



Received 30 September 2017

Accepted 26 October 2017

Edited by O. Büyükgüngör, Ondokuz Mayıs University, Turkey

**Keywords:** crystal structure; coordination polymer; barium; cobalt; macrocycle.**CCDC reference:** 1582341**Supporting information:** this article has supporting information at journals.iucr.org/e

# Crystal structure of the Ba<sup>II</sup>-based Co<sup>II</sup>-containing one-dimensional coordination polymer poly[[aqua- $\{\mu_4\text{-}2,2'\text{-}[(4,10\text{-dimethyl-}1,4,7,10\text{-tetraazacyclo-dodecane-}1,7\text{-diyl})\text{bis}(\text{methylidene})]\text{bis}(4\text{-oxo-}4H\text{-pyran-}3\text{-olato})\}\text{perchloratocobaltbarium}] perchlorate]$

**Paola Paoli,<sup>a</sup> Eleonora Macedi,<sup>a\*</sup> Patrizia Rossi,<sup>a</sup> Luca Giorgi,<sup>b</sup> Mauro Formica<sup>b</sup> and Vieri Fusi<sup>b</sup>**


<sup>a</sup>Department of Industrial Engineering, University of Firenze, via Santa Marta 3, I-50139 Firenze, Italy, and <sup>b</sup>Department of Pure and Applied Sciences, Lab of Supramolecular Chemistry, University of Urbino, via della Stazione, 4, I-61029 Urbino, Italy. \*Correspondence e-mail: eleonora.macedi@unifi.it

The title compound,  $\{[\text{Ba}(\text{H}_2\text{L1})(\text{ClO}_4)(\text{H}_2\text{O})]\text{ClO}_4\}_n$ , **L1** = 4,10-bis[(3-hydroxy-4-pyron-2-yl)methyl]-1,7-dimethyl-1,4,7,10-tetraazacyclododecane, is a one-dimensional coordination polymer. The asymmetric unit consists of a  $\{\text{Ba}[\text{Co}(\text{H}_2\text{L1})](\text{ClO}_4)(\text{H}_2\text{O})\}^+$  cationic fragment and a non-coordinating  $\text{ClO}_4^-$  anion. In the neutral  $[\text{Co}(\text{H}_2\text{L1})]$  moiety, the cobalt ion is hexacoordinated in a trigonal-prismatic fashion by the surrounding  $\text{N}_4\text{O}_2$  donor set. The  $\text{Ba}^{2+}$  ion is nine-coordinated and exhibits a distorted  $[\text{BaO}_9]$  monocapped square-anti-prismatic geometry, the six oxygen atoms coming from three distinct  $[\text{Co}(\text{H}_2\text{L1})]$  moieties, while the remaining three vertices are occupied by the oxygen atoms of a bidentate perchlorate anion and a water molecule. A barium- $\mu_2$ -oxygen motif develops along the *a* axis, connecting symmetry-related dinuclear  $\text{Ba}^{II}\text{-Co}^{II}$  cationic fragments in a wave-like chain, forming a one-dimensional metal coordination polymer. Non-coordinating  $\text{ClO}_4^-$  anions are located in the space between the chains. Weak C–H···O hydrogen bonds involving both coordinating and non-coordinating perchlorate anions build the whole crystal architecture. To our knowledge, this is the first example of a macrocyclic ligand forming a Ba<sup>II</sup>-based one-dimensional coordination polymer, containing Co<sup>II</sup> ions surrounded by a  $\text{N}_4\text{O}_2$  donor set.

## 1. Chemical context

Metal coordination polymers (CPs) have witnessed continuous growth, owing to their fascinating structural diversity in terms of architecture and topology and also their numerous potential applications, such as gas storage (Banerjee *et al.*, 2016; Fracaroli *et al.*, 2014; Sumida *et al.*, 2012; Suh *et al.*, 2012), chemical sensing (Campbell *et al.*, 2015; Hu *et al.*, 2014; Wang *et al.*, 2013; Kreno *et al.*, 2012), catalysis (Chughtai *et al.*, 2015; Mo *et al.*, 2014; Yoon *et al.*, 2012; Liu, Xuan *et al.*, 2010) and so forth. Recently, the interest in alkaline-earth metal ion-based CPs has been growing due to their unusual advantages such as low toxicity, wide distribution and low cost, which are of benefit for applications in the field of materials science (Raja *et al.*, 2014; Foo *et al.*, 2012, 2013; Xiao *et al.*, 2012).

According to a Cambridge Structural Database (CSD, Version 5.38, May 2017; Groom *et al.*, 2016) search, alkaline-




OPEN ACCESS

earth metal-based CPs are less common compared to the reported transition metal and rare-earth metal CPs (Cai *et al.*, 2017). Indeed, the study of alkaline-earth–metal systems is limited by challenges in the synthesis (Lian *et al.*, 2016; Douvali *et al.*, 2015; Mali *et al.*, 2015; Chakraborty *et al.*, 2014; Zhang, Huang *et al.*, 2012; Liu, Tsao *et al.*, 2010), the main reason being the variable coordination numbers (the most preferred coordination numbers are six for magnesium, six to eight for calcium, and six to twelve for strontium and barium), which lead to uncontrolled coordination geometries around the metal centre (Cai *et al.*, 2016; Feng *et al.*, 2015; Shi *et al.*, 2015; Zheng *et al.*, 2015; Jia *et al.*, 2014; Zhang, Yuan *et al.*, 2013; Smith *et al.*, 2013; Zhai *et al.*, 2013; Zhang, Guo *et al.*, 2013; Deng *et al.*, 2012; Foo *et al.*, 2012; Xiao *et al.*, 2012; Xie *et al.*, 2012; Zhang, Luo *et al.*, 2012; Jing *et al.*, 2010; Zhang *et al.*, 2010; Li *et al.*, 2009).

Besides, the ability of a system to bind alkaline-earth metal ions in aqueous solution is highly desirable and can be achieved thanks to the presence of oxygenated ligands and the preorganization of the receptor, which satisfies the need for a high coordination number without specific coordination requirements.

Ligand **L1** {4,10-bis[(3-hydroxy-4-pyron-2-yl)methyl]-1,7-dimethyl-1,4,7,10-tetraazacyclododecane} is a Maltol-based macrocycle (Amatori *et al.*, 2012) and is able to form discrete heteropolynuclear complexes. It has already proved to be able to form a Co<sup>II</sup> species (Borgogelli *et al.*, 2013) that is able to bind hard metal ions such as *Ln*<sup>III</sup> (*Ln* = Gd, Eu) and Na(I). In the case of *Ln*<sup>III</sup> ions, heterotrinuclear Co<sup>II</sup>–*Ln*<sup>III</sup>–Co<sup>II</sup> systems form, where the Co<sup>II</sup> cation preorganizes the system and two Co<sup>II</sup> species are involved in the coordination of one *Ln*<sup>III</sup> ion (Benelli *et al.*, 2013; Rossi *et al.*, 2017). In the case of the alkaline ion, a heterodinuclear complex forms, involving only one Co<sup>II</sup> species (Borgogelli *et al.*, 2013).

Herein we present a Ba<sup>II</sup>–Co<sup>II</sup> heterodinuclear metal coordination compound of **L1**, where a one-dimensional wave-like infinite array of barium ions bridges the [Co(H<sub>2</sub>**L1**)] moieties through a barium–μ<sub>2</sub>-oxygen motif. This is the first time that **L1** has proven able to form a coordination polymer and, to our knowledge, this is the first example of a macrocyclic ligand forming a Ba<sup>II</sup>-based 1D-CP containing Co<sup>II</sup> ions surrounded by an N<sub>4</sub>O<sub>2</sub> donor set.



## 2. Structural commentary

The title compound is the Ba<sup>II</sup>-based Co<sup>II</sup>-containing 1D-CP of **L1** of formula  $\{\text{Ba}[\text{Co}(\text{H}_2\text{L1})](\text{ClO}_4)(\text{H}_2\text{O})\}\cdot\text{ClO}_4\}_n$  and crystallizes in the monoclinic system in space group  $P2_1/n$ , with a  $\{\text{Ba}[\text{Co}(\text{H}_2\text{L1})](\text{ClO}_4)(\text{H}_2\text{O})\}^+$  cationic fragment (Fig. 1) and a  $(\text{ClO}_4)^-$  anion in the asymmetric unit.

In the neutral  $[\text{Co}(\text{H}_2\text{L1})]$  moiety, the Co<sup>2+</sup> ion is hexacoordinated and exhibits a distorted trigonal-prismatic geometry (Muettterties & Guggenberger, 1974), where the cobalt ion is surrounded by four nitrogen atoms of the macrocyclic base and two deprotonated hydroxyl oxygen atoms provided by both maltol rings of the ligand. In the distorted trigonal prism, the O1,N2,N3/O4,N1,N4 atoms define the two triangular faces, which are parallel within 12.51 (11)° (Fig. 2). The cobalt ion is displaced 1.0971 (5) Å above the mean plane described by the four nitrogen atoms of the tetraazamacrocyclic [maximum deviation of 0.068 (4) Å for N3] and falls, together with the Co–N(CH<sub>3</sub>) and Co–O bond distances (Table 1), in the expected range for Co-[12]aneN<sub>4</sub> complexes where the cobalt ion is hexacoordinated with a N<sub>4</sub>O<sub>2</sub> donor set (Fig. 3, left). The Co–N(Maltol) bond distances, instead, are longer (Table 1) than the Co–N(CH<sub>3</sub>) ones and longer with respect to those reported for other Co–**L1** complexes [Co–N(Maltol): range 2.26–2.44; Co–N(CH<sub>3</sub>): range: 2.13–2.19; Benelli *et al.*, 2013; Borgogelli *et al.*, 2013; Rossi *et al.*, 2017].

The conformation of the [12]aneN<sub>4</sub> macrocycle is the usual [3333]C-corners one (Meurant, 1987) with the *trans* nitrogen distances in agreement with those reported in the CSD for this conformation type, but the N2···N4 distance being longer than the N1···N3 one by 0.26 Å (Table 1), as found only in




Figure 1

The molecular structure of the  $\{\text{Ba}[\text{Co}(\text{H}_2\text{L1})](\text{ClO}_4)(\text{H}_2\text{O})\}^+$  cationic fragment, with the atom labelling and 30% probability displacement ellipsoids. Only one component of the disordered perchlorate anion and water molecule is shown. H atoms have been omitted for clarity. Symmetry codes: (i)  $-x + 1, -y, -z$ ; (ii)  $-x + 2, -y, -z$ .

Table 1

Selected bond lengths and angles ( $\text{\AA}$ ,  $^\circ$ ).

Co1—N1	2.199 (3)
Co1—N2	2.414 (3)
Co1—N3	2.220 (4)
Co1—N4	2.344 (3)
Co1—O1	2.044 (3)
Co1—O4	2.075 (3)
Ba1—O1	2.688 (3)
Ba1—O2	2.861 (3)
Ba1—O4	2.690 (3)
Ba1—O5	2.814 (3)
Ba1—O1W	2.774 (14)/2.75 (2)/2.972 (15) <sup>a</sup>
Ba1—O11	2.853 (19)/3.154 (13) <sup>b</sup>
Ba1—O12	2.955 (18)/2.863 (12) <sup>b</sup>
N1···N3	3.903 (5)
N2···N4	4.164 (5)
Ba1···Ba1 <sup>i</sup>	4.9123 (4)
Ba1···O2 <sup>i</sup>	2.860 (3)
O2···O2 <sup>i</sup>	2.932 (4)
Ba1···Ba1 <sup>ii</sup>	4.8443 (4)
Ba1···O5 <sup>ii</sup>	2.900 (3)
O5···O5 <sup>ii</sup>	3.033 (4)
Ba1 <sup>i</sup> ···Ba1 <sup>ii</sup>	8.8965 (4)
Ba1—O1—Co1	113.82 (12)
Ba1—O4—Co1	112.64 (11)
Ba1—O2—Ba1 <sup>i</sup>	118.34 (10)
Ba1—O5—Ba1 <sup>ii</sup>	115.92 (10)

Symmetry codes: (i)  $-x + 1, -y, -z$ ; (ii)  $-x + 2, -y, -z$ . Notes: (a) the values refer to O1WA/B/C atoms, respectively; (b) the values refer to the A/B oxygen atoms, respectively, of the disordered perchlorate anion (see Refinement section).

36% of cases. This is probably due to the fact that the Maltol units linked to atoms N2 and N4 are involved in chelate six-membered rings, which stiffen the system and force those nitrogen atoms to move farther apart.

The two maltolate rings are almost orthogonal to each other (dihedral angle between ring mean planes about  $71^\circ$ ); both rings form similar angles (about  $55^\circ$ ) with the mean plane N1,N2,N3,N4. The dimensions of the binding area defined by the four oxygen donor atoms of the ligand, as roughly estimated by the distances separating the opposite O1···O5 and O2···O4 atoms, are quite similar (about 4.5  $\text{\AA}$ ).




Figure 2

Coordination polyhedra around cobalt (left) and barium (right) ions. [Symmetry codes: (i)  $-x + 1, -y, -z$ ; (ii)  $-x + 2, -y, -z$ .]

Table 2

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots\cdot\cdot$	$D-\text{H}$	$\text{H}\cdots\cdot\cdot A$	$D\cdots\cdot\cdot A$	$D-\text{H}\cdots\cdot\cdot A$
C8—H8A···O21	0.99	2.91/2.62	3.877 (12)/3.60 (3)	165.3/172.3
C3—H3A···O22 <sup>iii</sup>	0.99	2.65/2.48	3.577 (11)/3.46 (3)	155.7/166.7
C8—H8B···O13 <sup>iv</sup>	0.99	2.49/2.59	3.46 (2)/3.524 (15)	168.5/156.6
C22—H22···O21 <sup>v</sup>	0.95	2.62/2.66	3.482 (11)/3.51 (3)	151.4/149.2
C22—H22···O23 <sup>v</sup>	0.95	2.57/2.54	3.455 (11)/3.39 (4)	155.1/149.9

Symmetry codes: (iii)  $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (iv)  $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (v)  $x - 1, y, z$ . Note: the first and second values for each entry refer to the A and B oxygen atoms, respectively, of the disordered perchlorate anion (see Refinement section).

The  $\text{Ba}^{2+}$  ion is nine-coordinated and exhibits a distorted  $[\text{BaO}_9]$  monocapped square-antiprismatic geometry (Guggenberger & Muettteries, 1976), Fig. 2, where the barium cation is surrounded by six oxygen atoms from three distinct  $[\text{Co}(\text{H}_2\text{L1})]$  moieties {four from two maltolate groups of a moiety and two from the carbonyl groups belonging to two distinct symmetry-related moieties, O2<sup>i</sup> and O5<sup>ii</sup> [symmetry codes: (i)  $-x + 1, -y, -z$ ; (ii)  $-x + 2, -y, -z$ ]}, an oxygen atom of a disordered water molecule and two oxygen atoms of a disordered perchlorate anion, the latter acting as a bidentate ligand (Fig. 1). In the distorted monocapped square antiprism, the O5 oxygen atom caps the O5<sup>ii</sup>, O4, O1W, O12 face (Fig. 2, right). All bond distances (Table 1) are in agreement with data found in the CSD.

The  $\text{Ba}^{2+}$  and  $\text{Co}^{2+}$  cations are located 3.9799 (7)  $\text{\AA}$  apart from each other, the line connecting them being normal to the mean plane described by the four nitrogen atoms of the macrocycle [angle value: 87.59 (7) $^\circ$ ; Fig. 1]. As for the bridged Co—O—Ba moiety (Fig. 3, right), while the Ba—O and Co—O bond distances and the Ba···Co distance are in agreement with those found in the CSD, the corresponding Ba—O—Co angles (Table 1) are outside the observed range (89.5–111.4 $^\circ$ ).

### 3. Supramolecular features

The title compound forms wave-like chains with a repeating unit comprising a dinuclear  $\text{Ba}^{II}-\text{Co}^{II}$  cationic fragment with associated coordinating water molecules and perchlorate ions (Fig. 4). Non-coordinating  $\text{ClO}_4^-$  anions are located in the space between the chains.

A barium- $\mu_2$ -oxygen motif develops along the  $a$  axis, the angle between the two mean planes formed by atoms Ba, O2,





Figure 3

Fragments searched in the CSD.


**Figure 4**

Wave-like one-dimensional  $\text{Ba}^{ii}$ -based coordination polymer that develops along the  $a$  axis. The oxygen and barium atoms belonging to the barium- $\mu_2$ -oxygen motif are depicted in ball and stick mode. Only one component of the disordered perchlorate anion and water molecule is shown. H atoms and the non-coordinating  $\text{ClO}_4^-$  anions have been omitted for clarity. [Symmetry codes: (i)  $-x + 1, -y, -z$ ; (ii)  $-x + 2, -y, -z$ .]

$\text{Ba}^i$  and  $\text{O}2^i$  and atoms  $\text{Ba}$ ,  $\text{O}5$ ,  $\text{Ba}^{ii}$  and  $\text{O}5^{ii}$  is about  $40^\circ$  [symmetry codes: (i)  $-x + 1, -y, -z$ ; (ii)  $-x + 2, -y, -z$ ], Fig. 4. The  $\text{Ba}-\text{O}$  bond distances, the  $\text{O}\cdots\text{O}$  and  $\text{Ba}\cdots\text{Ba}$  distances and the  $\text{Ba}-\text{O}-\text{Ba}$  angle values within each plane and the  $\text{Ba}^i\cdots\text{Ba}^{ii}$  distance (Table 1) are in agreement with data reported in the CSD.

Weak  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds (Desiraju & Steiner, 1999) involving both coordinating and non-coordinating perchlorate anions build the whole crystal architecture (Table 2). Distinct 1D-CPs are held together by weak  $\text{C}-\text{H}\cdots\text{O}$  interactions between the coordinating perchlorate anions belonging to a CP and methylene hydrogen atoms belonging to the adjacent CPs (Fig. 5).

The non-coordinating perchlorate anion connects, *via* a net of weak hydrogen bonds, three  $\{\text{Ba}[\text{Co}(\text{H}_2\text{L1})](\text{ClO}_4)(\text{H}_2\text{O})\}^+$


**Figure 5**

Adjacent CPs connected *via* hydrogen bonds involving the coordinating  $\text{ClO}_4^-$  anions as viewed along the  $b$  axis. The coordinating  $\text{ClO}_4^-$  anions are depicted in ball and stick mode. Hydrogen bonds are depicted as light-blue dotted lines. Only H atoms involved in the  $\text{C}-\text{H}\cdots\text{O}$  interactions and only one component of the disordered perchlorate anion and water molecule are shown.

cationic fragments belonging to two different 1D-CPs wave-like disposed along the  $b$  axis (Fig. 6).

#### 4. Database survey

Five structures containing **L1** were found in a search of the CSD (Version 5.38, May 2017; Groom *et al.*, 2016), three of them containing  $\text{Co}^{ii}$ : a hetero-trinuclear  $\text{Gd}^{iii}-\text{Co}^{ii}-\text{Gd}^{iii}$  dimer, a hetero-dinuclear  $\text{Na}^i-\text{Co}^{ii}$  complex and a  $\text{Co}^{ii}$  complex (Benelli *et al.*, 2013; Amatori *et al.*, 2012; Borgogelli *et al.*, 2013). In addition, our group recently published the

**Figure 6**

Crystal packing of the title compound as viewed along the  $a$  axis. The non-coordinating  $\text{ClO}_4^-$  anions are depicted in ball and stick mode. Hydrogen bonds involving the non-coordinating  $\text{ClO}_4^-$  anion and two  $\{\text{Ba}[\text{Co}(\text{H}_2\text{L1})](\text{ClO}_4)(\text{H}_2\text{O})\}^+$  cationic fragments on the same plane are depicted in light-blue dotted lines. Hydrogen bonds involving the non-coordinating  $\text{ClO}_4^-$  anion and a  $\{\text{Ba}[\text{Co}(\text{H}_2\text{L1})](\text{ClO}_4)(\text{H}_2\text{O})\}^+$  cationic fragment out of plane (symmetry operation  $x - 1, y, z$ ) are depicted in grey dotted lines. Only H atoms involved in the  $\text{C}-\text{H}\cdots\text{O}$  interactions and only one component of the disordered perchlorate anion and water molecule are shown.

corresponding hetero-trinuclear Eu–Co–Eu dimer (Rossi *et al.*, 2017).

A general search for structures containing both Co<sup>II</sup> and Ba<sup>II</sup> ions revealed 61 hits, 20 of which are polymeric structures formed by organic ligands containing both oxygen and nitrogen donor atoms and only two being 1D-CPs. It is noteworthy that none of the 20 structures contains either macrocyclic ligands or an N<sub>4</sub>O<sub>2</sub> donor set around the Co<sup>II</sup> ion. In eight out of those 20 polymeric structures, the Ba<sup>II</sup> and Co<sup>II</sup> ions are bridged by oxygen atoms and ten out of 20 show oxygen-bridged Ba<sup>II</sup> ions (only eight forming an infinite chain). Finally, only six out of the 20 polymeric structures contain both oxygen-bridged Ba<sup>II</sup> ions and oxygen-bridged Ba<sup>II</sup> and Co<sup>II</sup> ions.

All these data suggest that structures containing both oxygen-bridged Ba<sup>II</sup> ions and oxygen-bridged Ba<sup>II</sup> and Co<sup>II</sup> ions are not common and that no Ba<sup>II</sup>-based 1D-CPs formed by macrocyclic ligands and containing Co<sup>II</sup> ions surrounded by an N<sub>4</sub>O<sub>2</sub> donor set are present in the CSD.

## 5. Synthesis and crystallization

Compound **L1** was obtained following the synthetic procedure previously reported (Amatori *et al.*, 2012).

To obtain the Ba<sup>II</sup>-based Co<sup>II</sup>-containing 1D-CP of **L1**,  $\{\text{Ba}[\text{Co}(\text{H}_2\text{L1})](\text{ClO}_4)(\text{H}_2\text{O})\}\cdot\text{ClO}_4$ , 0.1 mmol of CoCl<sub>2</sub>·6H<sub>2</sub>O in water (10 mL) were added to an aqueous solution (20 mL) containing 0.1 mmol of **L1**·3HClO<sub>4</sub>·H<sub>2</sub>O. The solution was adjusted to pH 7 with 0.1 M N(CH<sub>3</sub>)<sub>4</sub>OH and then 0.05 mmol of BaCl<sub>2</sub>·2H<sub>2</sub>O were added. The solution was saturated with NaClO<sub>4</sub>. The Ba<sup>II</sup>–Co<sup>II</sup> 1D-CP of **L1** quickly precipitated as a microcrystalline pink solid. Crystals suitable for X-ray analysis were instead obtained by slow evaporation of a more diluted aqueous solution.

## 6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. All hydrogen atoms of the macrocycle were positioned geometrically and refined as riding with C–H = 0.95–0.99 Å with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C-methyl})$  and =  $1.2U_{\text{eq}}(\text{C})$  for other H atoms. Both perchlorate anions are disordered, all oxygen and chlorine atoms were set in double positions [anion 1: Cl1A/B, O11A/B, O12A/B, O13A/B, O14A/B, occupancy factor: 0.40 (3) and 0.60 (3); anion 2: Cl2A/B, O21A/B, O22A/B, O23A/B, O24A/B, occupancy factor: 0.78 (3) and 0.22 (3)]. The water molecule is disordered over three positions [SUMP command was used, occupancies 0.49 (3), 0.27 (3) and 0.24 (3)], the hydrogen atoms were not found in the Fourier-difference map and they were not introduced in the refinement. All non-hydrogen atoms were anisotropically refined: as for the disordered perchlorate anions, the SIMU instruction was used to restrain the anisotropic displacement parameters of the disordered atoms, while the ISOR instruction was used to model the disordered water oxygen atoms.

**Table 3**  
Experimental details.

Crystal data	[BaCo(C <sub>22</sub> H <sub>28</sub> N <sub>4</sub> O <sub>6</sub> )(ClO <sub>4</sub> )·(H <sub>2</sub> O)]·ClO <sub>4</sub>
$M_r$	857.67
Crystal system, space group	Monoclinic, $P2_1/n$
Temperature (K)	120
$a, b, c$ (Å)	8.8965 (2), 18.0995 (4), 19.0103 (6)
$\beta$ (°)	94.572 (2)
$V$ (Å <sup>3</sup> )	3051.34 (14)
$Z$	4
Radiation type	Mo $K\alpha$
$\mu$ (mm <sup>-1</sup> )	2.08
Crystal size (mm)	0.45 × 0.38 × 0.27
Data collection	
Diffractometer	Rigaku OD Xcalibur, Sapphire3
Absorption correction	Multi-scan ( <i>CrysAlis PRO</i> ; Rigaku OD, 2015). Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.
$T_{\min}, T_{\max}$	0.884, 1.000
No. of measured, independent and observed [ $I > 2\sigma(I)$ ] reflections	15496, 6966, 5315
$R_{\text{int}}$	0.037
(sin $\theta/\lambda$ ) <sub>max</sub> (Å <sup>-1</sup> )	0.682
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.041, 0.103, 1.06
No. of reflections	6966
No. of parameters	519
No. of restraints	133
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\max}, \Delta\rho_{\min}$ (e Å <sup>-3</sup> )	1.00, -0.72

Computer programs: *CrysAlis PRO* (Rigaku OD, 2015), *SIR2014* (Burla, 2015), *SHELXL2014/7* (Sheldrick, 2015).

## Acknowledgements

The authors acknowledge the CRIST (Centro di Cristallografia Strutturale, University of Firenze) where the data collection was performed.

## Funding information

Funding for this research was provided by: MIUR PRIN 2015.

## References

- Amatori, S., Ambrosi, G., Fanelli, M., Formica, M., Fusi, V., Giorgi, L., Macedi, E., Micheloni, M., Paoli, P., Pontellini, R. & Rossi, P. (2012). *J. Org. Chem.* **77**, 2207–2218.
- Banerjee, D., Simon, C. M., Plonka, A. M., Motkuri, R. K., Liu, J., Chen, X. Y., Smit, B., Parise, J. B., Haranczyk, M. & Thallapally, P. K. (2016). *Nat. Commun.* **7**, 11831.
- Benelli, C., Borgogelli, E., Formica, M., Fusi, V., Giorgi, L., Macedi, E., Micheloni, M., Paoli, P. & Rossi, P. (2013). *Dalton Trans.* **42**, 5848–5859.
- Borgogelli, E., Formica, M., Fusi, V., Giorgi, L., Macedi, E., Micheloni, M., Paoli, P. & Rossi, P. (2013). *Dalton Trans.* **42**, 2902–2912.
- Burla, M. C., Caliandro, R., Carrozzini, B., Cascarano, G. L., Cuocci, C., Giacovazzo, C., Mallamo, M., Mazzone, A. & Polidori, G. (2015). *J. Appl. Cryst.* **48**, 306–309.

- Cai, S.-L., He, Z.-H., Wu, W.-H., Liu, F.-X., Huang, X.-L., Zheng, S.-R., Fan, J. & Zhang, W. (2017). *CrystEngComm*, **19**, 3003–3016.
- Cai, S. L., Zheng, S. R., Fan, J., Zeng, R. H. & Zhang, W. G. (2016). *CrystEngComm*, **18**, 1174–1183.
- Campbell, M. G., Sheberla, D., Liu, S. F., Swager, T. M. & Dincă, M. (2015). *Angew. Chem. Int. Ed.* **54**, 4349–4352.
- Chakraborty, A. & Maji, T. K. (2014). *APL Mater* **2**, 124107-7.
- Chughtai, A. H., Ahmad, N., Younus, H. A., Laypkov, A. & Verpoort, F. (2015). *Chem. Soc. Rev.* **44**, 6804–6849.
- Deng, J. H., Zhong, D. C., Luo, X. Z., Liu, H. J. & Lu, T. B. (2012). *Cryst. Growth Des.* **12**, 4861–4869.
- Desiraju, G. R. & Steiner, T. (1999). *The Weak Hydrogen Bond in Structural Chemistry and Biology. IUCr Monographs on Crystallography*, Vol. 9. New York: Oxford University Press.
- Douvali, A., Papaefstathiou, G. S., Gullo, M. P., Barbieri, A., Tsipis, A. C., Malliakas, C. D., Kanatzidis, M. G., Papadas, I., Armatas, G. S., Hatzidimitriou, A. G., Lazarides, T. & Manos, M. J. (2015). *Inorg. Chem.* **54**, 5813–5826.
- Feng, X., Feng, Y. Q., Chen, J. J., Ng, S. W., Wang, L. Y. & Guo, J. Z. (2015). *Dalton Trans.* **44**, 804–816.
- Foo, M. L., Horike, S., Duan, J., Chen, W. & Kitagawa, S. (2013). *Cryst. Growth Des.* **13**, 2965–2972.
- Foo, M. L., Horike, S., Inubushi, Y. & Kitagawa, S. (2012). *Angew. Chem. Int. Ed.* **51**, 6107–6111.
- Fracaroli, A. M., Furukawa, H., Suzuki, M., Dodd, M., Okajima, S., Gándara, F., Reimer, J. A. & Yaghi, O. M. (2014). *J. Am. Chem. Soc.* **136**, 8863–8866.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst. B* **72**, 171–179.
- Guggenberger, L. J. & Muetterties, E. L. (1976). *J. Am. Chem. Soc.* **98**, 7221–7225.
- Hu, Z. C., Deibert, B. J. & Li, J. (2014). *Chem. Soc. Rev.* **43**, 5815–5840.
- Jia, H. L., Li, Y. L., Xiong, Z. F., Wang, C. J. & Li, G. (2014). *Dalton Trans.* **43**, 3704–3715.
- Jing, X. M., Zhang, L. R., Ma, T. L., Li, G. H., Yu, Y., Huo, Q. S., Eddaoudi, M. & Liu, Y. L. (2010). *Cryst. Growth Des.* **10**, 492–494.
- Kreno, L. E., Leong, K., Farha, O. K., Allendorf, M., Van Duyne, R. P. & Hupp, J. T. (2012). *Chem. Rev.* **112**, 1105–1125.
- Li, X., Wu, B. L., Niu, C. Y., Niu, Y. Y. & Zhang, H. Y. (2009). *Cryst. Growth Des.* **9**, 3423–3431.
- Lian, C., Liu, L., Guo, X., Long, Y., Jia, S., Li, H. & Yang, L. (2016). *J. Solid State Chem.* **233**, 229–235.
- Liu, H. K., Tsao, T. H., Lin, C. H. & Zima, V. (2010). *CrystEngComm*, **12**, 1044–1047.
- Liu, Y., Xuan, W. M. & Cui, Y. (2010). *Adv. Mater.* **22**, 4112–4135.
- Mali, G., Trebosc, J., Martineau, C. & Mazaj, M. (2015). *J. Phys. Chem. C*, **119**, 7831–7841.
- Meurant, G. (1987). *Stereochemical and Stereophysical Behaviour of Macrocycles*, edited by I. Bernal. Amsterdam; New York: Elsevier.
- Mo, K., Yang, Y. & Cui, Y. (2014). *J. Am. Chem. Soc.* **136**, 1746–1749.
- Muetterties, E. L. & Guggenberger, L. J. (1974). *J. Am. Chem. Soc.* **96**, 1748–1756.
- Raja, D. S., Luo, J. H., Yeh, C. T., Jiang, Y. C., Hsu, K. F. & Lin, C. H. (2014). *CrystEngComm*, **16**, 1985–1994.
- Rigaku OD (2015). *CrysAlis PRO*. Rigaku Oxford Diffraction Ltd., Abingdon, UK.
- Rossi, P., Ciattini, S., Formica, M., Fusi, V., Giorgi, L., Macedi, E., Micheloni, M. & Paoli, P. (2017). *Inorg. Chim. Acta*, doi: 10.1016/j.ica.2017.06.033.
- Sheldrick, G. M. (2015). *Acta Cryst. C* **71**, 3–8.
- Shi, B. B., Zhong, Y. H., Guo, L. L. & Li, G. (2015). *Dalton Trans.* **44**, 4362–4369.
- Smith, N. D., Roppe, J. R., Bonnefous, C., Payne, J. E., Zhang, H., Chen, X. H., Lindstrom, A. K., Duron, S. G., Hassing, C. A. & Noble, S. A. (2013). US patent 20080139558-A1.
- Suh, M. P., Park, H. J., Prasad, T. K. & Lim, D. W. (2012). *Chem. Rev.* **112**, 782–835.
- Sumida, K., Rogow, D. L., Mason, J. A., McDonald, T. M., Bloch, E. D., Herm, Z. R., Bae, T. H. & Long, J. R. (2012). *Chem. Rev.* **112**, 724–781.
- Wang, J. H., Li, M. & Li, D. (2013). *Chem. Sci.* **4**, 1793–1801.
- Xiao, D., Chen, H., Sun, D., He, J., Yan, S., Yang, J., Wang, X., Yuan, R. & Wang, E. (2012). *CrystEngComm*, **14**, 2849–2858.
- Xie, L. X., Hou, X. W., Fan, Y. T. & Hou, H. W. (2012). *Cryst. Growth Des.* **12**, 1282–1291.
- Yoon, M., Srirambalaji, R. & Kim, K. (2012). *Chem. Rev.* **112**, 1196–1231.
- Zhai, Q. G., Zeng, R. R., Li, S. N., Jiang, Y. C. & Hu, M. C. (2013). *CrystEngComm*, **15**, 965–976.
- Zhang, Y., Guo, B. B., Li, L., Liu, S. F. & Li, G. (2013). *Cryst. Growth Des.* **13**, 367–376.
- Zhang, X., Huang, Y.-Y., Zhang, M.-J., Zhang, J. & Yao, Y.-G. (2012). *Cryst. Growth Des.* **12**, 3231–3238.
- Zhang, F. W., Li, Z. F., Ge, T. Z., Yao, H. C., Li, G., Lu, H. J. & Zhu, Y. Y. (2010). *Inorg. Chem.* **49**, 3776–3788.
- Zhang, Y., Luo, X. B., Yang, Z. L. & Li, G. (2012). *CrystEngComm*, **14**, 7382–7397.
- Zhang, Y., Yuan, P., Zhu, Y. & Li, G. (2013). *Dalton Trans.* **42**, 14776–14785.
- Zheng, S. R., Wen, Z. Z., Chen, Y. Y., Cai, S. L., Fan, J. & Zhang, W. G. (2015). *Inorg. Chem. Commun.* **55**, 165–168.

# supporting information

*Acta Cryst.* (2017). E73, 1806-1811 [https://doi.org/10.1107/S2056989017015638]

## Crystal structure of the Ba<sup>II</sup>-based Co<sup>II</sup>-containing one-dimensional coordination polymer poly[[aqua{ $\mu_4$ -2,2'-[(4,10-dimethyl-1,4,7,10-tetraaza-cyclododecane-1,7-diyl)bis(methylidene)]bis(4-oxo-4H-pyran-3-olato)}perchloratocobaltbarium] perchlorate]


Paola Paoli, Eleonora Macedi, Patrizia Rossi, Luca Giorgi, Mauro Formica and Vieri Fusi

### Computing details

Data collection: *CrysAlis PRO* (Rigaku OD, 2015); cell refinement: *CrysAlis PRO* (Rigaku OD, 2015); data reduction: *CrysAlis PRO* (Rigaku OD, 2015); program(s) used to solve structure: *SIR2014* (Burla, 2015); program(s) used to refine structure: *SHELXL2014/7* (Sheldrick, 2015); molecular graphics: *SHELXL2014/7* (Sheldrick, 2015); software used to prepare material for publication: *SHELXL2014/7* (Sheldrick, 2015).

### Poly[[aqua{ $\mu_4$ -2,2'-[(4,10-dimethyl-1,4,7,10-tetraaza-cyclododecane-1,7-diyl)bis(methylidene)]bis(4-oxo-4H-pyran-3-olato)}perchloratocobaltbarium] perchlorate]

#### Crystal data



$M_r = 857.67$

Monoclinic,  $P2_{1}/n$

$a = 8.8965$  (2) Å

$b = 18.0995$  (4) Å

$c = 19.0103$  (6) Å

$\beta = 94.572$  (2) $^\circ$

$V = 3051.34$  (14) Å<sup>3</sup>

$Z = 4$

$F(000) = 1708$

$D_x = 1.867$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4989 reflections

$\theta = 2.2\text{--}27.7^\circ$

$\mu = 2.08$  mm<sup>-1</sup>

$T = 120$  K

Prism, pink

0.45 × 0.38 × 0.27 mm

#### Data collection

Rigaku OD Xcalibur, Sapphire3  
diffractometer

Radiation source: fine-focus sealed X-ray tube,  
Enhance (Mo) X-ray Source

Graphite monochromator

Detector resolution: 16.4547 pixels mm<sup>-1</sup>  
 $\omega$  scans

Absorption correction: multi-scan

(CrysAlisPro; Rigaku OD, 2015) Empirical  
absorption correction using spherical  
harmonics, implemented in SCALE3  
ABSPACK scaling algorithm.

$T_{\min} = 0.884$ ,  $T_{\max} = 1.000$

15496 measured reflections

6966 independent reflections

5315 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.037$

$\theta_{\max} = 29.0^\circ$ ,  $\theta_{\min} = 2.2^\circ$

$h = -12 \rightarrow 11$

$k = -22 \rightarrow 22$

$l = -22 \rightarrow 24$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.103$   
 $S = 1.06$   
 6966 reflections  
 519 parameters  
 133 restraints

Hydrogen site location: inferred from neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.037P)^2 + 2.9358P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 1.00 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.72 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ba1	0.75388 (3)	0.05421 (2)	0.01055 (2)	0.03237 (9)	
Co1	0.75301 (6)	0.26481 (3)	0.07072 (3)	0.02836 (14)	
O1	0.6344 (3)	0.19069 (15)	0.00703 (16)	0.0351 (7)	
N1	0.9308 (4)	0.33476 (19)	0.03199 (19)	0.0336 (8)	
C1	0.5130 (6)	0.3782 (3)	0.1078 (3)	0.0545 (14)	
H1A	0.4189	0.3897	0.1300	0.065*	
H1B	0.5870	0.4177	0.1210	0.065*	
O2	0.4340 (3)	0.07423 (16)	-0.00434 (18)	0.0400 (8)	
N2	0.6147 (4)	0.35215 (19)	-0.00548 (19)	0.0350 (8)	
C2	0.4817 (5)	0.3772 (3)	0.0300 (3)	0.0461 (12)	
H2A	0.3957	0.3438	0.0172	0.055*	
H2B	0.4530	0.4275	0.0133	0.055*	
O3	0.3318 (3)	0.26420 (17)	-0.11008 (16)	0.0374 (7)	
N3	0.5736 (4)	0.3060 (2)	0.1354 (2)	0.0389 (9)	
C3	0.7199 (5)	0.4133 (3)	-0.0140 (3)	0.0445 (11)	
H3A	0.7282	0.4440	0.0292	0.053*	
H3B	0.6818	0.4449	-0.0541	0.053*	
O4	0.8695 (3)	0.16663 (15)	0.09010 (15)	0.0340 (7)	
N4	0.8905 (4)	0.30110 (18)	0.17530 (19)	0.0324 (8)	
C4	0.8729 (5)	0.3827 (3)	-0.0275 (3)	0.0434 (11)	
H4A	0.8650	0.3537	-0.0718	0.052*	
H4B	0.9442	0.4239	-0.0330	0.052*	
O5	1.0650 (3)	0.05816 (16)	0.05118 (17)	0.0406 (8)	
C5	0.9983 (5)	0.3822 (2)	0.0890 (3)	0.0422 (11)	
H5A	0.9293	0.4240	0.0962	0.051*	
H5B	1.0941	0.4029	0.0746	0.051*	
O6	1.1536 (4)	0.16710 (18)	0.23669 (17)	0.0462 (8)	
C6	1.0284 (5)	0.3412 (3)	0.1570 (3)	0.0414 (11)	
H6A	1.1114	0.3055	0.1527	0.050*	
H6B	1.0601	0.3764	0.1952	0.050*	

C7	0.7907 (5)	0.3517 (3)	0.2110 (3)	0.0426 (11)
H7A	0.7868	0.4003	0.1870	0.051*
H7B	0.8308	0.3593	0.2606	0.051*
C8	0.6351 (5)	0.3191 (3)	0.2090 (3)	0.0442 (12)
H8A	0.6387	0.2719	0.2353	0.053*
H8B	0.5678	0.3533	0.2324	0.053*
C9	0.4532 (6)	0.2506 (3)	0.1367 (3)	0.0589 (15)
H9A	0.4106	0.2409	0.0885	0.088*
H9B	0.4947	0.2047	0.1576	0.088*
H9C	0.3738	0.2691	0.1650	0.088*
C10	1.0466 (5)	0.2848 (3)	0.0060 (3)	0.0500 (13)
H10A	1.0003	0.2536	-0.0319	0.075*
H10B	1.1274	0.3143	-0.0121	0.075*
H10C	1.0886	0.2536	0.0448	0.075*
C11	0.5658 (5)	0.3217 (2)	-0.0762 (2)	0.0395 (10)
H11A	0.5112	0.3604	-0.1049	0.047*
H11B	0.6556	0.3072	-0.1006	0.047*
C12	0.4661 (5)	0.2566 (2)	-0.0707 (2)	0.0318 (9)
C13	0.5033 (4)	0.1960 (2)	-0.0315 (2)	0.0316 (9)
C14	0.3993 (4)	0.1344 (2)	-0.0339 (2)	0.0325 (9)
C15	0.2580 (4)	0.1477 (2)	-0.0733 (2)	0.0310 (9)
H15	0.1815	0.1109	-0.0739	0.037*
C16	0.2308 (5)	0.2098 (3)	-0.1090 (2)	0.0367 (10)
H16	0.1358	0.2158	-0.1349	0.044*
C17	0.9321 (5)	0.2400 (2)	0.2241 (2)	0.0404 (11)
H17A	0.9863	0.2604	0.2673	0.048*
H17B	0.8390	0.2160	0.2380	0.048*
C18	1.0276 (5)	0.1841 (2)	0.1936 (2)	0.0354 (10)
C19	0.9936 (4)	0.1509 (2)	0.1305 (2)	0.0307 (9)
C20	1.0956 (5)	0.0936 (2)	0.1072 (2)	0.0364 (10)
C21	1.2276 (5)	0.0819 (3)	0.1537 (3)	0.0479 (12)
H21	1.3017	0.0479	0.1404	0.058*
C22	1.2496 (6)	0.1168 (3)	0.2147 (3)	0.0536 (14)
H22	1.3377	0.1056	0.2443	0.064*
Cl1A	0.800 (2)	0.1249 (9)	-0.1620 (10)	0.047 (2) 0.40 (3)
O11A	0.658 (2)	0.0849 (10)	-0.1333 (10)	0.047 (3) 0.40 (3)
O12A	0.9088 (18)	0.1007 (11)	-0.1140 (9)	0.052 (3) 0.40 (3)
O13A	0.860 (3)	0.0808 (10)	-0.2136 (12)	0.063 (4) 0.40 (3)
O14A	0.810 (3)	0.2057 (12)	-0.1686 (12)	0.074 (4) 0.40 (3)
Cl1B	0.7844 (14)	0.1280 (7)	-0.1622 (6)	0.0479 (16) 0.60 (3)
O11B	0.6498 (15)	0.1036 (8)	-0.1448 (7)	0.051 (2) 0.60 (3)
O12B	0.8977 (12)	0.1255 (8)	-0.1006 (6)	0.056 (2) 0.60 (3)
O13B	0.8144 (15)	0.0963 (8)	-0.2287 (6)	0.064 (3) 0.60 (3)
O14B	0.750 (2)	0.2000 (8)	-0.1816 (7)	0.076 (3) 0.60 (3)
Cl2A	0.6555 (9)	0.0780 (5)	0.3226 (4)	0.0424 (12) 0.78 (3)
O21A	0.5765 (11)	0.1464 (5)	0.3243 (6)	0.066 (2) 0.78 (3)
O22A	0.7030 (10)	0.0638 (6)	0.3946 (4)	0.062 (2) 0.78 (3)
O23A	0.5501 (11)	0.0220 (4)	0.2981 (6)	0.065 (2) 0.78 (3)

O24A	0.7821 (9)	0.0858 (7)	0.2801 (4)	0.081 (3)	0.78 (3)
Cl2B	0.661 (4)	0.073 (2)	0.317 (2)	0.060 (5)	0.22 (3)
O21B	0.610 (4)	0.1450 (16)	0.2989 (19)	0.053 (6)	0.22 (3)
O22B	0.733 (4)	0.0437 (17)	0.381 (2)	0.065 (5)	0.22 (3)
O23B	0.571 (4)	0.027 (2)	0.272 (2)	0.066 (5)	0.22 (3)
O24B	0.783 (3)	0.049 (2)	0.2882 (15)	0.071 (5)	0.22 (3)
O1WA	0.6542 (14)	0.0192 (9)	0.1411 (7)	0.079 (4)	0.493 (3)
O1WB	0.721 (2)	0.0144 (13)	0.1481 (12)	0.054 (5)	0.268 (3)
O1WC	0.842 (2)	-0.0011 (8)	0.1552 (8)	0.058 (4)	0.239 (3)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ba1	0.02904 (15)	0.02521 (14)	0.04255 (17)	0.00132 (10)	0.00089 (10)	-0.00540 (11)
Co1	0.0275 (3)	0.0241 (3)	0.0336 (3)	0.0011 (2)	0.0035 (2)	-0.0029 (2)
O1	0.0309 (15)	0.0251 (15)	0.0476 (19)	0.0022 (12)	-0.0083 (13)	-0.0040 (13)
N1	0.0362 (19)	0.0289 (19)	0.037 (2)	-0.0018 (15)	0.0072 (15)	-0.0018 (16)
C1	0.052 (3)	0.049 (3)	0.065 (4)	0.019 (2)	0.015 (3)	-0.007 (3)
O2	0.0332 (16)	0.0273 (16)	0.058 (2)	-0.0005 (12)	-0.0026 (14)	0.0005 (15)
N2	0.0336 (19)	0.031 (2)	0.040 (2)	0.0002 (15)	0.0013 (15)	-0.0011 (16)
C2	0.041 (3)	0.041 (3)	0.056 (3)	0.013 (2)	0.005 (2)	-0.007 (2)
O3	0.0348 (16)	0.0409 (18)	0.0354 (18)	0.0054 (13)	-0.0043 (13)	0.0019 (14)
N3	0.0327 (19)	0.045 (2)	0.040 (2)	0.0041 (17)	0.0069 (16)	-0.0030 (18)
C3	0.057 (3)	0.029 (2)	0.048 (3)	-0.001 (2)	0.000 (2)	0.008 (2)
O4	0.0332 (16)	0.0265 (15)	0.0403 (18)	0.0056 (12)	-0.0102 (12)	-0.0067 (13)
N4	0.0350 (19)	0.0272 (18)	0.035 (2)	0.0008 (15)	0.0057 (15)	-0.0068 (15)
C4	0.047 (3)	0.041 (3)	0.043 (3)	-0.011 (2)	0.008 (2)	0.007 (2)
O5	0.0372 (17)	0.0359 (18)	0.047 (2)	0.0083 (13)	-0.0064 (14)	-0.0124 (15)
C5	0.039 (3)	0.033 (2)	0.054 (3)	-0.008 (2)	0.002 (2)	-0.003 (2)
O6	0.051 (2)	0.0442 (19)	0.040 (2)	0.0051 (15)	-0.0151 (15)	-0.0055 (16)
C6	0.037 (2)	0.036 (2)	0.050 (3)	-0.005 (2)	-0.001 (2)	-0.005 (2)
C7	0.045 (3)	0.041 (3)	0.043 (3)	0.003 (2)	0.007 (2)	-0.015 (2)
C8	0.048 (3)	0.041 (3)	0.046 (3)	0.006 (2)	0.016 (2)	-0.008 (2)
C9	0.047 (3)	0.072 (4)	0.059 (4)	-0.016 (3)	0.016 (3)	-0.007 (3)
C10	0.043 (3)	0.051 (3)	0.059 (3)	0.002 (2)	0.020 (2)	-0.008 (3)
C11	0.045 (3)	0.034 (2)	0.039 (3)	0.000 (2)	0.001 (2)	0.002 (2)
C12	0.034 (2)	0.033 (2)	0.028 (2)	-0.0013 (18)	-0.0015 (17)	0.0001 (18)
C13	0.027 (2)	0.030 (2)	0.038 (2)	0.0033 (17)	-0.0002 (17)	-0.0061 (18)
C14	0.030 (2)	0.033 (2)	0.035 (2)	0.0061 (17)	0.0020 (17)	-0.0075 (19)
C15	0.025 (2)	0.037 (2)	0.029 (2)	-0.0004 (17)	-0.0038 (16)	-0.0079 (18)
C16	0.032 (2)	0.047 (3)	0.031 (2)	0.003 (2)	-0.0022 (17)	-0.003 (2)
C17	0.048 (3)	0.038 (3)	0.035 (3)	0.001 (2)	0.001 (2)	-0.003 (2)
C18	0.034 (2)	0.032 (2)	0.039 (3)	0.0022 (18)	-0.0061 (18)	0.0016 (19)
C19	0.031 (2)	0.026 (2)	0.034 (2)	-0.0013 (16)	-0.0025 (17)	-0.0028 (18)
C20	0.035 (2)	0.029 (2)	0.043 (3)	0.0009 (18)	-0.0058 (19)	-0.002 (2)
C21	0.040 (3)	0.037 (3)	0.064 (4)	0.010 (2)	-0.013 (2)	-0.011 (2)
C22	0.044 (3)	0.049 (3)	0.063 (4)	0.012 (2)	-0.024 (2)	-0.007 (3)
Cl1A	0.049 (4)	0.040 (4)	0.054 (4)	-0.001 (3)	0.010 (3)	0.013 (3)

O11A	0.047 (5)	0.039 (5)	0.055 (5)	-0.001 (4)	0.003 (4)	0.015 (4)
O12A	0.050 (5)	0.049 (5)	0.058 (5)	0.007 (4)	0.003 (4)	0.005 (4)
O13A	0.065 (7)	0.063 (6)	0.062 (7)	0.013 (5)	0.022 (5)	0.006 (5)
O14A	0.073 (8)	0.053 (7)	0.093 (8)	0.006 (7)	-0.019 (7)	0.024 (6)
Cl1B	0.049 (3)	0.057 (3)	0.037 (2)	0.0143 (18)	0.0014 (18)	0.0026 (18)
O11B	0.042 (3)	0.064 (5)	0.049 (4)	0.011 (4)	0.008 (3)	0.006 (4)
O12B	0.049 (3)	0.066 (4)	0.051 (4)	0.009 (4)	-0.006 (3)	0.001 (4)
O13B	0.058 (5)	0.100 (6)	0.035 (4)	0.024 (5)	0.006 (4)	-0.002 (4)
O14B	0.081 (7)	0.063 (5)	0.080 (6)	0.004 (6)	-0.018 (5)	0.025 (4)
Cl2A	0.0529 (19)	0.035 (2)	0.0386 (17)	-0.0050 (18)	-0.0023 (13)	0.0003 (14)
O21A	0.076 (5)	0.045 (3)	0.074 (5)	0.008 (3)	-0.011 (4)	-0.010 (4)
O22A	0.074 (4)	0.065 (5)	0.046 (4)	0.015 (3)	-0.001 (3)	0.007 (3)
O23A	0.082 (4)	0.034 (3)	0.074 (5)	-0.020 (3)	-0.022 (4)	0.006 (4)
O24A	0.097 (4)	0.070 (6)	0.083 (4)	-0.021 (4)	0.046 (3)	-0.011 (4)
Cl2B	0.075 (8)	0.034 (6)	0.069 (9)	0.014 (5)	0.002 (6)	-0.003 (5)
O21B	0.071 (10)	0.020 (8)	0.066 (12)	-0.001 (7)	-0.002 (9)	0.004 (8)
O22B	0.086 (10)	0.039 (9)	0.068 (10)	0.012 (7)	-0.003 (8)	-0.002 (7)
O23B	0.086 (9)	0.043 (9)	0.069 (11)	0.000 (7)	0.008 (8)	-0.007 (8)
O24B	0.091 (9)	0.046 (8)	0.076 (10)	0.009 (7)	0.012 (7)	-0.006 (7)
O1WA	0.100 (8)	0.095 (7)	0.042 (5)	0.000 (7)	0.005 (6)	-0.007 (5)
O1WB	0.067 (9)	0.056 (8)	0.038 (8)	-0.005 (8)	0.004 (8)	0.001 (6)
O1WC	0.077 (8)	0.045 (7)	0.052 (8)	-0.012 (6)	0.010 (6)	-0.012 (6)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Ba1—O1	2.688 (3)	C5—H5B	0.9900
Ba1—O4	2.690 (3)	O6—C22	1.339 (6)
Ba1—O1WB	2.75 (2)	O6—C18	1.370 (5)
Ba1—O1WA	2.774 (14)	C6—H6A	0.9900
Ba1—O5	2.814 (3)	C6—H6B	0.9900
Ba1—O11A	2.853 (19)	C7—C8	1.503 (6)
Ba1—O2 <sup>i</sup>	2.860 (3)	C7—H7A	0.9900
Ba1—O2	2.861 (3)	C7—H7B	0.9900
Ba1—O12B	2.863 (10)	C8—H8A	0.9900
Ba1—O5 <sup>ii</sup>	2.901 (3)	C8—H8B	0.9900
Ba1—O12A	2.955 (15)	C9—H9A	0.9800
Ba1—O1WC	2.972 (17)	C9—H9B	0.9800
Co1—O1	2.044 (3)	C9—H9C	0.9800
Co1—O4	2.075 (3)	C10—H10A	0.9800
Co1—N1	2.199 (3)	C10—H10B	0.9800
Co1—N3	2.220 (3)	C10—H10C	0.9800
Co1—N4	2.344 (4)	C11—C12	1.484 (6)
Co1—N2	2.414 (4)	C11—H11A	0.9900
O1—C13	1.330 (5)	C11—H11B	0.9900
N1—C5	1.472 (6)	C12—C13	1.351 (6)
N1—C10	1.484 (5)	C13—C14	1.448 (6)
N1—C4	1.485 (6)	C14—C15	1.432 (5)
C1—C2	1.484 (7)	C15—C16	1.326 (6)

C1—N3	1.492 (6)	C15—H15	0.9500
C1—H1A	0.9900	C16—H16	0.9500
C1—H1B	0.9900	C17—C18	1.471 (6)
O2—C14	1.252 (5)	C17—H17A	0.9900
O2—Ba1 <sup>i</sup>	2.860 (3)	C17—H17B	0.9900
N2—C3	1.466 (6)	C18—C19	1.353 (6)
N2—C2	1.479 (5)	C19—C20	1.469 (6)
N2—C11	1.486 (6)	C20—C21	1.429 (6)
C2—H2A	0.9900	C21—C22	1.320 (7)
C2—H2B	0.9900	C21—H21	0.9500
O3—C16	1.334 (5)	C22—H22	0.9500
O3—C12	1.366 (5)	Cl1A—O12A	1.35 (2)
N3—C9	1.470 (6)	Cl1A—O13A	1.40 (2)
N3—C8	1.481 (6)	Cl1A—O14A	1.47 (3)
C3—C4	1.511 (6)	Cl1A—O11A	1.59 (2)
C3—H3A	0.9900	Cl1B—O11B	1.342 (18)
C3—H3B	0.9900	Cl1B—O14B	1.383 (17)
O4—C19	1.325 (5)	Cl1B—O13B	1.433 (15)
N4—C17	1.471 (5)	Cl1B—O12B	1.484 (15)
N4—C7	1.477 (5)	Cl2A—O22A	1.423 (11)
N4—C6	1.490 (5)	Cl2A—O21A	1.425 (12)
C4—H4A	0.9900	Cl2A—O23A	1.434 (10)
C4—H4B	0.9900	Cl2A—O24A	1.444 (11)
O5—C20	1.254 (5)	Cl2B—O24B	1.33 (5)
O5—Ba1 <sup>ii</sup>	2.901 (3)	Cl2B—O23B	1.40 (4)
C5—C6	1.497 (6)	Cl2B—O21B	1.42 (4)
C5—H5A	0.9900	Cl2B—O22B	1.43 (5)
O1—Ba1—O4	56.75 (8)	N1—C4—H4A	109.6
O1—Ba1—O1WB	101.1 (5)	C3—C4—H4A	109.6
O4—Ba1—O1WB	74.2 (5)	N1—C4—H4B	109.6
O1—Ba1—O1WA	94.5 (3)	C3—C4—H4B	109.6
O4—Ba1—O1WA	78.8 (3)	H4A—C4—H4B	108.1
O1—Ba1—O5	111.22 (8)	C20—O5—Ba1	112.8 (3)
O4—Ba1—O5	60.20 (8)	C20—O5—Ba1 <sup>ii</sup>	127.8 (3)
O1WB—Ba1—O5	85.5 (4)	Ba1—O5—Ba1 <sup>ii</sup>	115.92 (10)
O1WA—Ba1—O5	97.9 (3)	N1—C5—C6	112.4 (4)
O1—Ba1—O11A	73.1 (4)	N1—C5—H5A	109.1
O4—Ba1—O11A	117.6 (4)	C6—C5—H5A	109.1
O1WA—Ba1—O11A	144.0 (5)	N1—C5—H5B	109.1
O5—Ba1—O11A	118.1 (4)	C6—C5—H5B	109.1
O1—Ba1—O2 <sup>i</sup>	121.15 (8)	H5A—C5—H5B	107.9
O4—Ba1—O2 <sup>i</sup>	146.50 (9)	C22—O6—C18	118.5 (4)
O1WB—Ba1—O2 <sup>i</sup>	73.8 (5)	N4—C6—C5	110.4 (4)
O1WA—Ba1—O2 <sup>i</sup>	67.9 (3)	N4—C6—H6A	109.6
O5—Ba1—O2 <sup>i</sup>	126.20 (8)	C5—C6—H6A	109.6
O11A—Ba1—O2 <sup>i</sup>	89.4 (4)	N4—C6—H6B	109.6
O1—Ba1—O2	59.51 (8)	C5—C6—H6B	109.6

O4—Ba1—O2	107.01 (8)	H6A—C6—H6B	108.1
O1WB—Ba1—O2	87.0 (5)	N4—C7—C8	109.4 (4)
O1WA—Ba1—O2	74.4 (3)	N4—C7—H7A	109.8
O5—Ba1—O2	166.61 (9)	C8—C7—H7A	109.8
O11A—Ba1—O2	70.2 (4)	N4—C7—H7B	109.8
O2 <sup>i</sup> —Ba1—O2	61.66 (10)	C8—C7—H7B	109.8
O1—Ba1—O12B	76.5 (3)	H7A—C7—H7B	108.2
O4—Ba1—O12B	84.3 (3)	N3—C8—C7	110.9 (4)
O1WB—Ba1—O12B	155.1 (5)	N3—C8—H8A	109.5
O5—Ba1—O12B	72.8 (2)	C7—C8—H8A	109.5
O2 <sup>i</sup> —Ba1—O12B	129.0 (3)	N3—C8—H8B	109.5
O2—Ba1—O12B	111.7 (2)	C7—C8—H8B	109.5
O1—Ba1—O5 <sup>ii</sup>	149.97 (9)	H8A—C8—H8B	108.1
O4—Ba1—O5 <sup>ii</sup>	123.92 (8)	N3—C9—H9A	109.5
O1WB—Ba1—O5 <sup>ii</sup>	107.8 (5)	N3—C9—H9B	109.5
O1WA—Ba1—O5 <sup>ii</sup>	115.4 (3)	H9A—C9—H9B	109.5
O5—Ba1—O5 <sup>ii</sup>	64.08 (10)	N3—C9—H9C	109.5
O11A—Ba1—O5 <sup>ii</sup>	83.3 (4)	H9A—C9—H9C	109.5
O2 <sup>i</sup> —Ba1—O5 <sup>ii</sup>	75.82 (9)	H9B—C9—H9C	109.5
O2—Ba1—O5 <sup>ii</sup>	128.99 (9)	N1—C10—H10A	109.5
O12B—Ba1—O5 <sup>ii</sup>	73.8 (3)	N1—C10—H10B	109.5
O1—Ba1—O12A	85.8 (4)	H10A—C10—H10B	109.5
O4—Ba1—O12A	93.0 (4)	N1—C10—H10C	109.5
O1WA—Ba1—O12A	169.9 (4)	H10A—C10—H10C	109.5
O5—Ba1—O12A	72.6 (3)	H10B—C10—H10C	109.5
O11A—Ba1—O12A	45.6 (5)	C12—C11—N2	111.3 (4)
O2 <sup>i</sup> —Ba1—O12A	120.5 (4)	C12—C11—H11A	109.4
O2—Ba1—O12A	114.1 (3)	N2—C11—H11A	109.4
O5 <sup>ii</sup> —Ba1—O12A	64.3 (4)	C12—C11—H11B	109.4
O1—Ba1—O1WC	114.0 (3)	N2—C11—H11B	109.4
O4—Ba1—O1WC	71.0 (3)	H11A—C11—H11B	108.0
O5—Ba1—O1WC	64.8 (3)	C13—C12—O3	123.3 (4)
O2 <sup>i</sup> —Ba1—O1WC	82.8 (3)	C13—C12—C11	124.2 (4)
O2—Ba1—O1WC	108.7 (3)	O3—C12—C11	112.5 (4)
O5 <sup>ii</sup> —Ba1—O1WC	91.4 (3)	O1—C13—C12	121.8 (4)
O1—Co1—O4	76.70 (11)	O1—C13—C14	119.4 (4)
O1—Co1—N1	122.08 (13)	C12—C13—C14	118.7 (4)
O4—Co1—N1	100.96 (12)	O2—C14—C15	123.7 (4)
O1—Co1—N3	100.86 (13)	O2—C14—C13	121.5 (4)
O4—Co1—N3	124.07 (13)	C15—C14—C13	114.8 (4)
N1—Co1—N3	124.08 (13)	C16—C15—C14	121.8 (4)
O1—Co1—N4	153.71 (12)	C16—C15—H15	119.1
O4—Co1—N4	82.53 (11)	C14—C15—H15	119.1
N1—Co1—N4	77.40 (13)	C15—C16—O3	122.6 (4)
N3—Co1—N4	77.62 (12)	C15—C16—H16	118.7
O1—Co1—N2	81.95 (12)	O3—C16—H16	118.7
O4—Co1—N2	152.96 (12)	C18—C17—N4	113.1 (4)
N1—Co1—N2	76.58 (12)	C18—C17—H17A	109.0

N3—Co1—N2	75.99 (13)	N4—C17—H17A	109.0
N4—Co1—N2	122.15 (12)	C18—C17—H17B	109.0
C13—O1—Co1	131.9 (2)	N4—C17—H17B	109.0
C13—O1—Ba1	114.1 (2)	H17A—C17—H17B	107.8
Co1—O1—Ba1	113.82 (11)	C19—C18—O6	123.0 (4)
C5—N1—C10	110.3 (4)	C19—C18—C17	124.1 (4)
C5—N1—C4	108.4 (3)	O6—C18—C17	112.9 (4)
C10—N1—C4	108.1 (4)	O4—C19—C18	122.3 (4)
C5—N1—Co1	110.6 (3)	O4—C19—C20	118.9 (4)
C10—N1—Co1	107.3 (3)	C18—C19—C20	118.7 (4)
C4—N1—Co1	112.1 (3)	O5—C20—C21	124.2 (4)
C2—C1—N3	112.0 (4)	O5—C20—C19	121.3 (4)
C2—C1—H1A	109.2	C21—C20—C19	114.5 (4)
N3—C1—H1A	109.2	C22—C21—C20	122.0 (4)
C2—C1—H1B	109.2	C22—C21—H21	119.0
N3—C1—H1B	109.2	C20—C21—H21	119.0
H1A—C1—H1B	107.9	C21—C22—O6	123.1 (4)
C14—O2—Ba1 <sup>i</sup>	124.5 (3)	C21—C22—H22	118.4
C14—O2—Ba1	111.2 (2)	O6—C22—H22	118.4
Ba1 <sup>i</sup> —O2—Ba1	118.33 (10)	O12A—Cl1A—O13A	89.8 (18)
C3—N2—C2	111.1 (4)	O12A—Cl1A—O14A	109.8 (13)
C3—N2—C11	108.9 (4)	O13A—Cl1A—O14A	118.6 (14)
C2—N2—C11	109.8 (3)	O12A—Cl1A—O11A	99.6 (12)
C3—N2—Co1	105.2 (3)	O13A—Cl1A—O11A	109.7 (13)
C2—N2—Co1	108.4 (3)	O14A—Cl1A—O11A	122.2 (15)
C11—N2—Co1	113.5 (3)	O12A—Cl1A—Ba1	52.5 (8)
N2—C2—C1	111.4 (4)	O13A—Cl1A—Ba1	121.1 (11)
N2—C2—H2A	109.3	O14A—Cl1A—Ba1	116.5 (12)
C1—C2—H2A	109.3	O11A—Cl1A—Ba1	51.0 (8)
N2—C2—H2B	109.3	Cl1A—O11A—Ba1	103.3 (10)
C1—C2—H2B	109.3	Cl1A—O12A—Ba1	106.3 (10)
H2A—C2—H2B	108.0	O11B—Cl1B—O14B	101.1 (12)
C16—O3—C12	118.6 (3)	O11B—Cl1B—O13B	108.6 (9)
C9—N3—C8	108.0 (4)	O14B—Cl1B—O13B	101.2 (12)
C9—N3—C1	111.1 (4)	O11B—Cl1B—O12B	111.0 (10)
C8—N3—C1	106.7 (4)	O14B—Cl1B—O12B	111.2 (9)
C9—N3—Co1	109.7 (3)	O13B—Cl1B—O12B	121.5 (12)
C8—N3—Co1	110.5 (2)	O11B—Cl1B—Ba1	61.1 (7)
C1—N3—Co1	110.8 (3)	O14B—Cl1B—Ba1	124.5 (8)
N2—C3—C4	109.5 (4)	O13B—Cl1B—Ba1	134.0 (9)
N2—C3—H3A	109.8	O12B—Cl1B—Ba1	50.2 (5)
C4—C3—H3A	109.8	Cl1B—O11B—Ba1	97.0 (7)
N2—C3—H3B	109.8	Cl1B—O12B—Ba1	106.3 (6)
C4—C3—H3B	109.8	O22A—Cl2A—O21A	104.2 (9)
H3A—C3—H3B	108.2	O22A—Cl2A—O23A	108.7 (6)
C19—O4—Co1	132.0 (2)	O21A—Cl2A—O23A	108.1 (7)
C19—O4—Ba1	115.2 (2)	O22A—Cl2A—O24A	111.8 (8)
Co1—O4—Ba1	112.64 (11)	O21A—Cl2A—O24A	109.6 (6)

---

C17—N4—C7	107.8 (3)	O23A—Cl2A—O24A	113.9 (9)
C17—N4—C6	110.0 (3)	O24B—Cl2B—O23B	90 (3)
C7—N4—C6	109.9 (3)	O24B—Cl2B—O21B	116 (3)
C17—N4—Co1	114.3 (3)	O23B—Cl2B—O21B	104 (3)
C7—N4—Co1	105.7 (3)	O24B—Cl2B—O22B	85 (3)
C6—N4—Co1	108.8 (3)	O23B—Cl2B—O22B	120 (3)
N1—C4—C3	110.4 (4)	O21B—Cl2B—O22B	131 (3)

---

Symmetry codes: (i)  $-x+1, -y, -z$ ; (ii)  $-x+2, -y, -z$ .