

ORIGINAL RESEARCH



A retrospective notes-based review of patients lost to follow-up from anti-retroviral therapy at Mulanje Mission Hospital, Malawi

Silas Webb¹, Joseph Hartland²

1. Croydon University Hospital and University of Bristol
2. Great Western Hospital and University of Bristol

Date Received: 20-March-2017
Revision Received: 08-Jan-2018
Date Accepted: 15-Jan-2018

Correspondence:
Silas Webb
(sw1167@my.bristol.ac.uk)

<https://dx.doi.org/10.4314/mmj.v30i2.4>

Abstract

Aim

To analyse patients with HIV who were lost to follow-up from anti-retroviral therapy (ART) at Mulanje Mission Hospital (MMH), Malawi.

Methods

All patients on adult antiretroviral combinations at MMH, who were classified as lost to follow-up (LTFU) according to the national guidelines (patients missing a scheduled follow-up visit by more than two months) over a 12-month period, were included in the study and compared against a control group who had never been lost. Variables compared were gender, age, months on ART, time of year, WHO clinical stage, ART regimen, reported side effects, number of doses missed in the previous 12 months, whether the patient has been followed up in the community and if so, the length of time elapsed before follow-up.

Results

In all, 136 patients had been LTFU over the previous 12 months at MMH. Of these, 43 had incomplete or missing ART cards, resulting in 93 LTFU patient's data that could be analysed. Patients were more likely to get LTFU if they were men ($p=0.03$), who had been on anti-retroviral therapy for a short duration ($p=0.06$) and the proportion of patients who missed more than 4 doses in the previous 12 months was higher among LTFU patients ($p=0.05$). Only 34.4% of those LTFU had been traced in the community at the time of analysis. Of those traced, 27% had moved to another area, 5.5% had died, 5.5% had the wrong documentation and 62% gave no reason as to why they had missed appointments.

Conclusion

This study in MMH has highlighted the importance and feasibility of comprehensive facility-level data-collection, both to identify local patient populations at risk of becoming lost to follow-up and to assess the follow-up measures in place to bring these lost to follow-up patients back into the programme. Even in the short time and with the small sample that was collected, there was evidence that patients most likely to get LTFU in MMH were young men, who had been on anti-retroviral therapy for a short duration and had missed over 4 doses in the last 12 months.

Introduction

More than three decades after the first cases of Human Immunodeficiency Virus (HIV) infections were recognized, the epidemic in sub-Saharan Africa (SSA) has claimed the lives of over 36 million people, and is still one of the world's most important public health issues¹. However, in recent years, significant advances have been made to address the epidemic, in line with the political intent of Millennium Development Goal 6, which was first to halve then begin to reverse the spread of HIV by 2015². Key to this success has been the expansion of antiretroviral therapy (ART), which has led to a 30-fold increase in ART coverage in SSA over the last decade³.

For patients accessing ART, Acquired Immune Deficiency Syndrome (AIDS) has been transformed from a life-limiting condition, with a median survival of one year from diagnosis, into a chronic disease⁴, one which requires life-long management⁵. However, the long-term retention of patients on ART has, itself, become a major challenge in SSA. In 2010, a systematic review of 39 papers found that 30% of patients were no longer taking their ARTs two years

after starting treatment⁶. In the same review, it was noted that there was a paucity of data as to why patients become lost to follow-up (LTFU) in SSA⁶.

This study will examine the current prevalence and reasons for patients becoming LTFU in resource-poor settings. The particular focus is on Mulanje Mission Hospital (MMH) in Malawi, serving one of the poorest regions of a nation with the lowest Gross Domestic Product (GDP) per capita in the world⁷.

Definitions

Throughout the literature, there is no universal definition for patients who are not retained in ART programmes. The World Health Organization (WHO) defines retention as "*patients known to be alive and receiving highly active ART (HAART) at the end of a follow-up period.*"⁸ This vague definition, which does not prescribe a specific follow-up period, has led to a wide disparity of retention periods being used in the studies: 30 days⁹, 3 months¹⁰, 6 months¹¹ and 1 year¹² are all quoted. Further confusion arises from the term 'LTFU' being used interchangeably with 'attrition' and 'defaulting' across studies⁵. In this report, LTFU has been utilized instead of

the two latter terms, which have been criticized for assigning blame to the patients as opposed to a “non-judgmental term that simply describes an outcome”¹².

Prevalence of patients LTFU from ART in SSA

The largest systematic review, which aimed to quantify the number of patients retained on ART in SSA (using 39 cohort studies with a total of 226,307 patients), found that retention was 70% after 24 months⁶. Although this study brought substantial attention to the issue, it did not differentiate the three distinct cohorts of patients who were not retained in care. One study which looked at samples of patients not retained in ART across 13 countries in SSA, including Malawi, found these results: 56% were LTFU, 40% had died and 4% were retained in care despite stopping ART¹³. In 2008, a more detailed single-cohort study in South Africa found that 31% of their non-retained patients had died, 25% had transferred to another facility and 44% were LTFU¹³. Despite differences in the absolute proportions across the different studies, the recurring finding is that patients LTFU make up the biggest share of patients not retained in ART programmes^{14,15}.

Reasons for patients LTFU in SSA

Adherence to drug programmes is affected by the complex interplay of self-care, community factors, care delivery and the drug itself¹⁵. Despite a paucity of qualitative research in developing countries, many reasons have been identified or hypothesized for the significant proportion of patients LTFU in ART programmes¹⁵. Several papers have identified financial constraints as a major cause of LTFU in SSA¹⁶. One study in Malawi found that 24% of patients originally recorded as LTFU, re-enrolled at the same site after ARTs were offered free of charge¹⁶. However, the variation in adherence rates in different areas of the country, irrespective of free access, suggests that other barriers to treatment are involved¹⁷. A qualitative study in Malawi's southern region of Chiradzulu found that fear and stigma were the main reasons for LTFU¹⁸. Of the 221 patients interviewed, 74% of ART patients had not disclosed their HIV status to household members¹⁸. In addition, the only published systematic review examining the concerns of HIV patients enrolled on ART programmes also found that fear of disclosure was the most consistently cited barrier to treatment¹⁹. As well as discussing structural and socio-cultural issues, some research has concentrated on identifying which patient groups are most likely to become LTFU. Younger age at treatment initiation²⁰, lower CD4 count at presentation⁶, type of ART regimen²¹ and the occurrence of side effects¹⁵ have all been found to be associated with LTFU in the early stages of treatment. Later on, a risk of LTFU arises from patients experiencing an improvement in health and subsequently assuming that treatment is no longer necessary; 28.3% of patients LTFU in rural Malawi cited improved health as their primary reason for leaving the programme¹⁸.

HIV and ART in Malawi

Malawi's HIV prevalence is one of the highest in the world, with the most recent Malawi Population-Based HIV Impact Assessment report estimating that 10.6% of the population (over 1 million people) are living with HIV²². The epidemic has affected all sectors of society, disproportionately targeting the youngest and most productive age groups, and has played a significant role in the country's life expectancy of only 54.8 years²³. This burden is compounded by a

shortage of health workers; the ratio of 1 physician per 50,000 people in Malawi is well below the WHO threshold of twenty¹⁹. Nevertheless, substantial efforts to reduce the burden of HIV in Malawi have been made over the last decade, bringing about significant reductions in mortality, morbidity and transmission. New infections have declined from 98,000 in 2005 to 34,000 in 2013, gaining 1.4 million life-years as well as a 67% reduction in mother-to-child transmission over the same period⁸. This has largely been attributed to remarkable increases in ART coverage across the country, with over 500,000 individuals initiated on ART since the introduction of free universal provision in 2004¹⁹. Prior to this, ART was only available at a cost of US\$25 for a month's supply¹⁷. However, ART adherence is variable. In 2012, only 79% of adults and children who started on ART in Malawi were taking them 12 months later, suggesting a significant proportion of patients LTFU⁷. National Reports in 2009, 2010 and 2012 have shown rates of LTFU of 12%, 15% and 21% respectively, which indicate that the problem is increasing^{8,19} and with it, the impact on individuals. Patients missing visits in the first year of treatment have a two-fold increase in long-term mortality²⁴. There is also a health risk to the general public posed by the likely increase in drug resistance²⁵. In response to this, in 2014, Malawi launched the 2015-2020 National HIV/AIDS Strategic Plan²⁶. A key objective involves “maintaining high levels of adherence and retention in ART programs”²³. This study is a contribution to this objective; as the number and demographics of patients LTFU from ART at MMH was not known, it provides this missing information and gives insight into the frequency and reasons patients may be LTFU locally.

Methods

Study design

This was a retrospective notes-based review.

Setting

MMH is a mission hospital located in southern Malawi, close to the border with Mozambique. The hospital serves a catchment of 650,000 people who mainly live in rural villages and are reliant on farming for subsistence⁷. The largest part of the hospital's workload is in the assessment and treatment of HIV/AIDS, with 10,714 HIV tests done and 74,136 ART clinic attendances during 2015 alone⁷. The most recent prevalence study in the area found that 18% of the population was living with HIV, significantly more than the national prevalence of 10.3%⁷.

Data collection

Using an existing clinical dataset, all patients on adult antiretroviral formulations at MMH, who were classified as ‘defaulters’ over the previous 12 months (2nd quarter 2015 – end of 1st quarter 2016), were included in the study. According to the local 2014 Malawian ART guidelines, defaulting was defined as “patients missing a scheduled follow-up visit by more than two months”²²; a time frame that has recently been shown by Rachlis et al to represent the period (>9 weeks) after which returning to clinic reduces substantially²⁷. Data was collected from the patient cards stored in the outpatient ART clinic. The following variables were included in the analysis: gender, age, months on ART, time of year LTFU, WHO clinical stage, ART regimen, reported side effects, number of doses missed in the previous 12 months, whether the patient has been followed up in the community

and if so, the length of time elapsed before follow up. To highlight comparisons amongst the studied population (the LTFU group), a control group was identified and their data analysed. The control group was selected by using all of the patients who attended the ART clinic between 18th and 22nd July 2016, and who had never defaulted.

Statistical analysis

Comparison data for LTFU and controls were analysed using the chi squared (χ^2) test, with alpha levels set at 0.05.

Ethical considerations

Ethical approval was sought and approved by the MMH medical director and hospital board. Confidentiality was maintained by coding the patient ART cards anonymously so that data was inputted and analysed separate from any patient identifiable information.

Results

In all, 136 patients had been LTFU over the previous 12 months at MMH. Of these, 43 patients had been missing ART cards, so the records of 93 patients LTFU were analysed. In the control group, 93 datasets were also analysed.

Table 1: Comparison data for LTFU and comparison groups

		Lost to Follow Up (n= 93)	Controls (n=93)
Gender	Male	42 (55%)	28 (30%)
	Female	51 (45%)	65 (70%)
Age (years)	≤20	3 (3.2%)	5 (5.4%)
	21-30	22 (23.7%)	13 (14.0%)
	31-40	36 (38.7%)	36 (38.7%)
	41-50	22 (23.7%)	23 (24.7%)
	>50	2 (2.2%)	7 (7.5%)
WHO Clinical Stage at Initiation of ART	1	39 (41.9%)	47 (50.5%)
	2	18 (19.4%)	15 (16.1%)
	3	30 (32.3%)	26 (28.0%)
	4	6 (6.5%)	5 (5.3%)
Duration of treatment on ART (months)	0-12	18 (19.4%)	10 (10.8%)
	13-24	20 (21.5%)	17 (18.3%)
	25-36	9 (9.7%)	11 (11.8%)
	37-48	18 (19.4%)	14 (15.1%)
	>48	28 (30.1%)	41 (44.1%)
Number of Doses of ART missed in last 12 months	0-3	48 (52%)	61 (66%)
	≥4	45 (48%)	32 (34%)




Figure 1: Number of Patients LTFU on ART in each quarter




Figure 2: Reasons for patients becoming LTFU (Recorded by expert clients) Pie Chart

had documented side effects in the previous 12 months (1 peripheral neuropathy and 1 hepatitis). The mean length of time on ART was 39.0 months in the LTFU group and 47.3 months in the control group, with a mean difference of 8.3 less months in the LTFU group ($p=0.06$).

The mean number of doses of ART missed in the previous 12 months was 13.15 in the LTFU group and 4.74 among controls, giving a mean difference of 8.41. When comparing the two groups, they were divided into those who had missed 0-3 doses and those who had missed 4 or more doses, following the criterion used by the clinic for non-compliance. It was found that the proportion of patients missing more than 4 doses was higher among the LTFU group compared to the controls ($p=0.05$).

Tracing

Of the LTFU patients 34.4% had been traced in the community at the time of analysis, with a mean length of time between clinic non-attendance and tracing of 13 weeks. According to the expert clients tracing the patients in the community, 27% had moved to another area, 5.5% had died, 5.5% had the wrong documentation (incorrect addresses documented in client files) and 62% gave no reason as to why they had missed appointments.

Discussion

The demographics of the MMH dataset echo previous findings in Malawi, in which male patients are more likely

to get LTFU on ART²⁷. The proportion of men in the LTFU group was 15.1% higher than in the control group, which was the only statistically significant finding of the study, with a chi-squared p -value of 0.03. The reasons for this cannot be definitely ascertained, since no interviews were undertaken with men who had been LTFU. However, previous studies in Malawi have shown that men are also less likely to know about their HIV status and to access treatment than women, suggesting that the gender disparity is one that encompasses all areas of HIV care¹⁷. The LTFU group also tended to be younger than the control group, with a mean difference of 3 years and a chi-squared value of 0.06. Fear of disclosure and stigma may be disproportionately affecting younger men, but more qualitative research is required to explore this. What this study, as well as a 2017 retrospective cohort study by Tweya et al, does illuminate is that both men and younger patients are at risk of LTFU in Malawi²⁸. Therefore, a recommendation is that interventions should be aimed specifically at men to improve their health outcomes.

Time of year LTFU

A higher proportion of those found LTFU were in the 4th quarter of 2015 and 1st quarter of 2016. These coincide with the rainy season in Mulanje District, which hinders the ability of patients to travel over long distances. Although data over several

years would be needed to confirm this trend, this is another matter warranting further research. Statistical confirmation could then be used to argue for more decentralised distribution of ART during the rainy season. This has been further evidenced by a recent retrospective cohort study in Neno District which found that ART decentralisation halved the average distance to health centres and was associated with a 70% increase in retention in care²⁹.

ART-regimen and length since initiation of ART

Two of the most instructive findings from the study came from an examination of the relationship between the length of time on ART and the likelihood of being LTFU. The mean length of time on medication was 8.3 months less in the LTFU group, with a p -value of 0.05, emphasizing the need for more intensive follow-up in the early stages of treatment, when patients are at the highest risk of LTFU. Patients were also more likely to get LTFU if they had missed 4 or more doses in the previous 12 months (p -value=0.05); so this represents another at-risk population who can be identified and could be given support before they get LTFU. The presence of side effects was not predictive of whether patients would get LTFU, with only 2.2% reporting them in the last 12 months. This is significantly lower than the figures quoted in trials looking at the rates of side effects in the 5A ART regimen, which is closer to 10%²⁶. This is possibly related to a failure in MMH documentation and so needs further research, especially in the light of a recent multi-country qualitative study which found that every additional self-reported side effect increased the odds of incomplete

ART adherence by 10%³⁰.

Tracing

Despite operating an expert client system in the clinic, attempting to trace and retrieve patients who miss their medication refills or appointments before they are officially classified as “defaulters”, only a third of those LTFU had received follow-up visits at the time of this study, suggesting that the community tracing is inconsistent. For those who were seen, the mean length of time between clinic non-attendance and tracing was 13 weeks. After taking into account the two-month delay that defines ‘defaulting’ in Malawi, this would mean a period of more than five months without medication for these patients, putting them at significant risk of adverse health consequences. When discussing this, one of the expert clients reported that a main barrier to seeing patients quickly was a lack of transport in the more remote areas, with their only bike currently broken and in need of servicing. Early follow-up has been shown to improve retention, with the largest ART provider in central Malawi finding that following up patients within 3 weeks of a missed appointment improved retention by 52%³¹. As only 45% of patients followed-up returned to clinic, this approach could be effective in MMH, but the expert clients will need to have access to better transport to try and achieve these results.

Another potential improvement to be made in the follow-up process is to take advantage of the growing number of mobile phone users in the developing world, with a project in Kenya finding that phone contacting improved ART adherence rates³². The expert clients also said that using phones would make for a more efficient follow-up. However, they were not currently provided with phone credit; so again this would need more funding to be made available.

Limitations

A significant limitation of this study was a reliance on the routine notes provided by the clinic, which often had large gaps and contained inconsistencies. The 43 initial missing records represent a potential selection bias if there are systematic differences between these patients and those analysed. Even when patient ART cards were available, some of the data was not entered: 8 ages were not recorded and for many patients, only their year of birth (not date) was documented. Absence of accurate data could have been a source of bias in age-related analyses.

Furthermore, there was no follow-up of two thirds of the LTFU patients; so their outcomes represented an unknown quantity. Without knowing if these patients had died, had moved elsewhere or had actively decided to stop treatment, very little can be inferred about their outcome. Of the patients who were followed up, 62% were documented as having “no reason” for leaving the programme. As their reasons are clearly more complex than that, these patients need to be interviewed in more depth to see whether they can be returned to the ART programme, with a view to reducing both morbidity and mortality. Ultimately, the biggest limitation was the study design; a case-control study design would be ideal. The way the control group was selected, which was by selecting patients attending over the course of a week, may not be truly comparable to the LTFU group, which was selected over the course of a year. This method was chosen as it was least disruptive to the running of the clinic in the short time available for data collection. However, as the control group was not randomly

selected over the same time period as the LTFU group, the data was susceptible to significant selection bias.

Conclusion

The global response to the challenge of scaling up access to ART in the developing world signifies one of the largest public health successes in recent history⁶. However, high rates of LTFU have the potential to reverse the gains that have been achieved thus far. Therefore, a focus of public health campaigns needs to be on long-term retention of patients on ART³³. The failure of patients to comply with their treatment should not be accepted as inevitable, but as a health injustice, affecting the poorest and most marginalized sectors of HIV patients and invariably worsening their health outcomes. This study at Mulanje Mission Hospital, Malawi, has highlighted the importance of comprehensive facility-level data-collection, both to identify local patient populations at risk of becoming LTFU and to assess the follow-up measures in place to bring these LTFU patients back into the programme. The authors found evidence that patients most likely to get LTFU in MMH were young men, who had been on ART for a short duration and had missed over 4 doses in the last 12 months. The increase in patients LTFU during the 1st and 4th quarters of the year needs further investigations and may highlight a need to provide larger quantities of ART or to further decentralize the clinic during these times. There was also evidence of serious inadequacies in the follow-up process, some of which could be remedied by modest expenditure on bikes and phone credits. In the immediate future, there is an urgent need to understand why patients are LTFU in MMH, to develop targeted interventions to prevent it. In order to further understanding, it is proposed that a qualitative study using semi-structured interviews with LTFU patients should be planned and undertaken in the area.

References

- LeVasseur M. T., Goldstein N. D., Welles S. A public health perspective on HIV/AIDS in Africa: Victories and unmet challenges. *Pathophysiology*. 2014; 21(3): 237-256. doi: <https://doi.org/10.1016/j.pathophys.2014.07.001>.
- UN. The Millennium Development Goals report 2010. United Nations, New York; 2010
- Rasschaert F., Koole O., Zachariah R., Lynen L., Manzi M., Van Damme W. Short and long term retention in antiretroviral care in health facilities in rural Malawi and Zimbabwe. *BMC Health Serv Res* 2012;12(1):1. doi: <https://doi.org/10.1186/1472-6963-12-444>
- Rosen S., Fox M. P., Gill C. J. Patient retention in antiretroviral therapy programs in sub-Saharan Africa: a systematic review. *PLoS One*. 2007 4(10) doi: [10.1371/journal.pmed.0040298](https://doi.org/10.1371/journal.pmed.0040298)
- Bisson G. P., Stringer J. S. ‘Lost but not forgotten—the economics of improving patient retention in AIDS treatment programs.’ *PLoS One*. 2009 6(10) doi: [10.1371/journal.pmed.1000174](https://doi.org/10.1371/journal.pmed.1000174)
- Fox M. P., Rosen S. Patient retention in antiretroviral therapy programs up to three years on treatment in sub-Saharan Africa, 2007–2009: systematic review. *Trop Med Int Health*. 2010;15(s1):1-15. doi: [10.1111/j.1365-3156.2010.02508.x](https://doi.org/10.1111/j.1365-3156.2010.02508.x)
- Shakespeare, R. Malawi: Challenges For Health [Lecture to University of Bristol SSC students]. Mulanje Mission Hospital. 8 July 2016.
- Joint United Nations Programme on HIV/AIDS (UNAIDS). (2014) ‘The Gap Report.’ Geneva: UNAIDS.
- Bisson G. P., Gaolathe T., Gross R., Rollins C., Bellamy S., Mogorosi

- M. 'Overestimates of survival after HAART: implications for global scale-up efforts.' *PloS Medicine*. 2008;3(3). doi: 10.1371/journal.pone.0001725
10. Bassett I. V., Wang B., Chetty S., Mazibuko M., Bearnot B., Giddy J. Loss to care and death before antiretroviral therapy in Durban, South Africa. *J Acquir Immune Defic Syndr*. 2009; 51(2):135-9
11. Geng E. H., Emenyonu N., Bwana M. B., Glidden D. V., Martin J. N. Sampling-based approach to determining outcomes of patients lost to follow-up in antiretroviral therapy scale-up programs in Africa. *JAMA*. 2008;300(5):506-507. doi: 10.1001/jama.300.5.506
12. Zachariah R., Harries A. D., Srinath S., Ram S. 'Language in tuberculosis services: can we change to patient-centred terminology and stop the paradigm of blaming the patients?' *Int J Tuberc Lung Dis*. 2012;16(6): 714-717. doi: 10.5588/ijtld.11.0635.
13. Dalal R. P., MacPhail C., Mqhayi M., Wing J., Feldman C., Chersich M. F. Characteristics and outcomes of adult patients lost to follow-up at an antiretroviral treatment clinic in Johannesburg, South Africa. *J Acquir Immune Defic Syndr*. 2008; 47(1):101-107. doi: 10.1097/QAI.0b013e31815b833a
14. Krebs D. W., Chi B. H., Mulenga, Y., Morris M., Cantrell R. A., Mulenga L. et al. Community-based follow-up for late patients enrolled in a district-wide programme for antiretroviral therapy in Lusaka, Zambia. *AIDS care*. 2008;20(3):311-317. doi: 10.1080/09540120701594776.
15. Miller C. M., Kethlapile M., Rybasack-Smith H., Rosen S. Why are antiretroviral treatment patients lost to follow-up? A qualitative study from South Africa. *Trop Med Int Health*. 2010;15(s1):48-54. doi:10.1111/j.1365-3156.2010.02514.x.
16. Hosseinipour M. C., Neuhaan F. H., Kanyama C. Lessons Learned From a Paying Antiretroviral Therapy Service in the Public Health Sector at Kamuzu Central Hospital, Malawi 1-Year Experience. *J Int Assoc Provid AIDS Care*. 2006;5(3):103-108. doi: 10.1177/1545109706288722
17. Muula A. S., Kataika E. (2008) Assessment of equity in the uptake of anti-retrovirals in Malawi. EQUINET Discussino Paper 58. University of Malawi, College of Medicine, Department of Community Health. [Online]. Available at: https://www.researchgate.net/publication/255594360_Assessment_of_equity_in_the_uptake_of_anti-retrovirals_in_Malawi (Accessed 18th March 2018)
18. McGuire M., Munyenembe T., Szumilin E., Heinzelmann A., Le Paih M., Bouithy N. et al. Vital status of pre-ART and ART patients defaulting from care in rural Malawi. *Trop Med Int Health*. 2010;15(s1):55-62. doi: 10.1111/j.1365-3156.2010.02504.x.
19. Rachlis B. S. Losses to Follow-up from an Antiretroviral Therapy (ART) Program in the Zomba District of Malawi. 2013. Doctoral dissertation, University of Toronto.
20. Chan A. K., Mateyu G., Jahn A., Schouten E., Arora P. Outcome assessment of decentralization of antiretroviral therapy provision in a rural district of Malawi using an integrated primary care model. *Trop Med Int Health*. 2010;15(s1): 90-97. doi: 10.1111/j.1365-3156.2010.02503.x.
21. Yiannoutsos C. T., Frangakis C. E., Musick, B. S. Sampling-based approaches to improve estimation of mortality among patient dropouts: experience from a large PEPFAR-funded program in Western Kenya. *PloS one*. 2008;3(12). doi: <https://doi.org/10.1371/journal.pone.0003843>
22. Ministry of Health, Malawi (2016) 'Summary Sheet Preliminary Findings: The Malawi Population-Based HIV Impact Assessment'
23. Ministry of Health, Malawi (2015) 'Malawi AIDS Response Progress Report.'
24. Mugavero M. J., Lin H. Y., Willig J. H., Westfall A. O., Ulett K. B. Missed visits and mortality among patients establishing initial outpatient HIV treatment. *Clin. Infect. Dis*. 2009;48(2):248-256. doi: 10.1086/595705.
25. Brennan, A. T., Maskew, M., Sanne, I., Fox, M. P. The importance of clinic attendance in the first six months on antiretroviral treatment: a retrospective analysis at a large public sector HIV clinic in South Africa. *J Int AIDS Soc*. 2010;13(1): 49. doi: 10.1186/1758-2652-13-49
26. Ministry of Health, Malawi (2014) 'Clinical Management of HIV in Children and Adults.'
27. Rachlis B., Ahmad F., van Lettow M., Muula A.S., Semba M., Cole D.C. Using concept mapping to explore why patients become lost to follow up from an antiretroviral therapy program in the Zomba District of Malawi. *BMC health services research*. 2013;13(1):210. doi: 10.1186/1472-6963-13-210.
28. Tweya H., Feldacker C., Heller T., Gugsu S, Ng'ambi W, Nthala O. et al. Characteristics and outcomes of older HIV-infected patients receiving antiretroviral therapy in Malawi: A retrospective observation cohort study. *PLoS One*. 2017;12(7) doi: <https://doi.org/10.1371/journal.pone.0180232>
29. Bilinski A., Birru E., Peckarsky M., Hecce M., Kalanga N., Neumann C., Bronson G et al. Distance to care, enrollment and loss to follow-up of HIV patients during decentralization of antiretroviral therapy in Neno District, Malawi: A retrospective cohort study. *PloS one*. 2017;12(10). doi: <https://doi.org/10.1371/journal.pone.0185699>
30. Renju J., Moshabela M., McLean E., Ddaaki W., Skovdal M., Odongo F. et al. 'Side effects' are 'central effects' that challenge retention in HIV treatment programmes in six sub-Saharan African countries: a multicountry qualitative study. *Sex Transm Infect*. 2017. doi: 10.1136/sextrans-2016-052971.
31. Tweya H., Gareta D., Chagwera F. Early active follow-up of patients on antiretroviral therapy (ART) who are lost to follow-up: the 'Back-to-Care' project in Lilongwe, Malawi. *Trop Med Int Health*. 2010;15(s1):82-89. doi: 10.1111/j.1365-3156.2010.02509.x.
32. Lester R. T. Effects of a mobile phone short message service on antiretroviral treatment adherence in Kenya (WelTel Kenya1): a randomised trial. *Lancet*. 2010;376(9755) :1838-1845. doi: [https://doi.org/10.1016/S0140-6736\(10\)61997-6](https://doi.org/10.1016/S0140-6736(10)61997-6)
33. Mberi M. N., Kuonza L. R., Dube N. M., Nattey C., Manda S. Determinants of loss to follow-up in patients on antiretroviral treatment, South Africa, 2004–2012: a cohort study. *BMC health services research*. 2015;15(1):1. doi: 10.1186/s12913-015-0912-2.