

RESEARCH ARTICLE

Open Access

Advanced practice registered nurses, physician assistants and cancer prevention and screening: a systematic review

Alexandria A Smith¹, Deanna Kepka^{2,3,4*} and K Robin Yabroff²

Abstract

Background: For more than two decades, integration of team-based approaches in primary care, including physicians, advanced practice registered nurses and physician assistants (APRN/PA), have been recommended for improving healthcare delivery, yet little is known about their roles in cancer screening and prevention. This study aims to review the current literature on the participation and roles of APRN/PAs in providing cancer screening and prevention recommendations in primary care settings in the United States.

Methods: We searched MEDLINE and CINAHL to identify studies published in 1990–2011 reporting on cervical, breast, and colorectal cancer screening and smoking cessation, diet, and physical activity recommendations by APRN/PAs in the United States. A total of 15 studies met all of our eligibility criteria. Key study, provider, and patient characteristics were abstracted as were findings about APRN/PA recommendations for screening and prevention.

Results: Most studies were cross-sectional, showed results from within a single city or state, had relatively small sample sizes, reported non-standardized outcome measures. Few studies reported any patient characteristics. APRN/PAs are involved in recommending cancer screening and prevention, although we found variation across screening tests and health behavior recommendations.

Conclusions: Additional research on the cancer prevention and screening practices of APRN/PAs in primary care settings using standardized outcome measures in relation to evidence-based guidelines may help strengthen primary care delivery in the United States.

Background

In the United States (US), cancer is the second leading cause of death [1]. According to current US Preventative Services Task Force (USPSTF) practice guidelines, cancer screening is associated with reduced cervical, breast, and colorectal cancer mortality and efforts to promote tobacco cessation [2-4], a healthful diet, and increased physical activity are associated with reduced cancer risk [5]. Yet in some populations with historically poor cancer outcomes, particularly the uninsured, low-income, and minorities [6,7], these evidence-based cancer control interventions have not been fully adopted [8]. The Affordable

Care Act will give approximately 32 million Americans greater access to healthcare and specifically the aforementioned cancer screening and prevention methods without any copayment by 2014 [9,10].

A recent Institute of Medicine Report (2011) called upon nurses to help meet the goals of the Affordable Care Act and outlined state and federal policy strategies to help ensure that nurses practice to the full extent of their education and training. For example, because APRN/PAs are not consistently able to see patients and provide medications without a physician's supervision, the report calls for reform of states' scope-of-practice laws. Furthermore, this report calls on nurses to serve as full partners with physicians and other healthcare professionals in the redesign of the US healthcare system and provides educational opportunities [11]. In addition,

* Correspondence: deanna.kepka@hci.utah.edu

²Division of Cancer Control and Population Sciences, National Cancer Institute, Bethesda, MD, USA

³College of Nursing, University of Utah, Salt Lake City, UT

Full list of author information is available at the end of the article

the Affordable Care Act provides for the Expansion of Physician Assistant Training Program [12], aims to increase student enrollment in primary care Physician Assistant (PA) programs. Since 1990 the number of APRN/PAs working in the US has risen from 50,000 to 250,000 in 2010, with midlevel providers conducting an increasing role in serving underserved populations and locations [13,14]. Approximately 55% APRNs/PAs work in a primary care setting [15]. The US Bureau of Labor Statistics reports that physician assistants within the United States are expected to grow by 30% between 2010 and 2020 [16]. Thus, this growing cadre of primary care health professionals is uniquely positioned to meet the growing demand for primary care resulting from the expansion of healthcare coverage by the Affordable Care Act.

As more individuals enter into the health care system, widespread implementation of evidence-based cancer screening and prevention interventions is critical for reducing cancer mortality and morbidity, but taxing on an already burdened healthcare system. In order to fulfill all of the current US Preventative Services Task Force recommendations for an average size patient panel, a primary care provider would exhaust an estimated 7.4 hours each day before providing diagnoses, treatments, or conducting administrative tasks [17]. Advanced practice nurses and physician assistants (APRN/PAs), including Nurse Practitioners (NPs), Physicians Assistants (PAs), and Certified Nurse Midwives (CNMs) are positioned to help meet this demand for recommended preventive services. For more than two decades, recommendations have been made to include APRN/PAs within primary care teams to improve chronic care delivery systems [18-21]. However, little is known about the roles of APRN/PAs in cancer screening and prevention, or how they might help meet an increasing demand for care. In this study, we conducted a systematic review of the recently published literature to assess the recommendations for and provision of USPSTF recommended cancer prevention and screening by APRN/PAs in primary care settings.

Methods

Study selection

We used the PubMed search mechanism for MEDLINE and CINAHL to identify English language studies on USPSTF recommended cancer prevention and screening recommendations among APRN/PAs in the United States published between January 1990 and December 2011. The search strategy used a combination of NIH librarian recommended terms that addressed APRN/PAs, cervical, breast, or colorectal cancer screening or recommendations for smoking cessation, or diet and physical activity. Using the following combination of search terms, a total of 594 studies were found: “Early Detection of Cancer” or “Vaginal Smears” or “Mammography”

or “Breast Neoplasms/prevention and control” or “Colorectal Neoplasms/prevention and control” or “Mass Screening” or “Papillomavirus Vaccines” and “Diet” or “Exercise” or “Motor Activity” or “Tobacco Use Cessation” and “Counseling” or “Nurse’s Role” or “Preventive Health Services” or “Patient Acceptance of Health Care” or “Patient Education as Topic” or “Health Knowledge, Attitudes, Practice” and “Nurse Practitioners” or “Physician Assistants” or “Nurse Midwives”. Organization of search terms and the number of published studies identified with each set of terms can be found in Table 1.

Abstracts were reviewed to identify the type of cancer screening or prevention, study design, sample size, and the country where the study was conducted. Inclusion criteria for this review were studies reporting: APRN/PAs conducting cancer prevention screening for cervical, breast or colorectal cancer, recommendations for smoking cessation, or diet and physical activity; United States

Table 1 Search terms used in med line and CINAHL

Group number	Search term	Number of articles medline	Number of articles CINAHL
Group 1	Nurse Practitioners OR Physician Assistants OR Nurse Midwives	22,644	3,779
Group 2	“Counseling”[Mesh:noexp] OR “Nurse’s Role”[Mesh] OR “Patient Acceptance of Health Care”[Mesh] OR “Patient Education as Topic”[Mesh] OR “Health Knowledge, Attitudes, Practice”[Mesh]	576,608	9,166
Group 3	“Diet”[Mesh] OR “Exercise”[Mesh] OR “Motor Activity”[Mesh] OR “Tobacco Use Cessation”[Mesh]	335,235	52,213
Group 4	“Early Detection of Cancer”[Mesh] OR “Vaginal Smears”[Mesh] OR “Mammography”[Mesh] OR “Breast Neoplasms/prevention and control”[Majr:noexp] OR “Colorectal Neoplasms/prevention and control”[Majr:noexp] OR “Mass Screening”[Mesh] OR “Papillomavirus Vaccines”[Mesh]	132,489	4,199
	Group 1 and Group 2 and (Group 3 or Group 4)	573	21

Limits: Humans, Journal Article, English, Publication Date from 1990/01/01 to 2011/12/31.

primary care setting; study publication dates between January 1990 and December 2011, quantitative data and at least 100 participants. We used a sample size eligibility requirement to ensure stable estimates of cancer prevention and screening as has been done [22,23]. Only 4 studies of more than 500 studies were excluded based solely on sample size. Reviews, editorials, letters, and essays were also excluded. Articles were initially reviewed by one author and any further decisions were discussed by all authors to determine if the study met all the eligibility criteria.

Ten studies met all of our eligibility criteria [24-33]. Because electronic searches may not identify all relevant studies [34], reference lists of the selected studies and published reviews of APRN/PAs and cancer screening or prevention were reviewed to identify other eligible studies. Five additional studies were identified in this manner [35-39]. A total of 15 studies are included in this paper.

Data abstraction

Data were abstracted from each paper using a standardized format in an excel spreadsheet. *Study characteristics included:* type of cancer screening test (Papanicolaou/Pap test, mammogram and/or any colorectal cancer screening test, including fecal occult blood test (FOBT), flexible sigmoidoscopy, or colonoscopy) or prevention recommendations (smoking cessation, diet and/or physical activity), geographic setting of sample (national, state, local), study design (cross-sectional, intervention and retrospective cohort), study size, delivery setting (single institution, network of institutions and multiple non-network institutions) and year of publication. *Type of APRN/PA* included NPs, CNMs, and PAs and was not a mutually exclusive category because some studies combined multiple types of APRN/PAs. Due to this and the limited number of studies, we reported on APRN/PAs as a combined unit unless the studies only focused on type of provider. The *comparison group* was recorded in three categories: physicians, other provider, and no comparison group. Where possible, comparisons were made between physicians and APRN/PAs. The *patient and provider characteristics* measured were age, gender, and race/ethnicity. The types of patient insurance coverage were also abstracted when reported. Cancer screening and prevention recommendations findings were abstracted as reported. Because some studies evaluated multiple screening tests and multiple types of prevention, these findings were abstracted and reported separately. The study cancer screening and prevention *outcome measures* were recorded in three categories: self-report, chart review, and biological samples. All study findings were abstracted as reported in the underlying study. This study

fully conforms to the PRISMA guidelines for systematic reviews (<http://www.prisma-statement.org>).

Results

Study characteristics

A total of seven studies reported outcomes on screening for cervical, breast or colorectal cancer [24,26,28-31,39], while ten studies measured cancer prevention recommendations for smoking cessation, diet, and physical activity (Table 2) [25-27,31-33,35-38]. We did not identify any studies of HPV vaccination or of post-treatment survivorship care. Three studies reported outcomes on more than one type of cancer screening [24,26,39], and four studies reported on more than one type of cancer prevention recommendations [26,27,31,37]. Most studies were cross-sectional, showed results from within a single city or state, had relatively small sample sizes (less than 500) and reported on the behavior of NPs. Only three studies reported on interventions, two focusing on tobacco and one on Pap test and mammograms [32,38,39]. The majority of studies presented self-reported data from providers about their own practice or their perceptions of APRN/PAs practice [24,25,28,31,35,37], while only a few presented self-reported data from patients [32,38]. Few studies reported receipt of services or documented changes in behavior as part of chart reviews [26,27,36,39]. A small number reported any patient characteristics. The response rates for studies varied, with those that presented self-reported data from providers ranged from 30% to 72% [24,25,28-31,35,37]. Many of the studies did not specify whether the APRN/PAs provided or recommended cancer screening.

Cancer screening

Cervical cancer screening

Of the five studies evaluating Pap tests, most showed that APRN/PAs provide or recommend Pap tests to patients (72% to 98%) and that physicians who currently work with APRN/PAs are amenable to APRN/PAs conducting Pap tests (Table 3) [24,26,29,31]. Physicians who practice in provider teams that include NPs and PAs are more supportive of NPs and PAs performing Pap tests than physicians who do not practice in provider teams that include NPs and PAs [29]. In addition, an intervention study compared NPs recommending and performing cervical cancer screening during routine visits to a provider reminder system. At follow-up there was a significant increase in the annual rate of women screened for cervical cancer by a NP at the intervention location (from 17.8% to 56.9%), while the annual rate of screening by physicians at the control location improved less (from 11.8% to 18.2%) during the study time period [39].

Table 2 Characteristics of studies of Advanced Practices Registered Nurses and Physician Assistants (APRN/PA) and cancer prevention and screening

Study characteristics	Study categories	Number of studies (N = 15)	Percentage of studies
Type of cancer prevention*	Diet/Physical activity	5	33%
	Smoking	9	60%
	Mammogram	3	20%
	Pap test	5	33%
	Colorectal cancer screening	4	27%
Study setting	National	6	40%
	State or regional	5	33%
	Local (City or Multiple Counties)	4	27%
Study design	Intervention	3	20%
	Retrospective cohort	2	13%
	Cross sectional	10	67%
Study date of publication	<1995	2	13%
	1995-1999	3	20%
	2000-2004	4	27%
	>= 2005	6	40%
Type of APRN/PA provider	Advanced practice nurse practitioner	13	87%
	Certified nurse midwife	2	13%
	Physician Assistant	3	20%
Comparison group	Physicians	5	33%
	None	5	33%
	Other	5	33%
Sample size	100-499	8	53%
	500-999	1	7%
	1,000-4,999	5	33%
	5,000-19,999	0	0%
	>20,000	1	7%
Healthcare delivery setting	Single institution or clinic	2	13%

Table 2 Characteristics of studies of Advanced Practices Registered Nurses and Physician Assistants (APRN/PA) and cancer prevention and screening (Continued)

	Network of institutions or clinics	3	20%	
	Multiple institutions (Not a Network)	10	67%	
Outcome measure	Chart review	3	20%	
	Self-Report by physicians	4	27%	
	Self-Report by APRN/PA providers	6	47%	
	Self-Report by patients	2	13%	
Patient characteristics	Insurance types*	Any medicare	2	13%
		Any medicaid	5	33%
		Private	6	40%
		Not reported	8	60%

*Measures not mutually exclusive.

Breast cancer screening

Of the three studies that studied breast cancer, two showed that a majority of patients who see NPs receive mammograms (69% to 91%) and that NPs recommend a similar amount of mammograms as physicians (Table 3) [24,26]. In the same NP intervention study mentioned previously for cervical cancer screening, the annual rate of mammography screening increased more among women seen at the NP screening recommendation site (18.3% to 40.0%) than at the cancer screening program using a provider reminder checklist on charts and referrals (18.0% at both time points) [39].

Colorectal cancer screening

Findings about APRN/PAs involvement in colorectal cancer screening are mixed and vary based on the screening modalities evaluated (Table 3). Of the four colorectal cancer screening studies, three showed a range of reported colorectal cancer screening provided or recommended by APRN/PAs (19% to 95%) [24,26,30]. This large variation in reported colorectal cancer screening is partially determined by the variation in reporting amongst the studies. The lowest percentage is based on chart review of an unspecified type of colorectal cancer screening among patients aged 50 and above [26] while the highest percentage is a self-reported survey answer from providers on how often they recommend FOBT to any patient (age not specified) [30]. We also found substantial variation in reporting on colorectal cancer screening modalities, with some studies reporting specific modalities [24] and others reporting whether an

Table 3 Studies of Advanced Practices Registered Nurses and Physician Assistants (APRN/PA) and breast, cervical and colorectal cancer screening

Study	Sample size	Outcome measure	Setting and study design	Pap findings	Mammogram findings	Colorectal cancer screening findings
Menees et al., [24]	Total providers: 336 OB/GYNs: 182	Self-Report of MDs and NPs	National survey of OB/GYNs and NPs	Pap test routinely provided:	Mammography routinely recommended:	CRC screening routinely recommended:
	NPs: 154		Cross sectional Survey	NPs: 94.8% OB/GYNs: 97.8% <i>P</i> < 0.01 Unadjusted	NPs: 90.9% OB/GYNs: 98.9% <i>P</i> < 0.01 Unadjusted	NPs: 61.7% OB/GYNs: 87.2% <i>P</i> < 0.01 Most common CRC screening recommended by either provider: FOBT: 76.2% Colonoscopy: 28.3% P-value not reported Ordered colonoscopy: NPs: 19.8% OB/GYNs: 37% <i>P</i> < 0.005 All measures unadjusted
Hopkins et al., [26]	Total patients: 1339	Chart review	Chart review in private practice	Receipt of Pap test:	Receipt of mammogram: (Aged 40+)	Patient receipt of colorectal screening: (Ages 50 +)
	PHCC NP patients: 755		and primary care health centers in New York City	PHCC NPs: 71.5% PHCC MDs: 53.8%	PHCC NPs: 69% PHCC MDs: 64.2%	PHCC NPs: 19.1% PHCC MDs: 45.7%
	MD patients: 441 PP NP patients: 143		Retrospective cohort	<i>P</i> < 0.001 Unadjusted	<i>P</i> = 0.240 Unadjusted	<i>P</i> < 0.001 Unadjusted
Sansbury et al., [28]	Total providers: 1900 PC MDs: 1235 NPC MDs: 665	Self-Report of MDs about APRN/PAs	National survey of MDs Cross sectional survey	NA	NA	Work with NP/PA to provide FOBT: MDs report working with a NP or PA to provide FOBT: 23.8% Of the 24% of physicians who work with NP/PA for FOBT, they reported frequency of supervising a NP or PA for FOBT: Supervised a NP: 75% Supervised a PA: 25% P-value not reported for all measures All measures unadjusted

Table 3 Studies of Advanced Practices Registered Nurses and Physician Assistants (APRN/PA) and breast, cervical and colorectal cancer screening (Continued)

Oliveria et al., [29]	Total providers: 1363 MDs: 1363	Self-Report of MDs about APRN/PAs	National survey of MDs Cross sectional survey	MDs amenable to NP/PA screening: Team Practice [^] : 89.6% Non-team Practice ^{^^} : 59.9% Team vs Non-team of amenable MDs: OR = 8.11 (95% CI: 5.80–11.35) P = 0.001 MDs reporting NP/PA screening: All MDs reporting frequency of NPs or PAs performing Pap tests: NPs: 33.5% PAs: 23.2% Team practice MDs reporting frequency of NP/PA performing Pap tests: NPs: 89.3% PAs: 82.7% P-value not reported for all other measures All measures unadjusted	NA	NA
Shaheen et al., [30]	Total providers: 1784 Total NPs: 526 Total PAs: 640 PC PAs: 322 PC NPs: 270	Self-Report of APRN/PAs	Survey of NPs and PAs in North Carolina Cross sectional survey	NA	NA	NP/PA who recommend/perform FOBT: Primary Care PA: 94.6% Primary Care NP: 92.1% NP/PA who recommend/perform flexible sigmoidoscopy: Primary Care PA: 76.1% Primary Care NP: 69.2% P-value not reported Unadjusted

Table 3 Studies of Advanced Practices Registered Nurses and Physician Assistants (APRN/PA) and breast, cervical and colorectal cancer screening (Continued)

Murphy, [31]	Total providers: 346 CNMs: 346	Self-Report of CNMs	National survey of CNMs Cross sectional survey	98% of CNMs report they routinely provide pap tests to 81-100% of their gynecologic patients P-value not reported Unadjusted	NA	NA
Mandelblatt et al., [39]	Total patients: 319 Intervention: 160 Control: 159 Total providers: Not Reported	Chart review	Two New York City study hospitals with NP led intervention and usual care control Intervention	Receipt of Pap test in intervention group: Baseline: 17.8% Post: 56.9% P < 0.01 Receipt of Pap test in control group: Baseline: 11.8% Post: 18.2% P-value not reported All measures unadjusted	Receipt of mammography in intervention group: Baseline: 18.3% Post: 40% P < 0.01 Receipt of mammography in control group: Baseline: 18.1% Post: 18.2% P-value not reported All measures unadjusted	NA

^ = (MDs who work with NPs or PAs).

^^ = (MDs who do not work with NPs or PAs).

NA = Not Applicable.

NPC = Non-Primary Care.

NP/PA = NP or PA.

PHCC = Primary Health Care Center.

PC = Primary Care.

PP = Private practice.

* = A higher number means the provider does the behavior more frequently.

CI = Confidence Interval.

unspecified type of colorectal cancer screening was offered [26]. In addition, two of the studies showed physicians reporting more colorectal cancer screening than APRN/PAs [24,26]. Only 24% of practicing primary care physicians reported working with APRN/PAs to provide FOBT in a nationally representative survey [28].

Recommendations for cancer prevention

Smoking cessation recommendations

Both physicians and APRN/PAs report frequently providing smoking cessation recommendations (Table 4). APRN/PAs self-reported a range of assessment of tobacco use and smoking cessation recommendations to patients (7% to 95%) in eight studies [25,26,31,32,36]. Of the nine studies that measured tobacco related outcomes, three showed that patients are more likely to receive recommendations for smoking cessation during visits with NPs than during visits without NPs (associations not always statistically significant) while another study showed MDs feel that they are more adequately trained to give smoking cessation counseling than NPs [27,32,37,38]. One smoking cessation intervention of NPs at a prenatal clinic visit compared to a usual care control prenatal clinic visit reported significantly increased abstinence among cigarette smokers at follow-up (19% vs. 0%) [38].

Diet and physical activity recommendations

The four studies that evaluated diet also evaluated physical activity, while one study only evaluated physical activity counseling (Table 4). These studies showed that while APRN/PAs do not frequently provide recommendations on diet and physical activity (12% to 52%), they do provide more recommendations related to diet and physical activity than their physician counterparts (3% to 15%) [26,27,31,37].

Discussion

In this paper, we reviewed the recent literature on the participation and roles of APRN/PAs in the delivery of cancer prevention and screening recommendations in US primary care settings. In the descriptive or intervention research we identified, only 15 studies during a 21 year period, APRN/PAs are involved in recommending cancer screening and prevention. The limited research is somewhat surprising, because a team approach, including physicians and APRN/PAs, has long been recommended for improving healthcare [18-21,40]. After receiving the appropriate training, APRN/PAs expect to provide or recommend Pap tests, mammograms and FOBT, while studies only reported on physicians working concurrently with APRN/PAs to screen for cervical cancer [29] and colorectal cancer [28]. With the enactment of the Affordable Care Act, millions of previously

uninsured or underinsured will gain access to healthcare. A better understanding of the potential roles of APRN/PAs in meeting this demand for cancer prevention and screening is critical.

The integration of more APRN/PAs into primary care can affect cancer screening and recommendations in several different ways. This integration has the potential to increase the overall percentage of the population ever receiving specific cancer prevention and screening recommendations, as was shown in an intervention study included in this review [39]. For example, colorectal cancer screening uptake in the US is substantially lower than for breast or cervical cancer screening [41]. The US Preventive Services Task Force (USPSTF) recommends any of three different tests for colorectal cancer (i.e., FOBT, flexible sigmoidoscopy, colonoscopy) [4]. These tests have different screening intervals, involvement of specialists, levels of invasiveness and other characteristics [4], potentially requiring detailed discussion to allow patients to make informed decisions about screening. Currently, less than 25% of physicians report actually working with APRN/PAs to provide colorectal cancer screening [28]. However, one challenge with moving forward with team-based health care is that physicians do not always want to work with nurse practitioners [42].

In a time constrained primary care setting, APRN/PAs might play a critical role in improving discussion about options and ultimately improving uptake of colorectal cancer screening. Alternatively, research featuring APRN/PAs might focus on improving all aspects of cancer control among specific populations, such as those previously uninsured or with key risk factors. Lack of health insurance and lack of prior screening has been consistently associated with late stage of disease at diagnosis for breast, cervical, and colorectal cancer [43-45]. Tobacco use and obesity are associated with many chronic diseases [46] and the role of APRN/PAs in encouraging healthy behaviors could improve a variety of health outcomes of the US population. Future research is needed that investigates that relationship between a visit with an APRN/PA and other primary care provider types within team-based primary care that oversamples racial and ethnic minorities and lower socioeconomic status populations.

We identified a number of methodological and reporting limitations in the studies included in this review related to study design and reporting of outcome measures and sample characteristics. Most of the studies were cross-sectional and did not assess cancer prevention or screening outcomes longitudinally. Surprisingly, only three studies reported results of interventions, therefore not allowing for a quantitative analysis of using APRN/PAs for cancer screening or prevention recommendations [32,38,39]. Few reported the type of APRN or PA provider separately, included comparison groups, or were based on well-described

Table 4 Studies of Advanced Practices Registered Nurses and Physician Assistants (APRN/PA) and diet, physical activity, and smoking cessation recommendations

Study	Sample size	Outcome measure	Setting	Findings for diet	Findings for physical activity	Findings for smoking cessation
Tompkins et al., [33]	Total providers: 398 NPs: 398	Self-report of NPs	Survey of NPs at Pacific Northwest Annual National Conference Cross sectional survey	NA	Physical activity counseling of appropriate patient in past week: 25% of NPs reported counseling 50% of appropriate patients 37.75% of NPs reported counseling 75% of appropriate patients 14.8% of NPs reported counseling 100% of appropriate patients Selected factors that facilitate physical activity counseling with patients: 69.2% of NPs reported length of patient visit 55.4% of NPs reported part of preventative health visit P-value not reported	NA
Patton et al., [25]	Total providers: 1802 Family physicians: 273 NPs: 294 Dentists: 584 Hygienists: 651	Self-report of MDs and NPs	Surveys of health professionals in North Carolina Cross sectional Survey	NA	NA	NPs report that they assess: Patient's past tobacco use: 95.1% Patient's present tobacco use: 97.9% Type and amount of tobacco: 92.3% P-value not reported Family MDs report that they assess: Patient's past tobacco use: 98.5% Patient's present tobacco use: 100% Type and amount of tobacco: 95.5% P-value not reported Adequately trained for smoking cessation NPs: 71.4% Family MD: 93.5% P-value not reported Physicians are significantly more likely to feel adequately trained to provide tobacco cessation compared to NPs

Table 4 Studies of Advanced Practices Registered Nurses and Physician Assistants (APRN/PA) and diet, physical activity, and smoking cessation recommendations (Continued)

						OR = 5.3 (3.2 - 8.6) P-value<.0001 All measures unadjusted CNMs reported that they always/usually: Document cigarette smoking use status at each visit: 73% Assess whether the patient is willing to make a quit attempt within the next 30 days: 66% Use counseling to help patients willing to make a quit attempt: 48% P-value not reported for all measures All measures unadjusted
Price et al., [35]	Total providers: 194 CNMs: 194 All patients pregnant women	Self-report of CNMs	Survey of CNMs in Ohio Cross sectional survey	NA	NA	
Running et al., [36]	Total patients: 400 NP patients: 200 Physician patients: 200	Chart review	Chart review of urgent care setting in HMO in the Southwest Retrospective cohort	NA	NA	Smoking cessation addressed among non-pharmacological interventions for sinusitis: NPs: 49% MDs: 31% Number of times smoking cessation is addressed for subjects in all categories NPs: .97 MDs: 1.95 P-value=.309 Unadjusted
Hopkins et al., [26]	Total Patients: 1339 Primary health care center (PHCC) NP patients: 755 MD patients: 441 Private practice NP patients: 143	Chart review	Chart review in private practice and primary care health centers in NY City Retrospective cohort	Receipt of assessment and counseling on nutrition and diet: PHCC NPs: 41.4% PHCC MDs: 14.7% P-value=0.000 Unadjusted	Receipt of assessment and counseling on physical activity: PHCC NPs: 15.8% PHCC MDs: 2.5% P-value=0.000 Unadjusted	Receipt of assessment and counseling on tobacco use: PHCC NPs: 79.2% PHCC MDs: 87.8% P-value=0.000 Unadjusted

Table 4 Studies of Advanced Practices Registered Nurses and Physician Assistants (APRN/PA) and diet, physical activity, and smoking cessation recommendations (Continued)

Lin et al., [27]	Total hospital outpatient department visits: 90,476 Visits with NP: 6,062 Visits without NP: 84,416	Chart review	National survey of hospital ambulatory settings (NAMCS) Cross sectional survey Odds ratio adjusted for patient age, sex, clinic type, metropolitan status, geographic region of hospital, and number of providers seen.	Received diet counseling at an OPD visit with a NP compared to one without a NP 32.6% vs. 22.9% Non-illness patients: 1.7 OR (95% CI OR: 1.2-2.5) P- value = 0.004 OPD visits for patients with chronic problems with a NP compared to one without a NP: 32.3% vs. 17.1% 2.5 OR (95% CI OR: 1.6-3.8) P-value = 0.001	Received physical activity counseling at an OPD visit with a NP compared to one without a NP 14.5% vs. 9.3% Non-illness patients: 1.8 OR (95% CI OR: 1.2-2.8) P- value = 0.007 OPD visits for patients with chronic problems with a NP compared to one without a NP: 20.2% vs. 8.9% 2.8 OR (95% CI OR: 1.6-5.1) P-value = 0.007	Received tobacco use counseling at an OPD visit with a NP compared to one without a NP 6.7% vs. 4.3% Non-illness patients: 1.7 OR (95% CI OR: 1.2-2.5) P- value = 0.004 OPD visits for patients with chronic problems with a NP compared to one without a NP: 4.7% vs.2.9% 1.8 OR (95% CI OR: 1.1-3.0) P-value = 0.01
Moody et al., [37]	Total Providers: 44 NPs: 44 Total patients: 680	Self-report of NPs	Survey of NPs in Tennessee Cross sectional survey	Provider report nutrition counseling: NPs: 19% MDs: 15% P-value not reported Unadjusted	Provider report physical activity counseling: NPs: 12% MDs: 7% P-value not reported Unadjusted	Provider report smoking cessation counseling: NPs: 7% MDs: 2.5% P-value not reported Unadjusted
Gebauer et al., [38]	Total patients: 178 Control patients: 94 Intervention patients: 84 All patients pregnant women who report smoking and intervention includes being seen by an advance practice NP	Self-report of patients and Salivary Cotinine Sample	Follow up survey at outpatient obstetric clinic - state not specified Intervention	NA	NA	Smoking rate/day at follow-up: Mean (SD) Control: 13.7 (14.1) Intervention: 7.8 (7.3) P =.008 Unadjusted Smoked any amount in past 7 days: Control Baseline: 94 participants Control Follow up: 94 participants Intervention Baseline: 83 participants Intervention Follow up: 70 participants Difference between groups =15.5% P-value<0.001 Unadjusted

Table 4 Studies of Advanced Practices Registered Nurses and Physician Assistants (APRN/PA) and diet, physical activity, and smoking cessation recommendations (Continued)

Murphy, [31]	Total providers: 346	Self-report of CNMs	National survey of CNMs	Nutritional counseling of gynecologic patients	Physical activity counseling of gynecologic patients	Smoking cessation counseling of gynecologic patients
	CNMs: 346		Cross sectional survey	52% of CNMs report counseling 81-100% of their patients P-value not reported Unadjusted	46% of CNMs report counseling 81-100% of their patients P-value not reported Unadjusted	72% of CNMs report counseling 81-100% of their patients P-value not reported Unadjusted
Zahnd et al, [32]	Total patients: 1217	Self-report of patients	Survey of patients from Four Kaiser Permanente Medical Centers in San Francisco Bay Area	NA	NA	Patients report discussing smoking cessation:
	NP patients: 269					NP Patients: 64%
	Physician patients: 948					MD Patients: 50%
	Total providers: 52					P-value<0.001
Physicians: 40	Intervention	Unadjusted	Independent predictors of counseling about smoking:			
NPs: 12			NP vs. Physician: OR 1.7			
			P-value=.0006			
			Adjusted for differences in patient characteristics			

*=A higher number means the provider does the behavior more frequently.

CI = Confidence Interval.

NA = Not Applicable.

NPC = Non-Primary Care.

NP/PA = NP or PA.

PHCC = Primary Health Care Center.

PC = Primary Care.

PP = Private practice.

samples (Tables 3 and 4) [24-28,36,38,39]. In addition, studies that did include comparison groups did not consistently report on statistical significance of comparisons. Inconsistencies in outcome measure reporting among these studies impacted our ability to compare guideline adherence and patient populations. Few studies evaluated whether screening recommendations were consistent with evidence-based guidelines for patient age at initiation or frequency [24,26]. This is particularly important because both overuse and underuse of screening can have adverse patient outcomes [47-49]. Most of the studies neglected to report patient demographics or key covariates, such as weight, body mass index, and comorbidities, hindering our ability to determine if either physicians or APRN/PAs are providing cancer screening based on guidelines.

Outcome measures were most commonly reported using either provider or patient self-reported data about recommendations and did not report on receipt of service or a documented change in behavior [24,25,30-33,35,37]. Even further removed from receipt of service, some studies reported what physicians perceived of APRN/PAs practice [28,29]. Self-reported and proxy-reported data may over or underestimate documented receipt of APRN/PA provider services [50]. Further, primary care addresses multiple preventive services, but only about half of the studies included more than one aspect of cancer control and no studies address post-treatment survivorship care [24,26,27,31,37,39]. Future research should address these limitations and be conducted in longitudinal cohorts with comparison groups of well-described provider types, document patient receipt of screening or prevention recommendations, and assess multiple cancer control recommendations. Use of standardized measures, including for patient characteristics associated with guideline recommendations, evaluation of guideline adherence and longer term patient outcomes will also be important.

Despite using a large number of search terms to identify published studies, manually reviewing all abstracts and relevant reference lists, it is possible we missed some relevant studies. The studies we identified were fairly heterogeneous in terms of patient populations, geographic region, provider type, and type of a comparison group. Additionally, included studies used a variety of approaches to measure cancer screening and prevention, such as physician, non-physician provider and patient self-report, as well as chart review. As a result, our synthesis of findings was descriptive rather than quantitative. Findings are generalizable only to the primary care setting.

Conclusion

In summary, further documentation of the role of APRN/PAs in recommending and providing cancer prevention and screening services in US primary care teams

is needed. Ensuring that future research measures cancer screening according to evidence-based USPSTF guidelines decreases variability among measures reporting and focuses on receipt of services. This will allow stakeholders to make more informed decisions on how best to utilize this growing workforce and provide cancer prevention and screening services to the US population.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AS reviewed articles for initial eligibility, created tables and drafted manuscript. AS, DK and RY reviewed final articles for eligibility, provided study analysis and edited the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to acknowledge Ms. Echo Warner, MPH (Research Analyst at the Huntsman Cancer Institute) for her assistance on the literature review for this study.

Author details

¹American Legacy Foundation, Washington, DC, USA. ²Division of Cancer Control and Population Sciences, National Cancer Institute, Bethesda, MD, USA. ³College of Nursing, University of Utah, Salt Lake City, UT. ⁴Cancer Control and Population Sciences, Huntsman Cancer Institute, Salt Lake City, UT.

Received: 16 August 2013 Accepted: 7 February 2014

Published: 12 February 2014

References

1. Addressing The Cancer Burden: At A Glance. <http://www.cdc.gov/chronicdisease/resources/publications/AAG/dcpch.htm>.
2. Screening for Cervical Cancer. <http://www.uspreventiveservicestaskforce.org/uspstf/uspsscerv.htm>.
3. Screening for Breast Cancer. <http://www.uspreventiveservicestaskforce.org/uspstf/uspssbrca.htm>.
4. Screening for Colorectal Cancer. <http://www.uspreventiveservicestaskforce.org/uspstf/uspsscolo.htm>.
5. Behavioral counseling to promote a healthful diet and physical activity for cardiovascular disease prevention in adults. <http://www.uspreventiveservicestaskforce.org/uspstf/uspssphys.htm>.
6. Ward E, Jemal A, Cokkinides V, Singh GK, Cardinez C, Ghafoor A, Thun M: Cancer disparities by race/ethnicity and socioeconomic status. *CA Cancer J Clin* 2004, **54**(2):78-93.
7. Freeman HP: Poverty, culture, and social injustice: determinants of cancer disparities. *CA Cancer J Clin* 2004, **54**(2):72-77.
8. Halle M, Lewis CB, Seshamani M: Health Disparities: A Case For Closing the Gap. In *U.S. Department of Health and Human Services. Government: Health Reform*; 2009.
9. The Affordable Care Act. <http://www.whitehouse.gov/healthreform/timeline>.
10. Take healthcare into your own hands. <http://www.healthcare.gov/prevention/index.html>.
11. IOM (Institute of Medicine): *The Future of Nursing: Leading Change, Advancing Health*. Washington, DC: The National Academies Press; 2011.
12. Division of Medicine and Dentistry, Bureau of Health Professions, Health Resources and Services Administration, U.S. Department of Health and Human Services: *Affordable Care Act: Expansion of Physician Assistant Training Program, Frequently Asked Questions*; 2010. Revised July 8, 2010, on the Internet at: <http://bhpr.hrsa.gov/grants/expansionfaq.pdf>.
13. Grumbach K, Hart LG, Mertz E, Coffman J, Palazzo L: Who is caring for the underserved? A comparison of primary care physicians and nonphysician clinicians in California and Washington. *Ann Fam Med* 2003, **1**(2):97-104.
14. Cooper RA: New directions for nurse practitioners and physician assistants in the era of physician shortages. *Acad Med* 2007, **82**(9):827-828.

15. Park M, Cherry D, Decker SL: **Nurse practitioners, certified nurse midwives, and physician assistants in physician offices.** *NCHS Data Brief* 2011, **69**:1–8.
16. Bureau of Labor Statistics, U.S. Department of Labor: **Occupational Outlook Handbook, 2012-2013 Edition, Physician Assistants.** on the Internet at <http://www.bls.gov/ooh/healthcare/physician-assistants.htm>.
17. Yarnall KS, Pollak KI, Ostbye T, Krause KM, Michener JL: **Primary care: is there enough time for prevention?** *Am J Public Health* 2003, **93**(4):635–641.
18. Wagner EH: **Chronic disease management: what will it take to improve care for chronic illness?** *Eff Clin Pract* 1998, **1**(1):2–4.
19. Wagner EH: **The role of patient care teams in chronic disease management.** *BMJ* 2000, **320**(7234):569–572.
20. Wagner EH, Austin BT, Von Korff M: **Organizing care for patients with chronic illness.** *The Milbank quarterly* 1996, **74**(4):511–544.
21. Coleman K, Austin BT, Brach C, Wagner EH: **Evidence on the chronic care model in the new millennium.** *Health Aff* 2009, **28**(1):75–85.
22. Yabroff KR, Kerner JF, Mandelblatt JS: **Effectiveness of interventions to improve follow-up after abnormal cervical cancer screening.** *Prev Med* 2000, **31**(4):429–439.
23. Mandelblatt J, Armetta C, Yabroff KR, Liang W, Lawrence W: **Descriptive review of the literature on breast cancer outcomes, through 2000.** *J Natl Cancer Inst Monogr* 1990, **2004**(33):8–44.
24. Menees SB, Patel DA, Dalton V: **Colorectal cancer screening practices among obstetrician/gynecologists and nurse practitioners.** *J Women's Health* 2009, **18**(8):1233–1238.
25. Patton LL, Ashe TE, Elter JR, Southerland JH, Strauss RP: **Adequacy of training in oral cancer prevention and screening as self-assessed by physicians, nurse practitioners, and dental health professionals.** *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2006, **102**(6):758–764.
26. Hopkins SC, Lenz ER, Pontes NM, Lin SX, Mundinger MO: **Context of care or provider training: the impact on preventive screening practices.** *Prev Med* 2005, **40**(6):718–724.
27. Lin SX, Gebbie KM, Fullilove RE, Arons RR: **Do nurse practitioners make a difference in provision of health counseling in hospital outpatient departments?** *J Am Acad Nurse Pract* 2004, **16**(10):462–466.
28. Sansbury LB, Klabunde CN, Mysliwiec P, Brown ML: **Physicians' use of nonphysician healthcare providers for colorectal cancer screening.** *Am J Prev Med* 2003, **25**(3):179–186.
29. Oliveria SA, Altman JF, Christos PJ, Halpern AC: **Use of nonphysician health care providers for skin cancer screening in the primary care setting.** *Prev Med* 2002, **34**(3):374–379.
30. Shaheen NJ, Crosby MA, O'Malley MS, Murray SC, Sandler RS, Galanko JA, Ransohoff DF, Klenzak JS: **The practices and attitudes of primary care nurse practitioners and physician assistants with respect to colorectal cancer screening.** *Am J Gastroenterol* 2000, **95**(11):3259–3265.
31. Murphy PA: **Primary care for women. Health assessment, health promotion, and disease prevention services.** *J Nurse-Midwifery* 1996, **41**(2):83–91.
32. Zahnd EG, Coates TJ, Richard RJ, Cummings SR: **Counseling medical patients about cigarette smoking: a comparison of the impact of training on nurse practitioners and physicians.** *Nurse Pract* 1990, **15**(3):10–13. 17–18.
33. Tompkins TH, Belza B, Brown MA: **Nurse practitioner practice patterns for exercise counseling.** *J Am Acad Nurse Pract* 2009, **21**(2):79–86.
34. Kawamoto K, Houlihan CA, Balas EA, Lobach DF: **Improving clinical practice using clinical decision support systems: a systematic review of trials to identify features critical to success.** *BMJ* 2005, **330**(7494):765.
35. Price JH, Jordan TR, Dake JA: **Perceptions and use of smoking cessation in nurse-midwives' practice.** *J Midwifery Womens Health* 2006, **51**(3):208–215.
36. Running A, Kipp C, Mercer V: **Prescriptive patterns of nurse practitioners and physicians.** *J Am Acad Nurse Pract* 2006, **18**(5):228–233.
37. Moody NB, Smith PL, Glenn LL: **Client characteristics and practice patterns of nurse practitioners and physicians.** *Nurse Pract* 1999, **24**(3):94–96. 99–100, 102–103.
38. Gebauer C, Kwo CY, Haynes EF, Wewers ME: **A nurse-managed smoking cessation intervention during pregnancy.** *J Obstet Gynecol Neonatal Nurs* 1998, **27**(1):47–53.
39. Mandelblatt J, Traxler M, Lakin P, Thomas L, Chauhan P, Matseoane S, Kanetsky P: **A nurse practitioner intervention to increase breast and cervical cancer screening for poor, elderly black women. The Harlem study team.** *J Gen Intern Med* 1993, **8**(4):173–178.
40. Newhouse RP, Stanik-Hutt J, White KM, Johantgen M, Bass EB, Zangaro G, Wilson RF, Fountain L, Steinwachs DM, Heindel L, et al: **Advanced practice nurse outcomes 1990–2008: a systematic review.** *Nurs Econ* 2011, **29**(5):230–250. quiz 251.
41. **Cancer Screening in the United States.** http://www.cdc.gov/cancer/colorectal/what_cdc_is_doing/screening_us.htm.
42. Donelan K, DesRoches CM, Dittus RS, Buerhaus P: **Perspectives of physicians and nurse practitioners on primary care practice.** *N Engl J Med* 2013, **368**(20):1898–1906.
43. Halpern MT, Ward EM, Pavluck AL, Schrag NM, Bian J, Chen AY: **Association of insurance status and ethnicity with cancer stage at diagnosis for 12 cancer sites: a retrospective analysis.** *Lancet Oncol* 2008, **9**(3):222–231.
44. Taplin SH, Ichikawa L, Yood MU, Manos MM, Geiger AM, Weinmann S, Gilbert J, Mouchawar J, Leyden WA, Altaras R, et al: **Reason for late-stage breast cancer: absence of screening or detection, or breakdown in follow-up?** *J Natl Cancer Inst* 2004, **96**(20):1518–1527.
45. Leyden WA, Manos MM, Geiger AM, Weinmann S, Mouchawar J, Bischoff K, Yood MU, Gilbert J, Taplin SH: **Cervical cancer in women with comprehensive health care access: attributable factors in the screening process.** *J Natl Cancer Inst* 2005, **97**(9):675–683.
46. **The Power to Prevent, The Call to Control: At A Glance.** 2009. <http://www.cdc.gov/chronicdisease/resources/publications/aag/chronic.htm>.
47. Korenstein D, Falk R, Howell EA, Bishop T, Keyhani S: **Overuse of health care services in the United States: an understudied problem.** *Arch Intern Med* 2012, **172**(2):171–178.
48. Kale MS, Bishop TF, Federman AD, Keyhani S: **Trends in the overuse of ambulatory health care services in the United States.** *JAMA Internal Medicine* 2013, **173**(2):142–148.
49. Chassin MR, Galvin RW: **The urgent need to improve health care quality.** Institute of Medicine National Roundtable on Health Care Quality. *JAMA* 1998, **280**(11):1000–1005.
50. Berry JA: **Nurse practitioners' use of clinical preventive services.** *J Am Acad Nurse Pract* 2009, **21**(8):454–460.

doi:10.1186/1472-6963-14-68

Cite this article as: Smith et al: Advanced practice registered nurses, physician assistants and cancer prevention and screening: a systematic review. *BMC Health Services Research* 2014 **14**:68.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

