

Research Article

The Relationship between VEGFC Gene Polymorphisms and Autoimmune Thyroiditis

Chaoqun Gao ^{1,2}, Jie Zhu,² Qiu Qin,² Xiaorong Yang,² Yanfei Jiang,² and Jinan Zhang ^{2,3}

¹Graduate School of Shanghai University of Traditional Chinese Medicine, Shanghai 201203, China

²Department of Endocrinology & Rheumatology, Shanghai University of Medicine & Health Sciences Affiliated Zhoupu Hospital, Shanghai 201318, China

³Shanghai University of Traditional Chinese Medicine, Shanghai 201203, China

Correspondence should be addressed to Jinan Zhang; zhangjinan@hotmail.com

Received 5 February 2022; Revised 6 May 2022; Accepted 17 May 2022; Published 12 July 2022

Academic Editor: Pei Jiang

Copyright © 2022 Chaoqun Gao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Autoimmune thyroid diseases (AITDs), representative autoimmune diseases, mainly consist of Graves' disease (GD) and Hashimoto's thyroiditis (HT). In this passage, we investigated the association between vascular endothelial growth factor C (VEGFC) gene polymorphisms and AITDs. **Methods.** A total of 1084 patients with AITDs and 794 healthy controls were tested for VEGFC gene genotypes in four single nucleotide polymorphisms (SNPs) by high-throughput sequencing, and the correlation between VEGFC gene polymorphisms and AITDs was statistically analyzed. **Results.** The genotype distribution of rs3775194 was statistically associated with AITDs compared with the control group. Rs3775194 was associated with AITDs under the overdominant model, both before and after adjusting for confounding factors, while the other three SNPs were not associated with GD and HT. There was a prominent discrepancy between male healthy controls and male AITD patients under overdominant model in rs3775194 and the recessive model in rs11947611. The genotype distribution of rs3775194 was statistically related to male HT. **Conclusion.** These results reveal the correlation between VEGFC mutation and AITD susceptibility.

1. Introduction

AITDs are thyroid-specific autoimmune diseases caused by the disorder of autoimmune mechanisms, among which GD and HT are the major subtypes [1]. GD and HT have different clinical manifestations and pathophysiological characteristics. GD shows elevated TSH receptor stimulating antibody (TRAb), accompanied with lymphocyte infiltration in the thyroid gland and hypertrophy of thyroid follicular epithelial cells. HT shows interstitial fibrous tissue hyperplasia and destruction of thyroid follicles, accompanied with elevated anti-thyroglobulin antibody (TgAb) and thyroid peroxidase antibody (TPOAb). According to related research, AITDs affect about 5% of the total population and are more popular in women than in men [2]. The pathogenesis of AITDs has not been clearly studied. Genetics, immunity, and environment may be involved in the occurrence and development of AITDs [3].

Vascular endothelial growth factor (VEGF) mainly involves in neovascular diseases such as malignant tumors and plays a role in increasing vascular permeability and mediating inflammation. It is affiliated with the platelet-derived growth factor family. Angiogenesis plays a role in a variety of autoimmune inflammatory diseases, including rheumatoid arthritis [4], systemic lupus erythematosus [5], and systemic sclerosis [6]. Inhibition of angiogenesis may be a promising treatment of these diseases. VEGFs are angiogenic factors which contain five members in mammals: VEGFA, VEGFB, VEGFC, VEGFD, and placental growth factor (PGF). VEGFC is encoded in humans by the VEGFC gene, which is located on chromosome 4q34. VEGFC is a ligand of VEGF-R3 and VEGF-R2, but it exerts effects mainly through VEGF-R3 [7]. It is widely believed that VEGFC plays a major role in lymphangiogenesis and promotes the survival, growth, and migration of lymphatic endothelial cells

TABLE 1: Clinical features and demographic statistics of AITD patients and controls.

Items	AITD (%)	GD (%)	HT (%)	Controls (%)
Number	1084	667	417	794
Age	41.7 ± 14.3	41 ± 14.6	42.8 ± 13.8	38.9 ± 10.5
Sex				
Females	828 (76.4)	476 (71.4)	352 (84.4)	477 (60.1)
Males	256 (23.6)	191 (28.6)	65 (15.6)	317 (39.9)
Ophthalmopathy (+)	/	99 (15.8)	/	/
Hypothyroidism (+)	/	/	174 (42.5)	/

(LECs) through its receptor VEGFR-3 [8]. We hypothesized that VEGFC gene is related to autoimmune thyroid diseases. Therefore, we investigated the association between VEGFC gene polymorphisms and AITD risk using a case-control approach.

2. Methods

2.1. Recruitment of the Participants. We recruited 1084 patients with AITD and 794 healthy controls from the Han Chinese population. AITD patients consisted of 256 men and 828 women. The normal healthy group was made up of 317 men and 477 women. To eliminate sampling error, all AITD patients were randomly recruited from the outpatient department of Shanghai Zhoupu Hospital, and the normal controls were consecutively enrolled from physical examination center of the same hospital. All AITD patients and normal controls were free of other autoimmune and inflammatory diseases. The participants were all from Shanghai, China. The study was permitted by the Ethics Committee of Shanghai University of Medicine & Health Sciences, and all the subjects in the study provided written informed consent.

In the AITD patients, there were 667 patients with GD (191 males and 476 females) and 417 patients with HT (65 males and 352 females). Patients with GD in the study were required to meet the following criteria: typical symptoms of hyperthyroidism, biochemical tests of hyperthyroidism, and positive TRAb [9]. Positive TPOAb or TgAb, and ultrasonographic findings of diffuse thyroid changes are the basis for the diagnosis of HT. TRAb, TgAb, TPOAb, and other serological parameters were detected by immunochemiluminescence kit (Roche Company, Switzerland) with high quality.

In order to explore the correlation between different clinical manifestations and genetic background, the relationship between SNP and various clinical subtypes was elaborated. In this study, the clinical manifestations such as Graves' ophthalmopathy (GO) in GD patients and hypothyroidism in HT patients were included. Table 1 summarizes the clinical characteristics of all the subjects. GO, also known as thyroid-related eye disease, is characterized by exophthalmos, excessive tearing, painful eye movement, and diplopia. We can make a diagnosis based on the criteria in Williams Textbook of Endocrinology.

2.2. Isolation and Extraction of DNA Samples. Two-milliliter peripheral venous blood was collected from each individual. Patients' DNA were extracted using Relax Gene Blood DNA System (Tiangen Biotech Co., Ltd., Beijing, China). The quality of genomic DNA was evaluated by Nano Drop 2000 spectrophotometer (Thermo Scientific Company, Waltham, USA) according to the manufacturer's guidelines.

2.3. SNP Selection and Genotyping. This study examined the four SNPs of VEGFC (i.e., rs7664413, rs11947611, rs2046463, and rs3775194). According to previous reports, these loci are associated with several autoimmune diseases. We hypothesized that these loci are also associated with AITD susceptibility. The four SNPs need to conform to Hardy-Weinberg equilibrium (HWE) with P value greater than 0.05 and minor allele frequency (MAF) > 0.05. Genotypes of DNA samples were tested with high-throughput SNP sequencing. In simple terms, the DNA sample was amplified in 10 microliter volume. The temperature condition was set to 95 °C for 15 minutes, followed by five cycles of 94 °C for 30s, 60 °C for 4 minutes, and 72 °C for 30 seconds and 10 cycles 94 °C for 30s, 60 °C for 1 minute, and 72 °C for 30s. The primer sequence we used is GAGT TTCTGTCTAGTTCTTTGTGG and GGAAAACATAC AAAAGGAAGATGC for rs7664413. For rs11947611, the primer sequences are AAACCTTGGCTTCTAACAATC TTC and CCTGAAAACATAAACCAAAAAGCC; TGTT TACGATACTCTCACTTTTGG and GGCCATGTAAA GAATAGTAGAACC are the primer sequences of rs2046463. TCACAGCTTAAGACTGAAATCAAC and ATTCTGTGACGATGTACTATAGGG are the primer sequences of rs3775194, respectively. In order to ensure the accuracy of the genotyping process, PCR was performed twice each time and negative control was performed using water as template.

2.4. Statistical Analysis. Statistical analysis included OR values, P values and 95% confidence intervals. We used SPSS (22.0, IBM, Chicago, USA) for our calculations. Measurement data are expressed as the mean plus or minus standard errors. The alleles and genotype frequency of SNP were analyzed by Chi-square test. By adjusting potential confounding factors such as thyroid function and gender, P and OR values before and after adjustment were calculated by multifactor logistic regression analysis to find

TABLE 2: Associations of rs7664413, rs11947611, rs2046463, and rs3775194 in VEGFC gene with AITD, GD, and HT.

		AITD <i>n</i> (%)	NC <i>n</i> (%)	P value AITD vs NC	GD <i>n</i> (%)	P value GD vs NC	HT <i>n</i> (%)	P value HT vs NC
rs7664413	CC	495 (45.7)	350 (44.1)	0.652	293 (43.9)	0.364	202 (48.4)	0.238
	TT	122 (11.3)	85 (10.7)		87 (13.0)		35 (8.4)	
	TC	467 (43.1)	359 (45.2)		288 (43.1)		180 (43.2)	
	C	1457 (67.2)	1059 (66.7)	0.739	874 (65.4)	0.470	584 (70.0)	0.095
rs11947611	T	711 (32.8)	529 (33.3)		462 (34.6)		250 (30.0)	
	AA	410 (37.8)	318 (40.1)	0.527	251 (37.6)	0.534	159 (38.1)	0.765
	GG	167 (15.4)	111 (14.0)		104 (15.6)		63 (15.1)	
	AG	507 (46.8)	365 (46.0)		313 (46.9)		195 (46.8)	
rs2046463	A	1327 (61.2)	1001 (63.0)	0.255	815 (61.0)	0.259	513 (61.5)	0.461
	G	841 (38.8)	587 (37.0)	0.600	521 (39.0)		321 (38.5)	
	AA	492 (45.4)	347 (43.7)		291 (43.6)	0.315	201 (48.2)	0.224
	GG	123 (11.3)	85 (10.7)		88 (13.2)		35 (8.4)	
rs3775194	AG	469 (43.3)	362 (45.6)	0.737	289 (43.3)	0.459	181 (43.4)	0.089
	A	1453 (67.0)	1056 (66.5)		871 (65.2)		583 (69.9)	
	G	715 (33.0)	532 (33.5)		465 (34.8)		251 (30.1)	
	CC	12 (1.1)	4 (0.5)	0.029	8 (1.2)	0.092	4 (1.0)	0.052
rs3775194	GG	906 (83.6)	636 (80.1)		553 (82.8)		354 (84.9)	
	CG	166 (15.3)	154 (19.4)		107 (16.0)		59 (14.1)	
	C	190 (8.8)	162 (10.2)	0.135	123 (9.1)	0.310	67 (8.0)	0.083
	G	1978 (91.2)	1426 (89.8)		1213 (90.9)		767 (92.0)	

TABLE 3: Associations of four polymorphisms models in VEGFC with AITD before and after adjusting for confounders.

Comparison models	Unadjusted estimates		Adjusted estimates	
	OR (95% CI)	<i>P</i> value	OR(95% CI)	<i>P</i> value
rs7664413				
Allele model	0.98 (0.85-1.12)	0.74	0.97 (0.84-1.12)	0.69
Dominant model	0.94 (0.78-1.13)	0.50	0.93 (0.77-1.12)	0.45
Recessive model	1.06 (0.79-1.42)	0.71	1.06 (0.78-1.43)	0.72
Overdominant model	0.92 (0.76-1.10)	0.36	0.91 (0.75-1.10)	0.32
rs11947611				
Allele model	1.08 (0.95-1.23)	0.26	1.06 (0.92-1.21)	0.43
Dominant model	1.10 (0.91-1.32)	0.33	1.07 (0.89-1.30)	0.47
Recessive model	1.12 (0.86-1.45)	0.39	1.08 (0.83-1.40)	0.59
Overdominant model	1.03 (0.86-1.24)	0.73	1.03 (0.86-1.24)	0.74
rs2046463				
Allele model	0.98 (0.85-1.12)	0.74	0.97 (0.84-1.12)	0.69
Dominant model	0.93 (0.78-1.12)	0.47	0.93 (0.77-1.12)	0.43
Recessive model	1.07 (0.80-1.43)	0.66	1.06 (0.79-1.43)	0.69
Overdominant model	0.91 (0.76-1.09)	0.32	0.90 (0.75-1.09)	0.29
rs3775194				
Allele model	0.85 (0.68-1.05)	0.14	0.83 (0.66-1.05)	0.12
Dominant model	0.79 (0.62-1.00)	0.05	0.79 (0.62-1.00)	0.05
Recessive model	2.21 (0.71-6.88)	0.15	1.91 (0.60-6.08)	0.25
Overdominant model	0.75 (0.59-0.96)	0.02	0.75 (0.59-0.96)	0.02

TABLE 4: Associations of four polymorphisms models in VEGFC with male AITD before and after adjusting for confounders.

Comparison models	Unadjusted estimates		Adjusted estimates	
	OR (95% CI)	<i>P</i> value	OR (95% CI)	<i>P</i> value
rs7664413				
Allele model	1.07 (0.83-1.37)	0.60	1.05 (0.82-1.34)	0.72
Dominant model	1.01 (0.72-1.40)	0.96	0.98 (0.70-1.37)	0.90
Recessive model	1.32 (0.78-2.23)	0.30	1.30 (0.77-2.21)	0.33
Overdominant model	0.90 (0.65-1.26)	0.54	0.88 (0.63-1.23)	0.46
rs11947611				
Allele model	1.26 (0.98-1.61)	0.07	1.25 (0.97-1.60)	0.08
Dominant model	1.20 (0.86-1.69)	0.28	1.16 (0.82-1.63)	0.40
Recessive model	1.71 (1.04-2.82)	0.04	1.79 (1.08-2.98)	0.02
Overdominant model	0.95 (0.68-1.32)	0.74	0.89 (0.64-1.24)	0.49
rs2046463				
Allele model	1.05 (0.82-1.35)	0.68	1.03 (0.81-1.33)	0.79
Dominant model	0.98 (0.71-1.37)	0.92	0.96 (0.68-1.34)	0.79
Recessive model	1.32 (0.78-2.23)	0.30	1.30 (0.77-2.21)	0.33
Overdominant model	0.88 (0.63-1.23)	0.45	0.86 (0.62-1.20)	0.38
rs3775194				
Allele model	0.72 (0.48-1.09)	0.12	0.74 (0.49-1.12)	0.15
Dominant model	0.65 (0.42-1.00)	0.05	0.66 (0.43-1.02)	0.06
Recessive model	—	0.03	—	0.02
Overdominant model	0.59 (0.38-0.92)	0.02	0.60 (0.38-0.94)	0.02

TABLE 5: Allele and genotype distributions of VEGF loci in subgroups of HT patients.

		Female controls <i>n</i> (%)	Female HT <i>n</i> (%)	Unadjust/ adjust <i>P</i> value		Male controls <i>n</i> (%)	Male HT <i>n</i> (%)	Unadjust/ adjust <i>P</i> value	
rs7664413	CC	207 (43.4)	170 (48.3)	0.208		143 (45.1)	32 (49.2)	0.831	
	TT	54 (11.3)	29 (8.2)			31 (9.8)	6 (9.2)		
	TC	216 (45.2)	153 (43.5)			143 (45.1)	27 (41.5)		
	C	493 (70.0)	630 (66.0)	0.082	0.07	429 (67.7)	91 (70.0)	0.600	0.49
	T	211 (30.0)	324 (34.0)			205 (32.3)	39 (30.0)		
rs11947611	AA	185 (38.8)	136 (38.6)	0.627		133 (42.0)	23 (35.4)	0.120	
	GG	80 (16.8)	51 (14.5)			31 (9.7)	12 (18.5)		
	AG	212 (44.4)	165 (46.9)			153 (48.3)	30 (46.1)		
rs2046463	A	582 (61.0)	437 (62.1)	0.660	0.69	419 (66.1)	76 (58.5)	0.091	0.19
	G	372 (39.0)	267 (37.9)			215 (33.9)	54 (41.5)		
	AA	206 (43.2)	169 (48.0)	0.212		141 (44.5)	32 (49.2)	0.780	
	GG	54 (11.3)	29 (8.2)			31 (9.8)	6 (9.2)		
	AG	217 (45.5)	154 (43.8)			145 (45.7)	27 (41.5)		
rs3775194	A	629 (65.9)	492 (69.9)	0.085	0.07	427 (67.4)	91 (70.0)	0.550	0.46
	G	325 (34.1)	212 (30.1)			207 (32.6)	39 (30.0)		
	CC	4 (0.8)	3 (0.9)	0.506		0 (0)	1 (1.5)	0.048	
rs3775194	GG	388 (81.3)	297 (84.4)			248 (78.2)	57 (87.7)		
	CG	85 (17.8)	52 (14.8)			69 (21.8)	7 (10.8)		
	C	93 (9.7)	58 (8.2)	0.290	0.23	69 (10.9)	9 (6.9)	0.140	0.17
	G	861 (90.3)	646 (91.8)			565 (89.1)	121 (93.1)		

out meaningful results. The correlation between VEGFC gene polymorphisms and AITD was further calculated under allele model, overdominant model, and recessive model. Linkage analysis was calculated using Haploview 4.2 software (Broad Institute, Cambridge, USA) with $p < 0.05$ deemed as positive.

3. Results

Table 2 shows the distribution of alleles and genotypes of VEGFC loci in AITD, GD, HT, and controls. No meaningful conclusions were found in the alleles and genotypes distributions of these four SNPs in GD and HT patients between cases and those in the control group. However, the genotype distribution of rs3775194 in AITD subjects (CC 1.1%, GG 83.6%, and CG 15.3%) was statistically different from that in the normal group (CC 0.5%, GG 80.1%, CG 19.4%) ($P = 0.029$). To further analyse the potential association between VEGFC gene polymorphisms and AITDs, we performed an analysis of four models before and after adjusting for confounding factors (age, sex). As demonstrated in Table 3, rs3775194 locus was strongly associated with AITD in overdominant model, both before and after adjusting for confounders ($P = 0.021$ and 0.024 , respectively). No positive results were found in our study comparing GD and HT with normal controls (data are not presented).

To further investigate whether VEGFC loci are associated with AITD in different genders, as presented in Table 4, we found that rs11947611 was associated with male AITD patients in a recessive model, both before and after adjustment for confounders ($P = 0.035$ and 0.024 , respectively). Rs3775194 was associated with male AITD patients in an overdominant model with P values of 0.017 and 0.023 before and after adjustment for confounding factors. Table 5 shows the genotypes and allele frequencies of these four loci in different sex subgroups of HT patients. The genotype distribution of rs3775194 in male HT patients (CC 1.5%, GG 87.7%, and CG 10.8%) was significantly different from that in male control group (GG 78.2% and CG 21.8%) ($P = 0.048$), but there were no positive results in allele frequency between the two groups. We have not yet found an association between VEGFC loci and female AITD patients (Table 6). No distinguishable difference was found between the female HT and the control group in four loci.

We further analyzed the correlation between four loci of VEGFC and some clinical phenotypes. It can be seen from Table 7 that there was no correlation between four polymorphisms of VEGFC and susceptibility to Graves' ophthalmopathy (GO). Table 8 shows that the distribution of these loci was not associated with hypothyroidism in HT patients, either.

Haploview software showed that rs2046463 and rs7664413 formed only one linkage disequilibrium (LD) region and three main haplotypes: CA, TG, and CG. However, we did not detect an association of VEGFC haplotypes CA, TG, and CG (Figure 1) with susceptibility to AITD, GD, and HT.

TABLE 6: Associations of four polymorphisms models in VEGFC with female AITD before adjusting for confounders.

	OR (95% CI)	<i>P</i> value
rs7664413		
Allele model	0.96 (0.83-1.12)	0.62
Dominant model	0.93 (0.76-1.13)	0.46
Recessive model	1.02 (0.74-1.39)	0.92
Overdominant model	0.92 (0.76-1.12)	0.42
rs11947611		
Allele model	1.08 (0.93-1.24)	0.31
Dominant model	1.09 (0.90-1.34)	0.38
Recessive model	1.11 (0.85-1.47)	0.44
Overdominant model	1.03 (0.85-1.25)	0.76
rs1485766		
Allele model	1.00 (0.88-1.15)	0.95
Dominant model	1.09 (0.88-1.36)	0.43
Recessive model	0.92 (0.73-1.15)	0.48
Overdominant model	1.14 (0.94-1.39)	0.18
rs2046463		
Allele model	0.97 (0.83-1.12)	0.64
Dominant model	0.93 (0.76-1.13)	0.46
Recessive model	1.03 (0.75-1.41)	0.85
Overdominant model	0.92 (0.75-1.12)	0.39
rs3775194		
Allele model	0.86 (0.68-1.09)	0.22
Dominant model	0.81 (0.63-1.04)	0.11
Recessive model	2.17 (0.67-7.08)	0.18
Overdominant model	0.77 (0.60-1.00)	0.05

4. Discussion

Epidemiological studies have confirmed that genetic factors play an important role in AITDs, but the known genes related to AITDs cannot fully explain the role of genetic factors in AITDs. In this study, we explored the relationship between VEGFC gene polymorphisms and AITDs using allele, dominant, recessive, and overdominant models as well as different subgroups of AITD. We found that the rs3775194 locus was associated with AITD patients, but not with GD or HT patients. We further found that rs3775194 was associated with male AITD patients under the overdominant model and rs3775194 was associated with the genotype distribution of male HT patients. Rs11947611 is associated with male AITD patients under the recessive model.

VEGF is a functional glycoprotein with high biological activity. It is also called vascular permeability factor due to its strong ability to promote the differentiation and proliferation of vascular endothelial cells. VEGF gene is involved in the occurrence and development of diabetic retinopathy and cancer [10, 11]. A study of 1 919 diabetic patients with gene polymorphisms found that three SNPs (rs17697419, rs17697515, and rs2333526) of VEGFC are associated with

TABLE 7: Allele and genotype distributions of VEGFC polymorphisms with ophthalmopathy in GD patients.

SNPs	Without (%)	With (%)	<i>P</i> value	OR (95% CI)
rs7664413				
CC	234 (44.4)	47 (47.5)	0.212	1.044 (0.759-1.437)
TT	64 (12.1)	17 (17.2)		
TC	229 (43.5)	35 (35.4)		
C	697 (66.1)	129 (65.2)		
T	357 (33.9)	69 (34.8)		
rs11947611				
AA	192 (36.4)	42 (42.4)	0.433	0.811 (0.59-1.113)
GG	84 (15.9)	12 (12.1)		
AG	251 (46.7)	45 (45.5)		
A	635 (60.2)	129 (65.2)		
G	419 (39.8)	69 (34.8)		
rs2046463				
AA	233 (44.2)	46 (46.5)	0.152	1.087 (0.791-1.493)
GG	64 (12.1)	18 (18.2)		
AG	230 (43.6)	35 (35.4)		
A	696 (66.0)	127 (64.1)		
G	358 (34.0)	71 (35.9)		
rs3775194				
CC	4 (0.8)	3 (3.0)	0.153	0.756 (0.462-1.237)
GG	440 (83.5)	80 (80.8)		
CG	83 (15.7)	16 (16.2)		
C	91 (8.6)	22 (11.1)		
G	963 (91.4)	176 (88.9)		

diabetic retinopathy. Rs17697515 is also specifically associated with diabetic macular edema in T2DM patients [12]. VEGFC gene also plays an important role in the development of various autoimmune diseases such as rheumatoid arthritis [4]. VEGF-C/sVEGFR-3 ratio is significantly lower in patients with Behcet's disease than in the control group and is correlated with the course of the disease [13]. In addition, serum VEGFC values are higher in adult-onset Still's disease, which may be a marker of disease activity [14]. On the other hand, VEGFC levels are low in patients with systemic sclerosis, and VEGFC may be a useful indicator for early prediction of pulmonary arterial hypertension in those patients [15]. Furthermore, VEGFC aggravates intestinal inflammation in mice with experimental colitis and is associated with inflammatory lymphatic formation [16]. Similarly, VEGFCs are highly expressed in salivary duct epithelial cells in patients with primary Sjogren's syndrome, and lymphangiogenesis is active in this syndrome [17]. Based on the facts above, it is reasonable to suspect that VEGFC is involved in AITD. Therefore, this study aimed to investigate the relationship between VEGFC gene polymorphisms and AITDs.

VEGFC gene is located on chromosome 4q34.3 and has many SNPs, including rs7664413, rs11947611, rs2046463, and rs3775194. All four of them are located

in the intron region. As we discussed previously, only the rs3775194 genotype distribution of the four SNPs was associated with AITD compared with the control group. Neither the allele nor genotype of four SNPs was involved in GD subgroup. Since AITDs are sex-specific diseases, we found significant differences in the genotype distribution of rs3775194 in male HT patients compared with controls. In addition, rs3775194 was significantly associated with AITD and male AITD patients under the overdominant model. Moreover, rs11947611 was associated with male AITD under the recessive model. We did not find any significant conclusions in the four SNPs of VEGFC comparing GO in GD and hypothyroidism in HT with corresponding controls. The lack of significant positive results might be due to the differences in the influence of environmental and genetic factors on the Han population and the limited number of sampled individuals. Further large population studies warrant further study.

In genomic DNA, changes in a single base may affect the amino acid sequence and ultimately affects susceptibility to diseases [18]. The phenotype is the result of environmental, genetic and other factors. Although these loci, such as rs3775194, are localized to noncoding regions, they may influence disease occurrence by regulating gene structure or expression [19]. VEGFC gene is highly expressed in

TABLE 8: Allele and genotype distributions of VEGFC polymorphisms with hypothyroidism in HT patients.

SNPs	Without (%)	With (%)	P value	OR (95% CI)
rs7664413				
CC	114 (48.5)	84 (48.3)	0.981	
TT	20 (8.5)	14 (8.0)		
TC	101 (43.0)	76 (43.7)	0.972	0.995 (0.735-1.346)
C	329 (70.0)	244 (70.1)		
T	141 (30.0)	104 (29.9)		
rs11947611				
AA	89 (37.9)	66 (37.9)	0.974	
GG	37 (15.7)	26 (14.9)		
AG	109 (46.4)	82 (47.1)	0.901	0.982 (0.739-1.305)
A	287 (61.1)	214 (61.5)		
G	183 (38.9)	134 (38.5)		
rs2046463				
AA	113 (48.1)	84 (48.3)	0.986	
GG	20 (8.5)	14 (8.0)		
AG	102 (43.4)	76 (43.7)	0.920	0.985 (0.728-1.332)
A	328 (69.8)	244 (70.1)		
G	142 (30.2)	104 (29.9)		
rs3775194				
CC	4 (1.7)	0 (0)	0.853	
GG	198 (84.3)	148 (85.1)		
CG	33 (14.0)	26 (14.9)	0.518	1.184 (0.709-1.975)
C	41 (8.7)	26 (7.5)		
G	429 (91.3)	322 (92.5)		




FIGURE 1: VEGFC linkage disequilibrium (LD) block in the Haploview 4.2.

thyroid tissue [20], whereas in patients with AITD, there is an increase in lymphocyte infiltration in thyroid tissue, which suggests lymphatic hyperplasia in thyroid tissue.

Haplotype analysis is a more powerful way to prove that a gene is associated with a disease. We found a strong link-

age disequilibrium in 2 SNPs between patients and controls. However, subsequent study showed no association between haplotypes and AITD, HT, or GD.

In conclusion, rs3775194 locus was associated with AITD, male AITD, and male HT patients, and rs11947611 was associated with male AITD patients. VEGFC loci is related to the immune system and may be a risk factor for AITDs.

Data Availability

The data generated during this study are available within the article and any further information can be made available upon request to the corresponding author.

Ethical Approval

Written informed consent was obtained from all participants. The present study was approved by the Ethics Committees of Shanghai University of Medicine & Health Sciences Affiliated Zhoupu Hospital (Shanghai, China).

Conflicts of Interest

The authors declare that they had no conflict of interest.

Authors' Contributions

Chaoqun Gao and Jie Zhu contributed equally to this work.

Acknowledgments

The present work was funded by the National Natural Science Foundation of China (No. 81873636), Pudong New Area Health Commission key sub-specialty (PWZy2020-12), Clinical Research Center of thyroid diseases of Shanghai Health Medical College (20MC20200002), Hospital Research Foundation from Shanghai University of Medicine & Health Sciences Affiliated Zhoupu Hospital (ZPXM-2019A-02), and the Project of Shanghai Medical Key Specialty (No. ZK2019C09).

References

- [1] T. J. Smith and L. Hegedus, "Graves' disease," *The New England Journal of Medicine*, vol. 375, no. 16, pp. 1552–1565, 2016.
- [2] A. T. Gabrielson, R. A. Sartor, and W. J. G. Hellstrom, "The impact of thyroid disease on sexual dysfunction in men and women," *Sexual Medicine Reviews*, vol. 7, no. 1, pp. 57–70, 2019.
- [3] Y. Tomer, "Mechanisms of autoimmune thyroid diseases: from genetics to epigenetics," *Annual Review of Pathology*, vol. 9, no. 1, pp. 147–156, 2014.
- [4] C. Dai, S. J. Kuo, S. L. Hu et al., "VEGF-C gene polymorphisms increase susceptibility to rheumatoid arthritis," *International Journal of Medical Sciences*, vol. 16, no. 10, pp. 1397–1403, 2019.
- [5] W. Tang, T. Zhou, Z. Zhong, and H. Zhong, "Meta-analysis of associations of vascular endothelial growth factor protein levels and -634G/C polymorphism with systemic lupus erythematosus susceptibility," *BMC Medical Genetics*, vol. 20, no. 1, p. 46, 2019.
- [6] A. Villa, G. Leodori, A. Iacolare, C. Pellicano, A. Gigante, and E. Rosato, "Renal parenchymal thickness is both related to vascular endothelial growth factor and intrarenal stiffness in systemic sclerosis," *Clinical and Experimental Rheumatology*, vol. 38, no. 3, Supplement 125, p. 171, 2020.
- [7] N. Ferrara, H. P. Gerber, and J. LeCouter, "The biology of VEGF and its receptors," *Nature Medicine*, vol. 9, no. 6, pp. 669–676, 2003.
- [8] K. Rauniyar, S. K. Jha, and M. Jeltsch, "Biology of vascular endothelial growth factor C in the morphogenesis of lymphatic vessels," *Frontiers in Bioengineering and Biotechnology*, vol. 6, p. 7, 2018.
- [9] F. Menconi, C. Marcocci, and M. Marino, "Diagnosis and classification of Graves' disease," *Autoimmunity Reviews*, vol. 13, no. 4-5, pp. 398–402, 2014.
- [10] L. Zhang, Y. Ding, X. Chen et al., "In Vivo fluorescence molecular imaging of the vascular endothelial growth factor in rats with early diabetic retinopathy," *Biomedical Optics Express*, vol. 12, no. 11, pp. 7185–7198, 2021.
- [11] M. C. Hsieh, H. T. Hsu, P. C. Hsiao et al., "Role of VEGF-C gene polymorphisms in susceptibility to hepatocellular carcinoma and its pathological development," *Journal of Clinical Laboratory Analysis*, vol. 28, no. 3, pp. 237–244, 2014.
- [12] G. Kaidonis, K. P. Burdon, M. C. Gillies et al., "Common sequence variation in the VEGFC gene is associated with diabetic retinopathy and diabetic macular edema," *Ophthalmology*, vol. 122, no. 9, pp. 1828–1836, 2015.
- [13] E. Sertoglu, Ç. Yücel, A. Omma et al., "Determination of Serum Vascular Endothelial Growth Factor (VEGF) and VEGF Receptor Levels with VEGF Gene Polymorphisms in Patients with Behcet's Uveitis," *Advances in Clinical and Experimental Medicine: Official Organ Wroclaw Medical University*, vol. 31, no. 3, pp. 231–240, 2021.
- [14] X. Chen, Q. Y. Hu, M. Wang et al., "Serum VEGF-C as an evaluation marker of disease activity in adult-onset Still's disease," *Rheumatology International*, vol. 42, no. 1, pp. 149–157, 2022.
- [15] H. Didriksen, Ø. Molberg, H. Fretheim et al., "Association of lymphangiogenic factors with pulmonary arterial hypertension in systemic sclerosis," *Arthritis & Rheumatology*, vol. 73, no. 7, pp. 1277–1287, 2021.
- [16] X. L. Wang, J. Zhao, L. Qin, and M. Qiao, "Promoting inflammatory lymphangiogenesis by vascular endothelial growth factor-C (VEGF-C) aggravated intestinal inflammation in mice with experimental acute colitis," *Brazilian Journal of Medical and Biological Research*, vol. 49, no. 5, article e4738, 2016.
- [17] A. Alunno, L. Ibba-Manneschi, O. Bistoni et al., "Mobilization of lymphatic endothelial precursor cells and lymphatic neovascularization in primary Sjogren's syndrome," *Journal of Cellular and Molecular Medicine*, vol. 20, no. 4, pp. 613–622, 2016.
- [18] A. Blauvelt and A. Chiricozzi, "The immunologic role of IL-17 in psoriasis and psoriatic arthritis pathogenesis," *Clinical Reviews in Allergy and Immunology*, vol. 55, no. 3, pp. 379–390, 2018.
- [19] B. Wang, X. Jia, Q. Yao et al., "CEP128 is a crucial risk locus for autoimmune thyroid diseases," *Molecular and Cellular Endocrinology*, vol. 480, pp. 97–106, 2019.
- [20] K. Tanaka, J. Kurebayashi, H. Sonoo et al., "Expression of vascular endothelial growth factor family messenger RNA in diseased thyroid tissues," *Surgery Today*, vol. 32, no. 9, pp. 761–768, 2002.