


Genomic Sequencing of Ranaviruses Isolated from Turbot (*Scophthalmus maximus*) and Atlantic Cod (*Gadus morhua*)

Ellen Ariel,^a Natalie K. Steckler,^b  Kuttichantran Subramaniam,^b Niels J. Olesen,^c Thomas B. Waltzek^b

College of Public Health, Medical and Veterinary Sciences, James Cook University, Townsville, Queensland, Australia^a; Department of Infectious Diseases and Pathology, College of Veterinary Medicine, University of Florida, Gainesville, Florida, USA^b; Technical University of Denmark, National Veterinary Institute, Frederiksberg C, Denmark^c

Ranaviruses have been isolated from Atlantic cod (*Gadus morhua*) and turbot (*Scophthalmus maximus*) in Denmark. Phylogenomic analyses revealed that these two ranaviruses are nearly identical and form a distinct clade at the base of the ranavirus tree branching off near other fish ranaviruses.

Received 19 October 2016 Accepted 21 October 2016 Published 15 December 2016

Citation Ariel E, Steckler NK, Subramaniam K, Olesen NJ, Waltzek TB. 2016. Genomic sequencing of ranaviruses isolated from turbot (*Scophthalmus maximus*) and Atlantic cod (*Gadus morhua*). *Genome Announc* 4(6):e01393-16. doi:10.1128/genomeA.01393-16.

Copyright © 2016 Ariel et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ellen Ariel, ellen.ariel@jcu.edu.au.

Cod iridovirus (CoIV) is the earliest isolation of a ranavirus in fish. It was isolated from wild Atlantic cod in Danish coastal waters in 1979 during an investigation of ulcers in cod (1). However, results of infection trials could not definitively link the CoIV to the cod ulcerative disease (2). *Ranavirus maximus* (Rmax) was isolated from clinically healthy turbot fry from an aquaculture facility during a screening for export certification in 1999 (3).

The Rmax and CoIV isolates were propagated in bluegill fry (BF-2) cells as previously described (4) and harvested at third and fourth passages, respectively. Cell culture supernatant was clarified at 3,000 × g for 20 min, and total nucleic acids were purified using a DNeasy blood and tissue kit (Qiagen). The DNA libraries were prepared using the Nextera XT DNA kit (Illumina), and sequencing was performed using a V3 chemistry 600-cycle kit on a MiSeq platform (Illumina). *De novo* assembly of the paired-end reads in SPAdes (5) produced contiguous consensus sequences of 115,510 bp with a G+C content of 55.19% and 114,865 bp with a G+C content of 56.99% for Rmax and CoIV, respectively.

The genomes of Rmax and CoIV were annotated using GATU (6) with epizootic hematopoietic necrosis virus (EHN, GenBank accession no. NC_028461) as the reference genome (7). Additional putative open reading frames (ORFs) were identified using GenemarkS (8), and gene functions were predicted based on BLASTp searches against the NCBI GenBank nonredundant (NR) protein sequence database. A total of 100 putative ORFs were predicted in Rmax and 98 in CoIV; compared to other closely related fish ranaviruses, 100 putative ORFs were predicted in EHN and 111 in short-finned eel ranavirus (SERV, GenBank accession no. KX353311) (4). Comparative genomic analyses revealed that Rmax and CoIV are nearly identical to each other, except for a nonsense mutation leading to an early stop codon in CoIV (ORF70 in Rmax, unannotated ORF located between ORFs 66 and 67 in EHN) and the absence of a gene in CoIV (ORF62 in Rmax, ORF58L in EHN). An analysis of locally collinear blocks in Mauve (9) revealed that the genomes of Rmax and CoIV are collinear with each other and nearly collinear to EHN and SERV, with the exception of a single minor inversion involving ORFs 95

and 96 in CoIV and ORFs 97 and 98 in Rmax (ORFs 95R and 96L in EHN, ORFs 108 and 109 in SERV). Maximum likelihood phylogenetic analyses based on the concatenated nucleotide sequences of the 26 *Iridoviridae* core genes (10) revealed that Rmax and CoIV form a distinct clade at the base of the ranavirus tree branching near other fish ranaviruses (e.g., EHN, SERV).

The isolation of nearly identical ranavirus strains from two different marine species, hundreds of kilometers and 20 years apart, indicates the persistence of a ranavirus strain of unknown virulence in the marine environment. With the exception of the grouper iridoviruses in Asia (11), the impact of ranaviruses on mariculture remains unclear. Experimental challenge studies are needed to determine whether CoIV or Rmax pose a risk to marine fisheries.

Accession number(s). The complete genome sequences of Rmax and CoIV have been deposited in GenBank under the accession numbers [KX574343](https://www.ncbi.nlm.nih.gov/nuccore/KX574343) and [KX574342](https://www.ncbi.nlm.nih.gov/nuccore/KX574342), respectively.

ACKNOWLEDGMENT

We thank Patrick Thompson for his technical assistance.

FUNDING INFORMATION

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

REFERENCES

- Jensen NJ, Bloch B, Larsen JL. 1979. The ulcer-syndrome in cod (*Gadus morhua*). III. A preliminary virological report. *Nord Vet Med* 31: 436–442.
- Jensen NJ, Larsen JL. 1982. The ulcer-syndrome in cod (*Gadus morhua*). IV. Transmission experiments with two viruses isolated from cod and *Vibrio anguillarum*. *Nord Vet Med* 34:136–142.
- Ariel E, Holopainen R, Olesen NJ, Tapiovaara H. 2010. Comparative study of ranavirus isolates from cod (*Gadus morhua*) and turbot (*Psetta maxima*) with reference to other ranaviruses. *Arch Virol* 155:1261–1271. <http://dx.doi.org/10.1007/s00705-010-0715-z>.
- Subramaniam K, Toffan A, Cappelozza E, Steckler NK, Olesen NJ, Ariel E, Waltzek TB. 2016. Genomic sequence of a ranavirus isolated from short-finned eel (*Anguilla australis*). *Genome Announc* 4(4): e00843-16. <http://dx.doi.org/10.1128/genomeA.00843-16>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov

- AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
6. Tcherepanov V, Ehlers A, Upton C. 2006. Genome annotation transfer utility (GATU): rapid annotation of viral genomes using a closely related reference genome. *BMC Genomics* 7:150. <http://dx.doi.org/10.1186/1471-2164-7-150>.
 7. Jancovich JK, Brémont M, Touchman JW, Jacobs BL. 2010. Evidence for multiple recent host species shifts among the ranaviruses (family *Iridoviridae*). *J Virol* 84:2636–2647. <http://dx.doi.org/10.1128/JVI.01991-09>.
 8. Besemer J, Lomsadze A, Borodovsky M. 2001. GeneMarkS: a self-training method for prediction of gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. *Nucleic Acids Res* 29:2607–2618. <http://dx.doi.org/10.1093/nar/29.12.2607>.
 9. Darling AC, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <http://dx.doi.org/10.1101/gr.2289704>.
 10. Eaton HE, Metcalf J, Penny E, Tcherepanov V, Upton C, Brunetti CR. 2007. Comparative genomic analysis of the family *Iridoviridae*: re-annotating and defining the core set of iridovirus genes. *Virol J* 4:11. <http://dx.doi.org/10.1186/1743-422X-4-11>.
 11. Duffus ALJ, Waltzek TB, Stöhr AC, Allender MC, Gotesman M, Whittington RJ, Hick P, Hines MK, Marschang RE. 2015. Distribution and host range of Ranaviruses, p 9–57. *In* Gray MJ, Chinchar VG (ed), *Ranaviruses: lethal pathogens of ectothermic vertebrates*. Springer, New York, NY.