



ELSEVIER

Contents lists available at ScienceDirect

## Data in Brief

journal homepage: [www.elsevier.com/locate/dib](http://www.elsevier.com/locate/dib)

## Data Article

## Deformed Wing virus absence/presence data across three genera on two Hawaiian Islands



Jessika Santamaria

University of Hawaii, Manoa, United States

## ARTICLE INFO

## Article history:

Received 17 November 2017

Received in revised form

4 December 2017

Accepted 7 December 2017

Available online 15 December 2017

## Keywords:

Deformed Wing virus

pollinator

*Apis mellifera**Ceratina smaragdula**Polistes*

Hawaii

## ABSTRACT

The data presented in this article relates to the research article, “Evidence of *Varroa*-mediated Deformed Wing virus spillover in Hawaii” (Santamaria et al., 2017) [3]. The article presents data collected throughout August 2014 to November 2015, on the two Hawaiian Islands of Oahu and Maui. *Apis* and non-*Apis* specimens – a total of four species – were collected and tested for Deformed Wing virus (DWV) absence or presence, only. Specific island locations are noted. This data is made publicly available to be analyzed or used in future relevant research.

© 2018 Published by Elsevier Inc. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

## Specifications Table

Subject area	Entomology, Virology
More specific subject area	Insect virology
Type of data	Table
How data was acquired	A stained SYBR Safe DNA gel stain (Invitrogen) was visualized under ultra-violet light. The positive samples were noted with “1” and negative samples with “0”.
Data format	Raw
Experimental factors	Samples were either from Oahu or Maui

DOI of original article: <https://doi.org/10.1016/j.jip.2017.11.008>

E-mail address: [jsantama@hawaii.edu](mailto:jsantama@hawaii.edu)

<https://doi.org/10.1016/j.dib.2017.12.020>

2352-3409/© 2018 Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Experimental features	Samples were collected on the two different islands, Oahu or Maui, across different sites. Samples were preserved in $-80^{\circ}\text{C}$ until they were tested for absence or presence of the Deformed Wing virus.
Data source location	Kahului, Hawaii Haleakala, Hawaii Iao Needle, Hawaii Makawao, Hawaii Ho'omaluhia, Hawaii Sandy Beach, Hawaii Pearl City, Hawaii Manoa, Hawaii Kaena Point, Hawaii Waimanalo, Hawaii
Data accessibility	Data is within this article

---

### Value of the data

- Data is useful as benchmark for viral presence in a pre-*Varroa* ecosystem
  - Data implicates how important it is to keep *Varroa* out of the *Varroa*-free islands of Hawaii
  - Potential future studies can explore pollinator-flower webs and how viruses could spread in these systems
- 

## 1. Data

The data in this article lists which species were collected, from which island, and island location they were collected from. The data shows whether each sample tested positive for Deformed Wing virus (DWV) or not; this is represented by “0” and “1” to indicate absence or presence, respectively. The raw data can be seen in [Table 1](#).

## 2. Experimental design, materials and methods

### 2.1. Specimen collection

Three species within two different Hymenoptera genera were selected as representatives of the local community of flower visitors: the introduced small carpenter bee *Ceratina smaragdula* (Apidae) which was first recorded in Hawaii in 1999 [2], and the introduced paper wasps *Polistes aurifer* and *Polistes exclamans* (Vespidae) first recorded in Hawaii in the 19th century and in 1952 respectively [1]. *Polistes* wasps collected on Oahu are *P. aurifer* and the specimens from Maui are *P. exclamans*. All samples were collected from either of the five sites on Oahu (*Varroa*-positive island), or the four sites on Maui, (*Varroa*-negative island). Collection sites on both islands consisted on a mix of agricultural fields, parks, gardens, and beach edge vegetation strips. The selected insect species are all relatively abundant and can be found in urban and agricultural environments, where they overlap in resource use with *A. mellifera*.

Samples were collected from August 2014 to November 2015. Insects were collected while they were foraging in fields or flower patches, via a hand-held net. Paper wasp samples were also collected from around their nests. Each insect was stored individually and kept on ice in the field until transferred to a  $-80^{\circ}\text{C}$  freezer for long term storage.

### 2.2. RNA extraction and reverse transcription PCR

Methods for RNA extraction and RT-PCR are fully described in Santamaria et al. [3].







**Table 1** (continued)

DWV	Species	Island	Location
0	<i>Polistes aurifer</i>	Oahu	Waimanalo
0	<i>Polistes aurifer</i>	Oahu	Waimanalo
0	<i>Polistes aurifer</i>	Oahu	Waimanalo
0	<i>Polistes aurifer</i>	Oahu	Waimanalo
0	<i>Polistes aurifer</i>	Oahu	Waimanalo
0	<i>Polistes aurifer</i>	Oahu	Waimanalo
0	<i>Polistes aurifer</i>	Oahu	Waimanalo
0	<i>Polistes aurifer</i>	Oahu	Waimanalo
0	<i>Polistes aurifer</i>	Oahu	Waimanalo
0	<i>Polistes aurifer</i>	Oahu	Waimanalo
1	<i>Polistes aurifer</i>	Oahu	Waimanalo
1	<i>Polistes aurifer</i>	Oahu	Waimanalo
1	<i>Polistes aurifer</i>	Oahu	Waimanalo
1	<i>Polistes aurifer</i>	Oahu	Waimanalo
1	<i>Polistes aurifer</i>	Oahu	Waimanalo
1	<i>Polistes aurifer</i>	Oahu	Waimanalo
1	<i>Polistes aurifer</i>	Oahu	Waimanalo
1	<i>Polistes aurifer</i>	Oahu	Waimanalo
1	<i>Polistes aurifer</i>	Oahu	Waimanalo
1	<i>Polistes aurifer</i>	Oahu	Waimanalo

## Acknowledgements

A huge thanks to all the beekeepers and farmers that allowed us access to their lands for the collection of this data. Particular thanks to Monstre Farm for their warmest hospitality.

## Transparency document. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2017.12.020>.

## References

- [1] J.R. Beggs, E.G. Brockerhoff, J.C. Corley, M. Kenis, M. Masciocchi, F. Muller, Q. Rome, C. Villemant, Ecological effects and management of invasive alien Vespidae, *BioControl* 56 (4) (2011) 505–526. <http://dx.doi.org/10.1007/s10526-011-9389-z>.
- [2] K.N. Magnacca, C.B.K. King, Assessing the Presence and Distribution of 23 Hawaiian Yellow-faced Bee Species on Lands Adjacent to Military Installations on O'ahu and Hawai'i Island. Technical Report No. 185, Pacific Cooperative Studies Unit, University of Hawai'i, Honolulu, Hawai'i, 2013, <http://scholarspace.manoa.hawaii.edu/bitstream/handle/10125/34064/v185.pdf>.
- [3] J. Santamaria, E.M. Villalobos, L. Brettell, S. Nikaido, J. Graham, S.J. Martin, Evidence of Varroa-mediated deformed Wing virus spillover in Hawaii, *J. Invertebr. Pathol.* (2017). <http://dx.doi.org/10.1016/j.jip.2017.11.008> (In Press).