

# Concurrent lung adenocarcinoma and bladder diffuse large B-cell lymphoma: a case report and literature review

Journal of International Medical Research

50(2) 1–8

© The Author(s) 2022

Article reuse guidelines:

[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)

DOI: 10.1177/03000605221081672

[journals.sagepub.com/home/imr](https://journals.sagepub.com/home/imr)

Feng Xu, Ying Liang, Wen-Bin Mo,  
Xiao-Jing Yan and Rui Zhang 

## Abstract

Lung adenocarcinoma is one of the most common solid tumors, and diffuse large B-cell lymphoma (DLBCL) is the most common histological subtype of adult non-Hodgkin's lymphoma. Although extra-nodular lesions are frequently observed in patients with DLBCL, urinary bladder involvement is rare. We report the case of a 77-year-old woman with lung adenocarcinoma who was diagnosed with a second primary bladder DLBCL, 9 months after treatment with molecular targeted drugs. Simultaneous therapies for her lymphoma with lenalidomide and rituximab and a tyrosine kinase inhibitor therapy for her lung cancer were both effective. This result was consistent with previous reports suggesting that patients unable to tolerate intensive chemotherapy could benefit from targeted therapies. Current research into the use of lenalidomide for the treatment of lymphomas and solid tumors is promising in terms of exploring immunotherapy as an alternative option for patients with concurrent solid tumors and lymphomas who have poor tolerance to radiotherapy and chemotherapy.

## Keywords

Diffuse large B-cell lymphoma, lung adenocarcinoma, rituximab, lenalidomide, concurrent lung cancer and lymphoma, treatment for concurrent lung cancer and lymphoma

Date received: 10 October 2021; accepted: 2 February 2022

## Introduction

Lung adenocarcinoma is the most common subtype of non-small cell lung cancer

The First Hospital of China Medical University, Shenyang 110001, Liaoning Province, China

### Corresponding author:

Rui Zhang, The First Affiliated Hospital of China Medical University 155 Nanjing North Street, Heping District, Shenyang, Liaoning 110001, China.

Email: [zhangruicmu1h@163.com](mailto:zhangruicmu1h@163.com)



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

(NSCLC), accounting for 83% of all lung cancers. It is considered as an advanced malignancy with a poor prognosis.<sup>1</sup> However, the recent identification of molecular targets has significantly improved the outcomes of patients with lung adenocarcinoma.<sup>2,3</sup> Mutations in the epidermal growth factor receptor gene (*EGFR*) are key drivers in NSCLC and are observed in up to 50% of lung adenocarcinomas.<sup>2-5</sup> Accordingly, *EGFR* tyrosine kinase inhibitors (TKIs) have been used in the treatment of NSCLC with *EGFR* mutations since 2000,<sup>6</sup> and have greatly improved the treatment response and long-term survival.<sup>7-9</sup>

Diffuse large B-cell lymphoma (DLBCL) is the most frequent histological subtype of adult non-Hodgkin's lymphoma (NHL),<sup>10,11</sup> and usually occurs as a primary B cell lymphoma or transforms from indolent lymphoma in rare cases. Although DLBCL can be cured by immunochemotherapy, its treatment remains challenging in older patients who more frequently present with a high international prognostic index (IPI) and are unable to tolerate standard chemotherapy. In addition to age and IPI, various other factors predict a poor response to immunochemotherapy, such as activated B cell (ABC) subtype,<sup>12</sup> double expression/double hit lymphoma,<sup>13-15</sup> and high-risk gene mutations, such as *TP53*.<sup>16</sup>

We present the case of an elderly female patient who developed DLBCL 9 months after a diagnosis of lung adenocarcinoma and during *EGFR*-TKI treatment. In addition to the TKIs for her lung adenocarcinoma, she received rituximab plus lenalidomide for the lymphoma.


## Case Report

This study was approved by the Ethics Committee of the First Hospital of China Medical University (Approval no.: [2020] 2020-277-2). Written informed consent

was signed by the patient. The study conformed to the CARE guidelines.<sup>17</sup>

A 77-year-old female patient was admitted to our hospital in April 2019 because of fatigue. A lung computed tomography scan revealed a mass lesion with a maximum diameter of 24 mm in the right upper lobe of the hilum (Figure 1a). Lung puncture biopsy was performed and histopathological analysis showed irregular distribution of heteromorphic cells with large nuclei and an imbalanced nuclear/cytoplasm ratio. Immunohistochemical assay results showed thyroid transcription factor-1 (+), carcinoembryonic antigen (+), p63 (scattered +), napsin-A (weak +), Ki-67 (30%), cytokeratin (CK)7 (+), and CK5/6 ( $\pm$ ) (Figure 1b, 1c). A diagnosis of lung adenocarcinoma was confirmed based on the above results. Furthermore, deletion of *EGFR* exon 19 was demonstrated by quantitative real-time polymerase chain reaction with a panel of NSCLC-related driver mutations (including *ALK*, *ROS1*, *RET*, *MET*, *KRAS*, *NRAS*, *HER2*, and *PIK3CA*). She was accordingly treated with the first-generation *EGFR*-TKI icotinib orally.

Nine months after initiation of icotinib treatment, the patient developed painless gross hematuria. Urinary bladder ultrasound revealed an 8.0  $\times$  3.4 cm lesion. Positron emission tomography-computed tomography (PETCT) showed an irregular mass with soft tissue density in the left posterior wall of the urinary bladder, involving the left ureter, with a maximum standardized uptake value (SUV) of 30 (Figure 1d, 1e). Increased metabolic activity was also observed in multiple soft tissue nodules in the lower lobe of the left lung, and hilar and mediastinal lymph nodes, with a maximum SUV of 10.7 (Figure 1f). There was no significant change in the size of the primary lesion in the right lung. Cystoscopy of the bladder lesion was performed. Histopathological analysis demonstrated diffuse distribution of atypical large



**Figure 1.** Lung adenocarcinoma and bladder diffuse large B-cell lymphoma (DLBCL) during treatment. (a) Lung computed tomography (CT) scan showed parahilar nodules in the upper lobe of the right lung with a maximum diameter of 24 mm before starting treatment. (b) Histopathological analysis of lung biopsy showed infiltration of heteromorphic cells (hematoxylin and eosin (H&E) staining,  $\times 200$ ). The nucleus was large and deeply stained and the nucleus/cytoplasm ratio was abnormal. (c) Histopathological analysis of lung biopsy showed positive expression of thyroid transcription factor-1. (d, e) Positron emission tomography (PET)-CT showed irregular soft tissue density mass in the left posterior wall of the bladder with left ureter involvement and a maximum standardized uptake value (SUV) of 30. (f) PET-CT showed a new nodule shadow in the left lower lung lobe with a maximum SUV of 10.7 after 9 months of treatment for lung adenocarcinoma. (g) Histopathological analysis of bladder biopsy showed diffuse distribution of atypical large lymphocytes (H&E staining,  $\times 200$ ) and positive expression of CD20 (h). (i, j) CT scan of the urinary system showed disappearance of the occupying lesion in the bladder after treatment. However, heterogeneous enhancement of the bladder wall could still be seen. (k, l) Lung CT showed that both lesions in the left lung and right lung were larger than before treatment with icotinib and R2.

lymphocytes with the following immunohistochemical features: CD20 (+), Bcl-2 (30% +), Bcl-6 (+), c-Myc (10% +), CD68 (–), multiple myeloma oncogene-1 (+), CD10 (–), Pax-5 (+), CD5 (–), CD3 (–), Ki-67 (80%), and Epstein–Barr encoding region *in situ* hybridization (–). DLBCL with ABC subtype was confirmed (Figure 1g, 1h). The prognosis was poor. Next-generation sequencing with a panel containing 82 DLBCL hotspot genes was carried out on the lymphoma tissue at Yuanqi Biomedical Technology Co. Ltd. (Shanghai, China). The results showed

mutations of *TET2*, *TNFAIP3*, *SGK1*, *IGLL5*, and *CREBBP*. However, no mutations in the *EGFR* gene, including exon 18-G719X, exon 19-del, exon 20-T90M, or exon 21L858R were identified in the lymphoma tissue by Droplet Digital PCR (Yuanqi Biomedical Technology Co., Ltd., Shanghai, China).

The patient declined radiotherapy and chemotherapy for their bladder lesion. Rituximab plus lenalidomide (R2 regimen) was administered, with rituximab 600 mg on day 1 and lenalidomide 25 mg on days 1 to 14 every 21 days. This regimen was

used in combination with continued icotinib for treatment of the lung adenocarcinoma. After the first course of treatment, the patient's gross hematuria completely resolved. Consent for the use of TKIs and R2 was acquired from the patient before treatment. After six courses of treatment, CT scan of the urinary system showed that the lymphoma had achieved complete remission (Figure 1i, 1j). The treatment was well tolerated by the patient. However, a lung CT scan showed that the lung cancer had progressed (Figure 1k, 1l). In light of the potential development of resistance to icotinib, the small molecule multi-target TKI anlotinib was added to the icotinib treatment. The size of the lung cancer lesion reduced significantly over the following 6 months of combination therapy. To date, the patient had been followed for 24 months with clinical evidence of continued remission of the urinary bladder lymphoma. However, her response to the TKIs was poor and her lung cancer continued to progress to a more advanced stage.

## Discussion

The occurrence of lymphomas as second tumors in patients with lung cancer is not rare. Although the exact cause of dual tumors is unknown, chemotherapy is considered to be a common cause of secondary lymphomas. However, evidence of secondary lymphomas after TKIs in the era of targeted therapy is uncommon.<sup>18–20</sup> Here we report a patient with a history of lung adenocarcinoma treated with TKIs who developed high-risk DLBCL with bladder involvement. Although DLBCL is curable with classical R-CHOP regimen,<sup>21</sup> the outcome in high-risk patients remains poor, especially in older, frail patients. The current patient was ineligible for standard therapy because of her age and poor disease condition. Although a driver *EGFR* mutation was detected as a potential target for

the treatment of her lung adenocarcinoma, the secondary lymphoma lacked indications for targeted drugs and was associated with several adverse factors for long-term survival, such as high IPI, extra-nodular disease, ABC subtype, dual expression of C-Myc and Bcl-2,<sup>21–24</sup> and *TET2*,<sup>25</sup> *TNFAIP3*,<sup>26</sup> and *SGKI* mutations.<sup>27</sup>

The incidence of concurrent lung cancer and lymphoma has increased in recent years. Both lung cancer and NHL are associated with higher risks of secondary malignancies of 36.7%<sup>28</sup> and 10.95%, respectively.<sup>29</sup> A search of the National Cancer Institute's SEER database ([www.seer.cancer.gov](http://www.seer.cancer.gov)) identified 14,665 patients with concurrent malignancies from 1975 to 2016, including 2607 patients who developed lymphoma secondary to their lung cancer. Of these, 44.5% had lung adenocarcinoma, and chronic lymphocytic leukemia or small cell lymphoma was the most common subtype of lymphoma (25.5%) secondary to the lung carcinoma. As the second most common subtype, DLBCL accounted for 20.2% of concurrent lymphomas in patients with lung cancer. There were significant differences in survival times among patients with different types of lung cancer who developed secondary DLBCL: the median survival of patients with lung adenocarcinoma and secondary DLBCL was 11 months, which was slightly poorer than that in patients with small cell lung cancer (Table 1). However, the relevance of the biological origins of the two types of tumors remains unclear. Evidence from the current case suggested that the concurrent malignancies originated separately because they had different mutation spectra. No *EGFR* mutations were found in the DLBCL specimens and there have been few reports of lymphoma secondary to *EGFR* TKIs,<sup>18,30</sup> indicating that the two tumors were likely to have different origins. It was therefore necessary to choose a

**Table 1.** Effects of various factors on survival time in patients with lymphoma secondary to lung cancer (data from the SEER database (www.seer.cancer.gov)).

	Number of people	Median survival (months)	Mean rank	Sum of rank	P
<b>Stage</b>					
I & II	307	16.5	299.34	91,897.00	<0.001*
III & IV	247	8	250.36	61,838.00	
Unknown	95	4.5	–	–	–
<b>Age, years</b>					
<60	57	12	336.88	19,202.00	0.616
≥60	592	9	323.86	191,723.00	
<b>Chemotherapy</b>					
Yes	400	15.5	368.46	147,384.00	<0.001*
No/unknown	249	4	255.18	63,541.00	
<b>Sex</b>					
Female	278	16	350.58	97,460.50	0.003*
Male	371	7.5	305.83	113,464.50	
Adenocarcinoma	309	11	236.49	$\chi^2 = 91.985$	<0.001*
Small-cell lung cancer	47	18	247.77		
Squamous cell carcinoma	200	7	374.89		

\*Significant difference between the two groups (P < 0.05).

**Table 2.** Influences of different indexes on the survival time of lymphoma secondary to lung cancer (data from the SEER database (www.seer.cancer.gov)).

	Number of people	Median survival (months)	Mean rank	Sum of rank	P
<b>Stage</b>					
I & II	136	18	140.18	19,064.50	0.015*
III & IV	122	10	117.59	14,346.50	
Unknown	51	6			–
<b>Age, years</b>					
<60	24	15.5	152.94	3670.50	0.906
≥60	285	11	155.17	44,224.50	
<b>Chemotherapy</b>					
Yes	195	16	172.54	33,646.00	<0.001*
No/unknown	114	5	124.99	14,249.00	
<b>Sex</b>					
Female	153	16	162.43	24,852.00	0.147
Male	156	11	147.71	23,043.00	

\*Significant difference between the two groups (P<0.05).

treatment regimen to take account of both types of tumor.

Chemotherapy has previously been the mostly widely used therapy for the treatment of lung cancer and lymphoma.

However, although results from the SEER database showed that chemotherapy partly prolonged survival of these patients (Table 2), the overall outcome was poor. The database reported 309 patients with



DLBCL secondary to adenocarcinoma from 1975 to 2016, of whom 46 patients had received chemotherapy for lung cancer before the occurrence of their secondary DLBCL. The median survival was 6 months in these patients. Except for a small number of patients who died due to complications and tumor progression leading to respiratory and circulatory deterioration,<sup>31,32</sup> most patients in the reported studies were not followed-up long-term.

In the current case, intensive chemotherapy was not considered appropriate because of the patient's age, prior history of lung cancer, and comorbid lung cancer and DLBCL, and a chemo-free regimen of R2, consisting of rituximab and lenalidomide, was considered in light of promising results in patients with B-cell lymphomas.<sup>33–36</sup> The therapeutic effects of lenalidomide are especially notable, showing that it enhance the proliferation and function of T cells, repair synapses of effector T cells, increase antibody-dependent cytotoxicity mediated by natural killer cells, up-regulate costimulatory molecules on the surface of tumor cells, and inhibit angiogenesis,<sup>37–41</sup> all of which could be relevant to the treatment of solid tumors<sup>42</sup> including NSCLC.<sup>32</sup> R2 accordingly produced a rapid and durable response in the present patient with an early-stage urinary bladder cancer.

In summary, the current challenging case involved an elderly patient who developed urinary bladder DLBLC during targeted treatment of her lung cancer. Previous analysis has demonstrated poor outcomes of traditional chemotherapy, especially for elderly patients who cannot tolerate chemotherapy. The patient's lymphoma was well-controlled by the R2 regimen and she tolerated the treatment well. However, further follow-up is needed to determine the long-term therapeutic effect and survival. Chemo-free therapy with novel agents may thus provide an alternative treatment choice in patients in

poor condition with concurrent lymphoma and lung cancer.


### Declaration of conflicting interest

The authors declare that there are no conflicts of interest.

### Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

### ORCID iD

Rui Zhang  <https://orcid.org/0000-0003-4731-9062>

### References

1. Miller KD, Nogueira L, Mariotto AB, et al. Cancer treatment and survivorship statistics, 2019. *CA Cancer J Clin* 2019; 69: 363–385.
2. Pao W, Miller V, Zakowski M, et al. EGF receptor gene mutations are common in lung cancers from “never smokers” and are associated with sensitivity of tumors to gefitinib and erlotinib. *Proc Natl Acad Sci USA* 2004; 101: 13306–13311.
3. Rosell R, Moran T, Queralt C, et al. Screening for epidermal growth factor receptor mutations in lung cancer. *N Engl J Med* 2009; 361: 958–967.
4. Shi Y, Au JS, Thongprasert S, et al. A prospective, molecular epidemiology study of EGFR mutations in Asian patients with advanced non-small-cell lung cancer of adenocarcinoma histology (PIONEER). *J Thorac Oncol* 2014; 9: 154–162.
5. Frol'Kis IV. [Cellular mechanisms of the activating effect of vasopressin on smooth vascular muscles]. *Dokl Akad Nauk SSSR* 1987; 294: 1004–1007.
6. Lynch TJ, Bell DW, Sordella R, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *New Engl J Med* 2004; 350: 2129–2139.
7. Pinsolle J, McLeer-Florin A, Giaj Levra M, et al. Translating systems medicine into clinical practice: examples from pulmonary

- medicine with genetic disorders, infections, inflammations, cancer genesis, and treatment implication of molecular alterations in non-small-cell lung cancers and personalized medicine. *Front Med* 2019; 6: 233.
8. Mok TS, Yi LW, Thongprasert S, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med* 2009; 361: 947.
  9. Zhou C, Wu YL, Chen G, et al. Erlotinib versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-positive non-small-cell lung cancer (OPTIMAL, CTONG-0802): a multicentre, open-label, randomised, phase 3 study. *Lancet Oncol* 2011; 12: 710–711.
  10. Morton LM, Wang SS, Devesa SS, et al. Lymphoma incidence patterns by WHO subtype in the United States, 1992–2001. *Blood* 2006; 107: 265.
  11. Jiang M, Bannani NN and Feldman AL. Lymphoma classification update: B-cell non-Hodgkin lymphomas. *Expert Rev Hematol* 2017; 10: 405–415.
  12. Alizadeh AA, Eisen MB, Davis RE, et al. Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. *Nature* 2000; 403: 503–511.
  13. Savage KJ, Johnson NA, Ben-Neriah S, et al. MYC gene rearrangements are associated with a poor prognosis in diffuse large B-cell lymphoma patients treated with R-CHOP chemotherapy. *Blood* 2009; 114: 3533–3537.
  14. Lin P, Dickason TJ, Fayad LE, et al. Prognostic value of MYC rearrangement in cases of B-cell lymphoma, unclassifiable, with features intermediate between diffuse large B-cell lymphoma and Burkitt lymphoma. *Cancer* 2012; 118: 1566–1573.
  15. Barrans S, Crouch S, Smith A, et al. Rearrangement of MYC is associated with poor prognosis in patients with diffuse large B-cell lymphoma treated in the era of rituximab. *J Clin Oncol* 2010; 28: 3360–3365.
  16. Peroja P, Pedersen M, Mantere T, et al. Mutation of TP53, translocation analysis and immunohistochemical expression of MYC, BCL-2 and BCL-6 in patients with DLBCL treated with R-CHOP. *Sci Rep* 2018; 8: 14814.
  17. Gagnier JJ, Kienle G, Altman DG, et al; CARE Group. The CARE guidelines: consensus-based clinical case reporting guideline development. *Headache* 2013; 53: 1541–1547.
  18. Cho BC, Chewaskulyong B, Lee KH, et al. Osimertinib versus standard of care EGFR TKI as first-line treatment in patients with EGFRm advanced NSCLC: FLAURA Asian Subset. *J Thorac Oncol* 2019; 14: 99–106.
  19. Wang L and Wang W. Safety and efficacy of anaplastic lymphoma kinase tyrosine kinase inhibitors in non-small cell lung cancer (Review). *Oncol Rep* 2021; 45: 13–28.
  20. Bauer TM, Felip E, Solomon BJ, et al. Clinical management of adverse events associated with lorlatinib. *Oncologist* 2019; 24: 1103–1110.
  21. Hans CP, Weisenburger DD, Greiner TC, et al. Confirmation of the molecular classification of diffuse large B-cell lymphoma by immunohistochemistry using a tissue microarray. *Blood* 2004; 103: 275–282.
  22. Johnson NA, Slack GW, Savage KJ, et al. Concurrent expression of MYC and BCL2 in diffuse large B-cell lymphoma treated with rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone. *J Clin Oncol* 2012; 30: 3452–3459.
  23. Green TM, Young KH, Visco C, et al. Immunohistochemical double-hit score is a strong predictor of outcome in patients with diffuse large B-cell lymphoma treated with rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone. *J Clin Oncol* 2012; 30: 3460–3467.
  24. Savage KJ, Slack GW, Mottok A, et al. The impact of dual expression of MYC and BCL2 by immunohistochemistry on the risk of CNS relapse in DLBCL. *Blood* 2016; 127: 2182.
  25. Kubuki Y, Yamaji T, Hidaka T, et al. TET2 mutation in diffuse large B-cell lymphoma. *J Clin Exp Hematol* 2017; 56: 145–149.
  26. Compagno M, Lim WK, Grunn A, et al. Mutations of multiple genes cause deregulation of NF-kappaB in diffuse large B-cell lymphoma. *Nature* 2009; 459: 717–721.
  27. Lu L, Zhu F, Li Y, et al. Inhibition of the STAT3 target SGK1 sensitizes diffuse large

- B cell lymphoma cells to AKT inhibitors. *Blood Cancer J* 2019; 9: 43.
28. Son C, Lee SK, Choi PJ, et al. Characteristics of additional primary malignancies in Korean patients with non-small cell lung cancer. *J Thorac Dis* 2013; 5: 737–744.
  29. Jiang S, Zhen H and Jiang H. Second primary malignancy in diffuse large B-cell lymphoma patients: A SEER database analysis. *Curr Probl Cancer* 2020; 44: 100502.
  30. Zhang W, Zhang Y, Zhao Q, et al. Long-term safety of icotinib in patients with non-small cell lung cancer: a retrospective, real-world study. *J Thorac Dis* 2020; 12: 639–650.
  31. Zheng JX, Li XZ, Xiang RL, et al. Synchronous primary pulmonary lymphoma presenting with pulmonary adenocarcinoma: A case report and literature review. *Indian J Cancer* 2015; 52: 37.
  32. Kim K, An S, Cha HJ, et al. Lenalidomide induces apoptosis and alters gene expression in non-small cell lung cancer cells. *Oncol Lett* 2013; 5: 588–592.
  33. Fowler NH, Davis RE, Rawal S, et al. Safety and activity of lenalidomide and rituximab in untreated indolent lymphoma: an open-label, phase 2 trial. *Lancet Oncol* 2014; 15: 1311–1318.
  34. Martin P, Jung SH, Pitcher B, et al. A phase II trial of lenalidomide plus rituximab in previously untreated follicular non-Hodgkin's lymphoma (NHL): CALGB 50803 (Alliance). *Ann Oncol* 2017; 28: 2806–2812.
  35. Becnel MR, Nastoupil LJ, Samaniego F, et al. Lenalidomide plus rituximab (R<sup>2</sup>) in previously untreated marginal zone lymphoma: subgroup analysis and long-term follow-up of an open-label phase 2 trial. *Br J Haematol* 2019; 185: 874–882.
  36. Fisher RI. Optimizing first-line therapy for follicular lymphoma: is it time for chemotherapy-free approaches? *J Natl Compr Canc Netw* 2019; 17: 1414–1416.
  37. Bartlett JB, Dredge K and Dalgleish AG. The evolution of thalidomide and its IMiD derivatives as anticancer agents. *Nat Rev Cancer* 2004; 4: 314–322.
  38. Gribben JG, Fowler N and Morschhauser F. Mechanisms of action of lenalidomide in B-cell non-Hodgkin lymphoma. *J Clin Oncol* 2015; 33: 2803–2811.
  39. Chang HD, Liu N, Klimek V, et al. Enhancement of ligand-dependent activation of human natural killer T cells by lenalidomide: therapeutic implications. *Blood* 108: 618–621.
  40. Wu L, Adams M, Carter T, et al. Lenalidomide enhances natural killer cell and monocyte-mediated antibody-dependent cellular cytotoxicity of rituximab-treated CD20+ tumor cells. *Clin Cancer Res* 14: 4650–4657.
  41. Mcdaniel JM, Pinilla-Ibarz J and Epling-Burnette PK. Molecular action of lenalidomide in lymphocytes and hematologic malignancies. *Adv Hematol* 2012; 2012: 513702.
  42. Segler A and Tsimberidou AM. Lenalidomide in solid tumors. *Cancer Chemother Pharmacol* 2012; 69: 1393–1406.