

ARTICLE

Estimation of brain receptor occupancy for trazodone immediate release and once a day formulations

Laura Oggianu¹ | Giorgio Di Dato¹ | Giorgina Mangano¹ | Maria Teresa Rosignoli¹ | Savannah McFeely² | Alice Ban Ke² | Hannah M. Jones² | Alessandro Comandini¹

¹Angelini Pharma S.p.A., Rome, Italy

²Certara UK, Simcyp Division, Sheffield, UK

Correspondence

Laura Oggianu, Angelini Pharma S.p.A., Viale Amelia, 70 – 00181, Rome, Italy.
Email: laura.oggianu@angelinipharma.com

Funding information

This work was supported by Angelini Pharma S.p.A.

Abstract

Trazodone is approved for the treatment of major depressive disorders, marketed as immediate release (IR), prolonged release, and once a day (OAD) formulation. The different formulations allow different administration schedules and may be useful to facilitate patients' compliance to the antidepressant treatment. A previously verified physiologically-based pharmacokinetic model based on in vitro and in vivo information on trazodone pharmacokinetics was applied, aiming at predicting brain receptor occupancy (RO) after single and repeated dosing of the IR formulation and repeated dosing of the OAD formulation in healthy subjects. Receptors included in the simulations were selected using static calculations of RO based on the maximum unbound brain concentration ($C_{\max, \text{brain}, u}$) of trazodone for each formulation and dosing scheme, resulting in 16 receptors being simulated. Seven receptors were simulated for the IR low dose formulation (30 mg), with similar t_{onset} and duration of coverage (range: 0.09–0.25 h and 2.1–>24 h, respectively) as well as RO (range: 0.64–0.92) predicted between day 1 and day 7 of dosing. The 16 receptors evaluated for the OAD formulation (300 mg) showed high RO (range: 0.97–0.84 for the receptors also covered by the IR formulation and 0.73–0.48 for the remaining) correlating with affinity and similar duration of time above the target threshold to the IR formulation (range: 2–>24 h). The dose-dependent receptor coverage supports the multimodal activity of trazodone, which may further contribute to its fast antidepressant action and effectiveness in controlling different symptoms in depressed patients.

Study Highlights

WHAT IS THE CURRENT KNOWLEDGE ON THE TOPIC?

The antidepressant efficacy of trazodone has been shown to be significantly correlated to its steady-state plasma levels, and previous work has shown some understanding of trazodone range of affinity for different receptors, at different doses, but without considering the different available formulations. Trazodone is commonly available as: immediate release (IR), prolonged release (PR), and once

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 Angelini Pharma S.p.A. *Clinical and Translational Science* published by Wiley Periodicals LLC on behalf of American Society for Clinical Pharmacology and Therapeutics.

a day (OAD) tablets. The IR formulation has a rapid onset and short duration of action, whereas the PR formulation is characterized by an absorption boost as soon as it is administered and has a comparatively delayed maximum concentration (C_{max}). Conversely, the OAD formulation provides a controlled release of trazodone over 24 h without the early high peak plasma concentration seen with the IR and PR formulations.

WHAT QUESTION DID THIS STUDY ADDRESS?

This work aims to identify the brain receptors reaching a threshold occupancy of 50% through static predictions and determine the occupancy versus time profile for those of interest following administration of short- and long-acting trazodone formulations.

WHAT DOES THIS STUDY ADD TO OUR KNOWLEDGE?

Brain receptor occupancy (RO) for key targets were predicted based on free drug concentrations, allowing for a physiologically relevant assessment of the different pathways affected by each formulation and dose.

HOW MIGHT THIS CHANGE CLINICAL PHARMACOLOGY OR TRANSLATIONAL SCIENCE?

The presented physiologically-based pharmacokinetic approach to assess RO can be used to guide formulation selection and dosing in clinical studies.

INTRODUCTION

Trazodone hydrochloride is a triazolopyridine derivative, defined as the first member of the class of serotonin antagonist/reuptake inhibitor (SARI) developed for the treatment of depression. Trazodone is currently approved for the treatment of major depressive disorder (MDD), with or without anxiety.¹

Trazodone acts as a potent antagonist of the serotonin (5-HT) receptors 5-HT_{2A} and shows moderate affinity to 5-HT_{1A}, 5-HT_{2C}, and 5-HT₇ receptors, acting as a weak agonist for the first one and as a weak antagonist for the following two receptors.² Trazodone also shows moderate affinity for the serotonin transporter SERT. Moreover, it was demonstrated that trazodone binds with high affinity to adrenergic receptors, where it acts blocking α_1 -adrenoceptors and moderately antagonizing α_2 -adrenoceptors.^{3,4} On the other hand, trazodone has very low affinity for acetylcholine muscarinic, dopaminergic, or GABA/benzodiazepine receptors, whereas there is no full consensus about trazodone affinity for the H₁-histaminic receptors.⁵

Trazodone can be defined as a multifunctional drug due to its dose-dependent pharmacological activity. Different clinical trials suggest that low doses of trazodone (i.e., 30 mg–50 mg per day) may be useful for controlling insomnia, probably due to the antagonism of 5-HT_{2A/2C} and α_1 that provides a hypnotic effect.⁶ When used at the proper antidepressant doses (e.g., starting from 100 to 150 mg per day, until 300 mg per day), trazodone is

able to exert additional pharmacological actions, such as the SERT blockade. This layered response allows for full antidepressant efficacy, with a complex mixture of pharmacological functions due to the simultaneous inhibition of the serotonin (5-HT) transporters (SERT), and 5-HT_{2A}, 5-HT_{2C}, and 5-HT₇ receptors, together with the partial agonism of 5-HT_{1A} receptors. The combination of these pathways allows for the antidepressant action of trazodone.¹ Considering the trazodone pharmacokinetic profile, and in particular the variability of the plasma concentration, to achieve therapeutically effective levels to manage major depressive disorder episodes, trazodone should be administered at the target dose of 300 mg/day.¹

Two formulations for trazodone are currently available—immediate-release (IR) tablets requiring multiple administrations daily, and, in Europe, prolonged-release tablets for twice-daily administration. Trazodone once a day (OAD) is a prolonged release formulation of trazodone for once daily administration. The IR formulation has a rapid onset and short duration of action, whereas the prolonged release (PR) formulation is characterized by an absorption boost as soon as it is administered and has a comparatively delayed maximum concentration (C_{max}). Conversely, the OAD formulation provides a controlled release of trazodone over 24 h without the early high peak plasma concentration seen with the IR and PR formulations.^{7,8}

To better support and clarify the multimodal mechanism of action of trazodone and its relevance in managing MDD symptoms, it is important to understand the

occupancy at the receptors indicated in MDD, the time at which a significant RO (t_{onset}) is reached, as well as the duration where the target RO is maintained. Therefore, the aim of this project was to simulate unbound brain concentrations in adults using an existing physiologically-based pharmacokinetic (PBPK) model for trazodone, refined with a series of human in vitro parameters, and apply a dynamic pharmacodynamic (PD) model to determine RO over time. The 30 mg IR (at single dose) and the 300 mg OAD (at steady-state) formulations, standing at the extremes of the available dosing range, will be assessed with the aim of simulating the widest possible range of exposures achievable by trazodone.

METHODS

PBPK modeling strategy and receptor occupancy determination

Development of trazodone PBPK models

The Simcyp Population-Based Simulator (version 18, release 2; Simcyp Ltd., Sheffield, UK) and Caucasian Healthy Volunteer population was used for all simulations. A PBPK model for IR trazodone has previously been developed.⁹ Plasma CL_{iv} was used directly in the model as input. A K_p scalar was incorporated to predict the observed volume of distribution at steady state (V_{ss}) value accurately.

For OAD trazodone, this model was updated to include the mechanistic absorption model, Advanced Dissolution Absorption and Metabolism (ADAM) model. The ADAM model within Simcyp has been described previously.¹⁰ In brief, the dissolution profile of OAD trazodone was estimated using the Weibull cumulative dissolution function parameters (F_{max} , α , and β , respectively, standing for maximum %Dissolved in vivo, dissolution profile scale factor, and dissolution profile shape factor) for each subject in the clinical trial (Equation 1). The mean deconvoluted in vivo dissolution profile was then compared with the optimized in vitro dissolution (Supplementary Methods).¹¹

$$F(t) = F_{\text{max}} * \left(1 - e^{-\frac{(t-\text{lag})^\beta}{\alpha}} \right) \quad (1)$$

The dissolution profile for the commercial formulation (manufactured by Aziende Chimiche Riunite Angelini Francesco S.p.A) was input as a discrete profile with a coefficient of variation (%CV) at each time point as estimated from in vivo deconvolution to capture population variability and a fixed value of %CV was also assumed. Simulations were then performed to reflect the reference

clinical study for a 300 mg single dose. The mean area under the curve (AUC)_{last}, AUC_{inf} , and C_{max} were calculated for each of the simulated subjects and then compared with those from the clinical trial (Supplementary Methods).¹¹

The updated model also includes a perfusion-limited, one-compartment brain model. Given that in vitro studies show trazodone is not a substrate for relevant transporters (P-gp,¹² BCRP, MRPs, OATPs, and OCTs) this model is valid (Table S1). Simulated total brain concentrations were corrected for tissue binding by human $f_{\text{u,brain}}$ values from the literature.¹³ A maximum effect (E_{max}) sigmoid model was applied to compute receptor occupancy using the simulated unbound brain concentration and the in vitro K_i values for the identified receptors of interest. The final model input parameters used are shown in Table 1.

Simulations for IR and OAD trazodone model verification

To verify the developed models for trazodone IR and OAD formulations, simulated plasma concentrations were compared with observed clinical data for a single 30 mg dose of IR trazodone and a single 300 mg dose of OAD trazodone. Simulation study design was matched to the clinical studies (number of subjects, demographics, etc.). For the IR formulation, 10 trials of 23 subjects per trial (age 22–54, 49% female) were simulated, whereas for the OAD formulation, a single trial of 43 subjects aged 18–56 years (46.5% women) was used. Model predictions were determined to be acceptable if the simulated parameters fell within 1.5-fold of the observed values. A comparison of the observed and predicted concentration-time profiles (visual check) for each formulation was also performed.

Target identification

Measured K_i values for several human molecular targets related to activity in the central nervous system (CNS) were available. These K_i values were determined using competitive radioligand binding assays using recombinant (CHO or HEK-293) cells expressing human receptors (58 targets, mainly GPCR and transporters, reported in Table S1). Trazodone was tested at seven log dilutions starting from 10 μM concentration. From the resulting competition curve, half-maximal inhibitory concentration (IC_{50}) values were determined by a nonlinear least square regression analysis. Inhibition constant (K_i) was calculated for each receptor according to the Cheng-Prusoff equation.¹⁴ Affinity toward the different receptors was determined in the range 10–1000 nM.

Parameter name	Value	Method/source
Physical chemistry and blood binding		
MW, g/mol	408.32	⁵
Log P	2.87	Calculated from experimental value of logD _{7.4} (=2.79) ¹
Compound type	Monoprotic base	¹
pK _a	6.61	Measured ¹
B/P	0.68	Calculated from measured E:P ratio of 0.2 (data on file)
f _{u,p}	0.0354	Measured by equilibrium dialysis (data on file)
Absorption		
Model	IR: first order OAD: ADAM with Solid Monolithic /Dissolution Profile Input (Table S1)	
F _a	IR: 0.98	Predicted from mean P _{app} (24.2 * 10 ⁻⁶ cm/s) obtained in Caco-2 cells and calibrated using metoprolol data (28.1 * 10 ⁻⁶ cm/s) ¹⁰
k _a (h ⁻¹)	IR/oral solution: 1.60 OAD: 0.07	IR: Predicted from mean P _{app} (24.2 * 10 ⁻⁶ cm/s) obtained in Caco-2 cells and calibrated using metoprolol data (28.1 * 10 ⁻⁶ cm/s) ¹⁰ OAD: fitting of concentration-time data following a single oral dose of 300 mg OAD trazodone ¹⁶
f _{u,gut}	1.0	Default value
Distribution		
Model	Full-PBPK	
V _{ss} (L/kg)	1.0	Predicted (Method 2) ¹⁷
Elimination		
CL _{iv} (L/h)	5	¹⁸
f _{m,CYP3A4} (%)	100	

TABLE 1 Input parameter values used to simulate the kinetics of trazodone

Abbreviations: ADAM, Advanced Dissolution Absorption and Metabolism; IR, immediate release; MW, molecular weight; OAD, once a day; PBPK, physiologically-based pharmacokinetic; V_{ss}, volume of distribution at steady state.

The list of targets was refined by identifying those with greater than or equal to 50% predicted receptor occupancy (RO; ≥ 0.5) at the maximum unbound brain concentration ($C_{\max, \text{brain}, u}$) for each dose/formulation level. This threshold was selected as it is the minimal SERT RO achieved for another multimodal drug, vortioxetine, in its clinical effective dose range.¹⁵ Free drug concentrations in the brain were used for final calculations as only protein-unbound drug concentrations are considered pharmacologically active. The RO was calculated using an E_{max} sigmoid model following the formula:

$$[\text{trazodone}]_{\text{brain}} / (K_i + [\text{trazodone}]_{\text{brain}}) \quad (2)$$

with K_i being the affinity constant for trazodone HCl on several molecular targets (Table S1). Trazodone maximum free base brain concentrations were taken from Simcyp V18r2

simulations for each dose (30 mg IR single dose, 30 mg IR at steady-state, and 300 mg OAD at steady-state). The reported literature value for human $f_{u, \text{brain}}$ of 0.077 was used to correct for the unbound concentration.¹³

Receptor occupancy simulations

The previously developed and validated trazodone model was used for all simulations.⁹ To determine the RO for the IR trazodone formulation, 10 virtual trials of 10 healthy subjects (50% women) aged 20–50 years receiving either a single oral dose or multiple oral doses (q.d., 8 days) of 30 mg IR trazodone were generated. Simulations for the OAD formulation were completed in the same manner, with 10 virtual trials of 10 healthy subjects (50% women) aged 20–50 years receiving multiple oral doses of OAD

trazodone (300 mg q.d.; 8 days). Each receptor was simulated independently due to limitations in applying the dynamic PD model across multiple targets.

Effect of in vitro variability on RO

As trazodone K_i values used for computations were the result of repeated in vitro assays on human receptors of interest, the effect of standard variability was evaluated on SERT and its RO value was determined for both formulations using two relative K_i values (160 and 280 nM; Table S2). The difference in the resulting RO, onset time, and duration of time at or above the target threshold between the two K_i values was assessed to determine the impact of the observed variability on RO predictions.

RESULTS

Model validation

The observed and simulated pharmacokinetic (PK) parameters for the 30 mg single dose IR trazodone and 300 mg single dose OAD trazodone are summarized in Table S3. The concentration-time profiles for the IR and OAD formulations are shown in Figure S1. The simulated values showed good agreement with the observed parameters, with the ratio of simulated to observed (S/O) for the AUC and C_{max} falling within 1.6-fold for both formulations (30 mg IR S/O range: 0.97–1.57; 300 mg OAD S/O range: 0.92–0.97).

Target identification

The simulated plasma and brain concentrations of trazodone following single and multiple oral doses of 30 mg

IR trazodone and multiple doses of 300 mg OAD trazodone are shown in Figures S2–S4. The PK parameters for all dosing intervals and formulations are summarized in Table 2.

Using predicted brain $C_{max,brain,u}$, static calculations were completed for previously identified targets to determine those reaching at least 50% occupancy for each dose and formulation of interest (Table 3). A total of 16 targets were found to be at or above the threshold value at least in one of the conditions evaluated (i.e., 300 mg trazodone OAD at steady-state) and were included in subsequent receptor occupancy modeling.

Immediate release formulation

Single dose receptor occupancy

Mean brain receptor occupancy following a single oral dose of 30 mg (IR formulation, adjusted to free base concentration) to healthy subjects (10 trials of 10 subjects) was simulated. RO was predicted for the seven targets through the application of the E_{max} sigmoid model using unbound brain concentrations. Simulations were used to estimate mean t_{onset} , that is the time to reach RO greater than or equal to 0.5, and duration of coverage at or above the target RO threshold for each target of interest.

Mean peak RO ranged from 0.59–0.91 for the targets of interest with an average t_{onset} of 0.12 h (Figure 1; Table 4). As would be expected, time at or above the target RO threshold (TAT) of 0.5 decreased with increasing K_i , with TAT of 27 h for the most potent (5-HT_{2A}, K_i = 14 nM) dropping to just over 2 h for the highest K_i (α_{1a} , K_i = 98 nM).

TABLE 2 Summary of total plasma and brain PK parameters for 30 mg IR trazodone following single and multiple oral doses and 300 mg OAD trazodone following multiple oral doses

Dose/formulation	Dosing frequency	C_{max} (ng/ml)	T_{max} (h)	AUC ^a (ng/ml × h)
Plasma				
30 mg IR	s.d.	442 (17%)	0.50	4992 (36%)
	q.d.	533 (21%)	0.50	4992 (40%)
300 mg OAD	q.d.	1632 (38%)	5.18	26651 (46%)
Brain				
30 mg IR	s.d.	702 (18%)	0.72	7841 (39%)
	q.d.	860 (23%)	0.65	8610 (41%)
300 mg OAD	q.d.	2814 (39%)	5.27	47882 (48%)

C_{max} and AUC are reported as geometric mean value, T_{max} as median, and variability as %CV.

Abbreviations: AUC, area under the curve; C_{max} , maximum concentration; IR, immediate release; OAD, once a day; PK, pharmacokinetic; T_{max} , time to maximum concentration.

^aAUC is AUC_{0-inf} following s.d. administration and AUC_{0-24h} following q.d. dosing.

TABLE 3 Calculated RO for proposed targets of interest by unbound brain concentration ($C_{\max, \text{brain}, u}$)

Target	System	K_i (nM)	RO		
			30 mg IR, s.d. ($C_{\max, \text{brain}, u} = 145$ nM/54.1 ng/ml)	30 mg IR, q.d. ($C_{\max, \text{brain}, u} = 178$ nM/66.3 ng/ml)	300 mg OAD, q.d. ($C_{\max, \text{brain}, u} = 583$ nM/217 ng/ml)
5-HT _{2A}	Serotonergic	14	0.912	0.927	0.977
α_{1B}	Adrenergic	15	0.906	0.922	0.975
5-HT _{1D}	Serotonergic	26	0.848	0.873	0.957
α_{1D}	Adrenergic	27	0.843	0.868	0.956
5-HT _{2B}	Serotonergic	48	0.752	0.788	0.924
5-HT _{1A}	Serotonergic	82	0.640	0.685	0.877
α_{1A}	Adrenergic	98	0.597	0.645	0.856
H ₁	Histaminergic	190	0.433	0.484	0.754
α_{2C}	Adrenergic	240	0.377	0.426	0.708
SERT	Serotonergic	280	0.342	0.389	0.675
5-HT _{2C}	Serotonergic	307	0.321	0.367	0.655
5-HT ₇	Serotonergic	406	0.264	0.305	0.589
α_{2A}	Adrenergic	430	0.253	0.293	0.575
α_{2B}	Adrenergic	460	0.240	0.279	0.559
D ₃	Dopaminergic	490	0.229	0.267	0.543
D ₂	Dopaminergic	560	0.206	0.241	0.510
5-HT _{1B}	Serotonergic	814	0.152	0.180	0.417
D ₁	Dopaminergic	929	0.135	0.161	0.385
D ₅	Dopaminergic	1200	0.108	0.129	0.327
5-HT _{5A}	Serotonergic	2340	0.058	0.071	0.199
NK-1	Tachykinin	3300	0.042	0.051	0.150
σ_1	Sigma	3980	0.035	0.043	0.128
β_2	Adrenergic	6000	0.024	0.029	0.089
5-HT ₄	Serotonergic	10,000	0.014	0.018	0.055
MOP	Opiate	15,436	0.009	0.011	0.036
M ₄	Cholinergic	30,000	0.005	0.006	0.019
M ₂	Cholinergic	34,127	0.004	0.005	0.017

For normalization from molar to ng/ml the molecular weight of the trazodone free-base was considered.

Values in bold indicate RO greater than or equal to 50%.

Abbreviations: $C_{\max, \text{brain}, u}$, unbound brain concentration; IR, immediate release; RO, receptor occupancy.

Multiple dose receptor occupancy

Ten trials of 10 healthy subjects were simulated for each of the seven targets showing static RO greater than or equal to 50% following multiple doses of 30 mg IR trazodone. The E_{\max} sigmoid model was applied to compute RO using simulated dynamic brain unbound concentration-time profiles at steady-state, day 7 of 8 days of dosing. The predicted mean peak receptor occupancies for the targets of interest were then used to estimate t_{onset} (relative to the first dose) and duration of time at or above the threshold, shown in Table 4 and Figure 2.

Mean peak RO at steady-state was similar to what was observed following a single IR dose, ranging from 0.64 to

0.92. Similarly, average t_{onset} to reach the target RO was 0.15 h (range: 0.09–0.25 h). Targets with high affinity showed TAT greater than 24 h (5-HT_{2A} and α_{1B} , $K_i = 14$ nM and 15 nM, respectively) with TAT ranging from 18.0 h to 2.10 h for the remaining receptors evaluated.

Once a day formulation

Multiple dose receptor occupancy

Ten trials of 10 healthy subjects were simulated for each of the 16 targets showing static RO greater than or equal




FIGURE 1 Simulated receptor occupancy vs. time profiles for 30 mg immediate release trazodone following a single oral dose in healthy subjects (mean – black line, 95th and 5th percentiles – grey lines, target receptor occupancy – horizontal dashed line)

to 50% following daily administration of 300 mg OAD trazodone. The E_{\max} sigmoid PD model was applied to compute receptor occupancies at steady-state (day 7), as shown in Figure 3. The predicted mean receptor occupancy, t_{onset} , and TAT for targets of interest are summarized in Table 4. At steady-state, the mean peak RO ranged from 0.48 to 0.97. The threshold value of RO greater than or equal to 0.5 was reached after the first

dose for 11 of 16 targets (t_{onset} ranging from 0.42 to 3.44 h). For the remaining four targets, t_{onset} was reached after the second dose (t_{onset} 27.1–28.4 h) for all except one receptor (D_3 , $t_{\text{onset}} = 52.4$ h) and one target identified in the static modeling, D_2 , that did not reach the RO threshold in the time simulated (mean peak RO = 0.48).

To determine if the average brain concentration at steady-state ($C_{\text{ave,brain,u}}$) could be used to determine those targets at

TABLE 4 Predicted geometric mean peak RO and, t_{onset} , and duration of TAT on day 1 and day 7 of 30 mg IR trazodone and on day 7 of 300 mg OAD trazodone administration

Target	K_i (nM)	Mean peak RO			t_{onset} (h)			TAT (h)		
		30 mg IR, s.d.	30 mg IR, q.d.	300 mg OAD	30 mg IR, s.d.	30 mg IR, q.d.	300 mg OAD	30 mg IR, s.d.	30 mg IR, q.d.	300 mg OAD
5-HT _{2A}	14	0.91	0.92	0.97	0.07	0.09	0.42	27.0	>24.0	>24.0
α_{1B}	15	0.90	0.92	0.97	0.07	0.09	0.42	26.0	>24.0	>24.0
5-HT _{1D}	26	0.84	0.87	0.95	0.07	0.09	0.50	18.3	18.2	>24.0
α_{1D}	27	0.84	0.86	0.95	0.10	0.09	0.50	17.7	17.7	>24.0
5-HT _{2B}	48	0.75	0.78	0.91	0.12	0.17	0.67	9.36	9.32	>24.0
5-HT _{1A}	81.8	0.63	0.68	0.86	0.19	0.25	0.76	3.29	3.11	>24.0
α_{1A}	98	0.59	0.64	0.84	0.22	0.25	0.84	2.18	2.1	>24.0
H ₁	190	-	-	0.73	-	-	1.43	-	-	16.6
α_{2C}	240	-	-	0.68	-	-	2.18	-	-	11.9
SERT	280	-	-	0.76	-	-	2.94	-	-	8.40
5-HT _{2C}	307	-	-	0.63	-	-	3.44	-	-	6.30
5-HT ₇	406	-	-	0.56	-	-	27.05	-	-	5.88
α_{2A}	430	-	-	0.55	-	-	27.64	-	-	4.28
α_{2B}	460	-	-	0.53	-	-	28.39	-	-	2.19
D ₃	490	-	-	0.52	-	-	52.42	-	-	2.01
D ₂	560	-	-	0.48	-	-	-	-	-	-

Abbreviations: IR, immediate release; OAD, once a day; RO, receptor occupancy; TAT, time above the RO threshold; t_{onset} , onset time.




FIGURE 2 Simulated mean receptor occupancy versus time following 7 days of dosing of immediate release trazodone (30 mg q.d.) in healthy subjects (mean – black line, 95th and 5th percentiles – grey lines, target receptor occupancy – horizontal dashed line)

or above the RO threshold, receptor occupancies were also computed using $C_{ave,brain,u}$. The simulated difference in RO between the two methods ranges from 10 to 17% for targets with relatively low binding affinity ($K_i > 100$ nM), whereas minimal difference was observed for those with high affinity ($\leq 5\%$; Table S4). It is important to note that the simulated difference in RO also reflects the difference between mean RO ($C_{ave,u}$ as inputs) and peak RO (dynamic brain concentration as inputs), both calculated at the seventh dosing day. For those targets with lower affinity, time to reach the target RO (t_{onset} , which is calculated starting from the first dose)

is reached after the second dose (i.e., >24 h), once the drug has accumulated in the brain. This is supported by the same targets not reaching the threshold value when mean steady-state concentration was used to determine occupancy and can be observed in the PD profiles presented in Figure S5.

In vitro variability

Ten virtual trials of 10 healthy subjects, as described previously, for each dose and formulation of interest were generated using




FIGURE 3 Simulated mean receptor occupancy following 7 days of dosing of once a day trazodone (300 mg q.d.; mean – black line, 95th and 5th percentiles – grey lines) in healthy subjects

two reported in vitro K_i values (160 nM and 280 nM) for SERT, summarized in Table S5 and Figure S6. For the IR formulation, a difference of ~ 1.4-fold in RO was observed for the low and high K_i values. There was no significant difference in RO between day 1 and day 7. A similar difference was seen for the OAD formulation, with a 1.2-fold difference in RO. For those formulations and dosing schemes reaching the target RO of 0.5, there was a slight increase in t_{onset} (1.18 h vs. 2.94 h) with a

significant decrease in the time at or above the threshold (20 h vs. 8.4 h) for the higher K_i value.

DISCUSSION

Using the updated trazodone PBPK model, brain receptor occupancy was determined for the 30 mg IR (single dose

and at steady-state) and 300 mg OAD (at steady-state) formulations. As it is difficult to determine brain concentrations, and therefore receptor occupancies *in vivo*, this approach allows for a better mechanistic understanding of the mechanism of action of trazodone when administered as an IR versus OAD formulation. This analysis demonstrated the utility of a PBPK-based approach for generating initial estimates of *in vivo* occupancy, particularly when clinical data are unavailable. Simulating the widest possible range of exposures for trazodone, 30 mg IR (SD) to 300 mg OAD (QD), allowed to further support determination of the optimal exposure-related differentiation of brain receptor activation by trazodone between formulations.

For the 30 mg IR formulation, seven targets (5-HT_{2A}, α_{1B} , 5-HT_{1D}, α_{1D} , 5-HT_{2B}, 5-HT_{1A}, and α_{1A}) were identified through static predictions from the unbound brain C_{max} to meet the threshold of 50% RO. Simulation shows that all targets exhibit rapid onset ($t_{onset} \leq 0.25$ h) after the first dose, with time above the target threshold (i.e., RO $\geq 50\%$) ranging from 2.4 to 26.9 h. It is important to note that, in the current study, the kinetics of association and dissociation of drugs binding to a specific target is not considered due to lack of data on k_{on} and k_{off} rates. Therefore, the duration of coverage at or above the target RO threshold may be overestimated. In this work, a “time above threshold” is reported as compared to “duration of occupancy” to address this difference. No significant differences were observed in the RO between day 1 and day 7.

In the current study, the PBPK simulations provided the mechanistic basis to support the clinically observed efficacy of trazodone IR to treat insomnia, already when given at the low dose of 30 mg, with predicted RO for 5-HT_{1A}, 5-HT_{1D}, 5-HT_{2A}, 5-HT_{2B}, α_{1A} , α_{1B} , and α_{1D} receptors of more than 0.63, 0.84, 0.91, 0.75, 0.59, 0.90, and 0.84, respectively.

Sixteen targets were identified from static modeling to reach 50% RO for the OAD formulation (300 mg q.d.). Simulations showed geometric mean peak RO at steady-state ranging from 0.48 to 0.97 with higher affinity targets ($K_i \leq 300$ nM) reaching the threshold occupancy after the first dose (t_{onset} range: 0.42–3.44 h). Targets with poor affinity did not reach RO greater than or equal to 0.5 until the second or third day of dosing, and in one case did not reach the threshold value at all, depending on the K_i value.

Comparing mean RO determined from the $C_{ave,brain,u}$ to the mean peak RO simulated using the dynamic brain unbound concentration-time profiles as PK inputs, differences of 6% or less in the predicted RO was observed for those targets with relatively tight binding affinity ($K_i < 100$ nM). The simulated difference in RO increased to 10–17% for targets with relatively poor binding affinity ($K_i > 100$

nM). Due to the similar results obtained from the two methods, it can be assumed that an adequate estimate of RO can be determined from average concentrations when dynamic concentration-time data is unavailable.

It is frequently noted that there are multiple reported values for parameters, such as K_i , with little information on how this effects subsequent clinical predictions. Here, the utility of PBPK for assessing the impact of this variability is shown through simulations with two reported K_i values for SERT—a key receptor in trazodone efficacy. This also serves to evaluate the effect of PD variability, as it is important to note that the simulator only accounts for variability in the PK simulation through built-in covariates (i.e., age, weight, sex, and plasma protein levels), not the PD calculations. Using different experimental K_i values, RO for a specific formulation and dosing scheme differed by an average of 1.4-fold. For repeat doses of 30 mg trazodone IR, this resulted in predicted ROs that reached the target threshold in one scenario and not in the other scenario (RO = 0.52 and 0.38 for the low and high K_i value, respectively). This difference appears to be more pronounced for borderline occupancy cases. In contrast, the difference in RO was significantly reduced in the case of OAD trazodone where SERT RO was generally higher (RO = 0.76 and 0.65, respectively).

Although previous work has been performed to determine brain RO for trazodone targets, the approach presented here differs in three key areas.¹⁶ First, a comprehensive evaluation of potential targets was completed using initial static calculations with over 50 targets evaluated compared to ~30 previously explored. Next, the RO calculations used here were based on unbound trazodone concentrations in the brain as compared to total brain trazodone concentrations, as only the free drug is able to bind with the receptors. Finally, RO was determined using a full PBPK model with a perfusion-limited brain compartment, as is supported by data showing no significant involvement of brain transporters, compared to a two-compartment model using a rodent $K_{p,uu}$ (i.e., a ratio of unbound brain to unbound plasma concentration) value to adjust concentrations. These key differences allow for an updated prediction of RO that better captures the processes involved *in vivo*.

It should be noted that the simulations presented here for the IR formulation do result in a 1.6-fold overprediction in trazodone AUC compared to the clinical value, which could affect the interpretation of the simulated RO. In model development for the IR and OAD formulations, two studies reporting trazodone clearance were identified with reported CL_{IV} values differing by approximately two-fold (5.3 L/h vs. 10.0 L/h).^{17,18} From the data presented, there is no obvious explanation for the discrepancy as the

only difference in study design appears to be PK sampling duration (24 h vs. 26 h). In the final model, the CL_{TV} has been set to 5.0 L/h for both formulations as to ensure the developed models can adequately capture the observed drug exposures for both formulations. Also, of importance for this discussion, is that the simulated F is different for both formulations—primarily due to the difference in the predicted value for F_a . For the IR formulation, F_a is ~ 1 in agreement with the findings from a human mass balance study in healthy volunteers, whereas the simulated F_a for the OAD formulation is 0.5, predicted from a mechanistic absorption model and in vivo dissolution profile. To improve the prediction of the IR formulation, a higher CL_{TV} or lower F_a would be required—the former resulting in poor prediction of the OAD formulation, and the latter not supported by the high permeability and reasonable solubility of the drug.

As described by Morgan et al., it is important to understand drug exposure at the site of action, target binding, and expression of functional pharmacological activity.¹⁹ In turn, achieving sufficient exposure, target binding, and pharmacology modulation for efficacy is key to clinical success. Although further work including imaging studies to visualize the in vivo occupancy during treatment would serve as confirmation for these results, this modeling exercise was able to simulate exposure at the site of action (brain) and occupancy at the relevant receptors. Overall, the results are consistent with the dose-dependent pharmacological activity of trazodone. The lower doses (i.e., the 30 mg of the IR formulation, both after single and repeated q.d. dosing), acts via the most potent functional properties—5-HT_{2A} and α_1 adrenergic antagonism and 5-HT_{1A} partial agonism and achieves reasonable levels of occupancy for these receptors (>0.59). The proper antidepressant dose of trazodone (i.e., 300 mg of the OAD formulation after repeated q.d. dosing) recruits additional pharmacological actions, such as the SERT blockade and antagonism of histaminergic H₁, 5-HT_{2C}, and 5-HT₇, and is able to achieve reasonable levels of occupancy at these additional receptors (>0.56). These findings form a strong foundation to further evaluate the multifunctional and multimodal mechanism of action for trazodone in achieving full antidepressant efficacy at target daily dose of 150–300 mg.

ACKNOWLEDGEMENTS

The authors wish to thank Matteo Vergassola and Parisa Foroutan (both part of Angelini Pharma S.p.A.) for their assistance in reviewing this work.

CONFLICT OF INTEREST

The authors declared no competing interests for this work.

AUTHOR CONTRIBUTIONS

L.O., M.T.R., A.C., G.D., S.J.M., A.K., and H.M.J. wrote the manuscript. L.O., G.M., M.T.R., A.C., G.D., S.J.M., A.K., and H.M.J. designed the research. S.J.M., A.K., and H.M.J. performed the research. L.O., G.M., M.T.R., A.C., G.D., S.M., A.K., and H.M.J. analyzed the data.

REFERENCES

- Herr J, Caspi A. Oleptro™ (trazodone hydrochloride) extended-release tablets. *PT*. 2011;36:2-18.
- Albert U, Lamba P, Stahl SM. Early response to trazodone once-a-day in major depressive disorder: review of the clinical data and putative mechanism for faster onset of action. *CNS Spectr*. 2021;26:232-242.
- Fagiolini A, Comandini A, Catena Dell'Osso M, Kasper S. Rediscovering trazodone for the treatment of major depressive disorder. *CNS Drugs*. 2012;26:1033-1049.
- Stahl SM. Mechanism of action of trazodone: a multifunctional drug. *CNS Spectr*. 2009;14:536-546.
- Angelini Pharma S.p.A. Trazodone Investigator's Brochure 8656B01; 2013.
- Schwasinger-Schmidt TE, Macaluso M. Other antidepressants. *Handb Exp Pharmacol*. 2019;250:325-355.
- Beasley CM Jr, Dornseif BE, Pultz JA, Bosomworth JC, Sayler ME. Fluoxetine versus trazodone: efficacy and activating-sedating effects. *J Clin Psychiatry*. 1991;52:294-299.
- Weisler RH, Johnston JA, Lineberry CG, et al. Comparison of bupropion and trazodone for the treatment of major depression. *J Clin Psychopharmacol*. 1994;14:170-179.
- Oggianu L, Ke AB, Chetty M, et al. Estimation of an appropriate dose of trazodone for pediatric insomnia and the potential for a trazodone-atomoxetine interaction. *CPT Pharmacometrics Syst Pharmacol*. 2020;9:77-86.
- Jamei M, Turner D, Yang J, et al. Population-based mechanistic prediction of oral drug absorption. *AAPS J*. 2009;11:225-237.
- Oggianu L, Pasculli G, Furlotti G, et al. Development and verification of population level IVIVC using mechanistic PBPK modeling for trazodone extended release formulations to support biowaiver for SUPAC changes to the formulation. *AAPS PharmSci* 2018;360. <https://www.eventscribe.net/2018/PharmSci360/fsPopUp.asp?efp=UUFSQIZZVFM1OTQ2&PosterID=166145&rnd=0.550835&mode=posterinfo>
- Gertz M, Harrison A, Houston JB, Galetin A. Prediction of human intestinal first-pass metabolism of 25 CYP3A substrates from in vitro clearance and permeability data. *Drug Metab Dispos*. 2010;38:1147-1158.
- Di LI, Umland JP, Chang G, et al. Species independence in brain tissue binding using brain homogenates. *Drug Metab Dispos*. 2011;39:1270-1277.
- Cheng Y, Prusoff WH. Relationship between the inhibition constant (K₁) and the concentration of inhibitor which causes 50 per cent inhibition (I₅₀) of an enzymatic reaction. *Biochem Pharmacol*. 1973;22:3099-3108.
- Sanchez C, Asin KE, Artigas F. Vortioxetine, a novel antidepressant with multimodal activity: review of preclinical and clinical data. *Pharmacol Ther*. 2015;145:43-57.
- Settimo L, Taylor D. Evaluating the dose-dependent mechanism of action of trazodone by estimation of occupancies for

- different brain neurotransmitter targets. *J Psychopharmacol*. 2018;32:96-104.
17. Nilsen OG, Dale O. Single dose pharmacokinetics of trazodone in healthy subjects. *Pharmacol Toxicol*. 1992;71:150-153.
 18. Greenblatt DJ, Friedman H, Burstein ES, et al. Trazodone kinetics: effect of age, gender, and obesity. *Clin Pharmacol Ther*. 1987;42:193-200.
 19. Morgan P, Van Der Graaf PH, Arrowsmith J, et al. Can the flow of medicines be improved? Fundamental pharmacokinetic and pharmacological principles toward improving Phase II survival. *Drug Discov Today*. 2012;17:419-424.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Oggianu L, Di Dato G, Mangano G, et al. Estimation of brain receptor occupancy for trazodone immediate release and once a day formulations. *Clin Transl Sci*. 2022;15:1417-1429. doi:[10.1111/cts.13253](https://doi.org/10.1111/cts.13253)