

Some Comments for Better Understanding of the Study Entitled “Reduced Vitamin D Levels are Associated with Stroke-Associated Pneumonia in Patients with Acute Ischemic Stroke” [Letter]

This article was published in the following Dove Press journal:
Clinical Interventions in Aging

Gulistan Bahat ¹
Serdar Ozkok ¹
Savaş Ozturk ²
Mehmet Akif Karan¹

¹Istanbul University, Istanbul Medical School, Department of Internal Medicine, Division of Geriatrics, Istanbul, Turkey;
²Istanbul Haseki Training and Research Hospital, Department of Internal Medicine, Istanbul, Turkey

Dear editor

We have read the article by Huang et al with great interest.¹ Authors aimed to reveal the relationship between serum vitamin D levels and stroke-associated pneumonia (SAP). They measured the serum vitamin D levels of the acute ischemic stroke (AIS) patients within 24 hrs of admission and divided them into three groups accordingly. Compared to the patients without SAP, patients with SAP had significantly lower vitamin D levels. The prevalence of SAP was higher in the vitamin D deficient group. In the logistic regression analysis, it was detected that vitamin D deficiency and insufficiency were independently associated with SAP.

This study is an important contribution to the literature and we congratulate the authors. We have some comments regarding the study analyses. Firstly, the authors mentioned about the anti-inflammatory effects of vitamin D and hypothesized that a reduction of vitamin D level might be associated with the occurrence of SAP due to its reduced anti-inflammatory effect. The anti-inflammatory effect of vitamin D is important for many autoimmune diseases or low-level inflammatory conditions like obesity or cardiovascular diseases.² In other words, sufficient levels of vitamin D would be helpful in regards to protection from the situations stated above. However, an increased likelihood of infections in vitamin D deficiency is basically caused by the decreased effect of vitamin D on promoting protective immunity. Vitamin D plays an important role in the innate antimicrobial response, and deficiency of it causes increased susceptibility to infections.³ Hence, one can suggest that a reduction of vitamin D level might be associated with the occurrence of SAP due to its “reduced effect on promoting protective immunity”, instead of “reduced anti-inflammatory effect”.

Secondly, dysphagia was one of the baseline clinical parameters stated in the methods section, but how it was detected is important and was not mentioned. Was it detected by a validated tool or by examination or just by asking about it? And was it present before stroke event or developed after? Actually, dysphagia history before admission might be a cause for malnutrition and vitamin D deficiency and, therefore may be an underlying factor for the relationship between vitamin D deficiency

Correspondence: Gulistan Bahat
Tel +90 212 414 20 00 Ext. 31478
Email gbahatozturk@yahoo.com

and SAP. This data should be clarified, and if there is no data on dysphagia history before admission, this should be stated as a limitation of the study. Furthermore, patients' nutrition status should also be mentioned. Malnutrition itself can be the cause of vitamin D deficiency and the factor related to SAP. Therefore, if the nutrition status of patients was evaluated, the data should be given, and if not, we suggest that it should also be stated as a limitation of the study.

In addition, the authors noted that the increase in the incidence of diabetes with the decrease of vitamin D level is an interesting finding. In fact, there is mounting evidence about the inverse relationship between vitamin D status and diabetes mellitus or insulin resistance.^{4,5} Therefore, the increasing prevalence of diabetes with decreasing vitamin D levels is an expected finding.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Huang GQ, Cheng HR, Wu YM, et al. Reduced vitamin D levels are associated with stroke-associated pneumonia in patients with acute ischemic stroke. *Clin Interv Aging*. 2019;14:2305–2314. doi:10.2147/CIA.S230255
2. Mousa A, Misso M, Teede H, et al. Effect of vitamin D supplementation on inflammation: protocol for a systematic review. *BMJ Open*. 2016;6:e010804. doi:10.1136/bmjopen-2015-010804
3. Aranow C. Vitamin D and the immune system. *J Investig Med*. 2011;59(6):881–886. doi:10.2310/JIM.0b013e31821b8755
4. Berridge MJ. Vitamin D deficiency and diabetes. *Biochem J*. 2017;474(8):1321–1332. doi:10.1042/BCJ20170042
5. Husemoen LL, Thuesen BH, Fenger M, et al. Serum 25(OH)D and type 2 diabetes association in a general population: a prospective study. *Diabetes Care*. 2012;35:1695–1700. doi:10.2337/dc11-1309

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Interventions in Aging 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Interventions in Aging editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Clinical Interventions in Aging

Dovepress

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier

Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>