



## Original Research

# Prediction of Prognosis Acute Pancreatitis with Inflammatory Markers and Patient Characteristics Compared to the Scoring System: Real-Life Data

Cagatay Ak,<sup>1</sup> Resul Kahraman,<sup>2</sup> Suleyman Sayar,<sup>2</sup> Ebru Tarikci Kilic,<sup>3</sup> Gupse Adali,<sup>2</sup> Kamil Ozdil<sup>2</sup>

<sup>1</sup>Department of Gastroenterology, Nigde Training and Research Hospital, Nigde, Türkiye

<sup>2</sup>Department of Gastroenterology, Health Sciences University, Umraniye Training and Research Hospital, Istanbul, Türkiye

<sup>3</sup>Department of Anesthesiology, Health Sciences University, Umraniye Training and Research Hospital, Istanbul, Türkiye

### Abstract

**Objectives:** Acute pancreatitis (AP) is an inflammatory disease with a high morbidity and mortality rate. It is one of the most common causes of hospitalization among gastrointestinal system diseases. Inflammatory and other factors that predict the severity of AP are very important for patient management. This study will analyze the factors associated with the severity of AP.

**Methods:** The sample consisted of 514 patients. Demographic characteristics, comorbid diseases, causes of AP, body mass index (BMI), tobacco use, blood at admission, amylase, lipase, leukocyte, neutrophil, lymphocyte, C-reactive protein (CRP), mean platelet volume, red cell distribution width, albumin, calcium, and CRP values at 48th h were recorded. The bedside index of severity in AP (BISAP), Ranson score, neutrophil-lymphocyte ratio (NLR), and platelet-to-lymphocyte ratio (PLR) values was calculated and recorded. The relationship between these parameters and the severity of AP was analyzed according to the Atlanta classification.

**Results:** Participants had a mean age of 55±17.8 years. More than half the participants were women (n=272, 52.9%). Biliary causes were the most common etiological causes (n=299, 58.2%). Most participants had mild pancreatitis (n=416, 80.9%). The severity of AP was associated with tobacco use, high BMI, thrombocytosis, high NLR, high PLR, high 48<sup>th</sup> h CRP, hypoalbuminemia, hypocalcemia, aspartate aminotransferase/alanine aminotransferase ratio (AST/ALT ratio), and high Ranson and BISAP scores.

**Conclusion:** Biochemical markers that give rapid results in the early period can provide information about the severity of AP. We may develop new scores by combining these parameters.

**Keywords:** Acute pancreatitis, de ritis ratio, neutrophil-lymphocyte ratio, platelet-lymphocyte ratio, prognostic markers

Please cite this article as "Ak C, Kahraman R, Sayar S, Tarikci Kilic E, Adali G, Ozdil K. Prediction of Prognosis Acute Pancreatitis with Inflammatory Markers and Patient Characteristics Compared to the Scoring System: Real-Life Data. Med Bull Sisli Etfal Hosp 2023;57(2):182–188".

Acute pancreatitis (AP) is an inflammatory disease of the pancreas with multifactorial pathogenesis that causes systemic and peripancreatic tissue inflammation. Enzyme activation plays a central role in local pancreatic injury.<sup>[1]</sup> AP mostly follows a mild course, resulting in rapid clinical improvement with fluid resuscitation and the symptomatic treatment of

pain and nausea. Population-based studies have shown the proportion of severe pancreatitis between 8% and 20%.<sup>[2]</sup> Despite treatment, AP can lead to complications, morbidity, and mortality. Predicting the prognosis related to disease severity in the earliest period is important in terms of providing intensive care support and other specific treatments for the patient.

**Address for correspondence:** Cagatay Ak, MD. Department of Gastroenterology, Nigde Training and Research Hospital, Nigde, Türkiye

**Phone:** +90 505 660 80 66 **E-mail:** cagatayak88@gmail.com

**Submitted Date:** July 07, 2022 **Revised Date:** December 11, 2022 **Accepted Date:** December 26, 2022 **Available Online Date:** June 20, 2023

©Copyright 2023 by The Medical Bulletin of Sisli Etfal Hospital - Available online at [www.sislietfaltip.org](http://www.sislietfaltip.org)

**OPEN ACCESS** This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).



Researchers have developed scoring systems, such as Ranson score,<sup>[3]</sup> Assessment of Acute Physiology and Chronic Health (APACHE-2),<sup>[4]</sup> and Bedside Index for Severity in AP (BISAP).<sup>[5]</sup> However, the Atlanta classification revised in 2012 is the most common classification system used to assess the severity of AP.<sup>[6]</sup> Researchers also focus on the relationship between new markers of inflammation (hemogram and biochemical tests, neutrophil/lymphocyte ratio [NLR], and platelet/lymphocyte ratio [PLR]) and severity of pancreatitis.<sup>[7-9]</sup> Obesity and tobacco use are associated with the severity of AP.<sup>[10,11]</sup> The most common causes of AP are gallstones, alcohol use, and idiopathic factors. However, its causes vary from region to region.<sup>[12]</sup> Identifying etiologic causes by geography and determining practical prognostic markers can be invaluable in patient management. This study will look into the associated with the severity of AP (according to the Atlanta classification) and demographic characteristics, etiological causes, tobacco use, body mass index (BMI), leukocytes, NLR, PLR, mean platelet volume (MPV), red cell distribution width (RDW), albumin, calcium values, BISAP, and Ranson scores. Another objective of this study is to determine, in which scores or parameters are most effective in predicting the severity of AP.

Methods

Seven hundred and ninety-six AP patients were admitted to the Health Sciences University Umraniye Training and Research Hospital Gastroenterology Clinic between January 01, 2015, and January 01, 2020. The sample consisted of 514 patients. Figure 1 shows the inclusion criteria. A patient was diagnosed with AP if he/she had two of the three parameters: 1 – amylase or lipase values more than 3 times

the upper limit of normal, 2 – typical abdominal pain, and 3 – typical radiological findings (According to Atlanta criteria).<sup>[6]</sup> Demographic characteristics, comorbid diseases, causes of AP, BMI, tobacco use, blood at admission, amylase, lipase, leukocyte, neutrophil, lymphocyte, hemoglobin, platelet, C-reactive protein (CRP), MPV, RDW, albumin, calcium, AST, ALT, and CRP values at 48<sup>th</sup> h were recorded. Ranson score, BISAP score, Atlanta score, AST/ALT ratio, and NLR and PLR values were calculated and recorded. In addition, the parameters evaluated in Ranson score, BISAP score, and Atlanta score are shown in Tables 1 and 2.<sup>[3,5,6]</sup>


Table 1. Ranson's criteria and BISAP score	
Ranson's criteria (1 point each)	BISAP score (1 point each)
On admission	BUN >25
WBC >16.000/μL	Impaired mental status
Age >55 years	SIRS
Glucose >200 mg/dL	Age >60 years
AST >250 IU/L	Pleural effusions
LDH >350 IU/L	
Within 48 h of admission	
HCT decrease >10%	
BUN increase >5	
Serum calcium <8 mg/dL	
Arterial pO <sub>2</sub> <60 mmHg	
Base deficit >4 mEq/L	
Fluid needs >6 L	

WBC: White blood cells; AST: Aspart aminotransferase; LDH: Lactate dehydrogenase; HCT: Hematocrit; BUN: Blood urea nitrogen; SIRS: Systemic inflammatory response syndrome.

Table 2. Revised atlanta classification	
A. Mild acute pancreatitis	
I. No organ failure	
II. No local or systemic complications	
B. Moderately severe acute pancreatitis	
I. Organ failure that resolves within 48 h (transient organ failure and/or	
II. Local or systemic complications without persistent organ failure	
C. Severe acute pancreatitis: Persistent organ failure (>48 h)	
I. Single organ failure	
II. Multiple organ failure	

Local complication: acute fluid collections; pancreatic necrosis; acute pseudocyst; and pancreatic abscess.

Organ failure and systemic complications: shock: SBP <90 mmHg; pulmonary insufficiency: PaO<sub>2</sub> ≤60 mmHg; renal failure: creatinine ≥170 μmol/L (≥2 mg/dL) after rehydration; gastrointestinal bleeding: 500 mL in 24 h; disseminated intravascular coagulation: platelets ≤100, 000/mm<sup>3</sup>, fibrinogen <1.0 g/L and fibrin-split products >80 μg/L; and severe metabolic disturbances: calcium ≤1.87 mmol/L or ≤7.5 mg/dL.



**Figure 1.** Inclusion and Exclusion Criteria.

Late admission: The patient's admission to the hospital one day after the onset of pain, or referral from another centre; Recurrent pancreatitis: Repeated hospitalizations due to acute pancreatitis.

All patients were hospitalized and followed up. All patients, except those with hypercalcemia etiology, were administered (IV) 5–10 mL/kg/h ringer lactate solution. Patients hospitalized for hypercalcemia were administered (IV) 5–10 mL/kg/h isotonic sodium chloride. Vital signs were monitored. Daily laboratory analysis was performed. Cross-sectional imaging was performed to detect local complications in patients whose abdominal pain did not regress and/or serum amylase lipase values did not regress 3 times below the reference limit at the 72<sup>nd</sup> h of treatment. Participants were classified as mild pancreatitis (MSAP), moderate pancreatitis (MAP), and SAP according to the 2012 revision of the Atlanta classification.<sup>[6]</sup> The general characteristics, inflammatory and laboratory indicators, and prognostic scores of the patients were compared according to the severity of AP.

This study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of Umraniye Training and Research Hospital (Date: March 18, 2020 and No: B.10.1.TKH.4.34.H.GP.0.01/70).

### Statistical Analysis

All statistical procedures were performed using SPSS software (version 25.0, SPSS Inc., Chicago, IL, institutionally registered software). The distribution of the data was found to be normal with the Kolmogorov–Smirnov test. While evaluating the study data, the descriptive statistical methods mean, standard deviation, frequency, and percentage were used. One-way ANOVA test was used to compare more than two groups for parametric data. Chi-square test was used when investigating the difference between groups for categorical data. Significance was evaluated at  $p < 0.05$  for all values.

### Results

The sample consisted of 514 patients. Participants had a mean age of  $55 \pm 17.8$  years. More than half the participants were women (52.9%). Our results showed that the three most common etiologic causes of AP were biliary in 299 (58.2%) patients, idiopathic in 98 (19.1%) patients, and alcohol-related causes in 44 (8.6%) patients. After these three etiological causes, drug use in 29 (5.6%) patients, post-ERCP in 20 (3.9%) patients, hypertriglyceridemia in 19 (3.7%) patients, hypercalcemia in 2 (0.4%) patients, autoimmune pancreatitis in 2 (0.4%) patients, and 1 (0.4%) patient had trauma-related AP. Table 3 shows the participants' general characteristics. Participants had MSAP (80.9%;  $n=416$ ), MAP (14.8%;  $n=76$ ), or SAP (4.3%;  $n=22$ ). Three participants (0.6%) died due to AP. BMI elevation and smoking were found to be statistically significantly different between MAP, MSAP, and SAP patient groups

**Table 3.** General characteristics and etiological causes

Parameters	n STD (%)
Age (years)	55±17.80
Gender	
Female	272 (52.9)
Male	242 (47.1)
Comorbidities	
Hypertension	141 (27.4)
Diabetes	68 (13.2)
Coronary artery disease	40 (7.8)
Congestive heart failure	17 (3.3)
Chronic renal failure	9 (1.8)
Chronic obstructive pulmonary disease	12 (2.3)
Asthma	19 (3.7)
Cerebrovascular diseases	9 (1.8)
Other diseases	84 (16.3)
Pancreatitis etiology	
Biliary	299 (58.2)
Alcohol	44 (8.6)
Idiopathic	98 (19.1)
Drug	29 (5.6)
Post-ERCP	20 (3.9)
Hypertriglyceridemia	19 (3.7)
Hypercalcemia	2 (0.4)
Autoimmune	2 (0.4)
Trauma	1 (0.2)
BMI	24.1±3.1
Hospitalization (day)	5.56±4.52

ERCP: Endoscopic retrograde cholangiopancreatography; BMI: Body mass index; Other diseases: Rheumatoid arthritis, Alzheimer, Dyslipidemia, Ankylosing spondylitis, Parkinson, Hypothyroidism, Dementia, Benign prostatic hyperplasia.

( $p=0.000$ ). There was no statistical difference between the patient groups in serum amylase and lipase levels at admission to the hospital ( $p=0.657$ ,  $p=0.524$ ). Platelet count was statistically significantly different between NLR and PLR, MAP, MSAP, and SAP patient groups ( $p=0.000$ ). There was no statistical difference between the CRP values of the patient groups at the time of admission to the hospital, but a statistically significant difference was found between the CRP values at the 48th h ( $p=0.000$ ). Albumin and calcium levels were statistically significantly different in MAP, MSAP, and SAP patient groups ( $p=0.000$ ). In addition, AST/ALT ratio, Ranson score, and BISAP score were statistically significantly different in MAP, MSAP, and SAP patient groups ( $p=0.000$ ). Table 4 shows the risk factors related with MSAP and SAP.

**Table 4.** Factors associated with acute pancreatitis severity

Parameters	MAP n=416 (80.9%) (Mean±SD)	MSAP n=76 (14.8) (Mean±SD)	SAP n=22 (4.3) (Mean±SD)	p*
Gender				
Female	222 (53.4)	41 (53.9)	9 (40.9)	0.512 <sup>†</sup>
Male	194 (46.6)	35 (46.1)	13 (59.1)	
Age	54.85±17.63	54.08±19.80	55.73±14.29	0.941 <sup>†</sup>
BMI	23.9±2.5	27.5±3.8	28.5±2.8	0.000*
Tobacco use	35 (8.4)	21 (27.6)	8 (36.4)	0.000 <sup>†</sup>
Hospitalization (day)	4.00±1.01	10.12±3.31	19.36±10.78	0.000 <sup>†</sup>
Etiology				
Biliary	255 (61.3)	34 (44.7)	10 (45.5)	0.000 <sup>†</sup>
Alcohol	22 (5.3)	18 (23.7)	4 (18.2)	0.002 <sup>†</sup>
Idiopathic	77 (18.5)	17 (22.4)	4 (18.2)	0.000 <sup>†</sup>
Amylase U/L	1433.11±962.7	1372.45±808.88	1578.91±933.76	0.657*
Lipase U/L	1686.17±1632.76	1743.92±1233.37	2084.32±2437.09	0.524*
Leukocyte u/L	9.69±3.6	10.26±4.01	11.35±3.55	0.069*
Hemoglobin g/dL	12.68±1.73	12.55±2.07	13.01±1.26	0.563*
Platelet u/L	218.30±69.44	304.84±97.53	332.31±115.70	0.000*
NLR	3.87±2.02	7.37±2.99	10.18±3.37	0.000*
PLR	130.19±74.30	287.39±140.25	360.97±157.58	0.000*
MPV fL	8.59±1.63	8.3±1.29	8.19±1.2	0.195*
RDW fL	13.53±1.22	13.67±1.16	13.75±1.59	0.491*
CRP mg/L	1.82±1.87	1.79±1.82	2.58±2.81	0.188*
48 <sup>th</sup> h CRP mg/L	7.26±3.23	14.77±6.26	18.96±5.9	0.000*
Albumin g/dL	3.71±0.41	3.3±0.37	3.08±0.44	0.000*
Calcium mg/dL	8.73±0.56	8.47±0.44	8.29±0.41	0.000*
AST IU/L	131.63±159.78	172.61±261.32	174.23±190.56	0.123*
ALT IU/L	158.42±183.45	119.55±170.45	119.09±146.93	0.158*
AST/ALT	1.02±0.62	1.48±0.58	1.64±0.73	0.000*
Ranson Score	1.08±1.09	1.67±1.06	2.91±1.19	0.000*
BISAP Score	0.86±0.80	1.74±0.87	3.77±0.75	0.000*

Data are shown as median (interquartile range) or n (%). BMI: body mass index; NLR: neutrophil-lymphocyte ratio; PLR: platelet-lymphocyte ratio; MPV: mean platelet volume; RDW: red cell distribution width; CRP: C-reactive protein; AST: aspartate aminotransferase; ALT: alanine aminotransferase. \*One-way ANOVA<sup>†</sup> Chi-square.

## Discussion

Eight in ten AP patients present with a mild edematous form and are discharged after a couple of days. However, two in ten AP patients develop a severe or complicated course of pancreatitis characterized by early or delayed systemic and local complications. Our results showed that eight in ten participants had MSAP (80.9%), which is consistent with the literature.<sup>[13]</sup>

Although there are various scoring systems (Ranson score, Balthazar score, SOFA score, APACHE II score, and Marshall score) to assess pancreatitis severity early (<24 h), they are not used very commonly in clinical practice.<sup>[13,14]</sup> It is more practical to use the BISAP score to diagnose SAP.<sup>[15]</sup> In addition, some clinical and laboratory parameters can provide

information about the prognosis of pancreatitis at the time of admission.<sup>[16]</sup> This study analyzed the relationship between clinical and laboratory parameters at admission and BISAP and Ranson scores and AP severity

Our results showed that the three most common etiologic causes of AP were biliary in 299 (58.2%) patients, idiopathic in 98 (19.1%) patients, and alcohol-related causes in 44 (8.6%) patients. Biliary causes were higher, while alcohol-related causes were lower than those reported by earlier studies.<sup>[17,18]</sup> This is probably because alcohol consumption is low in Turkey. A similar result was found in a study conducted in our country.<sup>[19]</sup> In their meta-analysis, Roberts et al. (2017)<sup>[20]</sup> have reported that the biliary/alcohol ratio is higher in Southern European countries (including Turkey) than in Northern European countries.



Obesity is an independent risk factor for SAP.<sup>[21]</sup> We found that SAP ( $28.5 \pm 2.8$ ) and MSAP ( $27.5 \pm 3.8$ ) participants had a higher mean BMI than MAP participants ( $23.9 \pm 2.5$ ). In their meta-analysis, Cruz-Monserrate, Conwell, and Krishna (2016) have also found that a BMI  $>25$  increases the risk for SAP.<sup>[22]</sup> There is a positive correlation between tobacco use and AP.<sup>[23-25]</sup> In this study, we found that tobacco use was statistically higher in MSAP and SAP than in MAP. Epigenetic changes or environmental stimuli combined with smoking can further advance pancreatic damage.<sup>[26]</sup>

CRP is one of the most important biomarkers related to the severity of AP. It is widely used despite the delayed increase peaking 72 h after the onset of symptoms.<sup>[27]</sup> We could not detect any difference in CRP levels between patient groups at admission. However, the SAP, MSAP, and MAP groups had a 48th-h CRP level of  $18.96 \pm 5.9$  mg/L,  $14.77 \pm 6.26$  mg/L, and  $7.26 \pm 3.23$  mg/L, respectively. CRP levels above 15 mg/L 48 h after admission help differentiate MSAP from SAP.<sup>[27]</sup>

NLR attracts the attention of many researchers because it has such advantages as rapid detection, high sensitivity, low cost, and non-invasiveness.<sup>[28]</sup> Furthermore, like increased NLR, PLR has been associated with inflammatory states and poor outcomes in SAP are explained by uncontrolled SIRS and its progression to multi-organ dysfunction syndrome.<sup>[29]</sup> We detected a positive association with NLR and PLR to AP severity, which is consistent with the literature.<sup>[30-32]</sup>

Inflammation reduces red blood cell (RBC) half-life, affects iron metabolism and erythropoiesis, and increases hemolysis, resulting in increased RBC size and heterogeneity. Therefore, RDW can be used as a non-specific inflammatory indicator.<sup>[33]</sup> There is a correlation between RDW and AP severity.<sup>[32]</sup> AP may increase platelet consumption due to pancreatitis region and distant organ inflammation and cause a decrease in MPV value.<sup>[34]</sup> We found that the SAP and MSAP groups had lower MPV but higher RDW than the MAP group. However, the difference was statistically insignificant. We excluded patients who were admitted to the clinic late. MPV and RDW may not change significantly early in the inflammatory process and may not be an early indicator of the severity of AP.

We detected that hypocalcemia and hypoalbuminemia were associated with the severity of AP, which is consistent with the literature.<sup>[35-37]</sup> Albumin maintains osmotic pressure, binds endogenous/exogenous substances with high vascular permeability, prevents coagulation, buffers acid-base status, and has antioxidant, anti-inflammatory, and anti-apoptosis effects.<sup>[38,39]</sup> Therefore, hypoalbuminemia may predict persistent organ failure and severe inflammation rather than being a marker of AP severity. We think that

AP causes hypoalbuminemia and that the higher the hypoalbuminemia, the more complicated the AP. The causes of early-phase hypocalcemia in AP are the self-digestion of mesenteric fat by pancreatic enzymes, the release of free fatty acids that form calcium salts, transient hypoparathyroidism, and hypomagnesemia.<sup>[40-42]</sup> Therefore, calcium levels may decrease with an increase in the severity of inflammation and may predict the severity of AP.

In our study, no correlation was found between AST and ALT levels and the severity of the disease, but it was found that the AST/ALT ratio related the severity of the disease. Fernando De Ritis (1957) was the first to define the ratio of serum activities of AST and ALT.<sup>[43]</sup> This ratio, also known as the De Ritis ratio, is used as a marker in the prognosis of many diseases.<sup>[44]</sup> A high De Ritis ratio in AP may be an early marker of severe inflammation and cell destruction resulting from organ failure. To the best of our knowledge, this is the first study to show associated with the severity of AP and De Ritis ratio. However, the low number of patients in the SAP group and the higher incidence of biliary etiology in the MAP group may have affected this result.

We detected a positive correlation between Ranson and BISAP scores and the severity of AP, which is consistent with the literature.<sup>[32]</sup> The main limitation of the Ranson criteria is that evaluation cannot be completed until 48 h after admission, which may miss an early therapeutic window and increase mortality. Both BISAP and Ranson scoring systems accurately predict the severity of AP, but the former is more effective because it is easier to use than the latter.

The mortality rate was 0.6% in this study. Research shows that the AP-related mortality rate ranges from 2.5% to 12.5%.<sup>[13,15,27,31]</sup> The low mortality rate in this study may be because we administered (IV) 5–10 mL/kg/h (aggressively) hydration on the 1<sup>st</sup> day and the breadth of patients' exclusion criteria from the study. We observed that the higher the AP severity, the longer the hospitalization. Determining the parameters of the severity of AP is effective in predicting morbidity and mortality rates and increased workload and costs. AP is one of the most common causes of hospitalization due to the gastrointestinal tract. Therefore, health-care professionals involved in patient management should carefully evaluate the parameters related to the severity of AP. In this study, it was determined that BISAP and Ranson scores were associated with the severity of AP. In addition, in this study, it was determined that laboratory parameters (platelet, NLR, PLR, albumin, calcium, 48<sup>th</sup> h CRP value, and AST/ALT ratio) that gave results in a short time were associated with the severity of AP. Predicting the severity of AP early may be effective in reducing morbidity and mortality with closer follow-up of the patient and more

aggressive treatment. Perhaps, with future studies, these laboratory parameters can be used in the development of new scoring systems that predict the severity of AP.

This study was conducted in one of the centers evaluating the highest number of AP patients in Turkey. The strengths of the study are the inclusion criteria, sample size, and the broad spectrum of parameters. The limitation of the study is that it was retrospective.

## Conclusion

High BMI, tobacco use, high NLR, high PLR, thrombocytosis, and high CRP (48<sup>th</sup> h) levels were found to be associated with the severity of AP. MPV and RDW may not predict the severity of AP in the early admission. Hypoalbuminemia, hypocalcemia, Ranson, and BISAP scores were found to be associated with the severity of AP.

## Disclosures

**Ethics Committee Approval:** This study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of Umranıye Training and Research Hospital (Date: 18/03/2020 & No: B.10.1.TKH.4.34.H.GP.01/70).

**Peer-review:** Externally peer-reviewed.

**Conflict of Interest:** None declared.

**Authorship Contributions:** Author Contributions: Concept – C.A., K.O.; Design – R.K., G.A.; Supervision – G.A., S.S.; Materials – C.A., E.T.K.; Data collection and/or processing – C.A., E.T.K.; Analysis and/or interpretation – E.T.K.; Literature search – C.A., S.S.; Writing – C.A.; Critical review – R.K., K.O.

## References

- Ruiz-Rebollo ML, Muñoz-Moreno MF, Mayo-Iscar A, Udaondo-Cascante MA, Nistal RB. Statin intake can decrease acute pancreatitis severity. *Pancreatol* 2019;19:807–12.
- Garg PK, Singh VP. Organ failure due to systemic injury in acute pancreatitis. *Gastroenterology* 2019;156:2008–23.
- Ranson JH, Rifkind KM, Roses DF, Fink SD, Eng K, Localio SA. Objective early identification of severe acute pancreatitis. *Am J Gastroenterol* 1974;61:443–51.
- Larvin M, McMahon MJ. APACHE-II score for assessment and monitoring of acute pancreatitis. *Lancet* 1989;2:201–5.
- Wu BU, Johannes RS, Sun X, Tabak Y, Conwell DL, Banks PA. The early prediction of mortality in acute pancreatitis: a large population-based study. *Gut* 2008;57:1698–703.
- Banks PA, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, Sarr MG, et al. Classification of acute pancreatitis--2012: revision of the Atlanta classification and definitions by international consensus. *Gut* 2013;62:102–11.
- Silva-Vaz P, Abrantes AM, Morgado-Nunes S, Castelo-Branco M, Gouveia A, Botelho MF, et al. Evaluation of prognostic factors of severity in acute biliary pancreatitis. *Int J Mol Sci* 2020;21:4300.
- Yarkaç A, Kose A, Bozkurt Babuş S, Ates F, Örekici Temel G, Ölmez A. The value of hematological parameters in acute pancreatitis. *Ulus Travma Acil Cerrahi Derg* 2019;25:453–60.
- Azab B, Jaglall N, Atallah JP, Lamet A, Raja-Surya V, Farah B, et al. Neutrophil-lymphocyte ratio as a predictor of adverse outcomes of acute pancreatitis. *Pancreatol* 2011;11:445–52.
- Aune D, Mahamat-Saleh Y, Norat T, Riboli E. High body mass index and central adiposity is associated with increased risk of acute pancreatitis: a meta-analysis. *Dig Dis Sci* 2021;66:1249–67.
- Ye X, Lu G, Huai J, Ding J. Impact of smoking on the risk of pancreatitis: a systematic review and meta-analysis. *PLoS One* 2015;10:e0124075.
- Zilio MB, Eyff TF, Azeredo-Da-Silva ALF, Bersch VP, Osvaldt AB. A systematic review and meta-analysis of the aetiology of acute pancreatitis. *HPB (Oxford)* 2019;21:259–67.
- Gliem N, Ammer-Herrmann C, Ellenrieder V, Neesse A. Management of severe acute pancreatitis: an update. *Digestion* 2021;102:503–7.
- Lankisch PG, Weber-Dany B, Hebel K, Maisonneuve P, Lowenfels AB. The harmless acute pancreatitis score: a clinical algorithm for rapid initial stratification of nonsevere disease. *Clin Gastroenterol Hepatol* 2009;7:702–5.
- Singh VK, Wu BU, Bollen TL, Repas K, Maurer R, Johannes RS, et al. A prospective evaluation of the bedside index for severity in acute pancreatitis score in assessing mortality and intermediate markers of severity in acute pancreatitis. *Am J Gastroenterol* 2009;104:966–71.
- Brown A, Baillargeon JD, Hughes MD, Banks PA. Can fluid resuscitation prevent pancreatic necrosis in severe acute pancreatitis? *Pancreatol* 2002;2:104–7.
- Lankisch PG, Assmus C, Lehnick D, Maisonneuve P, Lowenfels AB. Acute pancreatitis: does gender matter? *Dig Dis Sci* 2001;46:2470–4.
- Spanier BW, Dijkgraaf MG, Bruno MJ. Epidemiology, aetiology and outcome of acute and chronic pancreatitis: an update. *Best Pract Res Clin Gastroenterol* 2008;22:45–63.
- Tandoğan G, Coşkun BN, Eroğlu A, Avcı DN, Irak K, Kıyıcı M. Retrospective analysis of treatment approach of patients diagnosed with acute pancreatitis. *Sisli Etfal Hastan Tip Bul [Article in Turkish]* 2012;46:8–15.
- Roberts SE, Morrison-Rees S, John A, Williams JG, Brown TH, Samuel DG. The incidence and aetiology of acute pancreatitis across Europe. *Pancreatol* 2017;17:155–65.
- Ye C, Liu L, Ma X, Tong H, Gao J, Tai Y, et al. Obesity aggravates acute pancreatitis via damaging intestinal mucosal barrier and changing microbiota composition in rats. *Sci Rep* 2019;9:69.
- Cruz-Monserrate Z, Conwell DL, Krishna SG. The impact of obesity on gallstone disease, acute pancreatitis, and pancreatic cancer. *Gastroenterol Clin N Am* 2016;45:625–37.
- Tolstrup JS, Kristiansen L, Becker U, Gronbaek M. Smoking and risk of acute and chronic pancreatitis among women and men: a

- population-based cohort study. *Arch Intern Med* 2009;169:603–609.
24. Yadav D, Hawes RH, Brand RE, Anderson MA, Money ME, Banks PA, et al. Alcohol consumption, cigarette smoking, and the risk of recurrent acute and chronic pancreatitis. *Arch Intern Med* 2009;169:1035–45.
  25. Yang H, Wang L, Shi YH, Sui GT, Wu YF, Lu XH, et al. Risk factors of acute pancreatitis in the elderly Chinese population: a population-based cross-sectional study. *J Dig Dis* 2014;15:501–7.
  26. Thrower E. Pathologic cellular events in smoking-related pancreatitis. *Cancers (Basel)* 2015;7:723–35.
  27. Khanna AK, Meher S, Prakash S, Tiwary SK, Singh U, Srivastava A, et al. Comparison of Ranson, Glasgow, MOSS, SIRS, BISAP, APACHE-II, CTSI Scores, IL-6, CRP, and procalcitonin in predicting severity, organ failure, pancreatic necrosis, and mortality in acute pancreatitis. *HPB Surg* 2013;2013:367581.
  28. Kong W, He Y, Bao H, Zhang W, Wang X. Diagnostic value of neutrophil-lymphocyte ratio for predicting the severity of acute pancreatitis: a meta-analysis. *Dis Markers* 2020;2020:9731854.
  29. Suppiah A, Malde D, Arab T, Hamed M, Allgar V, Smith AM, et al. The prognostic value of the neutrophil-lymphocyte ratio (NLR) in acute pancreatitis: identification of an optimal NLR. *J Gastrointest Surg* 2013;17:675–81.
  30. Abaylı B, Gençdal G, Değirmencioğlu Ş. Correlation between neutrophil/lymphocyte ratio and Ranson score in acute pancreatitis. *J Clin Lab Anal* 2018;32:e22437.
  31. Li Y, Zhao Y, Feng L, Guo R. Comparison of the prognostic values of inflammation markers in patients with acute pancreatitis: a retrospective cohort study. *BMJ Open* 2017;7:e013206.
  32. Zhou H, Mei X, He X, Lan T, Guo S. Severity stratification and prognostic prediction of patients with acute pancreatitis at early phase: a retrospective study. *Medicine (Baltimore)* 2019;98:e15275.
  33. Orfanu AE, Popescu C, Leuştean A, Negru AR, Tilişcan C, Aramă V, et al. The importance of haemogram parameters in the diagnosis and prognosis of septic patients. *J Crit Care Med (Targu Mures)* 2017;3:105–10.
  34. Gasparyan AY, Ayvazyan L, Mikhailidis DP, Kitas GD. Mean platelet volume: a link between thrombosis and inflammation? *Curr Pharm Des* 2011;17:47–58.
  35. Hong W, Lin S, Zippi M, Geng W, Stock S, Basharat Z, et al. Serum albumin is independently associated with persistent organ failure in acute pancreatitis. *Can J Gastroenterol Hepatol* 2017;2017:5297143.
  36. Ammori BJ, Barclay GR, Larvin M, McMahon MJ. Hypocalcemia in patients with acute pancreatitis: a putative role for systemic endotoxin exposure. *Pancreas* 2003;26:213–7.
  37. Vatansever S, Doğru R, Pakoz ZB, Genç H, Ünsal B. Evaluation of laboratory findings and mortality in elderly patients with acute biliary pancreatitis. *Sisli Etfal Hastan Tip Bul* 2018;52:274–8.
  38. Viasus D, Garcia-Vidal C, Simonetti A, Manresa F, Dorca J, Guadiol F, et al. Prognostic value of serum albumin levels in hospitalized adults with community-acquired pneumonia. *J Infect* 2013;66:415–23.
  39. Vincent JL, Russell JA, Jacob M, Martin G, Guidet B, Wernerman J, et al. Albumin administration in the acutely ill: what is new and where next? *Crit Care* 2014;18:231. Erratum in: *Crit Care* 2014;18:630.
  40. Condon JR, Ives D, Knight MJ, Day J. The aetiology of hypocalcaemia in acute pancreatitis. *Br J Surg* 1975;62:115–8.
  41. McMahon MJ, Woodhead JS, Hayward RD. The nature of hypocalcaemia in acute pancreatitis. *Br J Surg* 1978;65:216–8.
  42. Ryzen E, Rude RK. Low intracellular magnesium in patients with acute pancreatitis and hypocalcemia. *West J Med* 1990;152:145–8.
  43. De Ritis F, Coltorti M, Giusti G. An enzymic test for the diagnosis of viral hepatitis; the transaminase serum activities. *Clin Chim Acta* 1957;2:70–4.
  44. Botros M, Sikaris KA. The de Ritis ratio: the test of time. *Clin Biochem Rev* 2013;34:117–30.