



Seminal plasma proteins and metabolites: effects on sperm function and potential as fertility markers

Arlindo Alencar Moura^{1,3}, Erdogan Memili², Antônia Moemia Rodrigues Portela¹, Arabela Guedes Viana¹, Ana Luiza Cazaux Velho¹, Maria Júlia Barbosa Bezerra¹, Fábio Róger Vasconcelos¹

¹Department of Animal Science, Federal University of Ceará, Fortaleza, CE, Brazil.

²Department of Dairy and Animal Sciences, Mississippi State University, MS, USA.

Abstract

Molecular components of sperm and in the media surrounding them influence male fertility. In this regard, seminal plasma proteins and metabolites modulate various reproductive events, including sperm motility and capacitation, cell protection, acrosome reaction, fertilization and embryonic development. Empirical associations between seminal proteins and metabolites and fertility indicate that these molecules are potential molecular markers of male reproductive status in cattle and other species.

Keywords: fertility, metabolites, proteins, seminal plasma, sperm.

Introduction

Pregnancy after artificial insemination (AI) is the best indicator of reproductive potential of sires. However, this information usually becomes available only after bulls are mature and have been selected for commercial use in the AI industry. Moreover, criteria such as sperm motility and morphology have limited associations with sire fertility, particularly in bulls selected by the AI industry (Killian *et al.*, 1993; Moura, 2005; Moura *et al.*, 2006; Oliveira *et al.*, 2013; Dogan *et al.*, 2015; Kaya and Memili, 2016). There can be substantial differences in fertility among bulls with normal semen parameters and those with non-compensable sperm defects may never achieve adequate fecundity (Oliveira *et al.*, 2013; Dogan *et al.*, 2015; Kaya and Memili, 2016). Therefore, mechanisms by which sperm molecular profiles influence bull fertility are not fully understood. In this context, there are efforts to identify molecular markers of gamete function in farm animals and humans. Candidate makers include sperm RNA, proteins and various molecules in reproductive fluids. These studies are based on the hypothesis that molecular components of sperm and/or from the surrounding media influence fertilizing capacity. In this regard, analysis of seminal plasma proteome and metabolome will provide information about mechanisms regulating sperm fertilizing potential and reproductive performance. Thus, the present review discusses the roles of selected seminal plasma proteins and metabolites and how their expression relates to fertility, especially in cattle.

Seminal plasma proteins

Proteins involved in sperm protection

Seminal plasma contains proteins that protect sperm in the epididymis (Hinton *et al.*, 1995; Kraus *et al.*, 2005), after ejaculation and in the female reproductive tract. Production of reactive oxygen species (ROS) is a component of sperm physiology (MacLeod, 1943); however, excessive ROS disturbs sperm homeostasis through formation of lipid peroxidation, reduction of enzymes that regulate calcium influx, and loss of ATP (Ohta *et al.*, 1989; Aitken *et al.*, 1993). To mitigate deleterious effects of excessive ROS, the epididymis secretes antioxidant enzymes (Hinton *et al.*, 1996), including glutathione S-transferase, tioredoxin peroxidase, superoxide dismutase, glutathione peroxidase (GSHPx) and catalase (Alvarez and Storey, 1983; Jeulin *et al.*, 1989; Fouchécourt *et al.*, 2000; Dacheux *et al.*, 2006). Of these, GSHPx catalyzes the reduction of hydrogen peroxide (Halliwell and Gutteridge, 1990), protecting sperm against excessive ROS (Perry *et al.*, 1992; Dacheux *et al.*, 2005). For example, increased GSHPx activity in ram semen is linked to maintenance of sperm viability (Casao *et al.*, 2010). Another seminal plasma molecule, acidic seminal fluid protein (aSFP), also controls oxidative stress in the bovine reproductive tract (Einspanier *et al.*, 1993; Schöneck *et al.*, 1996). This protein shares identity with molecules of the spermadhesin family (Romão *et al.*, 1997) and, in the bull, is secreted by the epididymis and accessory sex glands (Moura *et al.*, 2007a, 2010). Although aSFP binds to ejaculated sperm, it is lost after capacitation (Dostálová *et al.*, 1994). Therefore, unlike porcine spermadhesins (Caballero *et al.*, 2004, 2005), it appears that bovine aSFP does not participate in sperm-oocyte interactions. However, aSFP has been associated with survival of cryopreserved bull sperm (Jobim *et al.*, 2004).

Ion chelators in seminal plasma, such as lactoferrin (LF), also protect sperm against effects of lipid peroxidation (Ochsendorf, 1999). Lactoferrin sequesters ionic iron (Nozaki *et al.*, 2003) and adsorption to sperm during epididymal transit (Jin *et al.*, 1997) and ejaculation (Thaler *et al.*, 1990). In stallions, LF represents 41.2% of all proteins secreted by the epididymis (Fouchécourt *et al.*, 2000) and high

³Corresponding author: arlindo.moura@gmail.com

Received: February 15, 2018

Accepted: May 17, 2018



concentrations of LF in horse and dog seminal plasma relate to total number of sperm (Kikuchi *et al.*, 2003a, b). Seminal albumin, in turn, binds to lipid peroxides, contributing to sperm protection (Alvarez and Storey, 1983) and is positively correlated with percentage of morphologically normal sperm in bovine semen (Elzanaty *et al.*, 2007).

Clusterin, another seminal plasma molecule with protective roles, acts as a chaperone (Humphreys *et al.*, 1999) and inhibits cell lysis by complement-mediated mechanisms present in female secretions (Ibrahim *et al.*, 1999; Meri and Jarva, 2001). In the epididymis, clusterin affects maturation, lipid transport (Tenniswood *et al.*, 1992) and sperm membrane remodeling (Humphreys *et al.*, 1999). Clusterin chaperone activity is consistent with its ability to interact with various types of proteins *in vivo* (Carver *et al.*; 2003). *In silico* analysis of clusterin networking indicates potential interactions with proteases and protease inhibitors, such as plasminogen, alpha-2-macroglobulin, TIMP-1, alpha-2-antiplasmin precursor and plasminogen activator inhibitor 1. Clusterin also has putative links to fibronectins, which participate in cell adhesion, wound healing and maintenance of cell structure, including GTP protein-coupled receptors and modulators of cell growth. Seminal plasma clusterin is inversely related to percentage of sperm with intact membrane in peccaries (*Peccari tajacu* L.; Santos *et al.*, 2014) and with percentage of morphologically normal sperm in semen from Brahman bulls (Boe-Hansen *et al.*, 2015). In contrast, bull and ram sperm with morphologic defects have extensive clusterin binding (Ibrahim *et al.*, 2001a, b). This association probably occurs as a result of clusterin's ability to bind to damaged portions of hydrophobic regions of sperm membranes (Bailey and Griswold, 1999). Sertoli cell-secreted clusterin prevented apoptosis in rat testes subjected to hyperthermia (Matsushita *et al.*, 2016) and in humans, clusterin secreted in the fluid of the seminiferous epithelium has positive associations with fertility (Salehi *et al.*, 2013). In addition, seminal clusterin promotes immune tolerance to male antigens in humans, mitigating female immune reactions to male factors (Merlotti *et al.*, 2015). High levels of clusterin are associated with advanced physiopathological states, such as kidney diseases, neurodegenerative disorders, arteriosclerosis, heart attack and cancer (Wehrli *et al.*, 2001; Trougakos *et al.*, 2002; Pucci *et al.*, 2004; Calero *et al.*, 2005). We have characterized the seminal plasma proteome of several domestic and wild species, including bulls (Moura *et al.*, 2007a; Rego *et al.*, 2014, 2016; Menezes *et al.*, 2017), rams (Souza *et al.*, 2012), boars (González-Cadavid *et al.*, 2014), peccaries (Santos *et al.*, 2014), dogs (Aquino-Cortez *et al.*, 2017) and coatis (Silva *et al.*, 2018), among others. Clusterin is present in the semen of all these species, in moderate to high concentrations. Thus, many animals have a clusterin-based, conserved mechanism for sperm protection and regulation of immune reactions initiated by male gametes in the female reproductive tract.

Proteins associated with sperm motility

Seminal plasma contains various proteins associated with sperm motility (Baas *et al.*, 1983), e.g., kallikrein-cinins. In this case, kininogenin seminal plasma is a specific substrate for kalikrein (Fink *et al.*, 1989), an important stimulator of post-ejaculation sperm motility (Schill *et al.*, 1989). There is a positive correlation between seminal plasma kallikrein activity and sperm motility, with exogenous kallikrein enhancing bovine sperm motility (Somlev *et al.*, 1996). Angiotensin converting enzyme (ACE) is another seminal plasma component related to the kalikrein system (Hohlbrugger *et al.*, 1984). This enzyme catalyzes formation of angiotensin II and binds to receptors on sperm, enhancing motility (Vinson *et al.*, 1996). Furthermore, ACE activity in ram seminal plasma is positively correlated with sperm concentration and fertility (Métayer *et al.*, 2001; Gatti *et al.*, 2004). In contrast, inhibition of ACE activity in bovine seminal plasma decreases progressive motility and inhibits acrosome reaction after *in vitro* capacitation (Costa and Thundathil, 2012).

Proteins involved in sperm capacitation, acrosome reaction and fertilization

Phospholipid binding proteins belonging to the family of BSPs (Binder of Sperm Proteins) are present in seminal plasma of several species, including bulls, bucks (male goats and rabbits), rams, rodents, stallions and men (Moura *et al.*, 2007a; Manjunath *et al.*, 2009; Souza *et al.*, 2012; Plante *et al.*, 2016). BSPs comprise ~60% of all proteins of the accessory sex gland fluid (Moura *et al.*, 2007a) and seminal plasma (Manjunath and Sairam, 1987) of *Bos taurus* bulls and nearly the same amount in seminal plasma of *Bos indicus* bulls (Rego *et al.*, 2014). In cattle, BSP proteins are secreted as isoforms with 14-15 kDa (BSP1 and BSP3) and 30 kDa (BSP5). Both BSP1 and BSP5 have two fibronectin type II domains arranged in tandem and amino terminal extensions that are O-glycosylated at threonine residues. Such biochemical attributes allow BSP1 and BSP5 to interact with sperm and to modulate ligand-binding activities by similar mechanisms (Calvete *et al.*, 1996; Manjunath *et al.*, 2009), with functional similarities (Manjunath and Thérien, 2002). Bovine BSPs are typical accessory sex gland proteins (Manjunath and Thérien, 2002; Moura *et al.*, 2007a, 2010). BSPs bind to bull sperm at ejaculation (Manjunath and Thérien, 2002) and remain there after sperm contact oviductal secretions *in vitro* (Souza *et al.*, 2008), as well as in acrosome-reacted or frozen-thawed sperm (Rodríguez-Villamil *et al.*, 2016). BSPs also mediate the interaction between sperm and the oviduct epithelium (Gwathmey *et al.*, 2006; Suarez, 2016). The most studied role of BSPs is their ability to bind and remove phospholipids and cholesterol from the sperm membrane, an initial event of capacitation (Thérien *et al.*, 1999). Capacitating



effects of BSPs have been reported in other species, including mice (Plante and Manjunath, 2015) and humans (Plante *et al.*, 2014). However, while ruminant BSPs originate mainly from the accessory sex glands (Manjunath *et al.*, 2009; Souza *et al.*, 2012; van Tilburg *et al.*, 2014), they are expressed in the epididymis of mice and humans.

In addition to sperm capacitation, BSP1 affects *in vitro* fertilization and embryonic development. The study conducted by Rodriguez-Villamil *et al.* (2016) evaluated cumulus-oocyte complexes (COCs) incubated with frozen-thawed ejaculated sperm (18 h) in Fert-TALP medium containing: heparin and BSP1. With ejaculated sperm, cleavage rates were similar when Fert-TALP medium was incubated with heparin, 10 or 20 µg/ml BSP1. Day-7 blastocyst rates were equivalent after incubations with heparin or 10 µg/ml BSP1, but there were marked reductions in blastocyst formation after IVF media were supplemented with 20 or 40 µg/ml. Therefore, BSP1 is as efficient as heparin for inducing capacitation and fertilizing capacity of frozen-thawed ejaculated sperm. However, damage caused to embryo development may have been caused by BSP1 itself. High concentrations of and/or prolonged exposure to BSP proteins are harmful to cryopreserved sperm due to membrane destabilization and excessive phospholipid and cholesterol efflux (Thérien *et al.*, 1995, 1998; Manjunath and Thérien, 2002). Furthermore, content of BSP5 in accessory sex gland fluid has a quadratic association with bull fertility (Moura *et al.*, 2006), suggesting that excessive BSP was detrimental to sperm physiology and/or embryo development.

In the same study (Rodriguez-Villamil *et al.*, 2016), cleavage rates were higher after IVF using cauda epididymal sperm and any BSP1 concentration (10, 20 or 40 µg/ml) embryo development (day 8) was greater after inclusion of 20 or 40 µg/ml BSP1 in the IVF media, with or without heparin. Thus, we concluded that: 1) heparin has limited effect on cauda epididymal sperm *in vitro* (based on fertilization rates and blastocyst formation); 2) BSP1 has better effects on embryo growth than heparin; and 3) a combination of BSP1 with heparin does not enhance cleavage rates and embryo development beyond those obtained with BSP1. We also verified that SP1 did not cause reductions in bovine blastocyst growth after IVF with epididymal sperm, in contrast to results obtained with ejaculated sperm. Therefore, previous exposure of sperm or not to seminal plasma determines how sperm will respond to BSP *in vitro*. Additionally, combining heparin and BSP1 did not increase capacitation rates of ejaculated sperm. And both cleavage rates and blastocyst growth were similar after ejaculated sperm were incubated with heparin, BSP-1+heparin or BSP-1. With epididymal sperm, the best results or capacitation and blastocyst growth were obtained with BSP-1, when compared to heparin and heparin+BSP-1 (Rodriguez-Villamil *et al.*, 2018; Federal University of Ceara, Fortaleza, Brazil; unpublished data). Thus, BSP-1 is a potent capacitating

factor for bovine sperm and it increases fertilization rates, with no dependence on heparin.

Despite multiple beneficial roles of BSPs, these molecules can damage sperm during cryostorage, as they extract phospholipids and cholesterol from the membrane in a concentration- and time-dependent manner (Manjunath *et al.*, 2002; Plante *et al.*, 2016). Such deleterious effects occur when sperm are exposed for prolonged periods-and/or to excessive concentrations of BSPs. In this regard, extenders used for sperm preservation, such as egg-yolk (EY) and milk, contain components that associate with BSPs (Manjunath *et al.*, 2002). There are interactions between low-density lipoproteins in EY (Bergeron and Manjunath, 2006) or milk proteins; the latter can prevent excessive BSP binding to sperm and excessive phospholipid removal from the membrane, thereby protecting sperm during cryopreservation (Plante *et al.*, 2015). In goats, milk proteins (casein and β-lactoglobulin) bind to BSPs and reduces BSP interactions with sperm (Menezes *et al.*, 2016). Currently, BSPs are one of the most studied mammalian seminal plasma proteins and effects on ejaculated sperm, including capacitation, interaction with the oviduct epithelium and fertilization. That BSPs interact with components of semen extenders suggest that these proteins are potential targets for development of biomolecules that could enhance assisted reproductive technologies.

Seminal plasma phospholipase A2 (PLA2) participates in capacitation, acrosome reaction and sperm-oocyte membrane fusion (Soubeyrand *et al.*, 1997; Pietrobon *et al.*, 2005; Roldan and Shi, 2007), promotes release of fatty acids and phospholipids involved in final stages of gamete fusion (Roldan, 1998) and has antimicrobial effects. Furthermore, its expression in bovine seminal plasma is associated with fertility (Moura *et al.*, 2006). Osteopontin (OPN) concentrations in bovine seminal plasma were related to *in vivo* fertility of Holstein bulls (Killian *et al.*, 1993; Moura *et al.*, 2006) and to fertilizing capacity of cauda epididymal sperm treated with accessory sex gland fluid in IVF trials (Henault *et al.*, 1995; Moura *et al.*, 2007b). OPN is mainly secreted by the accessory sex glands and binds to sperm after ejaculation and after they contact secretions of the oviduct and are capacitated (Souza *et al.*, 2008). Also, OPN has a calcium binding site and a domain to link with heparin, consistent with its effects on sperm capacitation (Monaco *et al.*, 2009; Boccia *et al.*, 2013).

Alterations in the OPN gene reduce seminal plasma OPN concentration (Rorie *et al.*, 2016). Furthermore, in IVF studies, percentage of fertilized bovine oocytes was significantly reduced by addition of OPN antibodies to fertilization media and exposure of sperm or oocytes to antibodies against alpha V and alpha5 integrins before fertilization (Gonçalves *et al.*, 2007). Also, pre-treatment of bovine sperm and oocytes with OPN enhances both *in vitro* fertilization and early embryo development (Gonçalves *et al.*, 2008a, b). The



RGD amino acid sequence of osteopontin mediates its link with $\alpha 5$ and αv integrins (Denhardt, 2002; Wai and Kuo, 2004) and the ability of osteopontin to support cell adhesion is prevented when the RGD sequence is mutated (Liaw *et al.*, 1995; Xuan *et al.*, 1995). Treatment of sperm or oocytes with an RGD peptide, but not with an RGE sequence, reduced both the number of sperm bound to the zona pellucida and fertilization rates, similar to effects of anti-osteopontin antibodies. It appears that OPN interacts with sperm through integrins (Gonçalves *et al.*, 2007). Incubation of oocytes with osteopontin purified from bovine milk increased cleavage rates on day 4, blastocyst development on day 8 and hatched blastocysts on day 11 (Gonçalves *et al.*, 2007). Furthermore, OPN purified from milk improved sperm capacitation and addition of OPN to IVF media enhanced bovine blastocyst formation (Monaco *et al.*, 2009). Moreover, in an IVF system, using frozen-thawed bull semen, OPN improved fertilization rates and blastocyst development on day 8 (Gonçalves *et al.*, 2008a). In swine, supplementation of fertilization media with recombinant rat OPN enhanced fertilization rates by 41% and reduced polyspermy (Hao *et al.*, 2006). Exogenous OPN added to IVF media improved cleavage rates and swine embryo development, and inhibited apoptosis and DNA fragmentation (Hao *et al.*, 2008). Moreover, anti-OPN antibodies decreased rates of *in vitro* fertilization and blastocyst growth in mice (Liu *et al.*, 2015). Clearly, OPN affects fertilization and post-fertilization events.

Osteopontin is typically involved in cell adhesion, tissue and extracellular remodeling, inflammation and immune-mediated events (Denhardt, 2002; Wai and Kuo, 2004; Rittling and Singh, 2015; Boulefour *et al.*, 2016). Despite substantial knowledge regarding actions of osteopontin in several tissues, an understanding of its function in male reproduction is far from complete. There is general consensus that OPN secreted by the accessory sex glands binds to sperm during ejaculation through integrins and that the integrin-OPN complex interacts with the zona pellucida (D'Cruz, 1996). This model is supported by the presence of OPN in bovine oviductal fluid (Gabler *et al.*, 2003). Additionally, OPN binds to the CD44 receptor, which usually participates in cell adhesion (Cichy and Puré, 2003), and has been expressed in sperm (Bains *et al.*, 2002) and oocyte membranes (Schoenfelder and Einspanier, 2003). In the bull, OPN binds to the acrosome at ejaculation (Cancel *et al.*, 1999) and this sperm-OPN link is preserved after sperm contacts with oviductal fluid and undergoes an acrosome reaction *in vitro* (Souza *et al.*, 2008). In addition to sperm binding, OPN interacts with the zona pellucida and oolemma of bovine oocytes (Souza *et al.*, 2008). Consequently, we propose that OPN adheres to sperm and this complex connects to the zona pellucida or to OPN-zona pellucida, as OPN can form high-affinity bonds with other OPN molecules (Kaartinen *et al.*, 1999;

Goldsmith *et al.*, 2002). When entering the perivitelline space, OPN attached to the post-equatorial segment would mediate the interaction of sperm and oolema, also through integrins and/or CD44. Integrins (αv and $\alpha 5$) are present in bovine (Erikson *et al.*, 2008) and human sperm (Fusi *et al.*, 1996; Reddy *et al.*, 2003), as well as on human oolema (D'Cruz, 1996) and CD44 transmembrane glycoproteins are present in bovine sperm and oocytes. Interactions of sperm OPN with oocyte integrins and CD44 receptors could trigger intracellular signaling, as reported for other cell types (Wai and Kuo, 2004; Rangaswami *et al.*, 2006), and affect fertilization and early embryo development.

Metabolomics

Metabolites are the result of metabolic reactions associated with various biochemical pathways (Dunn *et al.* 2011). Many of these molecules have important roles in biological processes and represent potential biomarkers for predicting or detecting developmental states, physiological events, diseases or specific phenotypes. Therefore, metabolomics is used to understand networks of metabolites and have provided comprehensive identification and quantification of small molecules, including amino acids, peptides, vitamins, minerals, lipids, and carbohydrates in diverse cells, tissues, fluids, organs and organisms (Oliver *et al.*, 1998; Fiehn 2001, 2002; Dunn *et al.*, 2005).

Substantial progress has been made in the study and development of methodological strategies for using metabolomics, metabonomics, metabolic-fingerprinting, metabolite targeting, and metabolic profiling. Metabonomics is used to measure differences in levels of metabolites resulting from such factors as pathological or genetic changes, toxins, or use of drugs. Metabolic fingerprinting is a rapid method to evaluate and classify biologic samples or biopsies. Furthermore, metabolite target analysis is used to identify specific metabolic pathways of a limited number of metabolites, whereas metabolic profiling evaluates a cluster of metabolites that participate in a specific metabolic pathway (Dunn and Ellis, 2005; Hollywood *et al.*, 2006; Holmes *et al.*, 2008; Dunn *et al.*, 2011; Patti *et al.*, 2010). Metabolomics can be performed associated with other omics approaches, e.g. genomics, transcriptomics, and proteomics. For example, an Integrative Personal Omics Profile (iPOP) to identify markers for possible diseases affecting an individual could lead to an early diagnosis and perhaps prevention of certain diseases (Chen *et al.*, 2012). Although complementary to other omics, metabolomics provides identification and quantification of products from metabolism and its pathways, analyses of modifications in metabolic reactions, and characterization of phenotypes and identification of potential biomarkers for such phenotypes (Fiehn, 2001, 2002; Dunn *et al.*, 2005; Goodacre *et al.*, 2004; Hollywood *et al.*, 2006; Patti *et al.*, 2010).



Metabolomics and reproductive biology

Advanced and more sensitive methods are vital for addressing major questions in biology and biotechnology, including those related to assisted reproductive technologies (ART). It is well known that metabolites have critical roles in specific pathways related to fertilization, implantation and embryonic development. Some techniques used in metabolome analysis for studies of reproductive biology include proton nuclear magnetic resonance (¹H NMR), mass spectrometry (MS), fourier transform infrared spectroscopy (FTIR), near infrared (NIR) and Raman (Singh and Sinclair, 2007, Seli *et al.*, 2010a; Kovac *et al.*, 2013; Muñoz *et al.*, 2014a; b). Metabolomics methods have been used as noninvasive approaches to improve assessment of embryo quality (Singh and Sinclair, 2007; Bromer and Seli, 2008; Nagy *et al.*, 2008; Seli *et al.*, 2010a; Montag *et al.*, 2013). For example, ¹H NMR scans compared metabolomes in the culture media for human embryos before transfer. In that study, glutamate was associated with subsequent developmental potential (Seli *et al.*, 2008). In addition, NIR, Raman and ¹H NMR used for metabolome analysis of human embryo culture media were valuable for predicting successful implantation and pregnancy after IVF (Seli *et al.*, 2007, 2008, 2010b). FTIR metabolomics were used to determine gender of bovine embryos (Muñoz *et al.*, 2014a, b). This is also an effective and non-invasive method to determine embryo viability and the metabolic profile of blood plasma from recipient cows. Further, FTIR can be used to identify superior embryos and recipient females for optimum pregnancy outcome (Muñoz *et al.*, 2014b).

As indicated above, methods of conventional semen evaluation most often give only descriptive information and have limitations to predict in fertility. However, various molecular approaches, such as metabolomics, have provided more in depth understanding of mechanisms causing male infertility (Deepinder *et al.* 2007; Aitken, 2010). Metabolomics has promise in identifying potential biomarkers of male fertility and infertility (Gilany *et al.*, 2014; Goodacre *et al.*, 2004; Deepinder *et al.*, 2007; Kovac *et al.*, 2013). The presence or changes in specific metabolites could be related to male gamete functions, perhaps enabling evidence-based techniques to prevent or mitigate infertility (Aitken, 2010). Metabolomics approach using Raman spectroscopy to analyze human seminal plasma facilitated diagnosis of normospermic and asthenozoospermic men (Gilany *et al.*, 2014). Furthermore, ¹H NMR identified fertility-associated biomarkers in seminal plasma and serum of high- and low-fertility bulls. Metabolites, such as citrate,

tryptamine, taurine, and leucine were identified in seminal plasma, whereas asparagine, glycogen, citrulline, and isoleucine were present in serum (Kumar *et al.*, 2015). Using ¹H NMR, Hamamah *et al.* (1998) detected increased choline/citrate, choline/lactate, and glycerophosphorylcholine/choline ratios in seminal plasma of men afflicted with spermatogenic failure versus those with obstructive azoospermia. Several small molecular markers were identified in the urine of men with normozoospermic infertility using liquid chromatography-mass spectrometry (LC-MS) in combination with bioinformatics and multivariate analyses. In this research, leukotriene E₄, 3-hydroxypalmitoylcarnitine, aspartate, xanthosine, and methoxytryptophan were biomarkers of infertility (Zhang *et al.*, 2014). Clearly, metabolomics can be used to identify molecular markers of male fertility.

Conclusions

In recent decades, methods in proteomics and metabolomics have enabled detection of unprecedented numbers of molecules in the seminal plasma of farm animals, wild species and humans. This broadens our knowledge regarding roles of these molecules and their contributions to male fertility. Metabolomics can identify numerous classes of substances associated with metabolic pathways, leading to challenges in interpretation. Empirical associations exist between specific seminal proteins (Table 1), metabolites and fertility indexes. Experiments also confirm cause and effect relations between seminal plasma proteins (e.g. osteopontin and BSPs) and IVF and early embryo development, suggesting that seminal proteins have potential in animal biotechnology.

Studies to describe components of the seminal plasma are vital to construct comprehensive libraries of seminal plasma compounds. As many as 4,000 proteins have already been identified in human seminal plasma, although there may be up to 10,000 present (Gilany *et al.*, 2014). A human proteome atlas (<https://www.proteinatlas.org/>) of human tissues and organs is under development (Omenn *et al.*, 2017; Uhlén *et al.*, 2015; Thul *et al.*, 2017). Fundamental research sets the foundation of science and technology. However, investigations that use omics approaches and reproduction need to be focused on finding markers of traits that are important for livestock industry in different regions of the world. Lessons could be learned from translational research, where research is carried out by multidisciplinary teams, joining efforts from basic science, applied investigators and professionals in the front of technology.



Table 1. Functional groups and mechanism of actions of major seminal plasma proteins.

Major functional group	Protein	Mechanism of action	Major references
Proteins involved in sperm protection	Glutathione peroxidase	Catalyzes reduction of H ₂ O ₂ , protects sperm against excessive ROS.	(Halliwell and Gutteridge, 1990; rams); (Perry <i>et al.</i> , 1992; Dacheux <i>et al.</i> , 2005; bulls)
	Acidic seminal fluid protein	Sperm decapacitation, oxidative stress control, survival of cryopreserved sperm	(Einspanier <i>et al.</i> , 1993; Schöneck <i>et al.</i> , 1996; bull); (Dostálová <i>et al.</i> , 1994; bull); (Jobim <i>et al.</i> , 2004; bull)
	Lactoferrin	Ion chelators, protects sperm against effects of lipid peroxidation	(Ochsendorf, 1999; men)
	Albumin	Binds to lipid peroxides in sperm membrane, sperm protection	(Alvarez and Storey, 1983; rabbit)
Proteins associated with sperm motility	Clusterin	Chaperone and sperm membrane remodeling, protects against female reproductive tract immune response, binds to damage sperm membrane	(Humphreys <i>et al.</i> , 1999; men); (Ibrahim <i>et al.</i> , 1999; bull; Merlotti <i>et al.</i> , 2015; men); (Bailey and Griswold, 1999; rat)
	Kallikrein-cinins	Substrate for kallikrein, which enhance sperm motility	(Somlev <i>et al.</i> , 1996; bull)
Proteins involved in sperm capacitation, acrosome reaction and fertilization	Angiotensin converting enzyme	Catalyzes angiotensin II formation, Enhance motility.	(Vinson <i>et al.</i> , 1996; rat and men; Costa and Thundathil, 2012; bull)
	Binder of sperm proteins	Bind to sperm at ejaculation; mediate gametes interaction, phospholipids and cholesterol efflux from sperm membrane; enhance <i>in vitro</i> fertilization and embryonic development.	(Manjunath and Thérien, 2002; bull); (Gwathmey <i>et al.</i> , 2006; bull); (Thérien <i>et al.</i> , 1999; bull) (Rodriguez-Villamil <i>et al.</i> , 2016; bull); Manjunath <i>et al.</i> , 2002; bull; Plante <i>et al.</i> , 2016; bull)
	Phospholipase A2	Release of fatty acids and phospholipids involved in final stages of gamete fusion; antimicrobial effects.	(Soubeyrand <i>et al.</i> , 1997; bull; Pietrobon <i>et al.</i> , 2005; mouse; Roldan and Shi, 2007); Moura <i>et al.</i> , 2006; bull)
	Osteopontin	Binds to sperm acrosome after ejaculation; sperm capacitation; interacts with sperm through integrins, interacts with zona pellucida and oolemma; binds to sperm CD44 receptor; enhances <i>in vitro</i> fertilization and embryo development; reduces polyspermy and inhibits apoptosis and DNA fragmentation.	(Souza <i>et al.</i> , 2008; bull; Cancel <i>et al.</i> , 1999; bull); (Boccia <i>et al.</i> , 2013; buffalo; Monaco <i>et al.</i> , 2009; bull); (Gonçalves <i>et al.</i> , 2007; bull; D'Cruz, 1996; men); (Souza <i>et al.</i> , 2008; bull); (Cichy and Puré, 2003; men); (Gonçalves <i>et al.</i> , 2008a; bull); (Hao <i>et al.</i> , 2006; swine; Hao <i>et al.</i> , 2008; swine)

Acknowledgments

The present work has been supported by the Brazilian Federal Research Councils (CAPES and CNPq) and the Ceara State Foundation for Scientific Development and Research (FUNCAP).

References

- Aitken RJ, Harkiss D, Buckingham D. 1993. Relationship between iron catalysed lipid peroxidation potential and human sperm function. *J Reprod Fertil*, 98:257-265.
- Aitken RJ. 2010. Whither must spermatozoa wander? The future of laboratory seminology. *Asian J Androl* 12:99-103.
- Alvarez JG, Storey BT. 1983. Role of superoxide dismutase in protecting rabbit spermatozoa from O₂ toxicity due to lipid peroxidation. *Biol Reprod*, 28:1129-1136.
- Aquino-Cortez A, Pinheiro BQ, Lima DBC, Silva HVR, Mota Filho AC, Martins JAM, Rodriguez-



- Villamil P, Moura AA, Silva LDM. 2017. Proteomic characterization of canine seminal plasma. *Theriogenology*, 95:178-186.
- Baas JW, Molan PC, Shannon P. 1983. Factors in seminal plasma of bulls that affect the viability and motility of spermatozoa. *J Reprod Fertil*, 68:275-280.
- Bailey R, Griswold MD. 1999. Clusterin in the male reproductive system: localization and possible function. *Mol Cell Endocrinol*, 151:17-23.
- Bains R, Adeghe J, Carson J. 2002. Human sperm cells express CD44. *Fertil Steril*, 78:307-312.
- Bergeron A, Manjunath P. 2006. New insights towards understanding the mechanisms of sperm protection by egg yolk and milk. *Mol Reprod Dev*, 73:1338-1344.
- Boccia L, Di Francesco S, Neglia G, De Blasi M, Longobardi V, Campanil G, Gasparrini B. 2013. Osteopontin improves spermiacapacitation and in vitro fertilization efficiency in buffalo (*Bubalus bubalis*). *Theriogenology*, 80:212-217.
- Boe-Hansen GB, Rego JP, Crisp JM, Moura AA, Nouwens AS, Li Y, Venus B, Burns BM, McGowan MR. 2015. Seminal plasma proteins and their relationship with percentage of morphologically normal sperm in 2-year-old Brahman (*Bos indicus*) bulls. *Anim Reprod Sci*, 162:20-30.
- Boulefour W, Juignet L, Bouet G, Granito RN, Vanden-Bossche A, Laroche N, Aubin JE, Lafage-Proust MH, Vico L, Malaval L. 2016. The role of the SIBLING, Bone Sialoprotein in skeletal biology: contribution of mouse experimental genetics. *Matrix Biol*. 52/54:60-77.
- Bromer JG, Seli E. 2008. Assessment of embryo viability in assisted reproductive technology: shortcomings of current approaches and the emerging role of metabolomics. *Curr Opin Obstet Gynecol*, 20:234-241.
- Caballero I, Vazquez JM, Gil MA, Calvete JJ, Roca J, Sanz L, Parrilla I, Garcia EM, Rodriguez-Martinez H, Martinez EA. 2004. Does seminal plasma PSP-I/PSP-II spermadhesin modulate the ability of boar spermatozoa to penetrate homologous oocytes in vitro? *J Androl*, 25:1004-1012.
- Caballero I, Vazquez JM, Rodriguez-Martinez H, Gill MA, Calvete JJ, Sanz L, Garcia EM, Roca J, Martinez EA. 2005. Influence of seminal plasma PSP-I/PSP-II spermadhesin on pig gamete interaction. *Zygote*, 13:11-16.
- Calero M, Rostagno A, Frangione B, Ghiso J. 2005. Clusterin and Alzheimer's disease. *Subcell Biochem*; 38:273-298.
- Calvete JJ, Mann K, Sanz L, Raida M, Töpfer-Petersen E. 1996. The primary structure of BSP-30K, a major lipid-, gelatin-, and heparin-binding glycoprotein of bovine seminal plasma. *FEBS Lett*, 399:147-152.
- Cancel AM, Chapman DA, Killian GJ. 1999. Osteopontin localization in the Holstein bull reproductive tract. *Biol Reprod*, 60:454-460.
- Carver JA, Rekas A, Thorn DC, Wilson MR. 2003. Small heat-shock proteins and clusterin: intra- and extracellular molecular chaperones with a common mechanism of action and function? *IUBMB Life*, 55:661-668.
- Casao A, Cebrián I, Asumpção ME, Pérez-Pé R, Abecia JA, Forcada F, Cebrián-Pérez JA, Muño-Blanco T. 2010. Seasonal variations of melatonin in ram seminal plasma are correlated to those of testosterone and antioxidant enzymes. *Reprod Biol Endocrinol*, 11:59-66.
- Chen R, Mias GI, Li-Pook-Than J, Jiang LH, Lam HYK, Chen R, Miriami E, Karczewski KJ, Hariharan M, Dewey FE, Cheng Y, Clark M, Im H, Habegger L, Balasubramanian S, O'Huallachain M, Dudley JT, Hillenmeyer S, Haraksingh R, Sharon D, Euskirchen G, Lacroute P, Bettinger K, Boyle AP, Kasowski M, Grubert F, Seki S, Garcia M, Whirl-Carrillo M, Gallardo M, Blasco MA, Greenberg PL, Snyder P, Klein TE, Altman RB, Butte AJ, Ashley EA, Gerstein M, Nadeau KC, Tang H, Snyder M. 2012. Personal omics profiling reveals dynamic molecular and medical phenotypes. *Cell*, 148:1293-1307.
- Cichy J, Puré E. 2003. The liberation of CD44. *J Cell Biol*, 161:839-843.
- Costa DS, Thundathil JC. 2012. Characterization and activity of angiotensin-converting enzyme in Holstein semen. *Anim Reprod Sci*, 133:35-42.
- D'Cruz OJ. 1996. Adhesion molecules in human sperm-oocyte interaction: relevance to infertility. *Front Biosci*, 1:161-176.
- Dacheux JL, Castella S, Gatti JL, Dacheux F. 2005. Epididymal cell secretory activities and the role of proteins in boar sperm maturation. *Theriogenology*, 63:319-341.
- Dacheux JL, Belghazi M, Lanson Y, Dacheux F. 2006. Human epididymal secretome and proteome. *Mol Cell Endocrinol*, 250:36-42.
- Deepinder F, Chowdary HT, Agarwal A. 2007. Role of metabolomic analysis of biomarkers in the management of male infertility. *Expert Rev Mol Diagn*, 7:351-358.
- Denhardt DT. 2002. The third international conference on osteopontin and related proteins, San Antonio, Texas, May 10-12, 2002. *Calcif Tissue Int*, 74:213-219.
- Dogan S, Vargovic P, Oliveira R, Belser LE, Kaya A, Moura A, Sutovsky P, Parrish J, Topper E, Memili E. 2015. Sperm protamine-status correlates to the fertility of breeding bulls. *Biol Reprod*, 92:1-9. doi: 10.1095/biolreprod.114.124255.
- Dostálová Z, Calvete JJ, Sanz L, Hettel C, Riedel D, Schöneck C, Einspanier R, Töpfer-Petersen E. 1994. Immunolocalization and quantitation of acidic seminal fluid protein (aSFP) in ejaculated, swim-up, and capacitated bull spermatozoa. *Biol Chem Hoppe Seyler*, 375:457-461.
- Dunn B, Ellis DI. 2005. Metabolomics: current analytical platforms and methodologies. *TrAC-Trend Anal Chem*, 24:285-294.
- Dunn WB, Bailey NJC, Johnson HE. 2005. Measuring the metabolome: current analytical technologies. *Analyst*, 130:606-625.
- Dunn WB, Broadhurst DI, Atherton HJ, Goodacre R, Griffin JL. 2011. Systems level studies of mammalian metabolomes: the roles of mass



- spectrometry and nuclear magnetic resonance spectroscopy. *Chem Soc Rev*, 40:387-426.
- Einspanier R, Amselgruber W, Sinowatz F, Henle T, Ropke R, Schams D.** 1993. Localization and concentration of a new bioactive acetic seminal fluid protein (asfp) in bulls (*Bos taurus*). *J Reprod Fertil*, 98:241-244.
- Elzanaty S, Erenpreiss J, Becker C.** 2007. Seminal plasma albumin: origin and relation to the male reproductive parameters. *Andrologia*, 39:60-65.
- Erikson DW, Way AL, Chapman DA, Killian GJ.** 2008. Detection of osteopontin on Holstein bull spermatozoa, in cauda epididymal fluid and testis homogenates, and its potential role in bovine fertilization. *Reproduction*, 133:909-917.
- Fiehn O.** 2001. Combining genomics, metabolome analysis, and biochemical modelling to understand metabolic networks. *Compar Funct Genom*. 2:155-168.
- Fiehn O.** 2002. Metabolomics-the link between genotypes and phenotypes. *Plant Mol Biol*. 48:155-171.
- Fink E, Schill WB, Miska W.** 1989. Kinin-containing kininogen is present in human seminal plasma. *Adv Exp Med Biol*, 247:311-315.
- Fouchécourt S, Métayer S, Locatelli A, Dacheux F, Dacheux JL.** 2000. Stallion epididymal fluid proteome: qualitative and quantitative characterization; secretion and dynamic changes of major proteins. *Biol Reprod*, 62:1790-1803.
- Fusi FM, Tamburini C, Mangili F, Montesano M, Ferrari A, Bronson R.** 1996. The expression of alpha v, alpha 5, beta 1 and beta 3 integrin chains on ejaculated human spermatozoa varies with their functional state. *Mol Hum Reprod*, 2:169-175.
- Gabler C, Chapman DA, Killian GJ.** 2003. Expression and presence of osteopontin and integrins in the bovine oviduct during the estrous cycle. *Reproduction*, 126:721-729.
- Gatti J, Castella S, Dacheux F, Ecroyd H, Métayer S, Thimon V, Dacheux J.** 2004. Post-testicular sperm environment and fertility. *Anim Reprod Sci*, 82:321-339.
- Gilany K, Moazeni-Pourasil RS, Jafarzadeh N, Savadi-Shiraz E.** 2014. Metabolomics fingerprinting of the human seminal plasma of asthenozoospermic patients. *Mol Reprod Dev*. 81:84-86.
- Goldsmith HL, Labrosse JM, McIntosh FA, MaenpaaPH, Kaartinen MX, McKee MD.** 2002. Homotypic interactions of soluble and immobilized osteopontin. *Ann Biomed Eng*, 30:840-850.
- Gonçalves RF, Wolinetz CD, Killian GJ.** 2007. Influence of arginine-glycine-aspartic acid (RGD), integrins (αv and $\alpha 5$) and osteopontin on bovine sperm-egg binding, and fertilization in vitro. *Theriogenology*, 67:468-474.
- Gonçalves RF, Chapman DA, Bertolla RP, Eder I, Killian GJ.** 2008a. Pre-treatment of cattle semen or oocytes with purified milk osteopontin affects in vitro fertilization and embryo development. *Anim Reprod Sci*, 108:375-383.
- Gonçalves RF, Staros AL, Killian GJ.** 2008b. Oviductal fluid proteins associated with the bovine zona pellucida and the effect on *in vitro* sperm-egg binding, fertilization and embryo development. *Reprod Domest Anim*, 43:720-729.
- González-Cadavid V, Martins JA, Moreno FB, Andrade TS, Santo AC, Monteiro-Moreira AC, Moreira RA, Moura AA.** 2014. Seminal plasma proteins of adult boars and correlations with sperm parameters. *Theriogenology*, 82:697-707.
- Goodacre R, Vaidyanathan S, Dunn WB, Harrigan GG, Kell DB.** 2004. Metabolomics by numbers: acquiring and understanding global metabolite data. *Trends Biotechnol*, 22:245-252.
- Gwathmey TM, Ignatz GG, Mueller JL, Manjunath P, Suarez SS.** 2006. Bovine seminal plasma proteins PDC-109, BSP-A3, and BSP-30-kDa share functional roles in storing sperm in the oviduct. *Biol Reprod*, 75:501-507.
- Halliwell B, Gutteridge JMC.** 1990. *Free Radicals in Biology and Medicine*. 2nd. ed. Oxford, UK: Oxford University Press.
- Hao Y, Mathialagan N, Walters E, Mao J, Lai L, Becker D, Li W, Critser J, Prather RS.** 2006. Osteopontin reduces polyspermy during in vitro fertilization of porcine oocytes. *Biol Reprod*, 75:726-733.
- Hao Y, Murphy CN, Spate L, Wax D, Zhong Z, Samuel M, Mathialagan N, Schatten H, Prather RS.** 2008. Osteopontin improves in vitro development of porcine embryos and decreases apoptosis. *Mol Reprod Dev*, 75:291-298.
- Hamamah S, Seguin F, Bujan L, Barthelemy C, Mieuisset R, Lansac J.** 1998. Quantification by magnetic resonance spectroscopy of metabolites in seminal plasma able to differentiate different forms of azoospermia. *Hum Reprod*. 13:132-135.
- Henault MA, Killian GJ, Kavanaugh JF, Griel LC Jr.** 1995. Effect of accessory sex gland fluid from bulls of differing fertilities on the ability of cauda epididymal sperm to penetrate zona-free bovine oocytes. *Biol Reprod*, 52:390-397.
- Hinton BT, Palladino MA, Rudolph D, Labus JC.** 1995. The epididymis as protector of maturing spermatozoa. *Reprod Fertil Dev*, 7:731-745.
- Hinton BT, Palladino MA, Rudolph D, Lan ZJ, Labus JC.** 1996. The role of the epididymis in the protection of spermatozoa. *Curr Top Dev Biol*, 33:61-102.
- Hohlbrugger G, Pschorr J, Dahlheim H.** 1984. Angiotensin I-converting enzyme in the ejaculate of fertile and infertile men. *Fertil Steril*, 41:324-325.
- Hollywood K, Brison DR, Goodacre R.** 2006. Metabolomics: current technologies and future trends. *Proteomics*, 6:4716-4723.
- Holmes E, Wilson ID, Nicholso JK.** 2008. Metabolic phenotyping in health and disease. *Cell*, 134:714-717.
- Humphreys DT, Carver JA, Easterbrook-Smith SB, Wilson MR.** 1999. Clusterin has chaperone-like activity similar to that of small heat shock proteins. *J Biol Chem*. 274:6875-6881.
- Ibrahim NM, Troedsson MH, Foster DN, Loseth KJ, Farris JA, Blaschuk O, Crabo BG.** 1999. Reproductive tract secretions and bull spermatozoa contain different clusterin isoforms that cluster cells and inhibit complement-induced cytolysis. *J Androl*,



20:230-240.

- Ibrahim NM, Foster DN, Crabo BG.** 2001a. Localization of clusterin on freeze-preserved bull spermatozoa before and after glass wool-sephadex filtration. *J Androl*, 22:891-902.
- Ibrahim NM, Romano JE, Troedsson MH, Crabo BG.** 2001b. Effect of scrotal insulation on clusterin-positive cells in ram semen and their relationship to semen quality. *J Androl*, 22:863-877.
- Jeulin C, Soufir JC, Weber P, Laval-Martin D, Calvayrac R.** 1989. Catalase activity in human spermatozoa and seminal plasma. *Gamete Res*, 24:185-196.
- Jin YZ, Bannai S, Dacheux F, Dacheux JL, Okamura N.** 1997. Direct evidence for the secretion of lactoferrin and its binding to sperm in the porcine epididymis. *Mol Reprod Dev*, 47:490-496.
- Jobim MIM, Oberst ER, Salbego CG, Souza DO, Wald VB, Tramontina F, Mattos RC.** 2004. Two-dimensional polyacrylamide gel electrophoresis of bovine seminal plasma proteins and their relation with semen freezability. *Theriogenology*, 61:255-266.
- Kaartinen MT, Pirhonen A, Linnala-Kankkunen A, Mäenpää PH.** 1999. Cross-linking of osteopontin by tissue transglutaminase increases its collagen binding properties. *J Biol Chem*, 274:1729-1735.
- Kaya A, Memili E.** 2016. Sperm macromolecules associated with bull fertility. *Anim Reprod Sci*, 169:88-94.
- Kikuchi M, Mizoroki S, Kubo T, Ohiwa Y, Kubota M, Yamada N, Orino K, Ohnami Y, Watanabe K.** 2003a. Seminal plasma lactoferrin but not transferrin reflects gonadal function in dogs. *J Vet Med Sci*. 65:679-684.
- Kikuchi M, Takao Y, Tokuda N, Ohnami Y, Orino K, Watanabe K.** 2003b. Relationship between seminal plasma lactoferrin and gonadal function in horses. *J Vet Med Sci*, 65:1273-1274.
- Killian GJ, Chapman DA, Rogowski LA.** 1993. Fertility-associated proteins in Holstein bull seminal plasma. *Biol Reprod*, 49:1202-1207.
- Kovac JR, Pastuszak AW, Lamb DJ.** 2013. The use of genomics, proteomics, and metabolomics in identifying biomarkers of male infertility. *Fertil Steril*, 99:998-1007.
- Kraus M, Tichá M, Zelezná B.** 2005. Characterization of human seminal plasma proteins homologous to boar AQN spermadhesins. *J Reprod Immunol*, 65:33-46.
- Kumar A, Kroetsch T, Blondin P, Anzar M.** 2015. Fertility-associated metabolites in bull seminal plasma and blood serum: (1) H nuclear magnetic resonance analysis. *Mol Reprod Dev*, 82:123-131.
- Liaw L, Lindner V, Schwartz SM, Chambers AF, Giachelli CM.** 1995. Osteopontin and beta 3 integrin are coordinately expressed in regenerating endothelium *in vivo* and stimulate Arg-Gly-Asp-dependent endothelial migration *in vitro*. *Circ Res*, 77:665-672.
- Liu Q, Xie QZ, Zhou Y, Yang J.** 2015. Osteopontin is expressed in the oviduct and promotes fertilization and preimplantation embryo development of mouse. *Zygote*, 23:622-630.
- MacLeod J.** 1943. The role of oxygen in the metabolism and motility of the human spermatozoa. *Am J Physiol*, 138:512-518.
- Manjunath P, Sairam MR.** 1987. Purification and biochemical characterization of three major acidic proteins (BSP-A1, BSP-A2 and BSP-A3) from bovine seminal plasma. *Biochem J*, 241(3):685-692.
- Manjunath P, Nauc V, Bergeron A, Menard M.** 2002. Major proteins of bovine seminal plasma bind to the low-density lipoprotein fraction of hen's egg yolk. *Biol Reprod*, 67:1250-1258.
- Manjunath P, Therien I.** 2002. Role of seminal plasma phospholipid-binding proteins in sperm membrane lipid modification that occurs during capacitation. *J Reprod Immunol*, 53:109-119.
- Manjunath P, Lefebvre J, Jois PS, Fan J, Wright MW.** 2009. New nomenclature for mammalian BSP genes. *Biol Reprod*, 80:394-397.
- Matsushita K, Miyake H, Chiba K, Fujisawa M.** 2016. Clusterin produced by Sertoli cells inhibits heat stress induced apoptosis in the rat testis. *Andrologia*, 48:11-19.
- Menezes BM, van Tilburg M, Plante G, De Oliveira RV, Moura AA, Manjunath P.** 2016. Milk proteins interact with goat Binder of Sperm (BSP) proteins and decrease their binding to sperm. *Cell Tissue Res*, 366:427-442.
- Menezes EB, De Oliveira RV, Van Tilburg MF, Babosa EA, Nascimento NV, Velho ALMCS, Moren FB, Moreira RA, Monteiro-Moreira ACO, Carvalho GMC, Ramos AF, Memili E, Moura AA.** 2017. Proteomic analysis of seminal plasma from locally-adapted Curraleiro Pé-Duro bulls (*Bos taurus*): identifying biomarkers involved in sperm physiology in endangered animals for conservation of biodiversity. *Anim Reprod Sci*. 183:86-101.
- Meri S, Jarva H.** 2001. Complement regulatory proteins. In: Encyclopedia of Life Sciences (eLS). London: Nature Publ. pp. 1-7.
- Merlotti A, Dantas E, Remes Lenicov F, Ceballos A, Jancic C, Varese A, Rubione J, Stover S, Geffner J, Sabatte J.** 2015. Fucosylated clusterin in semen promotes the uptake of stress-damaged proteins by dendritic cells via DC-SIGN. *Hum Reprod*, 30:1545-1556.
- Métayer S, Dacheux F, Guérin Y, Dacheux JL, Gatti JL.** 2001. Physiological and enzymatic properties of the ram epididymal soluble form of germinal angiotensin I-converting enzyme. *Biol Reprod*, 65:1332-1339.
- Monaco E, Gasparrini B, Boccia L, De Rosa A, Attanasio L, Zicarelli L, Killian G.** 2009. Effect of osteopontin (OPN) on *in vitro* embryo development in cattle. *Theriogenology*, 71:450-457.
- Montag M, Toth B, Strowitzki T.** 2013. New approaches to embryo selection. *Reprod Biomed Online*. 27:539-546.
- Moura AA.** 2005. Seminal plasma proteins and fertility indexes in the bull: the case for osteopontin. *Anim Reprod*, 2:3-10.
- Moura AA, Koc H, Chapman DA, Killian GJ.** 2006. Identification of accessory sex gland fluid proteins as related to fertility indexes of dairy bulls: a proteomic approach. *J Androl*, 27:201-211.



- Moura AA, Chapman DA, Koc H, Killian GJ. 2007a. A comprehensive proteomic analysis of the accessory sex gland fluid from mature Holstein bulls. *Anim Reprod Sci*, 98:169-188.
- Moura AA, Chapman DA, Killian GJ. 2007b. Proteins of the accessory sex glands associated with the oocyte-penetrating capacity of cauda epididymal sperm from Holstein bulls of documented fertility. *Mol Reprod Dev*, 74:214-222.
- Moura AA, Souza CE, Stanley BA, Chapman DA, Killian GJ. 2010. Proteomics of cauda epididymal fluid from mature Holstein bulls. *J Proteomics*, 73:2006-2020.
- Muñoz M, Uyar A, Correia A, Diez, Fernandez-Gonzalez A, Caamano JN, Trigal B, Carrocera S, Seli E, Gomez E. 2014a. Non-invasive assessment of embryonic sex in cattle by metabolic fingerprinting of in vitro culture medium. *Metabolomics*, 10:443-451.
- Muñoz M, Uyar A, Correia E, Ponsart C, Guyader-Joly C, Martínez-Bello D, Marquant-Le Guienne B, Fernandez-Gonzalez A, Diez C, Caamano JN, Trigal B, Humblot P, Carrocera S, Martin D, Seli E, Gomez E. 2014b. Metabolomic prediction of pregnancy viability in superovulated cattle embryos and recipients with fourier transform infrared spectroscopy. *Biomed Res Int*. 2014:608579.
- Nagy Z, Sakkas, D, Behr B. 2008. Symposium: innovative techniques in human embryo viability assessment. Non-invasive assessment of embryo viability by metabolomic profiling of culture media ('metabolomics'). *Reprod Biomed Online* 17:502-507.
- Nozaki A, Ikeda M, Naganuma A, Nakamura T, Inudoh M, Tanaka K, Kato N. 2003. Identification of a lactoferrin-derived peptide possessing binding activity to hepatitis C virus E2 envelope protein. *J Biol Chem*, 278:10162-10173.
- Ochsendorf FR. 1999. Infections in the male genital tract and reactive oxygen species. *Hum Reprod Update*, 5:399-420.
- Ohta A, Mohri T, Ohyashiki T. 1989. Effect of lipid peroxidation on membrane-bound Ca₂-ATPase activity of the intestinal brush-border membrane. *Biochim Biophys Acta*, 984:151-157.
- Oliveira RV, Dogan S, Belser LE, Kaya A, Topper E, Moura A, Thibaudeau G, Memili E. 2013. Molecular morphology and function of bull spermatozoa linked to histones and associated with fertility. *Reproduction*, 146:263-272.
- Oliver SG, Winson MK, Kell DB, Baganz F. 1998. Systematic functional analysis of the yeast genome. *Trends Biotechnol*, 16:373-378.
- Omenn GS, Lane L, Lundberg EK., Overall CM, Deutsch EW. 2017. Progress on the HUPO draft human proteome: 2017 metrics of the human proteome project. *J Proteome Res*. 16:4281-4287.
- Patti GJ, Shriver LP, Wassif CA, Woo HK, Uritboonthai W, Apon J, Manchester M, Porter FD, Siuzdak G. 2010. Nanostructure-initiator mass spectrometry (NIMS) imaging of brain cholesterol metabolites in Smith-Lemli-Opitz syndrome. *Neuroscience*, 170:858-864.
- Perry AC, Jones R, Niang LS, Jackson RM, Hall L. 1992. Genetic evidence for an androgen-regulated epididymal secretory glutathione peroxidase whose transcript does not contain a selenocysteine codon. *Biochem J*, 285:863-870.
- Pietrobon EO, Soria M, Domínguez LA, MonclusMde L, Fornés MW. 2005. Simultaneous activation of PLA2 and PLC are required to promote acrosomal reaction stimulated by progesterone via G-proteins. *Mol Reprod Dev*, 70:58-63.
- Plante G, Thérien I, Lachance C, Leclerc P, Fan J, Manjunath P. 2014. Implication of the human Binder of Sperm Homolog 1 (BSPH1) protein in capacitation. *Mol Hum Reprod*, 20:409-421
- Plante G, Lusignan MF, Lafleur M, Manjunath P. 2015. Interaction of milk proteins and Binder of Sperm (BSP) proteins from boar, stallion and ram semen. *Reprod Biol Endocrinol*, 13:92-112.
- Plante G, Manjunath P. 2015. Murine binder of sperm protein homolog 1: a new player in HDL-induced capacitation. *Reproduction*, 149:367-376.
- Plante G, Prud'homme B, Fan J, Lafleur M, Manjunath P. 2016. Evolution and function of mammalian binder of sperm proteins. *Cell Tissue Res*, 363:105-127.
- Pucci S, Bonanno E, Pichiorri F, Angeloni C, Spagnoli LG. 2004. Modulation of different clusterin isoforms in human colon tumorigenesis. *Oncogene*, 23:2298-2304.
- Rangaswami H, Bulbule A, Kundu GC. 2006. Osteopontin: role in cell signaling and cancer progression. *Trends Cell Biol*, 16:79-87.
- Reddy VRK, Rajeev S, Gupta V. 2003. α6β1 integrin is a potential clinical marker for evaluating sperm quality in men. *Fertil Steril*, 79:1590-1596.
- Rego JP, Moura AA, Nouwens AS, McGowan MR, Boe-Hansen GB. 2014. Seminal plasma proteome of electroejaculated *Bos indicus* bulls. *Anim Reprod Sci*, 148:1-17.
- Rego JPA, Martins JM, Wolf CA, Van Tilburg M, Moreno F, Monteiro-Moreira AC, Moreira RA, Santos DO, Moura AA. 2016. Proteomic analysis of seminal plasma and sperm cells and their associations with semen freezability in Guzerat bulls. *J Anim Sci*, 94:5308-5320.
- Rittling S R, Singh R. 2015. Osteopontin in immune-mediated diseases. *J Dent Res*, 94:1638-1645.
- Rodríguez-Villamil P, Hoyos-Marulanda V, Martins JA, Oliveira AN, Aguiar LH, Moreno FB, Velho AL, Monteiro-Moreira AC, Moreira RA, Vasconcelos IM, Bertolini M, Moura AA. 2016. Purification of binder of sperm protein 1 (BSP1) and its effects on bovine in vitro embryo development after fertilization with ejaculated and epididymal sperm. *Theriogenology*, 85:540-554.
- Roldan ER.1998. Role of phospholipases during sperm acrosomal exocytosis. *Front Biosci*, 3:D1109-D1119.
- Roldan ER, Shi QX. 2007. Sperm phospholipases and acrosomal exocytosis. *Front Biosci*, 2:89-104.
- Romão MJ, Kolln I, Dias JM, Carvalho AM, Romero A, Varela PF, Sanz L, Topfer-Petersen E, CalveteJJ. 1997. Crystal structure of acidic seminal fluid protein (asfp) at 1.9 Å resolution: a bovine



- polypeptide of the spermadhesin family. *J Mol Biol*, 274:650-660.
- Rorie RW, Williams CL, Lester TD.** 2016. Association of osteopontin gene promoter single nucleotide polymorphisms with bull semen quality. *Adv Reprod Sci*, 4:1-7.
- Salehi M, Akbari H, Heidari MH, Molouki A, Murulitharan K, Moeini H, Novin MG, Aabed F, Taheri H, Fadaei F, Mohsenzadeh M, Jafari M, Pirouzi A, Heidari R.** 2013. Correlation between human clusterin in seminal plasma with sperm protamine deficiency and DNA fragmentation. *Mol Reprod Dev*, 80:718-724.
- Santos EA, Sousa PC, Martins JA, Moreira RA, Monteiro-Moreira AC, Moreno FB, Oliveira MF, Moura AA, Silva AR.** 2014. Protein profile of the seminal plasma of collared peccaries (*Pecari tajacu* Linnaeus, 1758). *Reproduction*, 147:753-764.
- Schill WB.** 1989. Significance of the kallikrein-kinin system in andrology. In: Fritz H, Schimidt I, Dietze G (Ed.). *The kallikrein-kinin System in Health and Disease*. Braunschweig: Limbach Verlag. pp. 171-203.
- Schoenfelder M, Einspanier R.** 2003. Expression of hyaluronan synthases and corresponding hyaluronan receptors is differentially regulated during oocyte maturation in cattle. *Biol Reprod*, 69:269-277.
- Schöneck C, Braun J, Einspanier R.** 1996. Sperm viability is influenced in vitro by the bovine seminal protein aSP: effects on motility, mitochondrial activity and lipid peroxidation. *Theriogenology*, 45:633-642.
- Seli E, Sakkas D, Scott R, Kwok SC, Rosendahl SM, Burns, DH.** 2007. Noninvasive metabolomic profiling of embryo culture media using Raman and near-infrared spectroscopy correlates with reproductive potential of embryos in women undergoing in vitro fertilization. *Fertil Steril*, 88:1350-1357.
- Seli E, Botros L, Sakkas D, Burns DH.** 2008. Noninvasive metabolomic profiling of embryo culture media using proton nuclear magnetic resonance correlates with reproductive potential of embryos in women undergoing in vitro fertilization. *Fertil Steril*, 90:2183-2189.
- Seli E, Robert C, Sirard MA.** 2010a. OMICS in assisted reproduction: possibilities and pitfalls. *Mol Hum Reprod*, 16:513-530.
- Seli E, Vergouw CG, Morita H, Botros L, Roos P, Lambalk CB, Yamashita N, Kato O, Sakkas D.** 2010b. Noninvasive metabolomic profiling as an adjunct to morphology for noninvasive embryo assessment in women undergoing single embryo transfer. *Fertil Steril*. 94:535-542.
- Silva HVR, Rodríguez-Villamil P, Magalhaes FF, Nunes TGP, Freitas LA, Ribeiro LR, Silva AR, Moura AA, Silva LDM.** 2018. Seminal plasma and sperm proteome of ring-tailed coatis (*Nasua nasua*, Linnaeus, 1766). *Theriogenology*, 111:34-42.
- Singh R, Sinclair KD.** 2007. Metabolomics: approaches to assessing oocyte and embryo quality. *Theriogenology*, 68:S56-62.
- Somlev B, Helili K, Karcheva V.** 1996. Tissue kallikrein activity in seminal plasma of bovine ejaculates with different quality. *Theriogenology*, 45:471-475.
- Soubeyrand S, Khadir A, Brindle Y, Manjunath P.** 1997. Purification of a novel phospholipase A2 from bovine seminal plasma. *J Biol Chem*, 272:222-227.
- Souza CEA, Moura AA, Monaco E, Killian GJ.** 2008. Binding patterns of bovine seminal plasma proteins A1/A2, 30 kDa and osteopontin on ejaculated sperm before and after incubation with isthmic and ampullary oviductal fluid. *Anim Reprod Sci*, 105:72-89.
- Souza CEA, Rego JP, Lobo CH, Oliveira JT, Nogueira FC, Domont GB, Fioramonte M, Gozzo FC, Moreno FB, Monteiro-Moreira AC, Figueiredo JR, Moura AA.** 2012. Proteomic analysis of the reproductive tract fluids from tropically-adapted Santa Ines rams. *J Proteomics*, 75:4436-4456.
- Suarez SS.** 2016. Mammalian sperm interactions with the female reproductive tract. *Cell Tissue Res*, 363:185-194.
- Tenniswood MP, Guenette RS, Lakins J, Mooibroek M, Wong P, Welsh JE.** 1992. Active cell death in hormone-dependent tissues. *Cancer Metastasis Rev*, 11:197-220.
- Thaler CJ, Vanderpuye OA, McIntyre JA, Faulk WP.** 1990. Lactoferrin binding molecules in human seminal plasma. *Biol Reprod*, 43:712-717.
- Thérien I, Bleau G, Manjunath P.** 1995. Phosphatidylcholine-binding proteins of bovine seminal plasma modulate capacitation of spermatozoa by heparin. *Biol Reprod*, 52:1372-1379.
- Thérien I, Moreau R, Manjunath P.** 1998. Major proteins of bovine seminal plasma and high-density lipoprotein induce cholesterol efflux from epididymal sperm. *Biol Reprod*, 59:768-776.
- Thérien I, Moreau R, Manjunath P.** 1999. Bovine seminal plasma phospholipid binding proteins stimulate phospholipid efflux from epididymal sperm. *Biol Reprod*, 61:590-598.
- Thul PJ, Åkesson L, Wiking M, Mahdessian D, Geladaki A, Ait Blal H, Alm T, Asplund A, Björk L, Breckels LM, Bäckström A, Danielsson F, Fagerberg L, Fall J, Gatto L, Gnann C, Hober S, Hjelmare M, Johansson F, Lee S, Lindskog C, Mulder J, Mulvey CM, Nilsson P, Oksvold P, Rockberg J, Schutten R, Schwenk JM, Sivertsson Å, Sjöstedt E, Skogs M, Stadler C, Sullivan DP, Tegel H, Winsnes C, Zhang C, Zwahlen M, Mardinoglu A, Pontén F, von Feilitzen K, Lilley KS, Uhlén M, Lundberg E.** 2017. A subcellular map of the human proteome. *Science*, 356:6340.
- Trougakos IP, Poulakou M, Stathatos M, Chalikia A, Melidonis A, Gonos ES.** 2002. Serum levels of the senescence biomarker clusterin/apolipoprotein J increase significantly in diabetes type II and during development of coronary heart disease or at myocardial infarction. *Exp Gerontol*, 37:1175-1187.
- Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C, Sjöstedt E, Asplund A, Olsson I, Lundberg E, Navani S, Szigartyo CA, Odeberg J, Djureinovic D, Takanen JO, Hober S, Alm T, Edqvist PH, Berling H, Tegel H, Mulder J, Rockberg J, Nilsson P, Schwenk JM, Hamsten M, von Feilitzen K, Forsberg**



- M, Persson L, Johansson F, Zwahlen M, von Heijne G, Nielsen J, Pontén F.** 2015. Proteomics. Tissue-based map of the human proteome. *Science*. 347:1260419.
- van Tilburg MF, Salles MG, Silva MM, Moreira RA, Moreno FB, Monteiro-Moreira AC, Martins JA, Cândido MJ, Araújo AA, Moura AA.** 2014. Semen variables and sperm membrane protein profile of Saanen bucks (*Capra hircus*) in dry and rainy seasons of the Northeastern Brazil (3 degrees S). *Int J Biometeorol*, 59:561-573.
- Vinson GP, Mehta J, Evans S, Matthews, S, Puddefoot, JR, Saridogan E, Holt WV, Djahanbakhch O.** 1996. Angiotensin II stimulates sperm motility. *Regul Pept*, 67:131-135.
- Wai PY, Kuo PC.** 2004. The role of osteopontin in tumor metastasis. *J Surg Res*, 121:228-241.
- Wehrli P, Charnay Y, Vallet P, Zhu G, Harmony J, Aronow B, Tschopp J, Bouras C, Viard-Leveugle I, French LE, Giannakopoulos P.** 2001. Inhibition of postischemic brain injury by clusterin overexpression. *Nat Med*, 7:977-979.
- Xuan JW, Hota C, Shigeyama Y, D'Errico JA, Somerman MJ, Chambers AF.** 1995. Site-directed mutagenesis of the arginine-glycine-aspartic acid sequence in osteopontin destroys cell adhesion and migration functions. *JCell Biochem*, 57:680-690.
- Zhang J, Mu X, Xia Y, Martin FL, Hang W, Liu L, Tian M, Huang Q, Shen H.** 2014. Metabolomic analysis reveals a unique urinary pattern in normozoospermic infertile men. *J Proteome Res*. 13:3088-3099.
-