

Meeting abstract

Identification of composite promoter modules in inflammation-regulated genes

A Weber*¹, O Dittrich-Breiholz², H Schneider², A Kel³, R Jauregui³, E Wingender³ and M Kracht¹

Address: ¹Rudolf-Buchheim-Institute of Pharmacology, Justus-Liebig-University, Giessen, Germany, ²Institute of Pharmacology, Medical School Hannover, Hannover, Germany and ³BIOBASE GmbH, Wolfenbuettel, Germany

* Corresponding author

from 12th Joint Meeting of the Signal Transduction Society (STS). Signal Transduction: Receptors, Mediators and Genes Weimar, Germany. 29–31 October 2008

Published: 26 February 2009

Cell Communication and Signaling 2009, **7**(Suppl 1):A105 doi:10.1186/1478-811X-7-S1-A105

This abstract is available from: <http://www.biosignaling.com/content/7/S1/A105>

© 2009 Weber et al; licensee BioMed Central Ltd.

In the context of the collaborative research centre SFB566 entitled "cytokine receptors and cytokine-dependent signalling pathways as therapeutic target structures" we have performed more than 1100 DNA microarray experiments under highly standardized experimental conditions with respect to RNA isolation, cRNA synthesis, cRNA labelling, cRNA hybridization and raw data acquisition. The expression of 90 inflammatory genes including 6 housekeeping genes were assessed in a large variety of human cell systems and stored in our microarray database CytoBASE <https://microarray.med.uni-giessen.de/base/index.phtml>. A set of 102 experiments comprising 18 human cell types, 59 experimental groups and treatment with 36 different stimuli/inhibitors in 47 different combinations was analyzed by hierarchical clustering using MultiExperiment Viewer (MEV, version 4.1, <http://www.tm4.org>) and Significance Analysis of Microarrays (SAM, <http://www-stat.Stanford.EDU/~tibs/SAM>). As a result 20 consistently coregulated inflammatory genes with different biological functions such as cytokines and chemokines (IL-6, CCL2, CXCL3, CXCL8, CXCL10), proteases (MMP-1, MMP-15), metabolic enzymes (COX-2, MnSOD) and intracellular signalling molecules (BIRC2, IκBα, IRF1, JUNB) were identified. By genome-wide microarray analysis we also compiled a control set of 48 genes that were not regulated under eight different inflammatory conditions. We searched 1.1 to 2.1 kB of the promoter regions of these genes for enriched transcription factor binding sites using F-Match <http://www.biobase.de>. We further identified

combinations of binding sites of enriched transcription factors that correlate with coregulation of inflammatory but not of control genes by using the composite module analysis tool <http://www.biobase.de>.

Several promoter modules containing 3 to 10 transcription factor binding sites within a promoter region of 200 bp were identified that distinguished the inducible 20 inflammatory from the 48 control genes. This information can be used in future experiments designed to reveal novel mechanisms of coregulation of different functional classes of inflammatory genes at the level of DNA.