

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

## 6-Methylsulfanyl-4H-pyrimido[1,6-a]-pyrimidin-4-one

Quinhua Huang,<sup>a</sup> Paul F. Richardson,<sup>a</sup> Eugene Rui,<sup>a</sup>  
Arnold L. Rheingold<sup>b</sup> and Alex Yanovsky<sup>a\*</sup>

<sup>a</sup>Pfizer Global Research and Development, La Jolla Labs, 10770 Science Center Drive, San Diego, CA 92121, USA, and <sup>b</sup>Department of Chemistry and Biochemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA  
Correspondence e-mail: alex.yanovsky@pfizer.com


Received 18 September 2009; accepted 23 September 2009

Key indicators: single-crystal X-ray study;  $T = 123$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.035;  $wR$  factor = 0.097; data-to-parameter ratio = 12.7.

Reaction of 2-(methylsulfanyl)pyrimidin-4-amine with the 5-(methoxyvinylidene) derivative of Meldrum's acid and subsequent heating of the product in Dowtherm fluid yielded the title compound,  $\text{C}_8\text{H}_7\text{N}_3\text{OS}$ , which was proven to contain a bicyclic 4H-pyrimido[1,6-a]pyrimidine system. All non-H atoms of the molecule are coplanar within 0.15 Å. The bond-length distribution in the bicyclic core shows localization of the double bonds. The geometry of the intramolecular  $\text{S}\cdots\text{O}$  1,5-contact [2.534 (2) Å] is consistent with the existence of an attractive interaction.

### Related literature

For the structure of a compound with a similar bicyclic carbon–nitrogen core, see: Olomucki *et al.* (1984). For statistical studies of the geometry of  $\text{S}\cdots\text{O}$  interactions, see: Rosenfield *et al.* (1977); Iwaoka *et al.* (2002).



### Experimental

#### Crystal data

$\text{C}_8\text{H}_7\text{N}_3\text{OS}$   
 $M_r = 193.23$   
Monoclinic,  $P2_1/c$   
 $a = 9.7621$  (8) Å  
 $b = 4.1725$  (3) Å  
 $c = 20.4092$  (16) Å  
 $\beta = 100.106$  (1)°  
 $V = 818.42$  (11) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.35$  mm<sup>-1</sup>  
 $T = 123$  K  
 $0.48 \times 0.14 \times 0.08$  mm

#### Data collection

Bruker APEXII CCD diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 2001)  
 $T_{\min} = 0.849$ ,  $T_{\max} = 0.972$   
6497 measured reflections  
1498 independent reflections  
1333 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.022$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$   
 $wR(F^2) = 0.097$   
 $S = 1.13$   
1498 reflections  
118 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.43$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.19$  e Å<sup>-3</sup>

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SIR2004 (Burla *et al.*, 2005); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-32 (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2363).

### References

- Bruker (2001). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giovacazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.  
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.  
Iwaoka, M., Takemoto, S. & Tomoda, S. (2002). *J. Am. Chem. Soc.* **124**, 10613–10620.  
Olomucki, M., Le Gall, J. Y., Colinart, S., Durant, F., Norberg, B. & Evrard, G. (1984). *Tetrahedron Lett.* **25**, 3471–3474.  
Rosenfield, R. E. Jr, Parthasarathy, R. & Dunitz, J. D. (1977). *J. Am. Chem. Soc.* **99**, 4860–4862.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

**supplementary materials**

*Acta Cryst.* (2009). E65, o2572 [ doi:10.1107/S1600536809038562 ]

## 6-Methylsulfanyl-4*H*-pyrimido[1,6-*a*]pyrimidin-4-one

Q. Huang, P. F. Richardson, E. Rui, A. L. Rheingold and A. Yanovsky

### Comment

The title compound was obtained by reaction of 2-(methylsulfanyl)pyrimidin-4-amine with 5-(methoxyvinylidene) derivative of Meldrum's acid and subsequent heating of the product in Dowtherm fluid (Fig. 1).

All non-hydrogen atoms of the molecule of the title compound (Fig. 2) are approximately coplanar; maximum deviation of the C1 atom from the least-squares plane is 0.143 (2) Å. The bond length values indicate localization of double bonds in the bicyclic core (N1–C2, 1.298 (3) Å; N3–C5, 1.308 (3) Å; C3–C4, 1.351 (3) Å; C6–C7, 1.355 (3) Å);, which is consistent with the fact that no resonance structures can be drawn. Surprisingly, to the best of our knowledge no other isolated bicyclic carbon-nitrogen system with the same positions of the N atoms was structurally studied. Similar bond lengths were observed in dihydropyrimido(1,2-*c*)purine derivative, where analogous bicyclic core makes up a part of the tricyclic system (Olomucki *et al.*, 1984).

The S1...O1 distance [2.534 (2) Å] and the O1...S1—C1 angle [177.5 (1)°] are consistent with the existence of the intramolecular attractive interaction (Rosenfield *et al.*, 1977; Iwaoka *et al.*, 2002).

### Experimental

A mixture of (2-methylsulfanyl)pyrimidin-4-ylamine (5.0 g, 35 mmol, 1.0 eq), 5-(methoxyvinylidene) derivative of Meldrum's acid (9.23 g, 49.6 mmol, 1.4 eq), and 2-PrOH (100 ml) was refluxed at 85°C for 1 h to reach reaction completion. The resulting suspension was cooled to 25°C, filtered, and washed with EtOH to afford 10 g of the displacement product as a yellow solid in 96% yield. LC—MS (APCI, *M*+1) 296.0; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ p.p.m. 1.69 (s, 6 H) 2.55 (s, 3 H) 7.43 (d, *J* = 5.54 Hz, 1 H) 8.57 (d, *J* = 5.54 Hz, 1 H) 9.20 (s, 1 H) 11.35 (s, 1 H).

The generated 5-[(2-methylsulfanyl)pyrimidin-4-ylaminovinylidene] derivative of Meldrum's acid (6.5 g, 22 mmol, 1.0 eq) was added portionwise to 130 ml of preheated Dowtherm A at 220°C (inner temperature). After being reacted at 220°C for 10 min, LC—MS indicated that reaction was complete. The reaction mixture was cooled to 25°C and diluted with 1 *L* heptane. The resulting suspension was left to stand overnight. The solid was filtered, washed with heptane, and dried under vacuum to afford 3.4 g of the target compound as a yellow solid in (80% yield). LC—MS (APCI, *M*+1) 194.0; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ p.p.m. 2.40 (s, 3 H) 6.35 (d, *J* = 6.55 Hz, 1 H) 7.04 (d, *J* = 6.29 Hz, 1 H) 8.08 (t, *J* = 6.55 Hz, 2 H). The product was recrystallized from EtOAc/hexane to yield single crystals suitable for X-ray diffraction studies.

### Refinement

All H atoms were placed in geometrically calculated positions (C—H = 0.95 Å and 0.98 Å for aromatic and methyl H atoms respectively) and included in the refinement in riding motion approximation. The *U*<sub>iso</sub>(H) were set to 1.2*U*<sub>eq</sub>(C) or 1.5*U*<sub>eq</sub>(C) for methyl H atoms.

## Figures




Fig. 1. Synthesis of the title compound.




Fig. 2. Molecular structure of the title compound, showing 50% probability displacement ellipsoids and atom numbering scheme. H atoms are drawn as circles with arbitrary small radius.

## 6-Methylsulfanyl-4H-pyrimido[1,6-a]pyrimidin-4-one

### Crystal data

$C_8H_7N_3OS$

$M_r = 193.23$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 9.7621 (8) \text{ \AA}$

$b = 4.1725 (3) \text{ \AA}$

$c = 20.4092 (16) \text{ \AA}$

$\beta = 100.1060 (10)^\circ$

$V = 818.42 (11) \text{ \AA}^3$

$Z = 4$

$F_{000} = 400$

$D_x = 1.568 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4134 reflections

$\theta = 3.2\text{--}25.4^\circ$

$\mu = 0.35 \text{ mm}^{-1}$

$T = 123 \text{ K}$

Blade, brown

$0.48 \times 0.14 \times 0.08 \text{ mm}$

### Data collection

Bruker APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 123 \text{ K}$

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(SADABS; Bruker, 2001)

$T_{\min} = 0.849$ ,  $T_{\max} = 0.972$

6497 measured reflections

1498 independent reflections

1333 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.022$

$\theta_{\max} = 25.4^\circ$

$\theta_{\min} = 2.1^\circ$

$h = -11 \rightarrow 11$

$k = -4 \rightarrow 5$

$l = -24 \rightarrow 24$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.035$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$wR(F^2) = 0.097$	$w = 1/[\sigma^2(F_o^2) + (0.045P)^2 + 0.5615P]$
$S = 1.13$	where $P = (F_o^2 + 2F_c^2)/3$
1498 reflections	$(\Delta/\sigma)_{\max} = 0.001$
118 parameters	$\Delta\rho_{\max} = 0.43 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.56563 (5)	0.91748 (12)	0.81678 (2)	0.02069 (19)
O1	0.80408 (15)	0.9008 (4)	0.78390 (7)	0.0276 (4)
N1	0.59517 (17)	0.5753 (4)	0.92701 (8)	0.0203 (4)
N2	0.79659 (16)	0.5851 (4)	0.87620 (7)	0.0172 (4)
N3	0.99408 (17)	0.2689 (4)	0.92351 (8)	0.0238 (4)
C1	0.3990 (2)	0.9169 (5)	0.84302 (11)	0.0280 (5)
H1A	0.3333	1.0474	0.8123	0.042*
H1B	0.4089	1.0064	0.8880	0.042*
H1C	0.3642	0.6966	0.8431	0.042*
C2	0.65797 (19)	0.6714 (5)	0.87950 (9)	0.0183 (4)
C3	0.6652 (2)	0.3762 (5)	0.97497 (10)	0.0209 (4)
H3B	0.6193	0.3094	1.0100	0.025*
C4	0.7959 (2)	0.2700 (5)	0.97507 (9)	0.0203 (4)
H4A	0.8392	0.1284	1.0089	0.024*
C5	0.8678 (2)	0.3717 (5)	0.92427 (9)	0.0186 (4)
C6	1.0583 (2)	0.3766 (5)	0.87347 (11)	0.0254 (5)
H6A	1.1495	0.3010	0.8719	0.030*
C7	0.9993 (2)	0.5853 (5)	0.82589 (9)	0.0199 (4)
H7A	1.0506	0.6512	0.7927	0.024*
C8	0.8652 (2)	0.7063 (5)	0.82424 (9)	0.0219 (4)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0177 (3)	0.0225 (3)	0.0219 (3)	0.00111 (19)	0.0038 (2)	0.0028 (2)

## supplementary materials

---

O1	0.0237 (8)	0.0363 (9)	0.0241 (7)	0.0002 (7)	0.0079 (6)	0.0095 (7)
N1	0.0196 (8)	0.0199 (9)	0.0220 (8)	-0.0009 (7)	0.0053 (7)	-0.0002 (7)
N2	0.0184 (8)	0.0176 (8)	0.0164 (8)	-0.0021 (6)	0.0050 (6)	-0.0017 (6)
N3	0.0199 (8)	0.0240 (9)	0.0278 (9)	0.0015 (7)	0.0047 (7)	-0.0014 (8)
C1	0.0201 (11)	0.0335 (12)	0.0312 (11)	0.0032 (9)	0.0068 (9)	0.0059 (10)
C2	0.0169 (9)	0.0178 (9)	0.0201 (10)	-0.0015 (8)	0.0026 (8)	-0.0037 (8)
C3	0.0211 (10)	0.0232 (10)	0.0190 (10)	-0.0025 (8)	0.0052 (8)	0.0004 (8)
C4	0.0233 (10)	0.0189 (10)	0.0184 (10)	-0.0007 (8)	0.0026 (8)	0.0006 (8)
C5	0.0179 (10)	0.0173 (9)	0.0200 (10)	-0.0016 (8)	0.0022 (8)	-0.0041 (8)
C6	0.0180 (10)	0.0264 (11)	0.0326 (11)	-0.0005 (9)	0.0068 (9)	-0.0070 (9)
C7	0.0253 (11)	0.0186 (10)	0.0183 (10)	-0.0042 (8)	0.0108 (8)	-0.0040 (8)
C8	0.0234 (10)	0.0245 (11)	0.0186 (10)	-0.0066 (9)	0.0059 (8)	-0.0026 (9)

### *Geometric parameters (Å, °)*

S1—C2	1.761 (2)	C1—H1B	0.9800
S1—C1	1.799 (2)	C1—H1C	0.9800
O1—C8	1.233 (2)	C3—C4	1.351 (3)
N1—C2	1.298 (3)	C3—H3B	0.9500
N1—C3	1.371 (3)	C4—C5	1.415 (3)
N2—C2	1.413 (2)	C4—H4A	0.9500
N2—C5	1.414 (2)	C6—C7	1.355 (3)
N2—C8	1.442 (2)	C6—H6A	0.9500
N3—C5	1.308 (3)	C7—C8	1.398 (3)
N3—C6	1.365 (3)	C7—H7A	0.9500
C1—H1A	0.9800		
C2—S1—C1	99.03 (9)	N1—C3—H3B	118.2
C2—N1—C3	118.60 (17)	C3—C4—C5	119.37 (18)
C2—N2—C5	119.08 (16)	C3—C4—H4A	120.3
C2—N2—C8	121.20 (16)	C5—C4—H4A	120.3
C5—N2—C8	119.70 (16)	N3—C5—N2	123.26 (18)
C5—N3—C6	117.29 (18)	N3—C5—C4	119.98 (18)
S1—C1—H1A	109.5	N2—C5—C4	116.76 (17)
S1—C1—H1B	109.5	C7—C6—N3	123.66 (19)
H1A—C1—H1B	109.5	C7—C6—H6A	118.2
S1—C1—H1C	109.5	N3—C6—H6A	118.2
H1A—C1—H1C	109.5	C6—C7—C8	121.85 (18)
H1B—C1—H1C	109.5	C6—C7—H7A	119.1
N1—C2—N2	122.57 (17)	C8—C7—H7A	119.1
N1—C2—S1	118.33 (15)	O1—C8—C7	126.55 (18)
N2—C2—S1	119.10 (14)	O1—C8—N2	119.26 (18)
C4—C3—N1	123.53 (18)	C7—C8—N2	114.19 (17)
C4—C3—H3B	118.2		

Fig. 1




Fig. 2

