

Clinical Commentary/Current Concept Review

Classification Based Treatment of Greater Trochanteric Pain Syndrome (GTPS) with Integration of the Movement System

Ashley E. Disantis^{1, a}, RobRoy L. Martin¹¹ Department of Physical Therapy, Duquesne University

Keywords: GTPS, classification, treatment, rehabilitation, exercise

<https://doi.org/10.26603/001c.32981>

International Journal of Sports Physical TherapyVol. 17, Issue 3, 2022

Greater trochanteric pain syndrome (GTPS) refers to pain in the lateral hip and thigh and can encompass multiple diagnoses including external snapping hip (coxa saltans), also known as proximal iliotibial band syndrome, trochanteric bursitis, and gluteus medius (GMed) or gluteus minimus (GMin) tendinopathy or tearing. GTPS presents clinicians with a similar diagnostic challenge as non-specific low back pain with special tests being unable to identify the specific pathoanatomical structure involved and do little to guide the clinician in prescription of treatment interventions. Like the low back, the development of GTPS has been linked to faulty mechanics during functional activities, mainly the loss of pelvic control in the frontal plane secondary to hip abductor weakness or pain with hip abductor activation. Therefore, an impairment-based treatment classification system is recommended in the setting of GTPS in order to better tailor conservative treatment interventions and improve functional outcomes.

Level of Evidence

Level V, clinical commentary

INTRODUCTION

Greater trochanteric pain syndrome (GTPS) refers to pain in the lateral hip and thigh and can encompass multiple diagnoses including external snapping hip (coxa saltans), also known as proximal iliotibial band syndrome, trochanteric bursitis, and gluteus medius (GMed) or gluteus minimus (GMin) tendinopathy or tearing.^{1,2} Although the symptoms associated with GTPS were previously thought to be primarily the result of trochanteric bursitis, multiple histologic and imaging studies have identified gluteal tendinopathy with or without bursitis as the primary source of pain and dysfunction.³⁻⁵ GTPS presents clinicians with a diagnostic challenge as no cluster of special test exists to identify the specific pathoanatomical structure involved or the severity of its involvement. Therefore, there is a need for an impairment-based classification system for GTPS to allow for tailored conservative treatment interventions and improved patient outcomes.

The use of the term GTPS is non-specific to pathology and refers to the many sources of pain in the lateral hip.

This is similar to the use of the term “non-specific low back pain”, which encompasses many different pathologies including but not limited to muscular strains, spinal stenosis, and intervertebral disc degeneration.² Instead of using special tests to identify the specific anatomical structure involved, which is common practice for many peripheral joints, examination procedures for the low back focus on the identification of impairments that lead to the development of pain.⁶ It has been suggested that identifying the specific anatomical structure involved in the onset of low back pain is not necessary to achieve successful resolution of pain.⁷⁻⁹ As a result, an impairment-based treatment classification approach is recommended in an attempt to tailor treatments to a particular subgroup depending on subjective and objective findings. GTPS presents clinicians with a similar diagnostic challenge as non-specific low back pain with special tests being unable to identify the specific pathoanatomical structure involved and do little to guide the clinician in prescription of treatment interventions. Like the low back, the development of GTPS has been linked to movement systems abnormalities during functional activities. The primary movement system finding is the loss

a Corresponding author:Ashley Disantis, PT, OCS
UPMC Children's Hospital of Pittsburgh
Pittsburgh, PA 15224
disantis@duq.edu
412-396-1811

Table 1. Differential Diagnoses: Key Subjective and Objective Findings

	Key Subjective Findings	Key Objective Findings
Lumbar Spine Pathology	<ul style="list-style-type: none"> • Pain that radiates below the knee • Low back pain that increases with sitting or walking • Pain that increases with movement of the lumbar spine 	<ul style="list-style-type: none"> • Pain, weakness, or altered sensation in dermatomal or myotomal pattern • Abnormal findings during lower extremity reflex testing • Pain reproduced with motion testing of the lumbar spine
Hip Joint Osteoarthritis	<ul style="list-style-type: none"> • Pain located in the anterior hip • Morning stiffness that resolves in < 1 hour • Increased pain and difficulty with weight bearing activities 	<ul style="list-style-type: none"> • Reproduction of pain during a FABER and/or Scour test • Decreased range of motion in a capsular pattern • Antalgic gait pattern
Femoral Stress Fracture	<ul style="list-style-type: none"> • Pain located in the anterolateral hip that worsens with weight bearing • Recent rapid increase in weight bearing activities 	<ul style="list-style-type: none"> • Painful and limited hip range of motion • Antalgic gait pattern
Avascular Necrosis of the Hip	<ul style="list-style-type: none"> • Pain located in the anterolateral hip that worsens with weight bearing • History of excessive alcohol or steroid use 	<ul style="list-style-type: none"> • Painful and limited hip range of motion • Antalgic gait pattern

of pelvic control in the frontal plane secondary to hip abductor weakness and/or pain.^{10,11} Therefore, conservative interventions should focus on the correction of these impairments to reduce pain and improve movement control in individuals with GTPS.

The purpose of this clinical commentary is to present an impairment-based classification system for GTPS to allow for tailored treatment interventions. Grimaldi & Fearon¹² presented a graded exercise program for the treatment of GTPS. Because the focus of their work was based solely on gluteal tendinopathy, further classification based on tissue irritability is warranted.¹² We propose a classification system based on findings identified during a standard physical examination coupled with findings from a movement system assessment. By addressing identified impairments, treatment will focus on correcting the faulty mechanics that lead to the development of GTPS rather than treatment being aimed towards a single anatomical structure.

GREATER TROCHANTERIC PAIN SYNDROME

GTPS has an incidence of 1.8 out of 1000 patients in the primary care setting, mainly affecting females in the 4th-6th decades of life.¹⁵⁻¹⁷ Multiple studies have demonstrated a higher incidence of GTPS with concomitant low back pain, hip osteoarthritis (OA), iliotibial band tenderness, and knee pain.^{18,19} While the majority of GTPS is observed in middle aged females, at a much lower rate runners, football players, and dancers have also shown to be affected due to high hip adduction moments during performance of their respective sports.^{17,20-22} In general, abnormal forces across the hip secondary to faulty biomechanics are hypothesized to lead to GTPS. Specifically, weakness or pain in the hip abductors results in a loss of pelvic control in the frontal plane, presenting as a Trendelenburg or compensated Trendelenburg position during tasks performed on a single leg. Pelvic obliquity increases the compressive forces through the lateral hip leading to degeneration of the GMed and GMin tendons and/or inflammation of the trochanteric bursae and proximal iliotibial band.²³

Individuals suffering from GTPS complain of pain at the lateral hip, with symptoms potentially radiating distally to the level of the buttock and lateral thigh. Pain generally worsens with single limb weight bearing activities and laying on the affected side. The greatest degree of tenderness should specifically be located on the lateral or posterior aspect of the greater trochanter.^{3,15,24,25} Activities that exacerbate symptoms include walking and ascending stairs, as well as other single leg activities, such as putting on pants or getting in/out of the bathtub. Due to the high likelihood of overlapping pathologies, particularly hip joint osteoarthritis and low back pain, it is important to first assess for the presence of intra-articular hip and/or lumbar spine pathology. It is also important to rule out the presence of a femoral stress fracture or avascular necrosis of the hip which would require a physician referral.¹⁹ Individuals complaining of radiating pain below the knee, pain in the low back that increases with sitting or walking, and pain specifically associated with movement of the lumbar spine should undergo a comprehensive lower quarter screen. This examination should assess for myotome related weakness, decreased sensation in dermatomal pattern, abnormal findings during lower extremity reflex testing, and lumbar range of motion.

For those with groin pain and morning stiffness hip osteoarthritis should be considered.²⁶ Intra-articular hip joint pathology should be considered with the reproduction of groin pain during a Flexion-Abduction-External Rotation (FABER) test and/or Scour test, decreased range of motion in a capsular pattern, and demonstration of an antalgic gait pattern.²⁷ With subjective findings including a rapid increase in weightbearing activities or excessive steroid or alcohol use, a femoral stress fracture or avascular necrosis of the hip, respectively, should be considered. If the objective examination is consistent with either of these diagnoses and an immediate referral to a physician is warranted. A summary of differential diagnoses can be found in [Table 1](#).

While special tests are frequently utilized in the clinical setting to identify GTPS, there is currently no test or cluster

of tests utilized to identify which *specific* pathoanatomical structures are involved or the severity to which they are involved. Some frequently utilized special tests include resisted hip abduction and external rotation, FABER, the resisted external de-rotational, and Trendelenburg tests. Ganderton et al.²⁸ found the FABER, resisted hip abduction, and the resisted external de-rotational tests to have the highest diagnostic test accuracy in confirming GTPS. Previous research on the Trendelenburg test and the resisted external rotation test has reported sensitivity and specificity values in individuals with GTPS of 73% and 77% and 88% and 97.3%, respectively.^{4,29} Multiple authors have also identified palpation of the greater trochanter and trochanteric bursae to be a useful tool in confirming the presence of GTPS.^{5,17,28} It should be noted, however, that no studies report sensitivity and specificity values for these palpatory tests in identifying the specific pathoanatomical structure involved. Therefore, it should be emphasized that these tests have minimal utility beyond confirming the presence or absence of GTPS and offer little to the clinician in prescribing appropriate treatment interventions.

Given the limited clinical utility of special tests, clinicians should focus on examination procedures that identify the impairments responsible for the development and persistence of GTPS. Previous research has established hip abductor weakness, loss of pelvic control in the frontal plane, and iliotibial band tightness and thickening as factors increasing compressive forces through the bursae and GMed and GMin tendons.^{10,11} Identifying impairments in hip flexibility and strength will assist with identifying individuals who will benefit from stretching and strengthening exercises and/or manual interventions. Along with standard range of motion, strength, and flexibility screens, a thorough analysis of the movement system should be performed. Movement system assessments are evaluations of biomechanics during functional activities providing insight into faulty movement patterns allowing clinicians to tailor treatment interventions to address identified impairments.³⁰

IMPAIRMENT BASED TREATMENT CLASSIFICATION

Since the mechanism for developing GTPS has been identified as strength and biomechanical deficits of and around the hip, an impairment-based treatment classification, similar to those described for the low back, may be useful to help appropriately tailor treatment interventions.^{10,11} We recommend classifying patients presenting with GTPS into two categories, contractile and/or non-contractile, and then further sub-classifying based on irritability.

CONTRACTILE

A patient with a contractile presentation would report a reproduction of pain with resisted hip abduction, lengthening or stretching of the hip abductors, palpation of the gluteal muscles and/or tendons, and would demonstrate faulty mechanics during functional activities. Any of these findings would suggest the gluteal muscles, proximal iliotibial band,

and/or tensor fascia latae are involved. However, it does not indicate which specific structure or the severity of involvement. Therefore, we recommend identifying the irritability of muscular involvement as high or low before prescribing interventions. Treatment for a contractile presentation should focus on muscle and tendon healing through soft tissue mobilization (STM) and graded loading exercises. In patients demonstrating severe pain, profound weakness, and a significant limited ability to walk without an assistive device, a physician referral is warranted to rule out a complete tear of the hip abductors.

HIGH IRRITABILITY

Highly irritable individuals in the contractile classification will be unable to abduct their hip against gravity and report moderate to severe reproduction of pain. The patient may also report severe difficulty with functional tasks involving single leg stance (SLS) and exhibit an antalgic gait. During this phase, STM to the gluteal muscle bellies, tensor fascia lata (TFL), and/or iliotibial band (ITB) are recommended to facilitate healing. STM of the gluteal tendons and trochanteric bursae should be avoided to limit further tissue irritation. GMed and GMin strengthening should be initiated through isometric contractions in gravity eliminated positions progressing to against gravity positions as tolerated. Low intensity isometrics have been shown to induce analgesic effects in individuals with patellar tendinopathy, however, these results are variable across populations.³¹ We therefore recommend a graded exercise program progressing from isometric contractions in the high irritability phase progressing to isotonic contractions in the low irritability phase. Core exercise should be initiated in supine and quadruped with a focus on proper activation of the lower abdominals. During this phase, all activities should be pain-free. Examples of gluteal and core exercise progressions for the high irritability phase can be found in Figures 1 and 2.

Along with a tailored exercise program, extensive education should be provided regarding activity and postural modifications with the goal of reducing load and compression through the lateral hip. Strategies should be given to help minimize time spent on a single leg during activities of daily living, such as sitting rather than standing to put on pants thereby reducing tensile load through the gluteal muscles. The importance of avoiding hip adduction should be stressed for sitting, standing, and sleeping to limit excessive compression through the lateral hip.¹² For example, education should be provided to avoid crossing legs in both sitting and standing and to sleep in supine with a pillow under their knees or in sidelying with a pillow between their legs (Figure 3).

LOW IRRITABILITY

Those with low irritability of the lateral hip musculature will be able to abduct their hip against gravity, but continue to complain of mild to moderate reproduction of pain. Treatment recommendations for this phase include initiation of pain-free stretching of the hip abductors, concentric and eccentric hip abductor strengthening, functional train-

ing focusing on single leg activities, and progression of core strengthening. STM should continue to be utilized in conjunction with pain-free stretching if tightness was identified through muscle length testing, such as an Ober's test. It should be noted that soft-tissue mobilization should be directed at the gluteal muscle bellies, TFL, and/or ITB, not at the trochanteric bursae or gluteal tendon insertions, to avoid further tissue irritation.¹⁹ GMed and GMin strengthening should be progressed to include isotonic contractions in the low irritability phase as the focus shifts from pain control and healing to restoration of function. Heavy slow resistance exercise has been shown to improve tendon quality and compared to eccentric exercise, heavy slow resistance exercise yielded similar clinical outcomes.³²⁻³⁴ We therefore recommend a combination of concentric and eccentric exercise as per pain tolerance. A large focus should be given to weight bearing hip abductor strengthening as weight bearing abductor exercises demonstrate significantly greater EMG activity than all non-weight bearing exercise, except sidelying hip abduction.³⁵ If tolerated, a resistance band should be incorporated to increase hip abductor activation and challenge lower extremity neuromuscular control. A GMed and GMin strengthening progression for the low irritability phase can be found in [Figure 4](#). Core exercises should be progressed to incorporate a plank progression as well as weight bearing functional tasks. Careful attention should be paid to frontal plane control of the pelvis during all weight bearing strengthening exercises.

The main goal of this phase is to restore appropriate pelvic control during functional tasks. Similar to the strengthening progression, functional training should begin in double leg stance progressing to the transition from double to SLS and finally to training on a single leg. In order to challenge neuromuscular control, dynamic functional training should be incorporated once the patient demonstrates mastery of static tasks. We recommend beginning the return to activity and sport progression with a walking program to incorporate progressive loading of the gluteal muscles as well as to improve cardiovascular endurance. Once the patient can ambulate 15-minutes with minimal pain and appropriate pelvic control, progression to sport-specific activities including running, jumping, hopping, and cutting should be initiated, if necessary. Progression should be dictated by pain and frontal plane control of the pelvis.

NON-CONTRACTILE

Individuals exhibiting pain during palpation of or around the greater trochanter, decreased strength of the hip abductors without reproduction of pain during resisted testing, and/or faulty mechanics during movement system evaluations would suggest a non-contractile source of GTPS. It should be emphasized that while these individuals may demonstrate weakness and functional deficiency of the hip abductors, these activities will not elicit pain. Clinicians should also evaluate for any cardinal signs of inflammation, including rubor, erythema, edema, and tenderness, which indicate involvement of the trochanteric bursae.^{36,37} Different from a contractile presentation, which focuses muscle and tendon healing through loading exercise, primary




Figure 1. Hip abductor isometric exercise (with or without resistance band) in a) short lever arm, gravity eliminated position and b) long lever arm, gravity eliminated position, c) short lever arm against gravity with towel roll to avoid hip abduction, and d) long lever arm against gravity with heel against the wall to avoid compensation.

interventions for a non-contractile presentation are aimed at reducing compressive forces over the greater trochanter and alleviating bursal irritation and overload of the lateral structures of the hip. Treatment should focus on decreasing inflammation, restoring flexibility of the lateral hip, particularly of the iliotibial band, improving strength of the hip abductors, and improving of functional movement patterns. We again recommend classifying the patient into either high or low irritability to better tailor treatment interventions.

HIGH IRRITABILITY

A patient with a highly irritable, non-contractile source of GTPS will exhibit severe signs inflammation. Modalities to decrease inflammation, including heat, ice, ultrasound, low-level laser therapy, shockwave therapy, and electrical stimulation can be utilized.^{38,39} Gentle soft-tissue mobilization of the structures surrounding the greater trochanter is useful in improving flexibility of the lateral hip. Clinicians should avoid any STM of or in close proximity to the trochanteric bursae as this would increase bursal irritation. Much like the highly irritable contractile presentation, hip abductor strengthening should begin with isometric exercise progressing to concentric exercise only as tolerated by pain. To progressively load the hip abductors and endure proper control of the pelvis, we recommend beginning abductor strengthening, core strengthening, and functional training in a non-weight bearing position and progressing to weight bearing as strength and movement patterns improve.

LOW IRRITABILITY

A patient with low-irritability and a non-contractile presentation will exhibit mild signs of inflammation, however, they may continue to report reproduction of pain with pal-



Figure 2. Transverse abdominis activation exercise with a) alternating upper extremity movement, b) bilateral upper extremity movement, and c) isometric lower extremity hold with alternating upper extremity movement.

pation of the greater trochanter and trochanteric bursae. Treatment principles should mirror those described for the low irritability, contractile presentation. Eccentric strengthening of the abductors, high-level core strengthening, and functional training including sport-specific tasks should be prescribed with the goal of returning the patient to their previous level of functioning.

ABNORMAL FRONTAL PLANE MOVEMENT SYSTEM DYSFUNCTION DURING SINGLE LEG TASKS

Development of GTPS is thought to result from faulty mechanics during functional activities, mainly the loss of pelvic control in the frontal plane secondary to hip abductor weakness or pain with hip abductor activation.^{10,11} The GMed and GMin are responsible for maintaining pelvic control in the frontal plane during single leg weight bearing activities and are the tendons most implicated in GTPS.⁴⁰ From the bottom up, these hip abductors also control the hip adduction moment resulting from the external ground reaction force medial to the hip.⁴¹ Any deviation in frontal plane control, due to either pain or weakness, increases the compressive forces of the GMed and GMin tendons on the greater trochanter and trochanteric bursae as well as increases the tension through the iliotibial band. Individuals with GTPS frequently complain of pain and difficulty while completing tasks requiring single leg support as these tasks stress the hip abductors. Given the large role faulty biomechanics plays in the development of GTPS, clinicians must thoroughly evaluate movement patterns through movement system dysfunction assessments. Movement system dysfunction assessments allow for identification of biomechanical deviations during functional activities. In the setting of GTPS, clinicians should evaluate pelvic control during a 30-second SLS, stair ascent, and the stance phase of the gait cycle.

SINGLE LEG STANCE

Assessing pelvic control during a 30-second SLS provides information on the function of the GMed and GMin as they are responsible for maintaining a level pelvis during single




Figure 3. Sleeping modifications in a) supine and b) sidelying




Figure 4. GMed and GMin strengthening progression (with or without resistance band) for the low irritability phase including a) bridge, b) single leg bridge, c) standing hip abduction, and d) side stepping.

leg activities.^{28,40,41} The SLS task should be broken down into the transition phase, going from double to SLS, and the stance phase, when only one foot is on the ground. The transition phase requires hip abductor activation to assist with the lateral translation of the pelvic girdle while stance phase requires the hip abductors to control pelvic tilt.⁴² In the setting of hip abductor weakness or pain with activa-

tion of the hip abductors, an individual may exhibit a Trendelenburg sign, meaning the contralateral hip will drop towards the floor, or a compensated Trendelenburg, meaning the patients trunk leans ipsilaterally to the stance leg (Figure 7).^{43,44} Allison et al.⁴² found individuals with gluteal tendinopathy exhibit greater hip adduction and ipsilateral pelvic shift in preparation for leg lift and greater hip adduction and less control over the pelvis during the stance phase. Other signs of hip abductor dysfunction include the need for upper or lower extremity support during the SLS task. To get a complete picture of SLS performance, a contralateral SLS should be assessed to highlight side-to-side differences in pelvic control.

STAIR ASCENT

The ability to negotiate stairs is critical for maintaining independence and is required to complete many activities of daily living. Compared to walking on level ground, the strength required to negotiate stairs is significantly higher.⁴⁵⁻⁴⁷ Individuals with GTPS frequently complain of pain during stair ascent or the inability to complete the stair task, greatly reducing their quality of life. During stair ascent, the hip abductors are responsible for maintaining a level pelvis as the body moves vertically to reach the next stair.⁴⁵ When compared to level ground walking, stair ascent requires greater hip flexion, hip adduction, and knee internal rotation moments which in turn increases the tension through the iliotibial band. Additionally, stair ascent requires activation of the tensor fascia lata, gluteus maximus, and the vastus lateralis, all which have attachments to the iliotibial band.⁴⁸⁻⁵⁰ This increase in tension on the iliotibial band leads to even greater compression over the greater trochanter.^{45,46} Allison et al.⁵¹ reported individuals with gluteal tendinopathy were 4.5 times more likely to exhibit a large hip adduction moment and greater pelvic translation at heel strike during stair ascent than healthy controls.⁵¹ Additionally, the presence of a contralateral trunk lean was observed in those with gluteal tendinopathy.⁵¹ When assessing the movement system during stair ascent, clinicians should evaluate the frontal plane mechanics during the stance phase, meaning when the lower extremity being assessed is on the ground. Poor performance of the hip abductors during stair ascent can manifest through a pelvic drop, pelvic translation, and/or trunk lean. The individual may also exhibit a step to gait pattern or excessive use of upper extremity support on a handrail.

GAIT

SLS comprises 40% of the gait cycle, potentially aggravating the symptoms of GTPS as there is a higher load through the hip abductor tendons compared to the swing phase.^{19,30} Compared to healthy controls, those with gluteal tendinopathy were found to have a greater hip adduction moment during the stance phase.⁵² A Trendelenburg gait pattern, with or without the complaint of pain, may be an indication of gluteal dysfunction. In the case of a mild gait deviation, a 30-second SLS test should also be performed in order to better assess the functional performance and endurance of the abductors. With severe gluteal dysfunction, a




Figure 5. Core progression for the low irritability including a) bird dog, b) transverse abdominis activation on physioball with alternating upper and lower extremity movement modified plank, c) modified plank, and d) modified side plank.




Figure 6. Functional squat progression (with or without resistance band from a) bilateral stance, b) staggered stance, and c) unilateral stance to improve lower extremity neuromuscular control and hip abductor strength.




Figure 7. Single leg stance with a) normal frontal plane mechanics, (b) a Trendelenburg, and (c) a compensated Trendelenburg.

compensatory trunk lean to balance out the center of gravity may be observed.⁵³

Table 2. Impairment-Based Treatment Classification for GTPS

CLASSIFICATION			
Contractile		Non-Contractile	
<ul style="list-style-type: none"> • Pain with palpation of the muscle belly or tendon of the GMed or GMin • Pain with palpation of the proximal iliotibial band • Pain and weakness during strength testing of the hip abductors • Decreased flexibility of the iliotibial band • Poor frontal plane control and reproduction of pain during single leg activities 		<ul style="list-style-type: none"> • Signs of inflammation including rubor, erythema, oedema, and tenderness • Pain with palpation of the greater trochanter or trochanteric bursae • Weakness of the hip abductors without reproduction of pain • Decreased flexibility of the iliotibial band • Poor frontal plane control during single leg activities without reproduction of pain 	
↓		↓	
High Irritability	Low Irritability	High Irritability	Low Irritability
<ul style="list-style-type: none"> • Unable to abduct hip against gravity with moderate to severe pain • Poor frontal plane control and difficulty completing single leg tasks due to moderate to severe pain • Antalgic gait secondary to pain 	<ul style="list-style-type: none"> • Able to abduct hip against gravity with mild to moderate pain • Poor frontal plane control and difficulty completing single leg tasks due to mild pain 	<ul style="list-style-type: none"> • Severe rubor, erythema, oedema, and tenderness • Unable to abduct hip against gravity with no reproduction of pain • Poor frontal plane control during single leg tasks 	<ul style="list-style-type: none"> • Mild rubor, erythema, oedema, and tenderness • Able to abduct hip against gravity with no reproduction of pain • Poor frontal plane control during single leg tasks
↓		↓	
TREATMENT			
Goal: Muscle and tendon healing		Goal: Reducing inflammation	
<ul style="list-style-type: none"> • STM of the hip abductors and iliotibial band to facilitate tendon healing • Submaximal isometric hip abductor exercises • Initiation of core strengthening 	<ul style="list-style-type: none"> • Heavy loading and eccentric strengthening of the hip abductors • Progression of core strengthening • Functional training and return to sport 	<ul style="list-style-type: none"> • Modalities to reduce inflammation • STM of the hip abductors and iliotibial band to improve flexibility • Stretching of the lateral hip 	<ul style="list-style-type: none"> • Pain free stretching of the lateral hip • Concentric and eccentric strengthening of the hip abductors • Core strengthening • Functional training and return to sport

In order to appropriately tailor treatment interventions, findings of the physical examination and movement assessments should be utilized to appropriately categorize the patient into a contractile or non-contractile presentation. Impairment-based treatment can then be utilized based on irritability to address muscle strength, flexibility deficits, and motor control impairments during functional activities. The goal of treatment is to improve motor patterns to reduce the compression of the gluteal tendons on the trochanteric bursae and greater trochanter and ultimately reduce pain and improve function. A summary of the recommended treatment algorithm for GTPS can be found in [Table 2](#).

CONCLUSION

GTPS is a term utilized to describe pain in the lateral hip and can include external snapping hip (coxa saltans), trochanteric bursitis, and gluteal tearing and tendinopathy.^{1,2} Individuals suffering from GTPS complain of lateral hip, thigh or buttock pain that worsens with single leg ac-

tivities and lying on the affected side. Currently, there are no tests or cluster of tests to identify the involved pathoanatomical structure or the severity of dysfunction. Therefore, an impairment-based treatment classification system is recommended, delineating patients into two categories: contractile or non-contractile. Once the patient is categorized into contractile or non-contractile based on physical examination findings and assessments of the movement system, impairment-based treatments can be utilized to appropriately tailor interventions to improve functional outcomes. Treatment of contractile sources of GTPS should focus on muscle and tendon healing through STM and loading exercises whereas treatment of non-contractile sources of GTPS should focus on reducing compressive forces over the greater trochanter and alleviating bursal irritation and overload of the lateral structures of the hip.

Submitted: October 19, 2021 CDT, Accepted: January 09, 2022 CDT



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

REFERENCES

1. Gordon EJ. Trochanteric bursitis and tendinitis. *Clin Orthop*. 1961;20:193-202.
2. Strauss EJ, Nho SJ, Kelly BT. Greater trochanteric pain syndrome. *Sports Med Arthrosc Rev*. 2010;18(2):113-119. doi:10.1097/JSA.0b013e3181e0b2ff
3. Barratt PA, Brookes N, Newson A. Conservative treatments for greater trochanteric pain syndrome: a systematic review. *Br J Sports Med*. 2017;51(2):97-104. doi:10.1136/bjsports-2015-095858
4. Bird PA, Oakley SP, Shnier R, Kirkham BW. Prospective evaluation of magnetic resonance imaging and physical examination findings in patients with greater trochanteric pain syndrome. *Arthritis Rheum*. 2001;44(9):2138-2145.
5. Grimaldi A, Mellor R, Nicolson P, Hodges P, Bennell K, Vicenzino B. Utility of clinical tests to diagnose MRI-confirmed gluteal tendinopathy in patients presenting with lateral hip pain. *Br J Sports Med*. 2017;51(6):519-524. doi:10.1136/bjsports-2016-096175
6. Hancock M. Approach to low back pain - physiotherapy. *Aust Fam Physician*. 2014;43(3):117-118.
7. Koes BW, van Tulder M, Lin CW, Macedo LG, McAuley J, Maher C. An updated overview of clinical guidelines for the management of non-specific low back pain in primary care. *Eur Spine J*. 2010;19(12):2075-2094. doi:10.1007/s00586-010-1502-y
8. van Tulder M, Becker A, Bekkering T, et al. Chapter 3. European guidelines for the management of acute nonspecific low back pain in primary care. *Eur Spine J*. 2006;15 Suppl 2:S169-91. doi:10.1007/s00586-006-1071-2
9. Hancock MJ, Maher CG, Latimer J, et al. Systematic review of tests to identify the disc, SIJ or facet joint as the source of low back pain. *Eur Spine J*. 2007;16(10):1539-1550. doi:10.1007/s00586-007-0391-1
10. Mellor R, Bennell K, Grimaldi A, et al. Education plus exercise versus corticosteroid injection use versus a wait and see approach on global outcome and pain from gluteal tendinopathy: prospective, single blinded, randomised clinical trial. *Br J Sports Med*. 2018;52(22):1464-1472. doi:10.1136/bjsports-2018-k1662rep
11. Khoury AN, Brooke K, Helal A, et al. Proximal iliotibial band thickness as a cause for recalcitrant greater trochanteric pain syndrome. *J Hip Preserv Surg*. 2018;5(3):296-300. doi:10.1093/jhps/hny025
12. Grimaldi A, Fearon A. Gluteal Tendinopathy: Integrating Pathomechanics and Clinical Features in Its Management. *J Orthop Sports Phys Ther*. 2015;45(11):910-922. doi:10.2519/jospt.2015.5829
13. Lievense A, Bierma-Zeinstra S, Schouten B, Bohnen A, Verhaar J, Koes B. Prognosis of trochanteric pain in primary care. *Br J Gen Pract*. 2005;55(512):199-204.
14. Long SS, Surrey DE, Nazarian LN. Sonography of greater trochanteric pain syndrome and the rarity of primary bursitis. *AJR Am J Roentgenol*. 2013;201(5):1083-1086. doi:10.2214/AJR.12.10038
15. Shbeeb MI, Matteson EL. Trochanteric bursitis (greater trochanter pain syndrome). *Mayo Clin Proc*. 1996;71(6):565-569. doi:10.4065/71.6.565
16. Tortolani PJ, Carbone JJ, Quartararo LG. Greater trochanteric pain syndrome in patients referred to orthopedic spine specialists. *Spine J*. 2002;2(4):251-254. doi:10.1016/s1529-9430(02)00198-5
17. Woodley SJ, Nicholson HD, Livingstone V, et al. Lateral hip pain: findings from magnetic resonance imaging and clinical examination. *J Orthop Sports Phys Ther*. 2008;38(6):313-328. doi:10.2519/jospt.2008.2685
18. Raman D, Haslock I. Trochanteric bursitis--a frequent cause of "hip" pain in rheumatoid arthritis. *Ann Rheum Dis*. 1982;41(6):602-603. doi:10.1136/ard.41.6.602
19. Segal NA, Felson DT, Torner JC, et al. Greater trochanteric pain syndrome: epidemiology and associated factors. *Arch Phys Med Rehabil*. 2007;88(8):988-992. doi:10.1016/j.apmr.2007.04.014
20. Anderson K, Strickland SM, Warren R. Hip and groin injuries in athletes. *Am J Sports Med*. 2001;29(4):521-533. doi:10.1177/03635465010290042501
21. Clancy WG. Runners' injuries. Part two. Evaluation and treatment of specific injuries. *Am J Sports Med*. 1980;8(4):287-289. doi:10.1177/036354658000800415

22. Del Buono A, Papalia R, Khanduja V, Denaro V, Maffulli N. Management of the greater trochanteric pain syndrome: a systematic review. *Br Med Bull*. 2012;102:115-131. doi:10.1093/bmb/ldr038
23. Grimaldi A, Mellor R, Hodges P, Bennell K, Wajswelner H, Vicenzino B. Gluteal tendinopathy: a review of mechanisms, assessment and management. *Sports Med*. 2015;45(8):1107-1119. doi:10.1007/s40279-015-0336-5
24. Fearon AM, Scarvell JM, Cook JL, Smith PN. Does ultrasound correlate with surgical or histologic findings in greater trochanteric pain syndrome? a pilot study. *Clin Orthop Relat Res*. 2010;468(7):1838-1844. doi:10.1007/s11999-009-1174-2
25. Silva F, Adams T, Feinstein J, Arroyo RA. Trochanteric bursitis: refuting the myth of inflammation. *J Clin Rheumatol*. 2008;14(2):82-86. doi:10.1097/RHU.0b013e31816b4471
26. Cibulka MT, White DM, Woehrle J, et al. Hip pain and mobility deficits--hip osteoarthritis: clinical practice guidelines linked to the international classification of functioning, disability, and health from the orthopaedic section of the American Physical Therapy Association. *J Orthop Sports Phys Ther*. 2009;39(4):A1-25. doi:10.2519/jospt.2009.0301
27. Takla A, O'Donnell J, Voight M, et al. The 2019 International Society of Hip Preservation (ISHA) physiotherapy agreement on assessment and treatment of femoroacetabular impingement syndrome (FAIS): an international consensus statement. *J Hip Preserv Surg*. 2020;7(4):631-642. doi:10.1093/jhps/hnaa043
28. Ganderton C, Semciw A, Cook J, Pizzari T. Demystifying the Clinical Diagnosis of Greater Trochanteric Pain Syndrome in Women. *J Womens Health (Larchmt)*. 2017;26(6):633-643. doi:10.1089/jwh.2016.5889
29. Lequesne M, Djian P, Vuillemin V, Mathieu P. Prospective study of refractory greater trochanter pain syndrome. MRI findings of gluteal tendon tears seen at surgery. Clinical and MRI results of tendon repair. *Joint Bone Spine*. 2008;75(4):458-464. doi:10.1016/j.jbspin.2007.12.004
30. Fearon AM, Cook JL, Scarvell JM, Neeman T, Cormick W, Smith PN. Greater trochanteric pain syndrome negatively affects work, physical activity and quality of life: a case control study. *J Arthroplasty*. 2014;29(2):383-386. doi:10.1016/j.arth.2012.10.016
31. Clifford C, Challoumas D, Paul L, Syme G, Millar NL. Effectiveness of isometric exercise in the management of tendinopathy: a systematic review and meta-analysis of randomised trials. *BMJ Open Sport Exerc Med*. 2020;6(1):e000760. doi:10.1136/bmjsem-2020-000760
32. Kongsgaard M, Kovanen V, Aagaard P, et al. Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy. *Scand J Med Sci Sports*. 2009;19(6):790-802. doi:10.1111/j.1600-0838.2009.00949.x
33. Beyer R, Kongsgaard M, Hougs Kjaer B, Ohlenschlaeger T, Kjaer M, Magnusson SP. Heavy slow resistance versus eccentric training as treatment for achilles tendinopathy: a randomized controlled trial. *Am J Sports Med*. 2015;43(7):1704-1711. doi:10.1177/0363546515584760
34. Kongsgaard M, Qvortrup K, Larsen J, et al. Fibril morphology and tendon mechanical properties in patellar tendinopathy: effects of heavy slow resistance training. *Am J Sports Med*. 2010;38(4):749-756. doi:10.1177/0363546509350915
35. Bolgla LA, Uhl TL. Electromyographic analysis of hip rehabilitation exercises in a group of healthy subjects. *J Orthop Sports Phys Ther*. 2005;35(8):487-494. doi:10.2519/jospt.2005.35.8.487
36. Shbeeb MI, O'Duffy JD, Michet, C. J., Jr., O'Fallon WM, Matteson EL. Evaluation of glucocorticosteroid injection for the treatment of trochanteric bursitis. *J Rheumatol*. 1996;23(12):2104-2106.
37. Williams BS, Cohen SP. Greater trochanteric pain syndrome: a review of anatomy, diagnosis and treatment. *Anesth Analg*. 2009;108(5):1662-1670. doi:10.1213/ane.0b013e31819d6562
38. Lustenberger DP, Ng VY, Best TM, Ellis TJ. Efficacy of treatment of trochanteric bursitis: a systematic review. *Clin J Sport Med*. 2011;21(5):447-453. doi:10.1097/ISM.0b013e318221299c
39. Reid D. The management of greater trochanteric pain syndrome: a systematic literature review. *J Orthop*. 2016;13(1):15-28. doi:10.1016/j.jor.2015.12.006
40. Al-Hayani A. The functional anatomy of hip abductors. *Folia Morphol (Warsz)*. 2009;68(2):98-103.
41. Yoon YS, Mansour JM. The passive elastic moment at the hip. *J Biomech*. 1982;15(12):905-910. doi:10.1016/0021-9290(82)90008-2

42. Allison K, Bennell KL, Grimaldi A, Vicenzino B, Wrigley TV, Hodges PW. Single leg stance control in individuals with symptomatic gluteal tendinopathy. *Gait Posture*. 2016;49:108-113. doi:10.1016/j.gaitpost.2016.06.020
43. Hardcastle P, Nade S. The significance of the Trendelenburg test. *J Bone Joint Surg Br*. 1985;67(5):741-746. doi:10.1302/0301-620X.67B5.4055873
44. Youdas JW, Mraz ST, Norstad BJ, Schinke JJ, Hollman JH. Determining meaningful changes in pelvic-on-femoral position during the Trendelenburg test. *J Sport Rehabil*. 2007;16(4):326-335. doi:10.1123/jsr.16.4.326
45. Nadeau S, McFadyen BJ, Malouin F. Frontal and sagittal plane analyses of the stair climbing task in healthy adults aged over 40 years: what are the challenges compared to level walking? *Clin Biomech*. 2003;18(10):950-959. doi:10.1016/s0268-0033(03)00179-7
46. Protopapadaki A, Drechsler WI, Cramp MC, Coutts FJ, Scott OM. Hip, knee, ankle kinematics and kinetics during stair ascent and descent in healthy young individuals. *Clin Biomech (Bristol, Avon)*. 2007;22(2):203-210. doi:10.1016/j.clinbiomech.2006.09.010
47. McFadyen BJ, Winter DA. An integrated biomechanical analysis of normal stair ascent and descent. *J Biomech*. 1988;21(9):733-744. doi:10.1016/0021-9290(88)90282-5
48. Birnbaum K, Pandorf T. Finite element model of the proximal femur under consideration of the hip centralizing forces of the iliotibial tract. *Clin Biomech*. 2011;26(1):58-64. doi:10.1016/j.clinbiomech.2010.09.005
49. Birnbaum K, Siebert CH, Pandorf T, Schopphoff E, Prescher A, Niethard FU. Anatomical and biomechanical investigations of the iliotibial tract. *Surg Radiol Anat*. 2004;26(6):433-446. doi:10.1007/s00276-004-0265-8
50. Dwek J, Pfirrmann C, Stanley A, Pathria M, Chung CB. MR imaging of the hip abductors: normal anatomy and commonly encountered pathology at the greater trochanter. *Magn Reson Imaging Clin N Am*. 2005;13(4):691-704, vii. doi:10.1016/j.mric.2005.08.004
51. Allison K, Vicenzino B, Bennell KL, Wrigley TV, Grimaldi A, Hodges PW. Kinematics and kinetics during stair ascent in individuals with Gluteal Tendinopathy. *Clin Biomech*. 2016;40:37-44. doi:10.1016/j.clinbiomech.2016.10.003
52. Allison K, Wrigley TV, Vicenzino B, Bennell KL, Grimaldi A, Hodges PW. Kinematics and kinetics during walking in individuals with gluteal tendinopathy. *Clin Biomech*. 2016;32:56-63. doi:10.1016/j.clinbiomech.2016.01.003
53. Gandbhir VN, Lam JC, Rayi A. Trendelenburg Gait. In: *StatPearls*. StatPearls Publishing; 2021.