

Original Article

Effect of footwear on standing balance in healthy young adult males

Ahmad H. Alghadir, Hamayun Zafar, Shahnawaz Anwer

Department of Rehabilitation Sciences, College of Applied Medical Sciences, King Saud University, Riyadh, Saudi Arabia

Abstract

Objective: The present study aimed to evaluate the effect of footwear on standing balance in healthy young adult males. **Methods:** Thirty healthy male participants aged 20-30 years were tested for standing balance on the Balance Master on three occasions, including wearing a sandal, standard shoe, or no footwear (barefoot). The tests of postural stability include; "Modified Clinical Test of Sensory Interaction on Balance" (mCTSIB), "Unilateral Stance" (US), and the "Limits of Stability" (LOS). The balance scores (mCTSIB, US, and LOS) was analyzed. **Results:** There was a significant effect between footwear conditions for mCTSIB with eye closed on a firm surface ($p=0.002$). There was a significant effect between footwear conditions for the US with eye open and closed ($p<0.05$). There was a significant effect between footwear conditions for LOS reaction time during forward movement ($p=0.02$). Similarly, there was a significant effect between footwear conditions for LOS reaction time during left side movement ($p=0.01$). **Conclusions:** Wearing sandals compared to bare feet significantly increased postural sway and reduced stability in healthy young adult males. However, wearing a standard shoe compared to bare feet did not significantly affect balance scores in standing.

Keywords: Balance, Footwear, Postural Sway, Barefoot, Stability

Introduction

Footwear has a vital role in improving well-being of any individuals. Previous study reported an association between footwear and disorders; therefore, considering footwear characteristics during management program is important¹. Footwear facilitates sensory information to the foot and control postural stability through the touch and proprioceptive system^{2,3}. The tactile stimulation is detected by the cutaneous mechanoreceptors of the plantar surface of the feet and gives information of plantar pressure distribution to the central nervous system². Previous study reported an impaired standing balance, if plantar surface afferents are not intact⁴. Whereas, stimulation of the cutaneous

mechanoreceptors can enhance postural stability⁵⁻⁷. The type of Footwear and their modifications, including foot orthoses, shoe inserts, and insoles can stimulate tactile as well as proprioceptive systems^{8,9}. In addition, previous study reported increased thresholds of plantar cutaneous vibrotactile particularly in the elderly^{7,10}.

The maintenance of static and dynamic balance is a vital to reduce risk of injuries in any sports. Type of footwear may influence the sensory feedback quality from the feet¹¹. Footwear often designed to give support and stability to the foot, therefore, potentially affecting balance and function of the foot¹. However, previous study reported 42% of fall in a group of 106 older people during wearing walking shoes¹². Another study reported postural instability in the elderly

The authors have no conflict of interest.

Corresponding author: Shahnawaz Anwer, MPT, Researcher, Department of Rehabilitation Sciences, CAMS, King Saud University, Riyadh, KSA
E-mail: anwer_shahnawazphysio@rediffmail.com

Edited by: F. Rauch

Accepted 7 August 2017

Abbreviations

mCTSIB = Modified Clinical Test of Sensory Interaction on Balance

US = Unilateral Stance

LOS = Limits of Stability

COG = Center of Gravity



Table 1. Comparison of balance scores between a sandal, standard shoe, or no footwear (barefoot) conditions.

Variables	Barefoot (BF) Mean ± SD	Standard shoe (SS) Mean ± SD	Sandal (S) Mean ± SD	ANOVA		BF vs SS	BF vs S	SS vs S
				F	p	p	p	p
mCTIB Firm - EO (deg/sec)	0.38 ± 0.19	0.41 ± 0.38	0.49 ± 0.14	1.784	0.17			
mCTIB Firm - EC (deg/sec)	0.31 ± 0.07	0.35 ± 0.18	0.52 ± 0.36	6.861	0.002	0.99	0.002	0.02
mCTIB Foam - EO (deg/sec)	0.67 ± 0.14	0.73 ± 0.18	0.78 ± 0.17	3.349	0.04	0.49	0.034	0.73
mCTIB Foam - EC (deg/sec)	1.29 ± 0.37	1.35 ± 0.40	1.62 ± 0.36	6.295	0.003	0.97	0.004	0.02
US - EO Left (deg/sec)	0.90 ± 0.21	0.98 ± 0.45	1.23 ± 0.65	3.904	0.02	0.89	0.03	0.14
US - EC Left (deg/sec)	2.46 ± 0.92	2.62 ± 0.80	3.08 ± 1.06	3.533	0.03	0.89	0.04	0.18
US - EO Right (deg/sec)	0.89 ± 0.25	1.08 ± 0.46	1.34 ± 0.88	4.293	0.02	0.70	0.01	0.27
US - EC Right (deg/sec)	2.54 ± 0.92	2.65 ± 0.94	3.17 ± 1.22	3.181	0.05	0.95	0.04	0.16
LOS - RT Forward (sec)	0.89 ± 0.30	0.89 ± 0.28	1.10 ± 0.36	4.344	0.02	0.88	0.04	0.04
LOS - RT Right (sec)	0.83 ± 0.29	0.87 ± 0.30	0.97 ± 0.28	2.028	0.14			
LOS - RT Back (sec)	0.77 ± 0.28	0.78 ± 0.32	0.90 ± 0.26	1.900	0.16			
LOS - RT Left (sec)	0.79 ± 0.27	0.88 ± 0.26	0.99 ± 0.28	4.429	0.01	0.56	0.01	0.31

mCTIB, Modified Clinical Test of Sensory Interaction on Balance; US, Unilateral Stance; EO, Eye open; EC, Eye closed; LOS, Limits of Stability; RT, Reaction time.

population who has poor footwear type and poor footwear characteristics¹³. In addition, Keegan et al.¹⁴ reported an increased risk of a foot fracture from a fall in individuals using slip-on shoes and sandals. Furthermore, Murphy et al.¹⁵ had suggested that the shoes were an important element in the development of human posture. Therefore, the present study aimed to evaluate the effect of footwear on standing balance in healthy young adult males.

Materials and methods

Participants

Thirty healthy adult male aged older than 18 years were recruited from the college of applied medical sciences, King Saud University, Riyadh. The present study was approved by the institution ethics committee, Rehabilitation Research Chair, King Saud University, Riyadh, Saudi Arabia. Each participant provided a written informed consent prior to the experiment. Individuals were excluded if they had a history of neurological disorders that can affect balance.

Procedures

Participants' demographic data such as age, height, weight, and body mass index (BMI) were recorded prior to testing. The Balance Master (NeuroCom Balance Master®; Natus Medical Incorporated, CA, USA) was used to measure the Postural stability¹⁶. The Balance Master utilizes a force platform to locate the center of gravity (COG) and provides various tasks for challenging balance. For all the tests, the participants were tested with three occasions, including wearing a sandal, standard shoe, and no footwear (barefoot). The tests of postural stability include;

“Modified Clinical Test of Sensory Interaction on Balance” (mCTSIB), “Unilateral Stance” (US), and the “Limits of Stability” (LOS). The mCTSIB includes 4 conditions: eye open on the firm surface, eye closed on the firm surface, eyes open on the foam, and eyes closed on the foam¹⁷. The mCTSIB tests measure subject's “sway velocity” (degree/sec) from 3 trials of 20 s duration for each of the four given conditions¹⁸. The US assessed the postural stability during single leg standing (right and left) in 2 conditions: eye open and eye closed. The US test measure subject's “sway velocity” (degree/sec) from 3 trials of 20 s duration for each of the given conditions^{19,20}. The LOS assessed the subject's stability limits without losing balance in a given directions. The LOS measures the “movement reaction time” (sec) and “movement velocity” (degrees/sec) from 3 trials of 20 s duration for given directions^{21,22}. In all the test conditions, the participants asked to stand in the marked position on the Balance master, with their hands positioned on the iliac crests²³. Participants had given a 15-second rest period between each trial and asked to perform a 5-m walk between each test condition. The order of all the test condition was randomized to minimize learning effect.

Statistical analysis

SPSS version 21 was used to analyze the data. The balance scores (mCTSIB, US, and LOS) between wearing a sandal, standard shoe, or no footwear (barefoot) was analyzed using the “repeated measures analysis of variance” with “Bonferroni's correction”. A $p < 0.05$ was considered as statistically significant.

Table 2. Comparison of balance scores between eyes open and closed conditions.

Footwear conditions	Balance outcome	Eyes open	Eyes closed	t-test	
				t	p
Barefoot (BF)	mCTIB on firm surface	0.38 ± 0.19	0.31 ± 0.07	1.948	0.06
	mCTIB on foam surface	0.67 ± 0.14	1.29 ± 0.37	-7.991	<0.001
	US on left leg	0.90 ± 0.21	2.46 ± 0.92	-9.970	<0.001
	US on right leg	0.89 ± 0.25	2.54 ± 0.92	-10.862	<0.001
Standard shoe (SS)	mCTIB on firm surface	0.41 ± 0.38	0.35 ± 0.18	0.734	0.47
	mCTIB on foam surface	0.73 ± 0.18	1.35 ± 0.40	-7.757	<0.001
	US on left leg	0.98 ± 0.45	2.62 ± 0.80	-13.895	<0.001
	US on right leg	1.08 ± 0.46	2.65 ± 0.94	-8.704	<0.001
Sandal (S)	mCTIB on firm surface	0.49 ± 0.14	0.52 ± 0.36	-0.394	0.69
	mCTIB on foam surface	0.78 ± 0.17	1.62 ± 0.36	-13.835	<0.001
	US on left leg	1.23 ± 0.65	3.08 ± 1.06	-11.765	<0.001
	US on right leg	1.34 ± 0.88	3.17 ± 1.22	-11.235	<0.001

mCTIB, Modified Clinical Test of Sensory Interaction on Balance; US, Unilateral Stance.

Results

The 30 male participants with mean (SD) of age, weight, height, and BMI were 23.1 (2.12), 71.1 (13.8), 1.69 (.05), and 24.69 (4.61), respectively. Table 1 presents the comparison of balance scores between a sandal, standard shoe, or no footwear (barefoot) conditions. Table 2 presents the comparison of balance scores between eyes open and closed conditions.

There were no significant effects between footwear conditions for mCTIB with eye open on firm surface ($p=0.17$). However, there were significant effects between footwear conditions for mCTIB with eye closed on firm surface ($p=0.002$). Post-hoc analyses indicate a significant differences in mCTIB between barefoot and sandal ($p=0.002$), and between standard shoe and sandal ($p=0.02$), but not between barefoot and standard shoe ($p=0.99$). In addition, there were significant effects between footwear conditions for mCTIB with eye open and closed on foam surface ($p=0.04$ and $p=0.003$, respectively). Post-hoc analyses indicate a significant differences in mCTIB with eye open between barefoot and sandal ($p=0.03$), but not between standard shoe and sandal ($p=0.73$), and between barefoot and standard shoe ($p=0.49$). However, there were significant differences in mCTIB with eye closed between barefoot and sandal ($p=0.004$), and between standard shoe and sandal ($p=0.02$), but not between barefoot and standard shoe ($p=0.97$).

There were significant effects between footwear conditions for US on left foot with eye open and closed ($p=0.02$ and $p=0.03$, respectively). Similarly, there were significant effects between footwear conditions for US on right foot with eye open and closed ($p=0.02$ and $p=0.05$, respectively). Post-hoc analyses indicate a significant differences in US on left foot with eye open and closed between barefoot and sandal

($p=0.03$ and 0.04 , respectively), but not between standard shoe and sandal ($p=0.14$ and $p=0.18$, respectively), and between barefoot and standard shoe ($p=0.89$ and $p=0.89$, respectively). Similarly, there were significant differences in US on right foot with eye open and closed between barefoot and sandal ($p=0.01$ and 0.04 , respectively), but not between standard shoe and sandal ($p=0.27$ and $p=0.16$, respectively), and between barefoot and standard shoe ($p=0.70$ and $p=0.95$, respectively).

There were significant effects between footwear conditions for LOS reaction time during forward movement ($p=0.02$). Post-hoc analyses indicate a significant differences in LOS reaction time during forward movement between barefoot and sandal ($p=0.04$), and between standard shoe and sandal ($p=0.04$), but not between barefoot and standard shoe ($p=0.88$). Similarly, there were significant effects between footwear conditions for LOS reaction time during left side movement ($p=0.01$). Post-hoc analyses indicate a significant differences in LOS reaction time during left side movement between barefoot and sandal ($p=0.01$), but not between standard shoe and sandal ($p=0.31$), and between barefoot and standard shoe ($p=0.56$). In addition, there were no significant effects between footwear conditions for LOS reaction time during backward and right side movement ($p=0.14$ and $p=0.16$, respectively).

There were insignificant differences in the postural sway measured by mCTIB on firm surface when tested in eyes closed condition compared to eyes open test conditions in all the three footwear conditions ($p>0.05$). There were significant differences in the postural sway measured by mCTIB on foam surface and US on left and right leg when tested in eyes closed condition compared to eyes open test conditions in all the three footwear conditions ($p<0.001$).

Discussion

The present study aimed to evaluate the differences between wearing a sandal, standard shoe, or no footwear (barefoot), in relation to postural stability measured by mCTSIB, US, and the LOS in healthy young adult males. The findings of the present study indicates that the wearing a sandal compared to bare feet significantly affect balance scores in mCTSIB with eye closed on firm surface, mCTSIB with eye open and closed on foam surface, US on left and right foot with eye open and closed, and the LOS reaction time during forward and left side movement. However, wearing a standard shoe compared to bare feet did not significantly affect balance scores in standing.

A previous study noted significantly increased medio-lateral and antero-posterior sway in eyes-closed testing in all footwear types compared to bare feet²⁴. Another study reported less sway in bare feet quiet standing compared to wearing own shoes and other footwear conditions²⁵. In addition, AP sway was increased in older adults wearing athletic footwear compared to bare feet²⁶. The tactile postural control mechanisms which transfer required updated sensory information to the CNS for the maintenance of balance can be altered by footwear²⁴.

In the present study, there were increased postural sway and reduced stability when tested in eyes closed condition compared to eyes open test conditions in all the three footwear conditions. Similarly, previous studies reported increased postural sway and reduced stability when tested in eyes closed condition compared to eyes open test conditions^{26,27}. The importance of vision in postural control is reported in the previous study²⁸.

In the preset study, there were increased postural sway and reduced stability when tested with standard shoes compared to sandal. Previous study reported that the poor footwear characteristics of sandal such as minimal heel counter stiffness and poor motion control resulting negative effect on balance²⁴.

The present study had some potential limitations. As this was a comparative study the small number of participants minimizes the generalizability of the results. The result of present study is limited to only healthy young adult males. Future research of footwear's effect on balance in young adult males who require stability (such as, acute ankle injuries, or lower extremity orthopedic conditions) would give further knowledge about the importance of footwear on balance in the younger adult population.

The present study concludes that the wearing a sandal compared to bare feet significantly increased postural sway and reduced stability in healthy young adult males. However, wearing a standard shoe compared to bare feet did not significantly affect balance scores in standing.

Acknowledgements

The authors would like to extend their sincere appreciation to the Deanship of Scientific Research at King Saud University for funding this research through the research group NO. RGP-VPP-209.

Ethics approval and consent to participate

The study was approved by the ethics committee of the College of Applied Medical Sciences, King Saud University. The participants were asked to sign a written informed consent form approved by the institution ethics committee of King Saud University.

Funding

This project was funded by the Deanship of Scientific Research, King Saud University through the research group NO. RGP-VPP-209. The funding body played no role in the study design, manuscript writing, or decision to submit the manuscript for publication.

Authors' contributions

SA: Corresponding author, participated in the design of the study, participated in the data collection, drafted the manuscript and finished the manuscript. AA: participated in the design of the study, helped in ethics applications and revised the manuscript critically. HZ: participated in the design of the study, developed the protocol, and revised the manuscript critically. All authors read and approved the final manuscript.

References

1. Barton CJ, Bonanno D, Menz HB. Development and evaluation of a tool for the assessment of footwear characteristics. *J Foot Ankle Res* 2009;2:10.
2. Hijmans JM, Geertzen JH, Dijkstra PU, Postema K. A systematic review of the effects of shoes and other ankle or foot appliances on balance in older people and people with peripheral nervous system disorders. *Gait Posture* 2007;25(2):316-23.
3. Perry SD, Radtke A, Goodwin CR. Influence of footwear midsole material hardness on dynamic balance control during unexpected gait termination. *Gait Posture* 2007; 25(1):94-8.
4. Meyer PF, Oddsson LI, De Luca CJ. The role of plantar cutaneous sensation in unperturbed stance. *Exp Brain Res* 2004;156(4):505-12.
5. Palluel E, Nougier V, Olivier I. Do spike insoles enhance postural stability and plantar-surface cutaneous sensitivity in the elderly? *Age (Dordr)* 2008;30(1):53-61.
6. Perry SD, Radtke A, McIlroy WE, Fernie GR, Maki BE. Efficacy and effectiveness of a balance-enhancing insole. *J Gerontol A Biol Sci Med Sci* 2008;63(6):595-602.
7. Priplata AA, Niemi JB, Harry JD, Lipsitz LA, Collins JJ. Vibrating insoles and balance control in elderly people. *Lancet* 2003;362(9390):1123-4.
8. Maki BE, Perry SD, Norrie RG, McIlroy WE. Effect of facilitation of sensation from plantar foot-surface boundaries on postural stabilization in young and older adults. *J Gerontol A Biol Sci Med Sci* 1999; 54(6):M281-7.
9. Rome K, Brown CL. Randomized clinical trial into the impact of rigid foot orthoses on balance parameters in excessively pronated feet. *Clin Rehabil* 2004; 18(6):624-30.
10. Galica AM, Kang HG, Priplata AA, D'Andrea SE, Starobinets OV, Sorond FA, et al. Subsensory vibrations

- to the feet reduce gait variability in elderly fallers. *Gait Posture* 2009;30(3):383-7.
11. Arnadottir SA, Mercer VS. Effects of footwear on measurements of balance and gait in women between the ages of 65 and 93 years. *Phys Ther* 2000;80(1):17-27.
 12. Frey C, Kubasak M. Faulty footwear contributes to why senior citizens fall. *Biomechanics* 1998;5:45-8.
 13. Brenton-Rule A, Bassett S, Walsh A, Rome K. The evaluation of walking footwear on postural stability in healthy older adults: an exploratory study. *Clin Biomech (Bristol, Avon)* 2011;26(8):885-7.
 14. Keegan TH, Kelsey JL, King AC, Quesenberry CP, Jr., Sidney S. Characteristics of fallers who fracture at the foot, distal forearm, proximal humerus, pelvis, and shaft of the tibia/fibula compared with fallers who do not fracture. *Am J Epidemiol* 2004;159(2):192-203.
 15. Murphy K, Curry EJ, Matzkin EG. Barefoot running: does it prevent injuries? *Sports Med* 2013;43(11):1131-8.
 16. Operators manual. Neurocom Balance Master, version 6.1. Neurocom International, Inc., 1998
 17. Cohen H, Blatchly CA, Gombash LL. A study of the clinical test of sensory interaction and balance. *Phys Ther* 1993;73(6):346-51; discussion 51-4.
 18. Wrisley DM, Whitney SL. The effect of foot position on the modified clinical test of sensory interaction and balance. *Arch Phys Med Rehab* 2004;85(2):335-8.
 19. Zouita Ben Moussa A, Zouita S, Dziri C, Ben Salah FZ. Single-leg assessment of postural stability and knee functional outcome two years after anterior cruciate ligament reconstruction. *Ann Phys Rehabil Med* 2009;52(6):475-84
 20. Ageberg E, Roberts D, Holmstrom E, Friden T. Balance in single-limb stance in healthy subjects-reliability of testing procedure and the effect of short-duration sub-maximal cycling. *BMC Musculoskelet Disord* 2003;4:14.
 21. Pickerill ML, Harter RA. Validity and reliability of limits-of-stability testing: a comparison of 2 postural stability evaluation devices. *J Athl Train* 2011;46(6):600-6.
 22. Vuillerme N, Burdet C, Isableu B, Demetz S. The magnitude of the effect of calf muscles fatigue on postural control during bipedal quiet standing with vision depends on the eye-visual target distance. *Gait Posture* 2006;24(2):169-72.
 23. Schneiders A, Gregory K, Karas S, Mündermann A. Effect of foot position on balance ability in single-leg stance with and without visual feedback. *J Biomech* 2016;49:1969-72.
 24. Brenton-Rule A, D'Almeida S, Bassett S, Carroll M, Dalbeth N, Rome K. The effects of sandals on postural stability in patients with rheumatoid arthritis: an exploratory study. *Clin Biomech* 2014;29(3):350-3.
 25. Lord SR, Bashford GM. Shoe characteristics and balance in older women. *J Am Geriatr Soc* 1996;44(4):429-33.
 26. Brenton-Rule A, Bassett S, Walsh A, Rome K. The evaluation of walking footwear on postural stability in healthy older adults: an exploratory study. *Clin Biomech* 2011;26(8):885-7.
 27. Menant JC, Steele JR, Menz HB, Munro BJ, Lord SR. Effects of footwear features on balance and stepping in older people. *Gerontology* 2008;54(1):18-23.
 28. Hytönen M, Pyykkö I, Aalto H, Starck J. Postural control and age. *Acta Otolaryngol* 1993;113(2):119-22.