

## A second monoclinic polymorph of *N*-[bis(morpholin-4-yl)phosphinoyl]-4-fluorobenzamide with the *P*2<sub>1</sub>/n space group

Atekeh Tarahhomī,<sup>a\*</sup> Mehrdad Pourayoubi,<sup>a</sup> Mojtaba Keikha,<sup>a</sup> Arnold L. Rheingold<sup>b</sup> and James A. Golen<sup>b</sup>

<sup>a</sup>Department of Chemistry, Ferdowsi University of Mashhad, Mashhad, Iran, and

<sup>b</sup>Department of Chemistry, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

Correspondence e-mail: tarahhomī\_at@yahoo.com


Received 18 June 2012; accepted 10 July 2012

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004 \text{ \AA}$ ;  $R$  factor = 0.056;  $wR$  factor = 0.160; data-to-parameter ratio = 15.7.

A second monoclinic polymorph of the title molecule,  $C_{15}H_{21}FN_3O_4P$ , is reported in the space group *P*2<sub>1</sub>/n and compared to the previously reported *C*2/c space group [Gholivand *et al.* (2006). *Polyhedron*, **25**, 711–721]. The asymmetric unit of the title compound consists of two independent molecules. The P atoms adopt a distorted tetrahedral environment. In the C(O)NHP(O) fragment, the P=O and the N—H groups are in a *syn* conformation with respect to each other and in the crystal, intermolecular N—H···O=P hydrogen bonds form dimeric aggregates.

### Related literature

For the monoclinic polymorph of the title molecule, in a *C*2/c space group, for bond lengths and angles and for preparation of the starting compound 4-F-C<sub>6</sub>H<sub>4</sub>C(O)NHP(O)Cl<sub>2</sub>, see: Gholivand *et al.* (2006). For related phosphoramides, see: Pourayoubi, Nečas & Negari (2012); Pourayoubi, Tarahhomī *et al.* (2012).



### Experimental

#### Crystal data

$C_{15}H_{21}FN_3O_4P$	$V = 3360.3 (2) \text{ \AA}^3$
$M_r = 357.32$	$Z = 8$
Monoclinic, <i>P</i> 2 <sub>1</sub> /n	Mo $K\alpha$ radiation
$a = 15.6093 (6) \text{ \AA}$	$\mu = 0.20 \text{ mm}^{-1}$
$b = 10.7114 (4) \text{ \AA}$	$T = 100 \text{ K}$
$c = 21.0045 (9) \text{ \AA}$	$0.18 \times 0.15 \times 0.10 \text{ mm}$
$\beta = 106.896 (2)^\circ$	

#### Data collection

Bruker APEXII CCD	25230 measured reflections
diffractometer	6912 independent reflections
Absorption correction: multi-scan	5355 reflections with $I > 2\sigma(I)$
(SADABS; Sheldrick, 2004)	$R_{\text{int}} = 0.037$
	$T_{\min} = 0.965, T_{\max} = 0.980$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$	H atoms treated by a mixture of
$wR(F^2) = 0.160$	independent and constrained
$S = 1.03$	refinement
6912 reflections	$\Delta\rho_{\max} = 1.45 \text{ e \AA}^{-3}$
439 parameters	$\Delta\rho_{\min} = -0.73 \text{ e \AA}^{-3}$
2 restraints	

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N4—H4N···O2 <sup>i</sup>	0.85 (2)	2.01 (2)	2.855 (3)	176 (3)
N1—H1N···O6 <sup>ii</sup>	0.87 (2)	2.04 (2)	2.870 (3)	159 (3)

Symmetry codes: (i)  $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXTL* and *enCIFer* (Allen *et al.*, 2004).

Support of this investigation by Ferdowsi University of Mashhad is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2145).

### References

- Allen, F. H., Johnson, O., Shields, G. P., Smith, B. R. & Towler, M. (2004). *J. Appl. Cryst.* **37**, 335–338.
- Bruker (2005). *SAINT* and *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Gholivand, K., Shariatinia, Z. & Pourayoubi, M. (2006). *Polyhedron*, **25**, 711–721.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Pourayoubi, M., Nečas, M. & Negari, M. (2012). *Acta Cryst. C* **68**, o51–o56.
- Pourayoubi, M., Tarahhomī, A., Karimi Ahmadabad, F., Fejsarová, K., van der Lee, A. & Dušek, M. (2012). *Acta Cryst. C* **68**, o164–o169.
- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

# supplementary materials

*Acta Cryst.* (2012). E68, o2444 [doi:10.1107/S1600536812031418]

## A second monoclinic polymorph of *N*-[bis(morpholin-4-yl)phosphinoyl]-4-fluorobenzamide with the *P*2<sub>1</sub>/*n* space group

**Atekeh Tarahhomí, Mehrdad Pourayoubi, Mojtaba Keikha, Arnold L. Rheingold and James A. Golen**

### Comment

The structure determination of a monoclinic polymorph of the title molecule, [4-F—C<sub>6</sub>H<sub>4</sub>C(O)NH]P(O)[NC<sub>4</sub>H<sub>8</sub>O]<sub>2</sub> (Fig. 1), in a *C*2/*c* space group with *Z* = 8 was investigated in ambient temperature, Gholivand *et al.* (2006). Here, we report on a second monoclinic polymorph in a space group *P*2<sub>1</sub>/*n* with *Z* = 8. Cell parameters for both polymorphs exhibit similar dimensions. The *C*2/*c* structure at 293 K has cell parameters of 15.732 (3), 10.740 (2), 21.553 (4) Å with  $\beta$  of 106.93 (3) $^\circ$  while those of *P*2<sub>1</sub>/*n* at 100 K have smaller cell dimensions indicative of a possible phase change.

The asymmetric unit consists of two independent molecules; in each molecule, the phosphoryl group adopts a *syn* orientation with respect to the N—H unit. In the first monoclinic modification of this compound (Gholivand *et al.*, 2006), the asymmetric unit is composed of one molecule involving disorder in one morpholin-4-yl moiety. The P atom is in a distorted tetrahedral environment as has been noted for other phosphoramides (Pourayoubi, Nečas & Negari, 2012 and Pourayoubi, Tarahhomí *et al.*, 2012).

The P=O, C=O and P—N bond lengths and P—N—C bond angles are within the expected values (Gholivand *et al.*, 2006).

In the crystal, two different intermolecular N—H···O(P) hydrogen bonds make dimeric aggregates. So, each hydrogen-bonded dimer is built of two symmetrically independent molecules (Fig. 2).

### Experimental

4-F—C<sub>6</sub>H<sub>4</sub>C(O)NHP(O)Cl<sub>2</sub> was prepared according to the literature method reported by Gholivand *et al.* (2006).


To a solution of 4-F—C<sub>6</sub>H<sub>4</sub>C(O)NHP(O)Cl<sub>2</sub> (0.723 g, 2.825 mmol) in dry acetonitrile (25 ml), a solution of morpholine (0.984 g, 11.300 mmol) in dry acetonitrile (5 ml) was added at 273 K. After 4 h stirring, the solvent was removed and the product was washed with distilled water and the resulting precipitate of 4-F—C<sub>6</sub>H<sub>4</sub>C(O)NHP(O)(NC<sub>4</sub>H<sub>8</sub>O)<sub>2</sub> was collected. Single crystals were obtained in a try on a reaction between 4-F—C<sub>6</sub>H<sub>4</sub>C(O)NHP(O)(NC<sub>4</sub>H<sub>8</sub>O)<sub>2</sub> and Sn(CH<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> in CH<sub>3</sub>OH under reflux, followed by slow evaporation of the filtered solution at room temperature.

### Refinement


All non-hydrogen atoms were refined anisotropically by full matrix least squares on F<sup>2</sup>. Hydrogen atoms H1N and H4N were found from a Fourier difference map and their N—H distances were fixed at 0.87 (2) Å and were allowed to refine isotropically with 1.20 *U*<sub>eq</sub> of parent N atoms. All other hydrogen atoms were placed in calculated positions and treated as riding on their parent C atoms with distances C—H = 0.990 Å (CH<sub>2</sub>) and 1.00 Å (CH) with 1.20*U*<sub>eq</sub> of their parent C atoms.

**Computing details**

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *enCIFer* (Allen *et al.*, 2004).

**Figure 1**

An *ORTEP*-style plot and atom labeling scheme for the asymmetric unit of the title molecule. Displacement ellipsoids are given at 50% probability level and H atoms are drawn as small spheres of arbitrary radii.

**Figure 2**

Crystal packing view showing  $\text{N}—\text{H}\cdots\text{O}(\text{P})$  hydrogen bond which are shown as dotted lines. The symmetrically independent molecules are shown as different colors and the H atoms not involved in hydrogen bonding have been omitted for the sake of clarity.

### *N*-[Bis(morpholin-4-yl)phosphinoyl]-4-fluorobenzamide

#### Crystal data

$\text{C}_{15}\text{H}_{21}\text{FN}_3\text{O}_4\text{P}$   
 $M_r = 357.32$   
Monoclinic,  $P2_1/n$   
Hall symbol: -P 2yn  
 $a = 15.6093 (6)$  Å  
 $b = 10.7114 (4)$  Å  
 $c = 21.0045 (9)$  Å  
 $\beta = 106.896 (2)$ °  
 $V = 3360.3 (2)$  Å<sup>3</sup>  
 $Z = 8$

$F(000) = 1504$   
 $D_x = 1.413 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 7455 reflections  
 $\theta = 2.3\text{--}26.4$ °  
 $\mu = 0.20 \text{ mm}^{-1}$   
 $T = 100 \text{ K}$   
Block, colourless  
 $0.18 \times 0.15 \times 0.10 \text{ mm}$

*Data collection*

Bruker APEXII CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
Detector resolution: 8.33 pixels mm<sup>-1</sup>  
 $\varphi$  and  $\omega$  scans  
Absorption correction: multi-scan  
(SADABS; Sheldrick, 2004)  
 $T_{\min} = 0.965$ ,  $T_{\max} = 0.980$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.056$   
 $wR(F^2) = 0.160$   
 $S = 1.03$   
6912 reflections  
439 parameters  
2 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H atoms treated by a mixture of independent  
and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0816P)^2 + 4.9555P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 1.45 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.73 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	0.47360 (4)	0.87432 (6)	0.15573 (3)	0.01350 (17)
P2	0.93848 (4)	0.36293 (6)	0.12376 (3)	0.01408 (17)
F1	0.07031 (11)	0.82774 (16)	0.31748 (9)	0.0255 (4)
F2	0.58872 (11)	0.32224 (15)	0.33946 (9)	0.0239 (4)
O1	0.30622 (14)	0.71437 (19)	0.12959 (10)	0.0250 (5)
O2	0.54401 (12)	0.95376 (17)	0.19942 (9)	0.0170 (4)
O3	0.26441 (14)	1.08121 (18)	0.01350 (10)	0.0263 (5)
O4	0.59087 (13)	0.54123 (18)	0.08540 (10)	0.0238 (5)
O5	0.76002 (14)	0.22559 (19)	0.10808 (10)	0.0246 (5)
O6	1.01711 (12)	0.43300 (18)	0.16535 (10)	0.0192 (4)
O7	0.74106 (13)	0.59113 (17)	-0.00982 (10)	0.0228 (4)
O8	1.08127 (14)	0.03799 (19)	0.09330 (11)	0.0294 (5)
N1	0.40849 (14)	0.8319 (2)	0.20362 (11)	0.0149 (5)
H1N	0.417 (2)	0.872 (3)	0.2410 (11)	0.018*
N2	0.41816 (15)	0.9499 (2)	0.08931 (11)	0.0165 (5)
N3	0.50455 (15)	0.7452 (2)	0.12658 (11)	0.0176 (5)

N4	0.87238 (15)	0.3430 (2)	0.17412 (11)	0.0157 (5)
H4N	0.8948 (19)	0.374 (3)	0.2123 (10)	0.019*
N5	0.88937 (14)	0.4404 (2)	0.05651 (11)	0.0164 (5)
N6	0.95166 (15)	0.2234 (2)	0.09645 (11)	0.0179 (5)
C1	0.27337 (18)	0.8749 (3)	0.27251 (14)	0.0186 (6)
H1A	0.3255	0.9257	0.2849	0.022*
C2	0.20861 (18)	0.8881 (3)	0.30509 (14)	0.0199 (6)
H2A	0.2152	0.9479	0.3396	0.024*
C3	0.13450 (17)	0.8125 (2)	0.28632 (14)	0.0172 (6)
C4	0.12203 (18)	0.7231 (3)	0.23765 (14)	0.0198 (6)
H4A	0.0707	0.6706	0.2269	0.024*
C5	0.18645 (17)	0.7119 (2)	0.20477 (14)	0.0173 (5)
H5A	0.1788	0.6520	0.1702	0.021*
C6	0.26243 (17)	0.7873 (2)	0.22151 (13)	0.0152 (5)
C7	0.32636 (18)	0.7737 (2)	0.18122 (13)	0.0168 (5)
C8	0.35807 (18)	0.8961 (3)	0.02858 (14)	0.0199 (6)
H8A	0.3567	0.8041	0.0330	0.024*
H8B	0.3805	0.9159	-0.0098	0.024*
C9	0.2642 (2)	0.9483 (3)	0.01647 (15)	0.0232 (6)
H9A	0.2249	0.9147	-0.0259	0.028*
H9B	0.2397	0.9213	0.0528	0.028*
C10	0.3181 (2)	1.1302 (3)	0.07509 (15)	0.0242 (6)
H10A	0.2939	1.1023	0.1114	0.029*
H10B	0.3160	1.2225	0.0735	0.029*
C11	0.4140 (2)	1.0871 (2)	0.08929 (15)	0.0226 (6)
H11A	0.4398	1.1206	0.0550	0.027*
H11B	0.4498	1.1193	0.1331	0.027*
C12	0.56485 (19)	0.7619 (3)	0.08400 (15)	0.0225 (6)
H12A	0.6268	0.7770	0.1122	0.027*
H12B	0.5454	0.8351	0.0546	0.027*
C13	0.5620 (2)	0.6472 (3)	0.04299 (15)	0.0249 (6)
H13A	0.5002	0.6333	0.0141	0.030*
H13B	0.6015	0.6582	0.0141	0.030*
C14	0.53270 (19)	0.5219 (3)	0.12520 (15)	0.0226 (6)
H14A	0.5529	0.4478	0.1538	0.027*
H14B	0.4715	0.5049	0.0960	0.027*
C15	0.53018 (19)	0.6333 (2)	0.16835 (14)	0.0204 (6)
H15A	0.4864	0.6186	0.1935	0.024*
H15B	0.5898	0.6459	0.2007	0.024*
C16	0.75142 (17)	0.3982 (2)	0.25373 (14)	0.0173 (5)
H16A	0.7956	0.4597	0.2545	0.021*
C17	0.70067 (17)	0.4068 (2)	0.29795 (14)	0.0177 (6)
H17A	0.7088	0.4741	0.3286	0.021*
C18	0.63820 (17)	0.3150 (2)	0.29609 (13)	0.0159 (5)
C19	0.62334 (16)	0.2167 (2)	0.25214 (14)	0.0158 (5)
H19A	0.5800	0.1546	0.2524	0.019*
C20	0.67331 (16)	0.2109 (2)	0.20749 (13)	0.0131 (5)
H20A	0.6630	0.1450	0.1759	0.016*
C21	0.73824 (17)	0.3000 (2)	0.20819 (13)	0.0142 (5)

C22	0.78981 (17)	0.2861 (2)	0.15893 (14)	0.0161 (5)
C23	0.81724 (17)	0.3917 (2)	0.00028 (13)	0.0171 (5)
H23A	0.8097	0.3012	0.0065	0.021*
H23B	0.8334	0.4033	-0.0415	0.021*
C24	0.73018 (18)	0.4588 (2)	-0.00470 (15)	0.0190 (6)
H24A	0.6832	0.4286	-0.0443	0.023*
H24B	0.7108	0.4401	0.0352	0.023*
C25	0.8061 (2)	0.6370 (3)	0.04805 (16)	0.0262 (7)
H25A	0.7871	0.6179	0.0880	0.031*
H25B	0.8112	0.7288	0.0449	0.031*
C26	0.89630 (19)	0.5771 (2)	0.05444 (16)	0.0244 (6)
H26A	0.9174	0.6018	0.0161	0.029*
H26B	0.9406	0.6070	0.0956	0.029*
C27	0.98143 (19)	0.2023 (3)	0.03698 (15)	0.0229 (6)
H27A	0.9799	0.2821	0.0129	0.027*
H27B	0.9401	0.1432	0.0068	0.027*
C28	1.0750 (2)	0.1502 (3)	0.05604 (18)	0.0305 (7)
H28A	1.0927	0.1334	0.0153	0.037*
H28B	1.1169	0.2127	0.0828	0.037*
C29	1.0563 (2)	0.0604 (3)	0.15248 (16)	0.0280 (7)
H29A	1.0983	0.1214	0.1806	0.034*
H29B	1.0610	-0.0184	0.1779	0.034*
C30	0.9619 (2)	0.1102 (3)	0.13704 (15)	0.0221 (6)
H30A	0.9191	0.0459	0.1129	0.027*
H30B	0.9479	0.1291	0.1791	0.027*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
P1	0.0156 (3)	0.0121 (3)	0.0144 (3)	0.0020 (2)	0.0069 (3)	0.0010 (3)
P2	0.0130 (3)	0.0138 (3)	0.0165 (4)	0.0014 (2)	0.0060 (3)	0.0030 (3)
F1	0.0196 (8)	0.0309 (9)	0.0323 (10)	0.0007 (7)	0.0173 (7)	-0.0010 (8)
F2	0.0236 (8)	0.0232 (8)	0.0321 (10)	-0.0011 (7)	0.0194 (7)	-0.0030 (7)
O1	0.0283 (11)	0.0296 (11)	0.0193 (11)	-0.0100 (9)	0.0104 (9)	-0.0112 (9)
O2	0.0146 (9)	0.0176 (9)	0.0180 (10)	0.0000 (7)	0.0034 (7)	0.0024 (8)
O3	0.0328 (11)	0.0195 (10)	0.0199 (11)	0.0081 (9)	-0.0026 (9)	-0.0024 (8)
O4	0.0260 (11)	0.0196 (10)	0.0305 (12)	0.0071 (8)	0.0155 (9)	0.0017 (9)
O5	0.0287 (11)	0.0272 (11)	0.0211 (11)	-0.0105 (9)	0.0124 (9)	-0.0097 (9)
O6	0.0142 (9)	0.0230 (10)	0.0201 (10)	-0.0004 (8)	0.0045 (8)	0.0065 (8)
O7	0.0235 (10)	0.0166 (9)	0.0240 (11)	0.0052 (8)	0.0002 (8)	0.0008 (8)
O8	0.0281 (11)	0.0238 (11)	0.0389 (13)	0.0130 (9)	0.0140 (10)	0.0058 (10)
N1	0.0151 (11)	0.0159 (10)	0.0157 (11)	-0.0010 (9)	0.0077 (9)	-0.0041 (9)
N2	0.0214 (11)	0.0114 (10)	0.0158 (11)	0.0024 (9)	0.0041 (9)	0.0001 (9)
N3	0.0260 (12)	0.0144 (11)	0.0162 (12)	0.0057 (9)	0.0120 (10)	0.0037 (9)
N4	0.0168 (11)	0.0171 (11)	0.0151 (11)	-0.0028 (9)	0.0075 (9)	-0.0033 (9)
N5	0.0145 (11)	0.0133 (10)	0.0198 (12)	0.0004 (8)	0.0026 (9)	0.0028 (9)
N6	0.0235 (12)	0.0165 (11)	0.0163 (11)	0.0052 (9)	0.0102 (9)	0.0030 (9)
C1	0.0153 (13)	0.0222 (14)	0.0196 (14)	-0.0039 (10)	0.0073 (11)	-0.0049 (11)
C2	0.0168 (13)	0.0239 (14)	0.0191 (14)	0.0007 (11)	0.0054 (11)	-0.0060 (11)
C3	0.0121 (12)	0.0205 (13)	0.0211 (14)	0.0046 (10)	0.0081 (11)	0.0060 (11)

C4	0.0133 (12)	0.0200 (13)	0.0259 (15)	-0.0025 (10)	0.0055 (11)	0.0016 (11)
C5	0.0154 (13)	0.0177 (13)	0.0169 (13)	-0.0003 (10)	0.0018 (10)	-0.0017 (11)
C6	0.0135 (12)	0.0150 (12)	0.0164 (13)	0.0011 (10)	0.0036 (10)	0.0008 (10)
C7	0.0205 (13)	0.0134 (12)	0.0168 (14)	-0.0010 (10)	0.0057 (11)	-0.0006 (10)
C8	0.0252 (14)	0.0183 (13)	0.0162 (14)	0.0028 (11)	0.0061 (11)	-0.0019 (11)
C9	0.0289 (15)	0.0178 (13)	0.0202 (15)	0.0035 (11)	0.0026 (12)	-0.0035 (11)
C10	0.0327 (16)	0.0163 (13)	0.0193 (15)	0.0074 (12)	0.0008 (12)	-0.0024 (11)
C11	0.0305 (16)	0.0130 (13)	0.0228 (15)	0.0008 (11)	0.0055 (12)	0.0020 (11)
C12	0.0262 (15)	0.0214 (14)	0.0241 (15)	0.0034 (12)	0.0140 (12)	0.0021 (12)
C13	0.0324 (16)	0.0238 (15)	0.0238 (16)	0.0070 (12)	0.0165 (13)	0.0015 (12)
C14	0.0240 (14)	0.0189 (13)	0.0280 (16)	0.0028 (11)	0.0122 (12)	0.0024 (12)
C15	0.0251 (14)	0.0175 (13)	0.0208 (14)	0.0038 (11)	0.0101 (12)	0.0047 (11)
C16	0.0127 (12)	0.0171 (13)	0.0229 (14)	-0.0039 (10)	0.0064 (10)	-0.0043 (11)
C17	0.0156 (13)	0.0187 (13)	0.0183 (14)	-0.0011 (10)	0.0042 (11)	-0.0068 (11)
C18	0.0124 (12)	0.0186 (13)	0.0190 (14)	0.0043 (10)	0.0084 (10)	0.0027 (11)
C19	0.0113 (12)	0.0129 (12)	0.0228 (14)	-0.0009 (9)	0.0043 (10)	0.0016 (10)
C20	0.0095 (11)	0.0120 (11)	0.0155 (13)	0.0020 (9)	-0.0002 (10)	0.0014 (10)
C21	0.0145 (12)	0.0131 (12)	0.0151 (13)	0.0003 (9)	0.0045 (10)	0.0017 (10)
C22	0.0180 (13)	0.0122 (12)	0.0187 (14)	-0.0033 (10)	0.0061 (11)	-0.0004 (10)
C23	0.0183 (13)	0.0167 (12)	0.0156 (13)	0.0021 (10)	0.0036 (11)	-0.0009 (10)
C24	0.0179 (13)	0.0159 (13)	0.0221 (14)	0.0012 (10)	0.0041 (11)	-0.0016 (11)
C25	0.0281 (15)	0.0127 (13)	0.0316 (17)	0.0043 (11)	-0.0010 (13)	-0.0011 (12)
C26	0.0226 (14)	0.0143 (13)	0.0307 (16)	-0.0031 (11)	-0.0011 (12)	0.0069 (12)
C27	0.0268 (15)	0.0242 (14)	0.0215 (15)	0.0043 (12)	0.0130 (12)	0.0026 (12)
C28	0.0291 (16)	0.0278 (16)	0.0419 (19)	0.0094 (13)	0.0221 (15)	0.0092 (14)
C29	0.0309 (16)	0.0234 (15)	0.0260 (16)	0.0081 (12)	0.0022 (13)	0.0091 (13)
C30	0.0296 (15)	0.0168 (13)	0.0210 (15)	0.0021 (11)	0.0091 (12)	0.0034 (11)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

P1—O2	1.4791 (19)	C9—H9B	0.9900
P1—N2	1.628 (2)	C10—C11	1.511 (4)
P1—N3	1.641 (2)	C10—H10A	0.9900
P1—N1	1.687 (2)	C10—H10B	0.9900
P2—O6	1.4866 (19)	C11—H11A	0.9900
P2—N5	1.626 (2)	C11—H11B	0.9900
P2—N6	1.635 (2)	C12—C13	1.495 (4)
P2—N4	1.692 (2)	C12—H12A	0.9900
F1—C3	1.356 (3)	C12—H12B	0.9900
F2—C18	1.357 (3)	C13—H13A	0.9900
O1—C7	1.217 (3)	C13—H13B	0.9900
O3—C10	1.422 (3)	C14—C15	1.505 (4)
O3—C9	1.425 (3)	C14—H14A	0.9900
O4—C14	1.417 (3)	C14—H14B	0.9900
O4—C13	1.432 (3)	C15—H15A	0.9900
O5—C22	1.220 (3)	C15—H15B	0.9900
O7—C25	1.426 (3)	C16—C17	1.388 (4)
O7—C24	1.435 (3)	C16—C21	1.396 (4)
O8—C28	1.422 (4)	C16—H16A	0.9500
O8—C29	1.427 (4)	C17—C18	1.378 (4)

N1—C7	1.380 (3)	C17—H17A	0.9500
N1—H1N	0.869 (17)	C18—C19	1.375 (4)
N2—C8	1.464 (3)	C19—C20	1.385 (4)
N2—C11	1.471 (3)	C19—H19A	0.9500
N3—C15	1.471 (3)	C20—C21	1.389 (3)
N3—C12	1.486 (3)	C20—H20A	0.9500
N4—C22	1.377 (3)	C21—C22	1.491 (4)
N4—H4N	0.846 (18)	C23—C24	1.514 (4)
N5—C26	1.470 (3)	C23—H23A	0.9900
N5—C23	1.470 (3)	C23—H23B	0.9900
N6—C30	1.465 (3)	C24—H24A	0.9900
N6—C27	1.471 (3)	C24—H24B	0.9900
C1—C2	1.383 (4)	C25—C26	1.517 (4)
C1—C6	1.397 (4)	C25—H25A	0.9900
C1—H1A	0.9500	C25—H25B	0.9900
C2—C3	1.373 (4)	C26—H26A	0.9900
C2—H2A	0.9500	C26—H26B	0.9900
C3—C4	1.373 (4)	C27—C28	1.504 (4)
C4—C5	1.381 (4)	C27—H27A	0.9900
C4—H4A	0.9500	C27—H27B	0.9900
C5—C6	1.392 (4)	C28—H28A	0.9900
C5—H5A	0.9500	C28—H28B	0.9900
C6—C7	1.491 (4)	C29—C30	1.512 (4)
C8—C9	1.519 (4)	C29—H29A	0.9900
C8—H8A	0.9900	C29—H29B	0.9900
C8—H8B	0.9900	C30—H30A	0.9900
C9—H9A	0.9900	C30—H30B	0.9900
O2—P1—N2	111.08 (11)	O4—C13—H13A	109.7
O2—P1—N3	118.04 (11)	C12—C13—H13A	109.7
N2—P1—N3	103.92 (11)	O4—C13—H13B	109.7
O2—P1—N1	104.63 (11)	C12—C13—H13B	109.7
N2—P1—N1	112.50 (12)	H13A—C13—H13B	108.2
N3—P1—N1	106.82 (11)	O4—C14—C15	111.9 (2)
O6—P2—N5	110.68 (11)	O4—C14—H14A	109.2
O6—P2—N6	120.07 (12)	C15—C14—H14A	109.2
N5—P2—N6	103.74 (12)	O4—C14—H14B	109.2
O6—P2—N4	104.22 (11)	C15—C14—H14B	109.2
N5—P2—N4	112.86 (12)	H14A—C14—H14B	107.9
N6—P2—N4	105.41 (11)	N3—C15—C14	109.7 (2)
C10—O3—C9	109.7 (2)	N3—C15—H15A	109.7
C14—O4—C13	110.1 (2)	C14—C15—H15A	109.7
C25—O7—C24	110.3 (2)	N3—C15—H15B	109.7
C28—O8—C29	110.2 (2)	C14—C15—H15B	109.7
C7—N1—P1	125.42 (19)	H15A—C15—H15B	108.2
C7—N1—H1N	115 (2)	C17—C16—C21	120.6 (2)
P1—N1—H1N	116 (2)	C17—C16—H16A	119.7
C8—N2—C11	112.1 (2)	C21—C16—H16A	119.7
C8—N2—P1	126.66 (18)	C18—C17—C16	118.1 (2)

C11—N2—P1	120.65 (19)	C18—C17—H17A	121.0
C15—N3—C12	110.3 (2)	C16—C17—H17A	121.0
C15—N3—P1	121.53 (18)	F2—C18—C19	118.5 (2)
C12—N3—P1	115.43 (18)	F2—C18—C17	118.5 (2)
C22—N4—P2	127.66 (19)	C19—C18—C17	123.1 (2)
C22—N4—H4N	120 (2)	C18—C19—C20	118.1 (2)
P2—N4—H4N	113 (2)	C18—C19—H19A	121.0
C26—N5—C23	112.1 (2)	C20—C19—H19A	120.9
C26—N5—P2	120.94 (19)	C19—C20—C21	121.0 (2)
C23—N5—P2	125.07 (18)	C19—C20—H20A	119.5
C30—N6—C27	111.0 (2)	C21—C20—H20A	119.5
C30—N6—P2	123.80 (19)	C20—C21—C16	119.1 (2)
C27—N6—P2	122.78 (19)	C20—C21—C22	117.6 (2)
C2—C1—C6	120.3 (2)	C16—C21—C22	123.3 (2)
C2—C1—H1A	119.9	O5—C22—N4	122.1 (2)
C6—C1—H1A	119.9	O5—C22—C21	121.0 (2)
C3—C2—C1	118.3 (3)	N4—C22—C21	116.9 (2)
C3—C2—H2A	120.9	N5—C23—C24	110.5 (2)
C1—C2—H2A	120.9	N5—C23—H23A	109.6
F1—C3—C2	118.0 (2)	C24—C23—H23A	109.6
F1—C3—C4	118.6 (2)	N5—C23—H23B	109.6
C2—C3—C4	123.4 (2)	C24—C23—H23B	109.6
C3—C4—C5	117.8 (2)	H23A—C23—H23B	108.1
C3—C4—H4A	121.1	O7—C24—C23	110.5 (2)
C5—C4—H4A	121.1	O7—C24—H24A	109.5
C4—C5—C6	120.9 (3)	C23—C24—H24A	109.5
C4—C5—H5A	119.5	O7—C24—H24B	109.5
C6—C5—H5A	119.5	C23—C24—H24B	109.5
C5—C6—C1	119.2 (2)	H24A—C24—H24B	108.1
C5—C6—C7	117.5 (2)	O7—C25—C26	110.0 (2)
C1—C6—C7	123.2 (2)	O7—C25—H25A	109.7
O1—C7—N1	120.9 (2)	C26—C25—H25A	109.7
O1—C7—C6	121.0 (2)	O7—C25—H25B	109.7
N1—C7—C6	118.1 (2)	C26—C25—H25B	109.7
N2—C8—C9	110.1 (2)	H25A—C25—H25B	108.2
N2—C8—H8A	109.6	N5—C26—C25	110.5 (2)
C9—C8—H8A	109.6	N5—C26—H26A	109.6
N2—C8—H8B	109.6	C25—C26—H26A	109.6
C9—C8—H8B	109.6	N5—C26—H26B	109.6
H8A—C8—H8B	108.2	C25—C26—H26B	109.6
O3—C9—C8	111.1 (2)	H26A—C26—H26B	108.1
O3—C9—H9A	109.4	N6—C27—C28	110.7 (2)
C8—C9—H9A	109.4	N6—C27—H27A	109.5
O3—C9—H9B	109.4	C28—C27—H27A	109.5
C8—C9—H9B	109.4	N6—C27—H27B	109.5
H9A—C9—H9B	108.0	C28—C27—H27B	109.5
O3—C10—C11	110.8 (2)	H27A—C27—H27B	108.1
O3—C10—H10A	109.5	O8—C28—C27	111.5 (2)
C11—C10—H10A	109.5	O8—C28—H28A	109.3

O3—C10—H10B	109.5	C27—C28—H28A	109.3
C11—C10—H10B	109.5	O8—C28—H28B	109.3
H10A—C10—H10B	108.1	C27—C28—H28B	109.3
N2—C11—C10	110.3 (2)	H28A—C28—H28B	108.0
N2—C11—H11A	109.6	O8—C29—C30	111.6 (2)
C10—C11—H11A	109.6	O8—C29—H29A	109.3
N2—C11—H11B	109.6	C30—C29—H29A	109.3
C10—C11—H11B	109.6	O8—C29—H29B	109.3
H11A—C11—H11B	108.1	C30—C29—H29B	109.3
N3—C12—C13	109.3 (2)	H29A—C29—H29B	108.0
N3—C12—H12A	109.8	N6—C30—C29	110.6 (2)
C13—C12—H12A	109.8	N6—C30—H30A	109.5
N3—C12—H12B	109.8	C29—C30—H30A	109.5
C13—C12—H12B	109.8	N6—C30—H30B	109.5
H12A—C12—H12B	108.3	C29—C30—H30B	109.5
O4—C13—C12	110.0 (2)	H30A—C30—H30B	108.1
O2—P1—N1—C7	169.2 (2)	P1—N2—C8—C9	119.3 (2)
N2—P1—N1—C7	48.5 (2)	C10—O3—C9—C8	-61.2 (3)
N3—P1—N1—C7	-64.9 (2)	N2—C8—C9—O3	56.1 (3)
O2—P1—N2—C8	165.5 (2)	C9—O3—C10—C11	61.6 (3)
N3—P1—N2—C8	37.6 (2)	C8—N2—C11—C10	52.5 (3)
N1—P1—N2—C8	-77.6 (2)	P1—N2—C11—C10	-119.2 (2)
O2—P1—N2—C11	-24.1 (2)	O3—C10—C11—N2	-57.0 (3)
N3—P1—N2—C11	-152.0 (2)	C15—N3—C12—C13	56.9 (3)
N1—P1—N2—C11	92.9 (2)	P1—N3—C12—C13	-160.6 (2)
O2—P1—N3—C15	76.8 (2)	C14—O4—C13—C12	61.5 (3)
N2—P1—N3—C15	-159.7 (2)	N3—C12—C13—O4	-60.0 (3)
N1—P1—N3—C15	-40.6 (2)	C13—O4—C14—C15	-59.7 (3)
O2—P1—N3—C12	-61.2 (2)	C12—N3—C15—C14	-54.3 (3)
N2—P1—N3—C12	62.3 (2)	P1—N3—C15—C14	165.87 (19)
N1—P1—N3—C12	-178.56 (19)	O4—C14—C15—N3	56.1 (3)
O6—P2—N4—C22	178.2 (2)	C21—C16—C17—C18	0.9 (4)
N5—P2—N4—C22	58.0 (3)	C16—C17—C18—F2	179.5 (2)
N6—P2—N4—C22	-54.5 (2)	C16—C17—C18—C19	-0.8 (4)
O6—P2—N5—C26	-25.8 (2)	F2—C18—C19—C20	179.2 (2)
N6—P2—N5—C26	-155.8 (2)	C17—C18—C19—C20	-0.5 (4)
N4—P2—N5—C26	90.6 (2)	C18—C19—C20—C21	1.6 (4)
O6—P2—N5—C23	171.3 (2)	C19—C20—C21—C16	-1.5 (4)
N6—P2—N5—C23	41.3 (2)	C19—C20—C21—C22	179.3 (2)
N4—P2—N5—C23	-72.3 (2)	C17—C16—C21—C20	0.2 (4)
O6—P2—N6—C30	79.2 (2)	C17—C16—C21—C22	179.3 (2)
N5—P2—N6—C30	-156.6 (2)	P2—N4—C22—O5	11.3 (4)
N4—P2—N6—C30	-37.8 (2)	P2—N4—C22—C21	-169.64 (19)
O6—P2—N6—C27	-81.7 (2)	C20—C21—C22—O5	22.0 (4)
N5—P2—N6—C27	42.5 (2)	C16—C21—C22—O5	-157.1 (3)
N4—P2—N6—C27	161.3 (2)	C20—C21—C22—N4	-157.1 (2)
C6—C1—C2—C3	0.5 (4)	C16—C21—C22—N4	23.8 (4)
C1—C2—C3—F1	-178.6 (2)	C26—N5—C23—C24	-52.1 (3)

C1—C2—C3—C4	1.1 (4)	P2—N5—C23—C24	112.1 (2)
F1—C3—C4—C5	177.7 (2)	C25—O7—C24—C23	−61.5 (3)
C2—C3—C4—C5	−2.0 (4)	N5—C23—C24—O7	55.9 (3)
C3—C4—C5—C6	1.3 (4)	C24—O7—C25—C26	61.8 (3)
C4—C5—C6—C1	0.2 (4)	C23—N5—C26—C25	52.7 (3)
C4—C5—C6—C7	−176.9 (2)	P2—N5—C26—C25	−112.2 (2)
C2—C1—C6—C5	−1.2 (4)	O7—C25—C26—N5	−57.1 (3)
C2—C1—C6—C7	175.8 (3)	C30—N6—C27—C28	−53.3 (3)
P1—N1—C7—O1	22.8 (4)	P2—N6—C27—C28	109.8 (3)
P1—N1—C7—C6	−156.19 (19)	C29—O8—C28—C27	−59.2 (3)
C5—C6—C7—O1	11.5 (4)	N6—C27—C28—O8	56.7 (3)
C1—C6—C7—O1	−165.6 (3)	C28—O8—C29—C30	58.9 (3)
C5—C6—C7—N1	−169.6 (2)	C27—N6—C30—C29	52.9 (3)
C1—C6—C7—N1	13.4 (4)	P2—N6—C30—C29	−110.0 (3)
C11—N2—C8—C9	−51.9 (3)	O8—C29—C30—N6	−56.0 (3)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N4—H4N···O2 <sup>i</sup>	0.85 (2)	2.01 (2)	2.855 (3)	176 (3)
N1—H1N···O6 <sup>ii</sup>	0.87 (2)	2.04 (2)	2.870 (3)	159 (3)

Symmetry codes: (i)  $-x+3/2, y-1/2, -z+1/2$ ; (ii)  $-x+3/2, y+1/2, -z+1/2$ .