

Correspondence and Personal Opinion

Influenza and pneumococcal vaccinations in dialysis patients

Viroj Wiwanitkit^{1,2}

¹Department of Laboratory Medicine, Chulalongkorn University, Bangkok, Thailand and ²Hainan Medical University, Haikou, China

Correspondence and offprint requests to: Viroj Wiwanitkit; E-mail: wviroj@yahoo.com

Sir, the report on influenza and pneumococcal vaccinations in dialysis patients is very interesting [1]. Wilmore *et al.* [1] noted that 'most dialysis patients were not up-to-date with both vaccinations.' Of interest, there are many problems preventing the success of vaccination programs for dialysis patients. The attitude of the practitioner seems to be very important [2]. Sometimes, the practitioners do not promote but act against vaccination. In addition, McGrath *et al.* recently reported that hospitalization and skilled nursing care were two important predictors of vaccination [3]. It was reported that 'patients with long hospitalizations or skilled nursing stays were less likely to be vaccinated suggesting evidence of the healthy-user effect [3]' To manage the problem of vaccination coverage, those factors should be carefully considered.

References

1. Wilmore SM, Philip KE, Cambiano V *et al.* Influenza and pneumococcal vaccinations in dialysis patients in a London district general hospital. *Clin Kidney J* 2014; 7: 27–32
2. Ridda I, Lindley IR, Gao Z *et al.* Differences in attitudes, beliefs and knowledge of hospital health care workers and community doctors to vaccination of older people. *Vaccine* 2008; 26: 5633–5640
3. McGrath LJ, Cole SR, Kshirsagar AV *et al.* Hospitalization and skilled nursing care are predictors of influenza vaccination among patients on hemodialysis: evidence of confounding by frailty. *Med Care* 2013; 51: 1106–1113

Received for publication: 28.1.14; Accepted in revised form: 4.2.14

Conflict of interest statement. None declared.