

Figure. De-escalation, escalation, and unchanged outcomes among 39,226 antibiotic admissions.

Rank	N (%)	N Antibiotics		
		Lower	Same	Higher
Lower	10551 (27)	1269 (3)	146 (<1)	
Same	1218 (6)	19703 (50)	3048 (8)	
Higher	110 (<1)	732 (2)	1349 (3)	

Note: De-escalation = green, Unchanged=yellow, Escalation=orange

Disclosures. All authors: No reported disclosures.

961. The Role of Negative Methicillin-Resistant *Staphylococcus aureus* Nasal Surveillance Swabs in Predicting the Need for Empiric Vancomycin Therapy
Darunee Chotipratsakul, MD, MPH¹; Pranita D. Tamma, MD, MHS²;

Avinash Gadala, MS, B.Pharm³; Sara E. Cosgrove, MS⁴; ¹Department of Medicine, Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand; ²Johns Hopkins University School of Medicine, Baltimore, Maryland; ³The Johns Hopkins Health System, Baltimore, Maryland; ⁴Johns Hopkins Medical Institutions, Baltimore, Maryland

Session: 123. Stewardship Tools
Friday, October 6, 2017: 8:30 AM

Background. The role of MRSA nasal surveillance swabs in guiding decisions about need for subsequent vancomycin therapy is unclear. Our objectives were to (1) determine the likelihood that patients with negative MRSA nasal swabs went on to develop MRSA infections during the same hospitalizations to assess if vancomycin therapy could be avoided once the nasal swab result returns negative, (2) assess days of vancomycin that potentially could be avoided, and (3) identify risk factors for having a negative MRSA nasal swab and developing an MRSA infection during the hospital stay.

Methods. This retrospective cohort study was conducted at six intensive care units (ICUs) at a tertiary care hospital in Baltimore from December 2013 to June 2015. MRSA nasal swabs are obtained at the time of admission and weekly thereafter for all ICU patients. The negative predictive value (NPV), defined as the ability of a negative MRSA nasal screening test to correctly predict no subsequent MRSA infection during the hospital stay, was calculated, accounting for the 3-day turnaround time of MRSA nasal surveillance swabs. Days of vancomycin therapy started or continued after 3 days from the first negative MRSA nasal swab were determined by chart review. A matched case-control study was performed to identify risk factors for patients with negative MRSA surveillance cultures who subsequently developed MRSA infections.

Results. Of 11,441 MRSA-nasal swab negative patients, the proportion of subsequent incident MRSA infections was 0.2%. Negative MRSA surveillance swabs had an NPV of 99.4% (95% CI 99.1-99.6%). Among 4,091 MRSA-negative patients receiving vancomycin, vancomycin was started or continued after 3 days since the first MRSA-negative nasal swab in 1,434 patients (35%), translating to 7,377 potentially avoidable vancomycin days. The matched case-control analysis did not identify risk factors associated with subsequent MRSA infection.

Conclusion. At our institution with robust infection control practices and low nosocomial MRSA transmission rates, patients with negative MRSA nasal swabs have a very low likelihood of subsequent MRSA infection during hospitalizations. MRSA nasal swabs can provide useful information when determining whether to initiate or stop empiric vancomycin.

Disclosures. All authors: No reported disclosures.

983. Doing the Same with Less: A Randomized, Multinational, Open-Label, Adjudicator-Blinded Trial of an Algorithm vs. Standard of Care to Determine Treatment Duration for Staphylococcal Bacteremia

Thomas L. Holland, MD¹; Helen W. Boucher, MD, FIDSA²; Issam Raad, MD³; Deverick J. Anderson, MD, MPH⁴; Sara E. Cosgrove, MD, MS⁵; Suzanne Aycocck, MSN⁶; John W. Baddley, MD⁷; Shein-Chung Chow, PhD¹; Vivian H. Chu, MD¹; Paul P. Cook, MD⁸; G. Ralph Corey, MD⁹; Jennifer S. Daly, MD⁹; Ray Y. Hachem, MD¹⁰; Anne-Marie Chaftari, MD¹⁰; James M. Horton, MD¹¹; Timothy C. Jenkins, MD¹²; Jiezhun Gu, PhD⁶; Donald P. Levine, MD¹³; Jose M Miro, MD, PhD^{14,15}; Paul Riska, MD¹⁶; Zachary A. Rubin, MD¹⁷; Mark E. Rupp, MD¹⁸; John Schrank Jr, MD¹⁹; Matthew Sims, MD, PhD²⁰; Dannah Wray, MD²¹; Marcus J. Zervos, MD²² and Vance Fowler Jr., MD, MHS¹; ¹Duke University Medical Center, Durham, North Carolina; ²Tufts Medical Center, Boston, Massachusetts; ³Department of Infectious Diseases, Infection Control and Employee Health, The University of Texas M.D. Anderson Cancer Center, Houston, Texas; ⁴Division of Infectious Diseases, Duke University Medical Center, Durham, North Carolina; ⁵Johns Hopkins Medical Institutions, Baltimore, Maryland; ⁶Duke Clinical Research Institute, Durham, North Carolina; ⁷University of Alabama at Birmingham, Birmingham, Alabama; ⁸Infectious Diseases, Brody School of Medicine at East Carolina University, Greenville, North Carolina; ⁹University of Massachusetts Medical School, Worcester, Massachusetts; ¹⁰University of Texas MD Anderson Cancer Center, Houston, Texas; ¹¹Medical Center, Charlotte, North Carolina; ¹²Denver Health, Denver, Colorado; ¹³Wayne State University, Detroit, Michigan; ¹⁴Hospital Clinic-IDIBAPS, Barcelona, Spain; ¹⁵Hospital Clinic-IDIBAPS, University of Barcelona, Barcelona, Spain; ¹⁶Albert Einstein College of Medicine, Bronx, New York; ¹⁷David Geffen School of Medicine/University of California, Los Angeles, Los Angeles, California; ¹⁸Internal Medicine, Division of Infectious Diseases, University of Nebraska Medical Center, Omaha, Nebraska; ¹⁹Greenville Health System, Greenville, South Carolina; ²⁰Beaumont Health System, Royal Oak, Michigan; ²¹Medical University of

South Carolina, Charleston, South Carolina; ²²Henry Ford Health System, Detroit, Michigan

Session: 132. Advances in Management of Bacteremia and Sepsis
Friday, October 6, 2017: 10:30 AM


Background. The appropriate duration of antibiotics for staphylococcal bloodstream infection (BSI) is unknown. An algorithm to identify patients with staphylococcal BSI who can be safely treated with shorter courses of therapy would improve care and reduce total antibiotic use.

Methods. Adult patients with staphylococcal BSI were randomized to treatment based on algorithm-based therapy (ABT) or to standard of care (SOC). Co-primary outcomes were clinical success, as determined by a blinded Adjudication Committee, and serious adverse event (SAE) rates. The prespecified secondary outcome measure was antibiotic days by treatment group, among patients without complicated BSI. Prespecified durations of therapy in ABT were: *S. aureus* BSI (SAB): uncomplicated: 14 days; complicated: 4-6 weeks. Coagulase-negative staphylococci BSI (CoNSB): simple (1 positive blood culture) (0-3 days), uncomplicated (>1 positive blood culture) (5 days), complicated (7-28 days). Outcomes were compared using intention-to-treat principles. The target sample size was 500 patients, to ensure 90% power for establishing noninferiority within a margin of 15%.

Results. Between April 2011 and March 2017, 509 adults with suspected staphylococcal BSI at 16 sites in the US and Spain were randomized to ABT (N = 255) or SOC (N = 254). There were 116 patients with SAB (23%) and 385 (76%) with CoNSB (Figure 1). Overall success rate in the ABT group was 82.0% vs. 81.5% in the SOC group, difference 0.5%, 95% CI -5.2% to 6.1%. SAEs were reported in 32.9% of ABT vs. 28.3% of SOC patients (OR 1.2, 95% CI 0.9 to 1.8). Among evaluable patients without complicated BSI, mean duration of therapy was 4.4 days in the ABT group vs. 6.4 days in the SOC group (difference -2.0 days, 95% CI -3.3 to -0.7, P = 0.003). Among patients with uncomplicated SAB, treatment durations were similar (15.3 days in ABT vs. 16.3 days in SOC, difference -1 day, 95% CI -3.89 to 1.91, P = 0.497), whereas for uncomplicated CoNSB, duration was shorter in the ABT group (5.3 days in ABT vs. 8.4 days in SOC, difference -3 days, 95% CI -4.87 to -1.34, P < 0.001).

Conclusion. The use of a treatment algorithm for staphylococcal BSI was associated with significant reductions in duration of antibiotic therapy in patients without complicated BSI, without significant differences in overall success or SAEs.

Figure 1. Schematic of Study Design



Disclosures. V. Fowler Jr., NIH: Investigator, Contract HHSN272200900023C

984. Induced Hypothermia in Patients with Septic Shock and Ventilator-demanding Respiratory Failure

Theis Skovsgaard Itenov, MD, PhD¹; Maria Egede Johansen, MD, PhD²; Morten Bestle, MD, PhD³; Katrin Thormar, MD⁴; Lars Hein, MD⁵; Louise Gyldensted, MD⁶; Anne Lindhardt, MD⁴; Henrik Christensen, MD⁷; Stine Estrup Damby, MD⁸; Henrik Planck Pedersen, MD⁹; Matthew Harmon, MD¹⁰; Uday Kant Soni, MD¹¹; Silvia Perez-Protto, MD, PhD¹²; Nikolaj Wesche, MD¹³; Rasmus Ehrenfried Berthelsen, MD¹³; Ulrik Skram, MD¹³; Asger Petersen, MD, PhD¹⁴; Thomas Mohr, MD, PhD¹⁵; Tina Waldau, MD, PhD¹⁵; Lone Museus Poulsen, MD⁸; Ditte Strange, MD¹⁴; Ole Christensen, MD¹⁶; Nicole P Juffermans, MD, PhD¹⁰; Daniel Sessler, MD, PhD¹²; Else Tonnesen, MD, DMSc¹⁷; Dennis Kristensen, MSc¹⁸; Alessandro Cozzi-Lepri, PhD¹⁸; Jens Lundgren, MD, DMSc, Professor¹⁹ and Jens Ulrik Stæhr Jensen, MD, PhD²⁰; ¹Chip, Rigshospitalet - University of Copenhagen, Copenhagen East, Denmark; ²Chip/Department of Infectious Diseases, Rigshospitalet, Copenhagen, Denmark; ³Anesthesia and Intensive Care, North Zealand Hospital, Hillerød, Denmark; ⁴Anesthesia and Intensive Care, Bispebjerg Hospital, Copenhagen NW, Denmark; ⁵Department of Anesthesia and Intensive Care, North Zealand Hospital, Hillerød, Denmark; ⁶Anesthesia and Intensive Care, Herlev & Gentofte Hospital, Hellerup, Denmark; ⁷Anesthesia & Intensive Care, Herlev & Gentofte Hospital, Hellerup, Denmark; ⁸Anesthesia and Intensive Care, Zealand University Hospital, Køge, Denmark; ⁹Anesthesia and Intensive Care, Roskilde Hospital, Roskilde, Denmark; ¹⁰Anesthesia and Intensive Care, Academic Medical Center, Amsterdam, the Netherlands; ¹¹Anesthesia and Intensive Care, Horsens Hospital, Horsens, Denmark; ¹²Outcomes Research, Cleveland Clinic, Cleveland, Ohio; ¹³Anesthesia and Intensive Care, North Zealand Hospital, Hillerød, Denmark; ¹⁴Anesthesia and Intensive Care, Bispebjerg Hospital, Copenhagen NW, Denmark; ¹⁵Anesthesia and Intensive Care, Gentofte Hospital, Hellerup, Denmark; ¹⁶Anesthesia and Intensive Care, Roskilde Hospital, Roskilde, Denmark; ¹⁷Anesthesia and Intensive Care, Aarhus University Hospital, Aarhus, Denmark; ¹⁸Chip/Department of Infectious Diseases, Rigshospitalet - University of Copenhagen,