



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

COVID-19 and social inequalities: a complex and dynamic interaction



COVID-19, in all its dimensions, including incidence, testing, and severity, is known to be associated with social inequalities.¹⁻⁶ Such inequalities are associated with differential exposure to the virus, greater susceptibility to infection, more frequent comorbidities associated with severe outcomes, and disparate access to care.

In *The Lancet Public Health*, Stéphanie Vandentorren and colleagues⁷ investigated the association between an area-based deprivation indicator and SARS-CoV-2 incidence, positivity and testing rates in France. To our knowledge, this is the first large-scale study to explore the issue of social inequalities relative to COVID-19 testing and the dynamics of the SARS-CoV-2 pandemic in France. The study benefited from the quality and exhaustiveness of national databases. The authors used data from the *Système d'Information de Dépistage Populationnel*, which records the results of all SARS-CoV-2 tests in France, for the period May 2020 to April 2021, during which almost 71 million SARS-CoV-2 tests were recorded, including 5 000 972 positive tests. Vandentorren and colleagues showed that SARS-CoV-2 incidence, positivity rates, and testing rates differed in the most socially deprived areas (measured by the European Deprivation Index [EDI]) when compared with the least deprived areas. A clear social gradient was identified, whereby individuals living in the most deprived areas had the highest risk of contracting SARS-CoV-2, but a concomitant lower likelihood of being tested. This social gradient was more pronounced in densely populated areas than in sparsely populated areas.

Vandentorren and colleagues analysed the weekly dynamic of SARS-CoV-2 incidence, positivity rates, and testing rates by deprivation and population density group, which showed that the measures implemented by the French Government to manage the pandemic, especially lockdowns, had a differential impact on these outcomes in the most and the least deprived areas, and that this impact varied according to population density. In less populated areas, during the second lockdown SARS-CoV-2 incidence, positivity rates, and testing rates were lower in the most deprived areas than the least

deprived areas. However, during the third lockdown, incidence and testing rates were higher in the most deprived areas, with a similar positivity rate. In densely populated municipalities, while testing rates were similar across deprivation quintiles during the second lockdown, they were lower in the most deprived than less deprived areas around the time of the third lockdown.

In this study, some area codes were excluded, therefore the authors compared the distribution of the total number of tests and the number of positive results by age and sex between the missing and included data. The differences, although statistically significant, were small. It would have been interesting to know how comparable the excluded areas were with regard to other socioeconomic variables, although it is unlikely that this issue biased the results. Moreover, the impact of each social variable in the EDI could not be studied, but it could be interesting to explore further. Socioeconomic data were not available at the individual level, therefore, the authors used an ecological measure of deprivation at the area level, precluding a detailed study of the mechanisms at work. Further studies are needed to investigate the complex spatiotemporal interactions suggested by the findings. Additionally, considering that access to COVID-19 vaccine might also be affected by socioeconomic inequalities, it would be of great interest to incorporate vaccination uptake into the analyses.

Vandentorren and colleagues have provided new information regarding the association between social inequalities and health in the context of a major health crisis. Importantly, the study also provided insights with regard to the dynamics involved, and suggest a worsening of the negative influence of socioeconomic inequalities on COVID-19 testing and incidence during the pandemic. Indeed, in densely populated municipalities, the testing rate decreased in the most deprived areas—signalling a deterioration in access to preventive measures. These results might also be associated with an increase in social inequalities resulting from the pandemic itself, since the crisis also increased income inequalities in France.⁸ In 2013, the French Government⁹ announced their intention to

Published Online
February 14, 2022
[https://doi.org/10.1016/S2468-2667\(22\)00033-0](https://doi.org/10.1016/S2468-2667(22)00033-0)
See [Articles](#) page e240

For more on the French Public Health Law see <http://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000028372809>

reduce social inequalities in health as part of the Public Health Law. However, doing so will require sustained and coordinated multisectoral initiatives, many of which have yet to be elaborated on and implemented. These challenges might explain the results described by Vandentorren and colleagues and findings in other national reports,⁸ including structural barriers to vaccination. Such studies might be useful for any future strategies to tackle such inequalities.

We declare no competing interests.

Copyright © 2022 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND 4.0 license.

**Catherine Quantin, Pascale Tubert-Bitter*
catherine.quantin@chu-dijon.fr

Service de Biostatistiques et d'Information Médicale (DIM), CHU Dijon Bourgogne, INSERM, Université de Bourgogne, Dijon 21079, France (CQ); Université Paris-Saclay, UVSQ, Université Paris-Sud, Inserm, High-Dimensional Biostatistics for Drug Safety and Genomics, CESP, Villejuif, France (CQ, PT-B)

1 Bamba C, Riordan R, Ford J, Matthews F. The COVID-19 pandemic and health inequalities. *J Epidemiol Community Health* 2020; **74**: 964–68.

2 Wilkins CH, Friedman EC, Churchwell AL, et al. A systems approach to addressing Covid-19 health inequities. *NEJM Catal Innov Care Deliv* 2021; **2**: 1–17.

3 Riou J, Panczak R, Althaus CL, et al. Socioeconomic position and the COVID-19 care cascade from testing to mortality in Switzerland: a population-based analysis. *Lancet Public Health* 2021; **6**: e683–91.

4 Green MA, García-Fiñana M, Barr B, et al. Evaluating social and spatial inequalities of large scale rapid lateral flow SARS-CoV-2 antigen testing in COVID-19 management: an observational study of Liverpool, UK (November 2020 to January 2021). *Lancet Reg Health Eur* 2021; **6**: 100107.

5 Lewis NM, Friedrichs M, Wagstaff S, et al. Disparities in COVID-19 incidence, hospitalizations, and testing, by area-level deprivation—Utah, March 3–July 9, 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**: 1369–73.

6 Burström B, Tao W. Social determinants of health and inequalities in COVID-19. *Eur J Public Health* 2020; **30**: 617–18.

7 Vandentorren S, Smaili S, Chatignoux E, et al. The effect of social deprivation on the dynamic of SARS-CoV-2 infection in France: a population-based analysis. *Lancet Public Health* 2022; published online Feb 14. [https://doi.org/10.1016/S2468-2667\(21\)00007-X](https://doi.org/10.1016/S2468-2667(21)00007-X).

8 Dubost C-L, Pollak C, Rey S. Les inégalités sociales face à l'épidémie de Covid-19: état des lieux et perspectives. <https://drees.solidarites-sante.gouv.fr/publications/les-dossiers-de-la-drees/les-inegalites-sociales-face-lepidemie-de-covid-19-etat-des> (accessed Jan 25, 2022).

9 Touraine M. Health inequalities and France's national health strategy. *Lancet* 2014; **383**: 1101–02.