




Draft Genome Sequences of 13 Plant-Associated Actinobacteria of the Family *Microbacteriaceae*

 Sergey V. Tarlachkov,^{a,b} Irina P. Starodumova,^a Lubov V. Dorofeeva,^a Natalia V. Prisyazhnaya,^a Semen A. Leyn,^{c,d} Jaime E. Zlamal,^e Sebastian Albu,^e Steven A. Nadler,^f Sergei A. Subbotin,^{e,f,g} Lyudmila I. Evtushenko^a

^aAll-Russian Collection of Microorganisms (VKM), G. K. Skryabin Institute of Biochemistry and Physiology of Microorganisms, Pushchino Scientific Center for Biological Research of the Russian Academy of Sciences, Pushchino, Russia

^bBranch of Shemyakin and Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences, Pushchino, Russia

^cInfectious and Inflammatory Disease Center, Sanford Burnham Prebys Medical Discovery Institute, La Jolla, California, USA

^dA. A. Kharkevich Institute for Information Transmission Problems, Russian Academy of Sciences, Moscow, Russia

^eCalifornia Department of Food and Agriculture, Sacramento, California, USA

^fDepartment of Entomology and Nematology, University of California, Davis, California, USA

^gCenter of Parasitology of A. N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia

ABSTRACT Draft genome sequences of 13 bacterial strains from the family *Microbacteriaceae* were generated using Illumina technology. The genome sizes varied from 3.0 to 4.8 Mb, and the DNA G+C content was 68.1 to 72.5%. The sequences obtained will contribute to the development of genome-based taxonomy and understanding of molecular interactions between bacteria and plants.

Members of the family *Microbacteriaceae* (class *Actinobacteria*) are widely distributed in various terrestrial and aquatic ecosystems and often occur in association with plants as endophytes and pathogens (1–3).

Novel strains of *Microbacteriaceae* were recovered from eight different plants of five families (Table 1) collected in various sites in California. *Rathayibacter* sp. strain VKM Ac-2835 was isolated from a *Malus* sp. with symptoms of bacterial wetwood disease by macerating several pieces of symptomatic superficial bark tissue in a sterile aqueous solution and then plating it onto *Pseudomonas* F agar (Becton, Dickinson, USA) amended with cycloheximide (100 mg/liter). The remaining strains were isolated from plants without visible symptoms of diseases, as described (3, 4), but Reasoner's 2A (R2A) agar (Fluka Analytical, USA) was used as the plating medium for isolation. *Rathayibacter agropyri* CA-4^T (=VKM Ac-2828^T) was kindly provided by T. D. Murray. For preservation, strains were grown on R2A agar and lyophilized using standard techniques. All strains were deposited in the All-Russian Collection of Microorganisms (VKM; <http://www.vkm.ru>).

Biomass for DNA extraction was grown in liquid peptone-yeast medium (5) inoculated with cells from a single colony, followed by cultivation at 28°C for 18 to 20 h on a rotary shaker. Genomic DNA was extracted using a QIAamp DNA minikit (Qiagen, Germany). DNA libraries for strains VKM Ac-2828^T, VKM Ac-2835, and VKM Ac-2836 were prepared in-house using a NEBNext Ultra II FS DNA library prep kit for Illumina (New England Biolabs, USA) following the protocol for use with inputs of ≥ 100 ng with modifications as described previously (6). Pooled DNA libraries were sequenced by Novogene Co., Ltd., on an Illumina HiSeq X Ten instrument to obtain 150-bp paired-end reads. For the remaining strains, DNA library construction and sequencing were conducted by Novogene Co., Ltd. Libraries were generated using a NEBNext DNA library prep kit for Illumina (New England Biolabs) following the manufacturer's recommendations. Pooled DNA libraries were sequenced on an Illumina NovaSeq 6000 instrument to obtain 150-bp paired-end reads.

Citation Tarlachkov SV, Starodumova IP, Dorofeeva LV, Prisyazhnaya NV, Leyn SA, Zlamal JE, Albu S, Nadler SA, Subbotin SA, Evtushenko LI. 2020. Draft genome sequences of 13 plant-associated actinobacteria of the family *Microbacteriaceae*. *Microbiol Resour Announc* 9:e00795-20. <https://doi.org/10.1128/MRA.00795-20>.

Editor David A. Baltrus, University of Arizona

Copyright © 2020 Tarlachkov et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sergey V.

Tarlachkov, sergey@tarlachkov.ru, or Sergei A. Subbotin, sergei.a.subbotin@gmail.com.

Received 13 July 2020

Accepted 27 August 2020

Published 17 September 2020

TABLE 1 Characteristics and DDBJ/ENA/GenBank accession numbers of the genomes

Organism	Associated plant (family)	No. of reads	Coverage (x)	No. of scaffolds	Scaffold N ₅₀ (bp)	Genome size (Mbp)	G+C content (%)	No. of proteins	SRA accession no.	GenBank accession no.
<i>Rathayibacter agropyri</i> VKM Ac-2828 ^T	<i>Paspopyrum smithii</i> (Poaceae)	14,362,336	578	25	656,029	3.0	68.1	2,835	SRX8466800	JABRPL0000000000
<i>Rathayibacter</i> sp. VKM Ac-2835	<i>Malus domestica</i> (Rosaceae)	12,953,304	410	6	1,292,012	4.3	72.2	3,849	SRX8466801	JABSNO0000000000
<i>Rathayibacter</i> sp. VKM Ac-2857	<i>Brachypodium distachyon</i> (Poaceae)	21,681,636	692	8	1,409,124	4.6	72.1	4,080	SRX8466811	JABMLE0000000000
<i>Rathayibacter</i> sp. VKM Ac-2856	<i>Brachypodium distachyon</i> (Poaceae)	20,040,560	689	9	783,576	4.3	72.5	3,806	SRX8466810	JABMLF0000000000
<i>Rathayibacter</i> sp. VKM Ac-2858	<i>Brachypodium distachyon</i> (Poaceae)	22,149,976	762	9	783,576	4.3	72.5	3,806	SRX8466812	JABMLD0000000000
<i>Curtobacterium</i> sp. VKM Ac-2852	<i>Avena fatua</i> (Poaceae)	19,012,470	724	7	1,020,126	3.9	70.8	3,580	SRX8466807	JABMLI0000000000
<i>Curtobacterium</i> sp. VKM Ac-2861	<i>Marah</i> sp. (Cucurbitaceae)	17,600,848	642	15	624,972	4.0	70.8	3,739	SRX8466804	JABMLA0000000000
<i>Frigoribacterium</i> sp. VKM Ac-2836	<i>Fragaria vesca</i> (Rosaceae)	28,426,948	1,100	10	1,120,753	3.3	70.4	3,010	SRX8466805	JABRPK0000000000
<i>Frigoribacterium</i> sp. VKM Ac-2859	<i>Brachypodium distachyon</i> (Poaceae)	14,404,054	638	5	1,688,707	3.3	71.3	3,027	SRX8466802	JABMLC0000000000
<i>Frigoribacterium</i> sp. VKM Ac-2860	<i>Brachypodium distachyon</i> (Poaceae)	14,248,970	632	5	1,688,688	3.3	71.3	3,029	SRX8466803	JABMLB0000000000
<i>Herbiconiux</i> sp. VKM Ac-2851	<i>Soliva sessilis</i> (Asteraceae)	18,251,960	623	9	1,342,178	4.3	70.7	4,032	SRX8466806	JABMLJ0000000000
<i>Microbacteriaceae</i> bacterium VKM Ac-2854	<i>Myosotis</i> sp. (Boraginaceae)	42,551,862	1,319	18	721,153	4.8	69.6	4,359	SRX8466808	JABMLH0000000000
<i>Microbacteriaceae</i> bacterium VKM Ac-2855	<i>Myosotis</i> sp. (Boraginaceae)	17,323,178	542	26	386,786	4.7	68.3	4,255	SRX8466809	JABMLG0000000000




FIG 1 Phylogenomic tree based on genomes of *Rathayibacter* strains sequenced in this work (in bold) and members of other validly published and some putative (6, 19, 20) *Rathayibacter* species. The tree was inferred by the balanced minimum evolution method using JolyTree 1.1.181205ac (21) with branch lengths scaled to the estimated number of substitutions per site. Branch support values (rate of elementary quartets) above 0.5 are indicated at the branch points. The genomic sequence of *Clavibacter michiganensis* subsp. *sepedonicus* ATCC 33113^T (GenBank accession numbers [AM849034.1](#) to [AM849036.1](#)) served as an outgroup (not shown) to root the tree.

Default parameters were used for all software unless otherwise specified. The quality of the reads was checked with FastQC 0.11.8 (7). Adapter sequences and low-quality regions in the raw reads were cut with Trimmomatic 0.39 (8) with the following options: ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10, SLIDINGWINDOW:4:15, and MINLEN:50. Trimmed reads were assembled using SPAdes 3.14.1 (9) with the following options: --cov-cutoff, auto; and --careful. The quality of assembly was assessed with QUASt 5.0.2 (10). Assemblies were annotated with NCBI PGAP (11) and the RAST Web server (12, 13). The pairwise similarity between the 16S rRNA gene sequences was determined using TaxonDC 1.3.1 (14). The average nucleotide identity (ANI) and digital DNA-DNA hybridization (dDDH) values were calculated using the JSpecies 1.2.1 (15) and GGDC 2.1 (16) tools, respectively.

Accession numbers and characteristics of the genomes are provided in Table 1. Figure 1 shows the phylogenomic positions of *Rathayibacter* strains sequenced here within the genus *Rathayibacter*. Four newly isolated strains clustered with *Rathayibacter festucae* but exhibited average nucleotide identity values (90.6 to 93.4%) and digital DNA-DNA hybridization levels (41.7 to 52.1%) to *R. festucae* DSM 15932^T not exceeding the thresholds for species delineation (17). No genome sequences of relevant type strains of the *Curtobacterium*, *Frigoribacterium*, and *Herbiconiux* species are available to precisely determine the phylogenomic positions of the remaining strains sequenced in this work (Table 1).

A BLAST search confirmed the presence of a genomic cluster comprising a complete suite of tunicaminylluracil-related biosynthetic genes in *R. agropyri* CA-4^T as already reported by Tancos et al. (18) for this strain. This gene cluster is not present in any other genomes sequenced in this work.

Further whole-genome sequencing of other *Microbacteriaceae* along with comparative genomic and phenotypic analyses of putative and known species with validly published names will result in valid descriptions of the revealed new taxa, contributing to the development of the genome-based taxonomy of prokaryotes.

Data availability. These whole-genome shotgun projects have been deposited in DDBJ/ENA/GenBank under the accession numbers listed in Table 1.

ACKNOWLEDGMENTS

We thank Andrei L. Osterman for his valuable help during work with this project, Timothy D. Murray for providing the *Rathayibacter agropyri* CA-4^T strain, and Dean Kelch and Robert Price for plant identification.

This work was supported by the U.S. Department of Agriculture Animal and Plant Health Inspection Service according to the research project AP18PPQS&T00C159 (18-0422-000-FR) "Enhancing Diagnostics of Plant Pathogenic Bacteria of the Genus *Rathayibacter*" (principal investigator [PI], S.A.S.).

REFERENCES

- Evtushenko LI. 2015. *Microbacteriaceae*. In Whitman WB, Rainey F, Kämpfer P, Trujillo M, Chun J, DeVos P, Hedlund B, Dedysh S (ed), *Bergey's manual of systematics of archaea and bacteria*. Wiley, Hoboken, NJ. <https://doi.org/10.1002/9781118960608.fbm00035>.
- Murray TD, Schroeder BK, Schneider WL, Luster DG, Sechler A, Rogers EE, Subbotin SA. 2017. *Rathayibacter toxicus*, other *Rathayibacter* species inducing bacterial head blight of grasses, and the potential for livestock poisonings. *Phytopathology* 107:804–815. <https://doi.org/10.1094/PHYTO-02-17-0047-RVW>.
- Dorofeeva LV, Starodumova IP, Krauzova VI, Prisyazhnaya NV, Vinokurova NG, Lysanskaya VY, Tarlachkov SV, Evtushenko LI. 2018. *Rathayibacter oskolensis* sp. nov., a novel actinobacterium from *Androsace koso-poljanskii* Ovcz. (Primulaceae) endemic to Central Russian Upland. *Int J Syst Evol Microbiol* 68:1442–1447. <https://doi.org/10.1099/ijsem.0.002681>.
- Dorofeeva LV, Krausova VI, Evtushenko LI, Tiedje JM. 2003. *Agromyces albus* sp. nov., isolated from a plant (*Androsace* sp.). *Int J Syst Evol Microbiol* 53:1435–1438. <https://doi.org/10.1099/ijms.0.02428-0>.
- Naumova IB, Kuznetsov VD, Kudrina KS, Bezzubenkova AP. 1980. The occurrence of teichoic acids in *Streptomyces*. *Arch Microbiol* 126:71–75. <https://doi.org/10.1007/BF00421893>.
- Tarlachkov SV, Starodumova IP, Dorofeeva LV, Prisyazhnaya NV, Leyn SA, Zlamal JE, Elane ML, Osterman AL, Nadler SA, Subbotin SA, Evtushenko LI. 2020. Complete and draft genome sequences of 12 plant-associated *Rathayibacter* strains of known and putative new species. *Microbiol Resour Announc* 9:e00316-20. <https://doi.org/10.1128/MRA.00316-20>.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <https://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
- Tarlachkov SV, Starodumova IP. 2017. TaxonDC: calculating the similarity value of the 16S rRNA gene sequences of prokaryotes or ITS regions of fungi. *J Bioinform Genom* 3:1–4. <https://doi.org/10.18454/jbg.2017.3.5.1>.
- Richter M, Rosselló-Móra R. 2009. Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A* 106:19126–19131. <https://doi.org/10.1073/pnas.0906412106>.
- Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14:60. <https://doi.org/10.1186/1471-2105-14-60>.
- Chun J, Oren A, Ventosa A, Christensen H, Arahal DR, da Costa MS, Rooney AP, Yi H, Xu XW, De Meyer S, Trujillo ME. 2018. Proposed minimal standards for the use of genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 68:461–466. <https://doi.org/10.1099/ijsem.0.002516>.
- Tancos MA, Sechler AJ, Davis EW, II, Chang JH, Schroeder BK, Murray TD, Rogers EE. 2019. The identification and conservation of tunicaminyluracil-related biosynthetic gene clusters in several *Rathayibacter* species collected from Australia, Africa, Eurasia, and North America. *Front Microbiol* 10:2914. <https://doi.org/10.3389/fmicb.2019.02914>.
- Starodumova IP, Tarlachkov SV, Prisyazhnaya NV, Dorofeeva LV, Ariskina EV, Chizhov VN, Subbotin SA, Evtushenko LI, Vasilenko OV. 2017. Draft genome sequence of *Rathayibacter* sp. VKM Ac-2630 isolated from leaf gall induced by the knapweed nematode *Mesoanguina picridis* on *Acropilton repens*. *Genome Announc* 5:e00650-17. <https://doi.org/10.1128/genomeA.00650-17>.
- Vasilenko OV, Starodumova IP, Tarlachkov SV, Dorofeeva LV, Avtukh AN, Evtushenko LI. 2016. Draft genome sequence of "*Rathayibacter tanacetii*" strain VKM Ac-2596 isolated from *Tanacetum vulgare* infested by a foliar nematode. *Genome Announc* 4:e00512-16. <https://doi.org/10.1128/genomeA.00512-16>.
- Crisuolo A. 2019. A fast alignment-free bioinformatics procedure to infer accurate distance-based phylogenetic trees from genome assemblies. *Res Ideas Outcomes* 5:e36178. <https://doi.org/10.3897/rio.5.e36178>.