



Does Raising the Arms Modify Head Tremor Severity in Cervical Dystonia?

BRIEF REPORT

ELIZABETH CISNEROS

JEANNE P. VU

HA YEON LEE

QIYU CHEN

CASEY N. BENADOF

ZHENG ZHANG

EMILY A. PETTITT

SUBHAGYA K. JOSHI

RICHARD L. BARBANO

JOSEPH JANKOVIC

HYDER A. JINNAH

JOEL S. PERLMUTTER

BRIAN D. BERMAN

ABHIMANYU MAHAJAN

CHRISTOPHER G. GOETZ

GLENN T. STEBBINS

CYNTHIA L. COMELLA

DAVID A. PETERSON

*Author affiliations can be found in the back matter of this article

]u[ubiquity press

ABSTRACT

Background: A defining characteristic of dystonia is its position-dependence. In cervical dystonia (CD), sensory tricks ameliorate head tremor (HT). But it remains unknown whether raising the arms alone has the same impact.

Methods: We analyzed data collected from patients enrolled by the Dystonia Coalition. For 120 patients with HT, we assessed how raising their arms without touching their head changed their HT severity.

Results: Forty-eight out of 120 patients exhibited changes in HT severity when raising their arms. These patients were more likely to exhibit decreases in HT severity ($N = 35$) than increases ($N = 13$, $\chi^2(1, N = 48) = 10.1$, $p = 0.002$). Demographic factors and sensory trick efficacy were not significant predictors of whether HT severity changed when raising their arms.

Discussion: Raising the arms without touching the head is a posture that can reduce HT severity in some CD patients. Our results extend the concept of position-dependent motor symptoms in CD to include the position of the arms.

Highlights

Head tremor (HT) is a prevalent symptom of cervical dystonia (CD) that can often be disabling. This study demonstrates that raising the arms without touching the head is a posture that can reduce HT severity in some CD patients. Our findings also identify a novel form of position-dependence in CD.

CORRESPONDING AUTHOR:

David A. Peterson, PhD

CNL-S, Salk Institute for Biological Studies, 10010 N. Torrey Pines Rd, La Jolla, CA 92037, USA

dap@salk.edu

KEYWORDS:

head tremor; cervical dystonia; dystonic tremor; posture

TO CITE THIS ARTICLE:

Cisneros E, Vu JP, Lee HY, Chen Q, Benadof CN, Zhang Z, Pettitt EA, Joshi SK, Barbano RL, Jankovic J, Jinnah HA, Perlmutter JS, Berman BD, Mahajan A, Goetz CG, Stebbins GT, Comella CL, Peterson DA. Does Raising the Arms Modify Head Tremor Severity in Cervical Dystonia? *Tremor and Other Hyperkinetic Movements*. 2021; 11(1): 21, pp. 1–6. DOI: <https://doi.org/10.5334/tohm.623>

INTRODUCTION

Cervical dystonia (CD) is a common form of focal dystonia characterized by muscle spasms that lead to abnormal head posture accompanied by head tremor (HT) in approximately 60% of patients [1, 2]. HT is often a disabling aspect of CD that has highly variable responses to conventional treatments [3]. Many patients develop a sensory trick (alleviating maneuver), which is a maneuver, such as touching the cheek, chin, or other area on the face or head, used to normalize abnormal head posture [4]. A sensory trick can also decrease HT severity [5]. Among CD patients with HT who report having a trick, about half exhibit reduced HT severity during execution of the trick while the arms were raised prior to the hand touching the face to execute the trick [5]. This suggests that the widely recognized position-dependent nature of CD motor symptoms [6] includes the position of the arms. In this study, our aim was to determine whether raising the arms alone without touching any part of the head has the potential to decrease HT severity.

METHODS

We analyzed data collected from 206 patients with isolated CD enrolled across 10 sites in the Dystonia Coalition's previous rating scale validation study (<https://clinicaltrials.gov/ct2/show/NCT01373424>). All patients provided informed consent prior to their participation in the study. The protocols for original data collection and subsequent analyses were approved by the Human Research Protection Offices at the Washington University School of Medicine (WUSM), Rush University Medical Center (RUMC), and the University of California, San Diego (UCSD; protocol 111255X). All patients were assessed three or more months after their last botulinum neurotoxin (BoNT) injections, and were excluded if they had prior DBS surgery. Patients were on a variety of medications in the following categories: Benzodiazepines, GABAergic drugs (Baclofen, pregabalin), beta adrenergic blockers, and dopamine receptor drugs. We calculated the duration patients had CD by subtracting their reported age of onset from their age at the time the protocol was administered.

Videos of patients were recorded using a standardized examination protocol. The examination included two segments: 1) a baseline segment where patients were instructed to let their head drift to its natural dystonic position, and 2) an "arms up" maneuver segment where patients were instructed to raise their arms into a series of horizontal positions including arms held straight out in

pronation, straight out in supination, and in an elbows-bent winged posture. Video segments were annotated using ELAN software version 4.9.4. All video recordings were reviewed by a movement disorders neurologist (CLC) with expertise in dystonia who assessed HT presence and severity during the baseline and arms up segments. During the arms up segment, changes in HT severity were categorized as whether or not they changed by at least an estimated 30% in comparison to the baseline segment, and were scored as decreased, unchanged, or increased.

The examination also included a segment where patients were instructed to demonstrate their most effective sensory trick or, if unaware of a trick, prompted to touch their right cheek, left cheek, and back of their head. The sensory trick was scored using the Toronto Western Spasmodic Torticollis Rating Scale (TWSTRS-2) [7] as complete (0), moderate [1], mild [2], minimal [3], or no improvement of posture [4] by one or more tricks. Because of challenges with multi-rater scoring [2, 8], the sensory trick was scored in a separate assessment by the same movement disorders neurologist (CLC), blind to the patients' original rating.

We tested the null hypothesis that no patients exhibited change in HT with arms up using a binomial test. For those exhibiting change in HT, we determined the likelihood of HT severity increasing or decreasing compared to baseline during the arms up segment using a chi-squared test. For all subsequent analyses, we examined several factors to determine whether or not they were associated with either a change in HT severity and/or whether they were more likely to be associated with an increase or decrease in HT severity. We examined the effect of type of head posture (anterocollis, retrocollis, laterocollis, rotation), the presence of hand tremor, gender, age at onset, and disease duration (and the interaction term between age at onset and disease duration) using a nominal logistic regression. For those patients who reported having a sensory trick, we examined the effect of the trick's efficacy using a nominal logistic regression. All statistical analysis was performed with SAS's John's Macintosh Project (JMP) version Pro 15. We used an alpha level of 0.05 to determine significance.

RESULTS

Of the 206 patients, 84 were excluded due to the lack of HT, and two were excluded because they were not prompted to perform the arms up segment, leaving 120 patients. Demographics, TWSTRS-2 total motor severity scores, and type of predominant head posture for the patient cohort are provided (**Table 1**).

A proportion of patients significantly greater than zero (48 out of 120) exhibited change in HT severity during the arms up segment ($p < 0.0001$; **Figure 1**). Of those that exhibited change in HT severity, it was more likely for them to exhibit a decrease in HT severity ($N = 35$ patients) than an increase ($N = 13$, $\chi^2(1, N = 48) = 10.1$, $p = 0.002$; **Figure 1**).

Type of head posture, the presence of hand tremor, gender, age at onset, disease duration, and the interaction term between age at onset and disease duration were not found to be significant predictors of whether there would be change or no change in HT severity ($\chi^2(9, N = 120) =$

4.34 , $p = 0.887$) or an increase or decrease in HT severity ($\chi^2(9, N = 48) = 14.9$, $p = 0.0953$).

Out of 120 patients, 106 reported having a sensory trick. The sensory trick score was not found to be a significant predictor of whether there would be change or no change in HT ($\chi^2(1, N = 106) = 0.08$, $p = 0.776$), or an increase or decrease in HT severity ($\chi^2(1, N = 42) = 0.131$, $p = 0.718$).

DISCUSSION

In this study, we assessed the effect of the arms up postural maneuver on CD HT severity. We found that HT severity changed for 40% of patients with HT. Of those whose HT severity changed with arms up, almost three times as many exhibited decreased HT severity compared to those exhibiting increased HT severity. Our results show that raising the arms alone, without touching the head, is a posture that can reduce HT severity in some patients with CD. Our finding is consistent with the prior observation that about half of CD patients with HT and a sensory trick exhibit reduced HT severity during execution of the sensory trick while the arms were being raised but prior to the hand touching the face [5]. Our findings also extend the widely recognized position-dependent nature of CD motor symptoms [6] to include the position of the arms. We also found that sensory trick efficacy was not a significant predictor of the effect of arms up on HT severity. Although tricks and raising the arms share elements of proprioceptive dynamics, our results suggest that they have at least partly independent influences on CD motor symptoms.

It remains unclear *how* the arms up maneuver reduces HT severity. It could be due to the role of proprioception, motor programming, biomechanical interactions, or

DEMOGRAPHICS	MEAN (SD)
Age at Onset (y)	42.9 (12.3)
Age at Exam (y)	61.1 (9.83)
Disease Duration (y)	18.3 (11.8)
Gender (F/M)	(91/29)
TWSTRS-2 Motor Total	17.4 (5.27)
Range (out of possible 0–48)	3–29
CERVICAL DYSTONIA POSTURE TYPE	COUNT
Rotation	111
Laterocollis	107
Retrocollis	47
Anterocollis	38

Table 1 Patient Characteristics.

Note: SD = standard deviation; TWSTRS-2 = Toronto Western Spasmodic Torticollis Rating Scale; each CD posture type category represents patients that have at least that posture type, however, most patients have a mixture of the four and are counted multiple times.




Figure 1 (A) Frequency of patients that exhibited change or no change in HT severity (B) Percentage of patients that exhibited an increase or decrease in HT severity. (A) Those with HT are more likely to experience no change in HT severity during the arms up segment. **(B)** Those who exhibited change are more likely to have a decrease in HT severity during the arms up segment.

some interaction thereof. Proprioceptive sensory input changes when raising the arms. Patients with CD have abnormal head proprioception evidenced by abnormal response to neck muscle vibration [9]. In addition, while performing out-of-sight upper limb reaching movement towards a target, CD patients showed significant trajectory abnormalities in comparison to normal controls [10]. One study showed that although tactile abnormalities were present in CD patients with or without tremor, proprioceptive dysfunction was observed in only those with tremor [11]. Motor programming has been suggested as a potential mechanism for the sensory trick because an effective sensory trick relates to not only increased parietal and bilateral visual cortex activation but also decreased activation in contralateral supplementary motor area and primary sensorimotor cortices [12]. Finally, biomechanical interactions between muscles involved in raising the arms and those producing head tremor may also play a role in how raising the arms decreases HT severity. Raising the arms involves levator and trapezius muscles, which are also implicated in head positioning. Although the arms up maneuver and sensory tricks share elements of upper limb proprioception, motor programming, and biomechanics, the arms up maneuver in our study has not been documented as a self-initiated maneuver that patients use and is therefore not considered a conventional sensory trick.

This study has a few limitations. First, we measured changes in HT severity categorically instead of numerically. More precise measures of HT severity may be more sensitive for detecting these changes. Second, we do not have data on how the sensory trick affected the severity of HT and head posture separately or how the arms up maneuver affected head posture. This information could be used to determine whether the arms up maneuver and sensory tricks have differential effects on HT even for individual patients. Third, although in our cross-sectional design we found that the likelihood that raising arms decreases HT severity is independent of demographic variables, HT likelihood in CD is higher for older age at onset [13, 14], longer disease duration [14, 15], and female gender [1, 14]. Longitudinal studies could determine whether individual patients exhibit changes over time in the relationship of HT to the arms up maneuver.

Future studies on the role of proprioception and motor programming in CD could benefit from experimental designs and analyses that go beyond this study. In terms of experimental design, one could passively move patients arms up or have them imagine moving their arms up. This would help determine if, like the sensory trick, imagined motor programming can modulate HT severity. In terms of new analyses, more fine-grained methods to capture

change in HT – such as with computer vision-based software that can quantify HT severity from videos – will also help us determine how proprioception and motor programming influence their CD symptoms.

DATA ACCESSIBILITY STATEMENT

Original patient data available from the Dystonia Coalition upon reasonable request.

ETHICS AND CONSENT

This study was performed in accordance with the ethical standards detailed in the Declaration of Helsinki. The authors' institutional ethics committee has approved this study and all patients have provided written informed consent.

ACKNOWLEDGEMENTS

We gratefully acknowledge Laura Wright and Matt Hicks (WUSM) for assistance with providing data access. We would also like to acknowledge the Max Planck Institute for Psycholinguistics in Nijmegen for the development of ELAN and the SAS Institute in Cary, NC for the development of JMP.

FUNDING INFORMATION


The work was supported in part by the Dystonia Coalition, a consortium of the Rare Diseases Clinical Research Network (RDCRN) organized by the Office of Rare Diseases Research (ORDR) at the National Center for Advancing Clinical and Translational Studies (U54 TR001456) in collaboration with the National Institute for Neurological Diseases and Stroke (U54 NS065701 and U54 NS116025). This work was also supported by the Office of the Assistant Secretary of Defense for Health Affairs, through the Peer Reviewed Medical Research Program under Award No. W81XWH-17-1-0393. Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the Department of Defense.

COMPETING INTERESTS

The authors have no competing interests to declare.

AUTHOR AFFILIATIONS

Elizabeth Cisneros  orcid.org/0000-0001-9466-9116
Institute for Neural Computation, University of California, San Diego,
La Jolla, CA, USA

Jeanne P. Vu  orcid.org/0000-0003-2665-2763
Institute for Neural Computation, University of California, San Diego,
La Jolla, CA, USA

Ha Yeon Lee
Institute for Neural Computation, University of California, San Diego,
La Jolla, CA, USA

Qiyu Chen
Institute for Neural Computation, University of California, San Diego,
La Jolla, CA, USA

Casey N. Benadof
Institute for Neural Computation, University of California, San Diego,
La Jolla, CA, USA


Zheng Zhang
Institute for Neural Computation, University of California, San Diego,
La Jolla, CA, USA

Emily A. Pettitt
Institute for Neural Computation, University of California, San Diego,
La Jolla, CA, USA


Subhagya K. Joshi
Institute for Neural Computation, University of California, San Diego,
La Jolla, CA, USA

Richard L. Barbano
Department of Neurology, University of Rochester, Rochester, NY,
USA

Joseph Jankovic
Parkinson's Disease Center and Movement Disorders Clinic,
Department of Neurology, Baylor College of Medicine, Houston,
TX, USA

Hyder A. Jinnah  orcid.org/0000-0003-1531-3969
Departments of Neurology and Human Genetics, Emory University,
Atlanta, GA, USA

Joel S. Perlmutter
Department of Neurology, Washington University School of
Medicine, St. Louis, MO, USA;
Departments of Radiology, Neuroscience, Physical Therapy, and
Occupational Therapy, Washington University School of Medicine,
St. Louis, MO, USA


Brian D. Berman  orcid.org/0000-0002-0602-9942
Department of Neurology, Virginia Commonwealth University,
Richmond, VA, USA

Abhimanyu Mahajan  orcid.org/0000-0001-8807-6672
Department of Neurological Sciences, Rush University Medical
Center, Chicago, IL, USA

Christopher G. Goetz  orcid.org/0000-0003-4457-5511
Department of Neurological Sciences, Rush University Medical
Center, Chicago, IL, USA

Glenn T. Stebbins
Department of Neurological Sciences, Rush University Medical
Center, Chicago, IL, USA

Cynthia L. Comella
Department of Neurological Sciences, Rush University Medical
Center, Chicago, IL, USA

David A. Peterson  orcid.org/0000-0001-8151-4078
Institute for Neural Computation, University of California, San
Diego, La Jolla, CA, USA;
Computational Neurobiology Laboratory, Salk Institute for
Biological Studies, La Jolla, CA, USA

REFERENCES

1. **Pal PK, Samii A, Schulzer M, Mak E, Tsui JK.** Head tremor in cervical dystonia. *Can J Neurol Sci.* 2000; 27(2): 137–42. DOI: <https://doi.org/10.1017/S0317167100052240>
2. **Shaikh AG, Beylergil SB, Scorr L, Kilic-Berkmen G, Freeman A, Klein C,** et al. Dystonia & tremor: A cross-sectional study of the dystonia coalition cohort. *Neurology*; 2020. DOI: <https://doi.org/10.1212/WNL.0000000000011049>
3. **Fasano A, Bove F, Lang AE.** The treatment of dystonic tremor: a systematic review. *J Neurol Neurosurg Psychiatry.* 2014; 85(7): 759–69. DOI: <https://doi.org/10.1136/jnnp-2013-305532>
4. **Patel N, Hanfelt J, Marsh L, Jankovic J, Coalition motD.** Alleviating manoeuvres (sensory tricks) in cervical dystonia. *J Neurol Neurosurg Psychiatry.* 2014; 85(8): 882–4. DOI: <https://doi.org/10.1136/jnnp-2013-307316>
5. **Wissel J, Müller J, Ebersbach G, Poewe W.** Trick maneuvers in cervical dystonia: investigation of movement- and touch-related changes in polymyographic activity. *Mov Disord.* 1999; 14(6): 994–9. DOI: [https://doi.org/10.1002/1531-8257\(199911\)14:6<994::AID-MDS1013>3.0.CO;2-K](https://doi.org/10.1002/1531-8257(199911)14:6<994::AID-MDS1013>3.0.CO;2-K)
6. **Frucht SJ.** The definition of dystonia: current concepts and controversies. *Mov Disord.* 2013; 28(7): 884–8. DOI: <https://doi.org/10.1002/mds.25529>
7. **Comella CL, Fox SH, Bhatia KP, Perlmutter JS, Jinnah HA, Zurowski M,** et al. Development of the Comprehensive Cervical Dystonia Rating Scale: Methodology. *Mov Disord Clin Pract.* 2015; 2(2): 135–41. DOI: <https://doi.org/10.1002/mdc3.12131>
8. **Cisneros E, Stebbins GT, Chen Q, Vu JP, Benadof CN, Zhang Z,** et al. It's tricky: Rating alleviating maneuvers in cervical dystonia. *J Neurol Sci.* 2020; 419: 117205. DOI: <https://doi.org/10.1016/j.jns.2020.117205>
9. **Bove M, Brichetto G, Abbruzzese G, Marchese R, Schieppati M.** Neck proprioception and spatial orientation in cervical dystonia. *Brain.* 2004; 127(Pt 12): 2764–78. DOI: <https://doi.org/10.1093/brain/awh291>
10. **Pelosin E, Bove M, Marinelli L, Abbruzzese G, Ghilardi MF.** Cervical dystonia affects aimed movements of nondystonic segments. *Mov Disord.* 2009; 24(13): 1955–61. DOI: <https://doi.org/10.1002/mds.22693>
11. **Avanzino L, Cherif A, Crisafulli O, Carbone F, Zenzeri J, Morasso P,** et al. Tactile and proprioceptive dysfunction differentiates cervical dystonia with and without tremor. *Neurology.* 2020; 94(6): e639–e50. DOI: <https://doi.org/10.1212/WNL.0000000000008916>

12. **Naumann M, Magyar-Lehmann S, Reiners K, Erbguth F, Leenders KL.** Sensory tricks in cervical dystonia: perceptual dysbalance of parietal cortex modulates frontal motor programming. *Ann Neurol.* 2000; 47(3): 322–8. DOI: [https://doi.org/10.1002/1531-8249\(200003\)47:3<322::AID-ANA7>3.0.CO;2-E](https://doi.org/10.1002/1531-8249(200003)47:3<322::AID-ANA7>3.0.CO;2-E)
13. **Chan J, Brin MF, Fahn S.** Idiopathic cervical dystonia: clinical characteristics. *Mov Disord.* 1991; 6(2): 119–26. DOI: <https://doi.org/10.1002/mds.870060206>
14. **Merola A, Dwivedi AK, Shaikh AG, Tareen TK, Da Prat GA, Kauffman MA,** et al. Head tremor at disease onset: an ataxic phenotype of cervical dystonia. *J Neurol.* 2019; 266(8): 1844–51. DOI: <https://doi.org/10.1007/s00415-019-09341-w>
15. **Chen Q, Vu JP, Cisneros E, Benadof CN, Zhang Z, Barbano RL,** et al. Postural Directionality and Head Tremor in Cervical Dystonia. *Tremor Other Hyperkinet Mov (N Y).* 2020; 10. DOI: <https://doi.org/10.5334/tohm.526>

TO CITE THIS ARTICLE:

Cisneros E, Vu JP, Lee HY, Chen Q, Benadof CN, Zhang Z, Pettitt EA, Joshi SK, Barbano RL, Jankovic J, Jinnah HA, Perlmutter JS, Berman BD, Mahajan A, Goetz CG, Stebbins GT, Comella CL, Peterson DA. Does Raising the Arms Modify Head Tremor Severity in Cervical Dystonia? *Tremor and Other Hyperkinetic Movements.* 2021; 11(1): 21, pp. 1–6. DOI: <https://doi.org/10.5334/tohm.623>

Submitted: 29 March 2021 Accepted: 03 June 2021 Published: 23 June 2021

COPYRIGHT:

© 2021 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

Tremor and Other Hyperkinetic Movements is a peer-reviewed open access journal published by Ubiquity Press.