

## Behavioral Biomarkers of Schizophrenia in High Drinker Rats: A Potential Endophenotype of Compulsive Neuropsychiatric Disorders

Silvia V. Navarro<sup>1</sup>, Roberto Alvarez<sup>1</sup>, M. Teresa Colomina<sup>2,3</sup>, Fernando Sanchez-Santed<sup>1</sup>, Pilar Flores<sup>1</sup>, and Margarita Moreno<sup>\*,1</sup>

<sup>1</sup>Department of Psychology, University of Almería, Campus de Excelencia Internacional Agroalimentario (CEIA3), Almería, Spain;

<sup>2</sup>Department of Psychology and Research Center for Behavior Assessment (CRAMC), Universitat Rovira i Virgili, Tarragona, Spain;

<sup>3</sup>Research in Neurobehavior and Health (NEUROLAB), Universitat Rovira i Virgili, Tarragona, Spain

\*To whom correspondence should be addressed; Department of Psychology, University of Almería, Carretera de Sacramento s/n, 04120 Almería, Spain; tel: +34-950-214624, fax: +34-950-015471, e-mail: [mjmoreno@ual.es](mailto:mjmoreno@ual.es)

**Psychogenic polydipsia, which is compulsive, non-regulatory fluid consumption, is present in 6%–20% of chronic psychiatric patients and frequently associated with the schizophrenia diagnosis. In the present study, we investigated the relation between schizophrenia-like symptoms and biomarkers with a compulsive drinking behavior phenotype in rats. Rats that were selected for low drinking vs high drinking behavior following schedule-induced polydipsia (SIP) were assessed in a latent inhibition (LI) paradigm using tone and electrical foot shock and in a spatial reversal learning task to evaluate behavioral inflexibility. We also analyzed the myelin basic protein in different brain areas of high drinker (HD) and low drinker (LD) rats. The HD rats, which were characterized by a compulsive drinking behavior on SIP, had a reduced level of LI effect and increased behavioral inflexibility in the spatial reversal learning task in comparison to the LD group. Moreover, HD rats showed less myelination in the center of the corpus callosum, striatum, and amygdala in comparison to LD rats. These findings strengthen the validity of HD rats that were selected by SIP as a possible phenotype of compulsive neuropsychiatric disorders, as evidenced by the existence of behaviors and biological markers that are related to schizophrenia and obsessive-compulsive disorder, including a reduced LI effect, behavioral inflexibility and reduced brain myelination. Future studies could contribute to the elucidation of the mechanisms underlying the compulsive phenotype of HD rats and its relation to vulnerability to schizophrenia.**

*Key words:* compulsivity/schizophrenia/  
schedule-induced polydipsia/reversal learning/latent inhibition/myelination

### Introduction

Schizophrenia is a highly debilitating neuropsychiatric disorder that affects 1% of the world population.<sup>1</sup> Psychotic symptoms affect almost all aspects of mental activity, including perception, attention, memory, and emotion.<sup>2</sup> One of the core symptoms that is observed in schizophrenia is engaging in repetitive thoughts and behaviors, which fosters a comorbidity with other neuropsychiatric disorders that are characterized by compulsive behaviors, such as obsessive-compulsive disorder (OCD) or autism.<sup>3,4</sup> Moreover, schizophrenic patients are also characterized by behavioral signs, such as the attenuation or absence of latent inhibition (LI),<sup>5,6</sup> which is a neural mechanism that allows for irrelevant stimuli to be ignored,<sup>7</sup> and behavioral inflexibility, which is defined as the lack of ability to adjust and respond to changes in the environment.<sup>8,9</sup>

Currently, the antipsychotic medication has shown a reduced efficacy for the treatment of neurocognitive and social dysfunction symptoms.<sup>10</sup> A proposed underlying neurobiological mechanism includes myelin abnormalities in the corpus callosum, which is hypothesized to be a central biomarker of the disorder.<sup>11–13</sup> Moreover, OCD patients have also shown myelin abnormalities, in terms of white matter volume and microstructure, primarily located within the fronto-striato-thalamo-cortical circuit.<sup>14–16</sup> Thus, the relationship between compulsivity and white matter abnormalities appear to be important behavioral and biological markers of vulnerability to compulsive spectrum disorders.

The animal model of scheduled-induced polydipsia (SIP), which is defined as excessive drinking behavior in food-deprived animals that is not result of physiological needs but rather a response to intermittent reinforcing

program,<sup>17,18</sup> has been shown to be a useful tool for exploring the neurobiology of compulsivity in different neuropsychiatric disorders.<sup>19,20</sup> These include schizophrenia,<sup>21–23</sup> OCD,<sup>24–27</sup> and alcohol abuse.<sup>28–31</sup> Indeed, psychogenic polydipsia (PP), which is defined as excessive non-physiologic drinking, is present in 6%–20% of psychiatric patients with schizophrenia and other compulsive spectrum disorders.<sup>32–34</sup>

SIP is a model that is sensitive to individual differences, where rats can be selected according to their compulsive drinking pattern.<sup>20,35,36</sup> Previous studies that were performed in our laboratory have shown relevant differences between high drinker (HD) and low drinker (LD) rats that were selected according to SIP acquisition. For example, the behavioral response to a challenge with dopaminergic drugs, such as D-amphetamine and cocaine,<sup>37</sup> diverges in the availability of dopamine D<sub>1</sub> and D<sub>2</sub> receptors in mesolimbic brain areas<sup>38</sup>; lack of inhibitory control in the 5-choice serial reaction time task (5-CSRTT); increased serotonin and noradrenalin activity in the amygdala in post-mortem brain analyses<sup>39</sup>; and a reduction of compulsive drinking by the 5-HT<sub>2A/C</sub> serotonin agonist DOI in HD rats.<sup>40</sup> HD rats selected by SIP have shown behavioral and neurochemical alterations that are characteristic of inhibitory control deficits; however, the knowledge of the possible co-occurrence with other behavioral biomarkers of schizophrenia remains unknown.

In the present study, we investigated if the compulsive rats, HD selected by SIP, showed possible differences, compared with LD rats, in behavioral and biological markers of schizophrenia. The HD rats showed: a reduced effect in a LI paradigm, considered as a behavioral model of schizophrenia<sup>7,41–43</sup>; an increased behavioral inflexibility in a spatial reversal learning task,<sup>44,45</sup> compared with LD rats. Finally, according the brain myelin abnormalities found in schizophrenia and OCD patients, we assessed the myelin basic protein (MBP) staining in the corpus callosum, striatum and basolateral amygdala. Our results showed a reduced MBP staining in HD compared with LD rats.

## Methods

### Subjects

Sixty-one male Wistar rats (Harlan Ibérica), weighing approximately 275–300 g at the start of the experiments, were housed 3 per cage (50 × 15 × 25 cm) at 22°C and under a 12-hour light–dark cycle (light off at 08:00 AM), with food and water available ad libitum. After 10 days of habituation, rats were gradually reduced to 85% of their free-feeding body weight by controlled feeding and maintained at this level of deprivation throughout the experiment. Food was available by daily feedings of lab chow approximately 30 minutes after each experimental session. All testings were conducted between 09:00 AM and 02:00 PM. All procedures were performed in accordance with Spanish RD1201/2005 on the protection of experimental animals and the European Community Council Directives (86/609/EEC).

### Experimental Procedures

**Schedule-Induced Polydipsia.** Two cohorts of rats (cohort 1:  $N = 32$  rats and cohort 2:  $N = 29$  rats) were tested in 13 operant SIP chambers (32 × 25 × 34 cm) (MED Associates). After 20 daily sessions of SIP, the animals were separated into 2 specific populations, HD and LD, according to whether their rates of drinking were above or below the group median, respectively.<sup>20,40</sup> The following measures were recorded for each rat: (a) total amount of water (milliliters) removed from the bottle, (b) total licks to the bottle and (c) total entries into the food magazine ([supplementary methods](#)).

The order of screening and testing was the following: for the first cohort, SIP and LI were tested; for the second cohort, SIP and the Reversal Learning task were tested.

**Latent Inhibition.** After the separation of HD ( $n = 16$ ) and LD ( $n = 16$ ) rats by the SIP procedure in the first cohort, selective attention was tested by a LI paradigm. Animals were formerly trained in the lever press. Then, 1 group was pre-exposed to a tone, and finally, all animals were tested in conditioned suppression of the lever press.<sup>46,47</sup> The variable measured was the suppression ratio ([supplementary methods](#)).<sup>46</sup>

**Reversal Learning.** After the SIP procedure, the second cohort of HD ( $n = 14$ ) and LD ( $n = 15$ ) rats was tested for behavioral inflexibility by the Spatial Reversal Learning task. The animals were required to learn the initial spatial location of a reinforced lever and to then press the correct lever when the position was reversed. The task procedure was based on Boulougouris et al.<sup>48</sup> and previous studies.<sup>45,49,50</sup> The variables measured were: the number of trials to reach the criterion, number of incorrect responses, perseverative errors, learning errors, latency to collect the food pellet, latency to respond to levers, and omissions ([supplementary methods](#)).

**MBP Immunohistochemistry.** After termination of behavioral experiments, 8 non-pre-exposed rats from the HD ( $n = 4$ ) and LD ( $n = 4$ ) groups of the first cohort (LI) were used to study myelination according to the method of Fuentes et al.<sup>51</sup> Different brain regions were selected according to the rat brain atlas of Paxinos and Watson ([supplementary methods](#)).<sup>52</sup> The results of the measure of each animal are expressed as the average density of both brain hemispheres.


### Statistical Analyses

Data were analyzed using ANOVA or Student's *t* test when appropriate ([supplementary methods](#)). Data in all figures are shown as the mean ± standard error of the mean (SEM). Statistical significance was set at  $P < .05$ .

## Results

### HD and LD Selected by SIP

The mean of water intake, licks and magazine entries of HD and LD rats on SIP are shown in [figure 1](#) and [supplementary table S1](#). In the 2 cohorts, HD rats increased water intake (interaction SIP session  $\times$  group effects: group 1  $F_{19,570} = 18.16$ ,  $P < .0001$ ; group 2  $F_{19,513} = 9.29$ ,  $P < .0001$ ) and licks (interaction SIP session  $\times$  group effect: cohort 1  $F_{19,570} = 6.83$ ,  $P < .0001$ ; cohort 2  $F_{19,513} = 14.03$ ,  $P < .0001$ ) across SIP sessions compared with the LD group ([figures 1A](#) and [1B](#)). Post hoc analysis indicated a significant increase in water intake on SIP of HD in comparison to LD rats starting at session 3 in cohort 1 ( $P < .01$ ) and at session 5 in cohort 2 ( $P < .0001$ ). Furthermore, compared to the first session, HD animals significantly increased their consumption of water during session 3 for cohort 1 ( $P < .05$ ) and during session 4 in cohort 2 ( $P < .01$ ), reaching stable levels of water intake at session 11 in cohort 1 and at session 10 in cohort 2. In addition, LD animals did not show a significant increase



**Fig. 1.** Schedule-induced Polydipsia, the mean ( $\pm$ SEM) water intake in FT-60s across 20 sessions of high drinker (HD) and low drinker (LD) groups, cohort 1 in panel A (HD,  $n = 16$ ; LD,  $n = 16$ ) and cohort 2 in panel B (HD,  $n = 14$ ; LD,  $n = 15$ ). \* $P < .05$  indicates significant differences between HD and LD groups from that session onward. # $P < .05$  indicates significant differences in comparison to session 1 on schedule-induced polydipsia (SIP) in the same group.

in their consumption of water across SIP sessions. In the number of total licks, HD rats showed a significant increase in comparison to LD rats that started during session 2 in cohort 1 ( $P < .01$ ) and during session 1 in cohort 2 ( $P < .01$ ). There were no significant differences between the HD and LD rats in the magazine entries measured on SIP ([supplementary table S1](#)).


### Latent Inhibition

**Training Sessions.** The means of lever pressing for HD and LD rats in the VI30 s and VI60 s reinforcement schedules are shown in [supplementary figure S1](#). In the VI60 s schedule, HD rats revealed a significant increase in lever pressing in comparison with LD rats ( $t(30) = -2.92$ ;  $P < .05$ ). No differences were observed in VI30 between HD and LD rats ( $t(30) = -1.15$ ,  $P = .25$ ).

**Conditioning.** The mean of suppression ratios of HD and LD rats in the LI paradigm are shown in [figure 2](#). In the LD group ([figure 2A](#)), the pre-exposure affected the conditioning phase, as shown by an increased suppression ratio response (interaction pre-exposure  $\times$  session:  $F_{3,42} = 2.88$ ,  $P < .05$ ; pre-exposure effect:  $F_{1,14} = 18.12$ ,  $P < .001$ ; session effect:  $F_{3,42} = 6.63$ ,  $P < .001$ ). Post hoc analysis indicated that PRE rats had an increased response in comparison to no-PRE rats from session 1 to 3 ( $P < .001$ ), while they displayed a trend towards still maintaining significant differences at session 4 ( $P = .056$ ). In the HD group ([figure 2B](#)), the pre-exposure affected the conditioning phase, as shown by an increased suppression ratio response (interaction pre-exposure  $\times$  session:  $F_{3,42} = 4.72$ ,  $P < .001$ ; session effect:  $F_{3,42} = 19.56$ ,  $P < .0001$ ; no pre-exposure effect:  $F_{1,14} = 1.86$ ,  $P = .19$ ). However, post hoc analysis revealed that there was a reduced pre-exposure effect in HD rats that showed differences in the suppression ratio response of PRE rats in comparison to no-PRE rats only at sessions 1 and 2 ( $P < .05$ ,  $P < .01$ ), but not at sessions 3 and 4 ( $P = .73$ ,  $P = .90$ ), during the conditioning phase. [Figure 2C](#) shows the different performance by the mean of the suppression ratio for session 3 and 4 under the pre-exposure ( $t(14) = 2.28$ ,  $P < .05$ ) vs the no pre-exposure condition ( $P > .05$ ) in the HD and LD rats.

### Reversal Learning

**Trials to Criterion.** The mean of the number of trials to reach the criterion of the Reversal Learning task are shown in [figures 3A](#) and [3B](#). During retention phase, HD rats had an increased in number of trials to complete the criterion in comparison to LD rats (interaction group  $\times$  retention:  $F_{3,81} = 5.31$ ,  $P < .05$ ; group effect:  $F_{1,27} = 5.45$ ,  $P < .05$ ; retention phase effect:  $F_{3,81} = 9.63$ ,  $P < .0001$ ). Post hoc analysis revealed that there were no differences in the spatial discrimination retention between HD and LD rats, however, HD rats had a significant increase in



**Fig. 2.** Latent inhibition, suppression ratio for the last day of pre-exposure and the 4 sessions of conditioning in the low drinker (LD) group in panel A (LD NO-PRE  $n = 8$ ; LD PRE  $n = 8$ ) and high drinker (HD) group in panel B (HD NO-PRE,  $n = 8$ ; HD PRE,  $n = 8$ ). Panel C, suppression ratio mean for the 2 last conditioning sessions (3 and 4). Data are expressed as the mean  $\pm$  SEM. In panel A and B,  $*P < .05$ ,  $**P < .01$  and  $***P < .001$  indicate significant differences between the pre-exposed and not pre-exposed groups. In panel C,  $*P < .05$  indicate significant differences between HD and LD groups.


the number of trials to reach the criterion in the retention of reversal 1 in comparison to the LD group ( $P < .0001$ ). These differences did not remain during the retention of reversals 2 and 3 (figure 3A). During reversal, HD rats needed an increased number of trials to reach the criterion in comparison with LD rats (interaction group  $\times$  reversal:  $F_{3,81} = 3.81$ ,  $P < .01$ ; group effect:  $F_{3,81} = 30.65$ ,  $P < .0001$ ). Post hoc analysis revealed that HD rats needed significantly more trials to reach the criterion during reversal 1 ( $P < .01$ ) and also during reversal 2, with a trend towards a significant difference ( $P = .09$ ) in comparison to LD rats (figure 3B).

**Incorrect Responses to Criterion.** The mean of the incorrect responses to the criterion on the Reversal Learning task are shown in figures 3C and 3D. In the retention phase, HD rats had an increase in incorrect responses (interaction group  $\times$  retention:  $F_{3,81} = 2.96$ ,  $P < .05$ ; retention effect:  $F_{3,81} = 4.55$ ,  $P < .01$ ). Post hoc analysis of retention phase revealed no significant differences in spatial discrimination retention between HD and LD rats, however indicated that HD rats made significantly more incorrect responses during retention of reversal 1 than LD rats ( $P < .01$ ) (figure 3C). During the reversal phase, HD rats needed an increased number of incorrect


responses to reach the criterion (interaction group  $\times$  reversal:  $F_{3,81} = 2.91$ ,  $P < .05$ ; session effect:  $F_{3,81} = 56.85$ ,  $P < .0001$ ; no group effect  $F_{1,27} = 2.91$ ,  $P = .14$ ). Post hoc analysis revealed that HD rats made significantly more incorrect responses during reversal 1 ( $P < .01$ ) and also during reversal 2, with a trend towards a significant difference ( $P = .06$ ) in comparison to LD rats (figure 3D).

**Perseverative and Learning Errors.** The mean of perseverative and learning errors in the reversal learning task are shown in figures 4A and 4B. During reversals, HD rats showed a significantly increased number of perseverative errors in comparison to LD rats (interaction group  $\times$  session effect:  $F_{3,78} = 2.55$ ,  $P < .01$ ; session effect:  $F_{3,78} = 55.26$ ,  $P < .001$ ; no group effect  $F_{1,26} = 1.82$ ,  $P = .18$ ). Post hoc analyses revealed that HD rats had a significantly increased number of perseverative errors during reversal 1 ( $P < .01$ ) in comparison to LD rats (figure 4A). In contrast, during reversal, no significant differences were observed in learning errors measure (figure 4B).

**Omissions and Latencies to Respond.** No significant differences were found between HD and LD rats in either omission (supplementary table S2) or in latency



**Fig. 3.** Reversal learning task, number of trials to criterion and incorrect responses to criterion through (A, C) the retention phase as well as the (B, D) reversal phase of the experiment. Abbreviations, SpDisRet: Spatial Discrimination Retention; RetRev1: retention reversal 1; RetRev2: retention reversal 2; RetRev3 Ret: retention reversal 3. Data are expressed as the mean  $\pm$  SEM. \*\* $P < .01$  and \*\*\* $P < .001$  indicates significant differences between the high drinker (HD;  $n = 14$ ) and low drinker (LD;  $n = 15$ ) rats.



**Fig. 4.** Reversal learning task, mean error scores  $\pm$  SEM of the high drinker (HD) and low drinker (LD) groups: (A) perseveration ( $\leq 6$  consecutive presses on the wrong lever) and (B) learning ( $> 6$  lever presses on the wrong lever). \*\* $P < .01$  indicates significant differences in the number of perseverative errors between the HD and LD rats.

to response (supplementary table S3) at any stage of the experiment.

#### Myelination in HD and LD Rats

The analyses of MBP revealed significant differences in the optical densities of myelinated fibers between HD and LD rats (Figure 5). HD rats showed reduced myelination within the corpus callosum ( $t(35) = -3.73$ ,  $P < .001$ ), striatum ( $t(15) = -2.51$ ,  $P < .05$ ) and basolateral amygdala ( $t(62) = -3.44$ ,  $P < .001$ ) in comparison to LD rats.


#### Discussion

In the present study, we demonstrated that an animal model of compulsivity, HD rats selected by SIP, showed

the existence of other behaviors and biological markers that were related to schizophrenia and OCD. The HD rats, which were characterized by a compulsive drinking behavior on SIP, showed less LI effect, as indicated by a reduced effect of pre-exposure to stimuli in a conditioned emotional response task; and increased behavioral inflexibility in a spatial reversal learning task in comparison to LD rats. Moreover, HD rats showed less myelination in the center of the corpus callosum, striatum, and basolateral amygdala compared with LD rats.

#### Latent Inhibition

During training previous to the pre-exposure phase, we found that HD compulsive rats selected by SIP showed significantly increased lever presses than LD rats under



**Fig. 5.** Expression of myelin basic protein (MBP) in the corpus callosum, striatum, and basolateral amygdala in the high drinker (HD;  $n = 4$ ) and low drinker (LD;  $n = 4$ ) rats selected by schedule-induced polydipsia (SIP). Data are expressed as the mean  $\pm$  SEM. \* $P < .05$  and \*\*\* $P < .001$  indicate significant differences between the HD and LD rats.

the VI60 s schedule. Previous studies have posited the idea that training under intervals may induce habit formation responses. Lever pressing is a response that is used in different animal models of compulsivity,<sup>53,54</sup> such as the reward omission procedure<sup>55,56</sup> and the post-training signal attenuation model (PTSA).<sup>57,58</sup> Our results in the VI60 s schedule are coherent with the above-cited studies, because HD rats continue pressing without earning additional pellets. Future studies based in the stimulus-response habit formation should explore if HD rats might be a possible sensitive population.

HD rats had a reduced LI effect in comparison with LD rats. In HD rats, the pre-exposed group only showed differences during session 1 and 2 compared with the pre-exposed group, while in the LD group, the LI effect was maintained until session 4. The disruption of the LI effect through different procedures has been used as a model of schizophrenia.<sup>43</sup> Lesion studies in rats have demonstrated the implication of different brain structures in this phenomenon: the nucleus accumbens shell,<sup>59,60</sup> basolateral amygdala, entorhinal cortex,<sup>61,62</sup> and ventral hippocampus.<sup>63</sup> Moreover, the pharmacological manipulation of NMDA receptors, which is as an animal model of schizophrenia, also induced LI disruption.<sup>64</sup> Likewise, recent experiments have shown that the administration of a NMDA receptor antagonist, MK-801, in rats increased water intake on SIP.<sup>21</sup> The idea that a compulsive trait could also be accompanied with an abnormal LI has also been observed in human studies with contradictory results. Two studies found enhanced LI in OCD patients,<sup>65,66</sup> whereas other studies have shown deficits in LI depending upon the OCD subtype.<sup>67</sup>

### Reversal Learning

The HD compulsive rats that were selected by SIP, showed behavioral inflexibility, as reflected by their significant poorer performance in all measures of the spatial reversal learning task, in comparison to LD rats. Thus, during reversal 1, HD rats needed more trials to reach the criterion as well as had more incorrect and perseverative

responses than LD rats. In terms of behavioral inflexibility, the failure in the trials to criterion, although perseverative in nature, reveals that HD rats might have a deficit in the initial reversal learning phase, thus in the detection of the contingency changes. This deficit was also accompanied by a lack of the inhibitory response revealed by the increased incorrect perseverative responses. In contrast, there were no differences in the number of learning errors or the acquisition and retention of the spatial discrimination between HD and LD rats, which supports the notion that the differences were not due to a deficit in the acquisition or retention of previous stimulus reward contingencies. Moreover, the finding that the significant differences are only found during the reversal 1 are in accordance with previous literature<sup>45,48-50</sup>, and might be explained by the large number of trials to criterion done in this reversal, that could benefit the performance in the next reversals. The relation between compulsive behaviors, thus behavioral inflexibility and repetitive drinking behavior on SIP, could be established by comparing several studies. First, animal lesion studies have previously demonstrated the implication of the orbitofrontal cortex (OFC) in impaired reversal learning.<sup>44,48,68</sup> Moreover, in OCD patients and their clinically unaffected relatives, there is an abnormally reduced activation of OFC during reversal learning.<sup>69</sup> According to these results, recent c-fos protein activation studies have also implicated the abnormal activity of OFC and prefrontal areas in compulsive drinking on SIP<sup>23</sup> (Merchan et al, unpublished). Second, the systemic administration of the D2/D3 receptors agonist quinpirole, which impairs reversal learning,<sup>50</sup> have also been shown to increase non-regulatory water intake in rats by contra-free loading, a proposed model of psychotic polydipsia.<sup>70-72</sup> Third, serotonin depletion has been shown to increase behavioral inflexibility.<sup>68,73</sup> Furthermore, a low tryptophan diet has shown to increase compulsive drinking in HD rats on SIP (Merchan et al, unpublished). Finally, in a recent study, Barlow et al<sup>74</sup> found that highly perseverative rats in a spatial-discrimination serial reversal learning showed reductions in both serotonin 5-HT<sub>2A</sub> receptor binding in the OFC and tryptophan hydroxylase

levels in the dorsal raphe nucleus, thereby suggesting that the serotonergic tone might be an endophenotype that predisposes animals to behavioral inflexibility and other forms of compulsive behavior. Similarly, HD compulsive rats on SIP have also shown reduced serotonin 5-HT<sub>2A</sub> receptor binding and alterations in 5-HT tone in comparison to LD rats.<sup>39</sup>

#### *Myelination in HD Compulsive Rats*

HD rats, selected by SIP, showed less myelination in the corpus callosum, striatum and basolateral amygdala in comparison with LD animals. In experimental studies, chronic treatment with the NMDA receptor antagonist MK-801 in mice have shown demyelination of the corpus callosum with schizophrenia-like behaviors, including hyperlocomotor activity and anxiety.<sup>75,76</sup> This pharmacological model of schizophrenia has also been shown to increase compulsive drinking in rats on SIP.<sup>21</sup> Moreover, animal models of schizophrenia, based on maternal immune activation, have also shown a reduction in myelinated fibers in the BLA amygdala, as well as long-lasting prepulse inhibition (PPI) deficits, locomotor hyperactivity and working memory deficits.<sup>77</sup> Reduced myelination in the corpus callosum and reduced oligodendrocyte density in the basolateral amygdala have been observed in schizophrenic patients.<sup>78–82</sup> A possible explanation, for the less myelination observed in the HD rats, could be related with the neuroadaptation induced by the hyponatremic state observed in human PP. However, this might be discarded because: first, the findings of reduced myelination in schizophrenic patients cited above did not report PP; and second HD rats did not show symptoms associated with hyponatremia, such as disorientation, ataxia or seizures after SIP; moreover the regulatory levels of drinking in HD rats are not different from LD rats in the home-cage.<sup>83</sup> In contrast, the compulsive water ingestion by SIP in rats has demonstrated to induce neuroadaptation by the increase in the spine density in the dorsolateral striatum neurons,<sup>84</sup> pointing towards a neuronal modification on habit formation brain structures after SIP. Further research is needed to clarify the relation between polydipsia, schizophrenia, and reduced brain myelin, where polydipsia could be acting as a symptom or maybe as a vulnerability factor in psychotic episodes.

In conclusion, HD rats that were selected by SIP showed a reduced LI effect, behavioral inflexibility and reduced brain myelination, which are well known behavioral and biological markers of schizophrenia. Our data confirm previous findings that propose that SIP is an animal model of compulsive behaviors associated with schizophrenia and extends knowledge of the possible compulsive phenotype of HD rats that are selected by SIP. Future studies on the compulsive phenotype of HD rats that are selected by SIP can contribute to elucidating possible explanations for the comorbidity of different compulsive behaviors and

schizophrenia-like deficits in the same individual. This will improve our understanding of the mechanisms of vulnerability in compulsive spectrum disorders.

#### **Supplementary Material**

Supplementary data are available at *Schizophrenia Bulletin* online.

#### **Funding**

Ministerio de Economía y Competitividad, Spanish Government (PSI2012-31660, PSI2014-55785-C2-1-R and PSI2015-70037-R).

#### **Acknowledgment**

The authors thank Luis Ruedas for his invaluable and continuous help. The authors have declared that there are no conflicts of interest in relation to the subject of this study.

#### **References**

1. Insel TR. Rethinking schizophrenia. *Nature*. 2010;468:187–193.
2. Lindenmayer JP, Khan A. Psychopathology. In: Lieberman JA, Stroup TS, Perkins DO, eds. *Essentials of Schizophrenia*. Washington, DC: American Psychiatric Publishing, Inc.; 2003:11–56.
3. Cederlöf M, Lichtenstein P, Larsson H, et al. Obsessive-compulsive disorder, psychosis, and bipolarity: a longitudinal cohort and multigenerational family study. *Schizophr Bull*. 2015;41:1076–1083.
4. Peng ZW, Xu T, Miao GD, et al. Neurological soft signs in obsessive-compulsive disorder: the effect of co-morbid psychosis and evidence for familiarity. *Prog Neuropsychopharmacol Biol Psychiatry*. 2012;39:200–205.
5. Brenner CA, Kieffaber PD, Clementz BA, et al. Event-related potential abnormalities in schizophrenia: a failure to “gate in” salient information? *Schizophr Res*. 2009;113:332–338.
6. Vlcek P, Bob P, Raboch J. Sensory disturbances, inhibitory deficits, and the P50 wave in schizophrenia. *Neuropsychiatr Dis Treat*. 2014;10:1309–1315.
7. Lubow R. A short history of latent inhibition research. In: Lubow R, Weiner I, eds. *Latent Inhibition. Cognition, Neuroscience and Applications to Schizophrenia*. New York, NY: Cambridge University Press; 2010:1–22.
8. Leeson VC, Robbins TW, Matheson E, et al. Discrimination learning, reversal, and set-shifting in first-episode schizophrenia: stability over six years and specific associations with medication type and disorganization syndrome. *Biol Psychiatry*. 2009;66:586–593.
9. Waltz JA, Gold JM. Probabilistic reversal learning impairments in schizophrenia: further evidence of orbitofrontal dysfunction. *Schizophr Res*. 2007;93:296–303.
10. Millan MJ, Fone K, Steckler T, Horan WP. Negative symptoms of schizophrenia: clinical characteristics, pathophysiological substrates, experimental models and prospects for improved treatment. *Eur Neuropsychopharmacol*. 2014;24:645–692.

11. Bernstein HG, Steiner J, Bogerts B. Glial cells in schizophrenia: pathophysiological significance and possible consequences for therapy. *Expert Rev Neurother*. 2009;9:1059–1071.
12. Bernstein HG, Ernst TS, Lendeckel U, Dobrowolny H, Steiner J, Bogerts B. What protects patients with schizophrenia from developing Alzheimer pathology? In: Ritsner MS, ed. *Brain Protection in Schizophrenia, Mood and Cognitive Disorders*. New York, NY: Springer; 2010:45–66.
13. Dwork AJ, Mancevski B, Rosoklija G. White matter and cognitive function in schizophrenia. *Int J Neuropsychopharmacol*. 2007;10:513–536.
14. Atmaca M, Onalan E, Yildirim H, Yuce H, Koc M, Korkmaz S. The association of myelin oligodendrocyte glycoprotein gene and white matter volume in obsessive-compulsive disorder. *J Affect Disord*. 2010;124:309–313.
15. Fan Q, Yan X, Wang J, et al. Abnormalities of white matter microstructure in unmedicated obsessive-compulsive disorder and changes after medication. *PLoS One*. 2012;7:e35889.
16. Fields RD. White matter in learning, cognition and psychiatric disorders. *Trends Neurosci*. 2008;31:361–370.
17. FALK JL. Production of polydipsia in normal rats by an intermittent food schedule. *Science*. 1961;133:195–196.
18. Falk JL. The motivational properties of schedule-induced polydipsia. *J Exp Anal Behav*. 1966;9:19–25.
19. Platt B, Beyer CE, Schechter LE, Rosenzweig-Lipson S. Schedule-induced polydipsia: a rat model of obsessive-compulsive disorder. *Curr Protoc Neurosci* 2008;Chapter 9:Unit 9.27. doi:10.1002/0471142301.ns0927s43.
20. Moreno M, Flores P. Schedule-induced polydipsia as a model of compulsive behavior: neuropharmacological and neuroendocrine bases. *Psychopharmacology (Berl)*. 2012;219:647–659.
21. Hawken ER, Delva NJ, Reynolds JN, Beninger RJ. Increased schedule-induced polydipsia in the rat following subchronic treatment with MK-801. *Schizophr Res*. 2011;125:93–98.
22. Hawken ER, Delva NJ, Beninger RJ. Increased drinking following social isolation rearing: implications for polydipsia associated with schizophrenia. *PLoS One*. 2013;8:e56105.
23. Gregory JG, Hawken ER, Banasikowski TJ, Dumont EC, Beninger RJ. A response strategy predicts acquisition of schedule-induced polydipsia in rats. *Prog Neuropsychopharmacol Biol Psychiatry*. 2015;61:37–43.
24. Rosenzweig-Lipson S, Sabb A, Stack G, et al. Antidepressant-like effects of the novel, selective, 5-HT<sub>2C</sub> receptor agonist WAY-163909 in rodents. *Psychopharmacology (Berl)*. 2007;192:159–170.
25. Schechter LE, Lin Q, Smith DL, et al. Neuropharmacological profile of novel and selective 5-HT<sub>6</sub> receptors agonists: WAY-181187 and WAY-208466. *Neuropharmacology* 2008;33:1323–1335.
26. Toscano CA, Kameyama M, Garcia-Mijares M, Silva MT, Santarem EM. Relationship between ethanol and sucrose self-administration and schedule-induced polydipsia. *Pharmacol Biochem Behav*. 2008;90:586–589.
27. Woods-Kettelberger A, Kongsamut S, Smith CP, Winslow JT, Corbett R. Animal models with potential applications for screening compounds for the treatment of obsessive-compulsive disorder. *Expert Opin Investig Drugs*. 1997;6:1369–1381.
28. Wayner MJ. Craving for alcohol in the rat: adjunctive behavior and the lateral hypothalamus. *Pharmacol Biochem Behav*. 2002;73:27–43.
29. Mittleman G, Van Brunt CL, Matthews DB. Schedule-induced ethanol self-administration in DBA/2J and C57BL/6J mice. *Alcohol Clin Exp Res*. 2003;27:918–925.
30. Mittleman G, Call SB, Cockroft JL, Goldowitz D, Matthews DB, Blaha CD. Dopamine dynamics associated with, and resulting from, schedule-induced alcohol self-administration: analyses in dopamine transporter knockout mice. *Alcohol*. 2011;45:325–339.
31. Gilpin NW, Badia-Elder NE, Elder RL, Stewart RB. Schedule-induced polydipsia in lines of rats selectively bred for high and low ethanol preference. *Behav Genet*. 2008;38:515–524.
32. de Leon J, Verghese C, Tracy JI, Josiassen RC, Simpson GM. Polydipsia and water intoxication in psychiatric patients: a review of the epidemiological literature. *Biol Psychiatry*. 1994;35:408–419.
33. de Leon J, Tracy J, McCann E, McGrory A. Polydipsia and schizophrenia in a psychiatric hospital: a replication study. *Schizophr Res*. 2002;57:293–301.
34. Iftene F, Bowie C, Milev R, et al. Identification of primary polydipsia in a severe and persistent mental illness outpatient population: a prospective observational study. *Psychiatry Res*. 2013;210:679–683.
35. Dantzer R, Terlouw C, Tazi A, et al. The propensity for schedule-induced polydipsia is related to differences in conditioned avoidance behaviour and in defense reactions in a defeat test. *Physiol Behav*. 1988;43:269–273.
36. Mittleman G, Jones GH, Robbins TW. Effects of diazepam, FG 7142, and RO 15-1788 on schedule-induced polydipsia and the temporal control of behavior. *Psychopharmacology (Berl)*. 1988;94:103–109.
37. López-Grancha M, Lopez-Crespo G, Sanchez-Amate MC, Flores P. Individual differences in schedule-induced polydipsia and the role of gabaergic and dopaminergic systems. *Psychopharmacology (Berl)*. 2008;197:487–498.
38. Pellón R, Ruíz A, Moreno M, Claro F, Ambrosio E, Flores P. Individual differences in schedule-induced polydipsia: neuroanatomical dopamine divergences. *Behav Brain Res*. 2011;217:195–201.
39. Moreno M, Gutiérrez-Ferre VE, Ruedas L, Campa L, Suñol C, Flores P. Poor inhibitory control and neurochemical differences in high compulsive drinker rats selected by schedule-induced polydipsia. *Psychopharmacology (Berl)*. 2012;219:661–672.
40. Navarro SV, Gutiérrez-Ferre V, Flores P, Moreno M. Activation of serotonin 5-HT<sub>2A</sub> receptors inhibits high compulsive drinking on schedule-induced polydipsia. *Psychopharmacology (Berl)*. 2015;232:683–697.
41. Lubow RE. *Latent Inhibition and Conditioned Attention Theory*. Cambridge, UK: Cambridge University Press; 1989.
42. Lubow RE. Construct validity of the animal latent inhibition model of selective attention deficits in schizophrenia. *Schizophr Bull*. 2005;31:139–153.
43. Weiner I. The “two-headed” latent inhibition model of schizophrenia: modeling positive and negative symptoms and their treatment. *Psychopharmacology (Berl)*. 2003;169:257–297.
44. Boulougouris V, Robbins TW. Enhancement of spatial reversal learning by 5-HT<sub>2C</sub> receptor antagonism is neuroanatomically specific. *J Neurosci*. 2010;30:930–938.
45. Castañé A, Theobald DE, Robbins TW. Selective lesions of the dorsomedial striatum impair serial spatial reversal learning in rats. *Behav Brain Res*. 2010;210:74–83.
46. Hall G, Channell S. Latent inhibition and conditioning after preexposure to the training context. *Lear Motiv*. 1985;16:381–397.
47. Killcross AS, Robbins TW. Differential effects of intra-accumbens and systemic amphetamine on latent inhibition



- using an on-baseline, within-subject conditioned suppression paradigm. *Psychopharmacology (Berl)*. 1993;110:479–489.
48. Boulougouris V, Dalley JW, Robbins TW. Effects of orbitofrontal, infralimbic and prelimbic cortical lesions on serial spatial reversal learning in the rat. *Behav Brain Res*. 2007;179:219–228.
  49. Boulougouris V, Glennon JC, Robbins TW. Dissociable effects of selective 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> receptor antagonists on serial spatial reversal learning in rats. *Neuropsychopharmacology*. 2008;33:2007–2019.
  50. Boulougouris V, Castañé A, Robbins TW. Dopamine D<sub>2</sub>/D<sub>3</sub> receptor agonist quinpirole impairs spatial reversal learning in rats: investigation of D<sub>3</sub> receptor involvement in persistent behavior. *Psychopharmacology (Berl)*. 2009;202:611–620.
  51. Fuentes S, Colomina MT, Vicens P, Franco-Pons N, Domingo JL. Concurrent exposure to perfluorooctane sulfonate and restraint stress during pregnancy in mice: effects on postnatal development and behavior of the offspring. *Toxicol Sci*. 2007;98:589–598.
  52. Paxinos G, Watson C. *The Rat Brain in Stereotaxic Coordinates*. New York, NY: Academic Press; 1998.
  53. Dickinson A, Nicholas DJ, Adams CD. The effect of the instrumental training contingency in susceptibility to reinforcer devaluation. *Q J Exp Psychol B*. 1983;35:35–51.
  54. Boulougouris V, Chamberlain SR, Robbins TW. Cross-species models of OCD spectrum disorders. *Psychiatry Res*. 2009;170:15–21.
  55. Derusso AL, Fan D, Gupta J, Shelest O, Costa RM, Yin HH. Instrumental uncertainty as a determinant of behavior under interval schedules of reinforcement. *Front Integr Neurosci*. 2010;28:17.
  56. Yin HH, Ostlund SB, Knowlton BJ, Balleine BW. The role of the dorsomedial striatum in instrumental conditioning. *Eur J Neurosci*. 2005;22:513–523.
  57. Joel D, Avisar A. Excessive lever pressing following post-training signal attenuation in rats: a possible animal model of obsessive compulsive disorder? *Behav Brain Res*. 2001;123:77–87.
  58. Joel D, Ben-Amir E, Doljansky J, Flaisher S. ‘Compulsive’ lever-pressing in rats is attenuated by the serotonin re-uptake inhibitors paroxetine and fluvoxamine but not by the tricyclic antidepressant desipramine or the anxiolytic diazepam. *Behav Pharmacol*. 2004;15:241–252.
  59. Gal G, Schiller D, Weiner I. Latent inhibition is disrupted by nucleus accumbens shell lesion but is abnormally persistent following entire nucleus accumbens lesion: The neural site controlling the expression and disruption of the stimulus preexposure effect. *Behav Brain Res*. 2005;162:246–255.
  60. Nelson AJ, Thur KE, Marsden CA, Cassaday HJ. Dopamine in nucleus accumbens: salience modulation in latent inhibition and overshadowing. *J Psychopharmacol*. 2011;25:1649–1660.
  61. Coutureau E, Galani R, Gosselin O, Majchrzak M, Di Scala G. Entorhinal but not hippocampal or subicular lesions disrupt latent inhibition in rats. *Neurobiol Learn Mem*. 1999;72:143–157.
  62. Coutureau E, Blundell PJ, Killcross S. Basolateral amygdala lesions disrupt latent inhibition in rats. *Brain Res Bull*. 2001;56:49–53.
  63. Ouhaz Z, Ba-M’hamed S, Bennis M. Haloperidol treatment at pre-exposure phase reduces the disturbance of latent inhibition in rats with neonatal ventral hippocampus lesions. *C R Biol*. 2014;337:561–570.
  64. Pouzet B, Zhang WN, Weiner I, Feldon J, Yee BK. Latent inhibition is spared by N-methyl-D-aspartate (NMDA)-induced ventral hippocampal lesions, but is attenuated following local activation of the ventral hippocampus by intracerebral NMDA infusion. *Neuroscience*. 2004;124:183–194.
  65. Kaplan O, Dar R, Rosenthal L, Hermesh H, Fux M, Lubow RE. Obsessive-compulsive disorder patients display enhanced latent inhibition on a visual search task. *Behav Res Ther*. 2006;44:1137–1145.
  66. Swerdlow NR, Hartston HJ, Hartman PL. Enhanced visual latent inhibition in obsessive-compulsive disorder. *Biol Psychiatry*. 1999;45:482–488.
  67. Lee HJ, Telch MJ. Differences in latent inhibition as a function of the autogenous-reactive OCD subtype. *Behav Res Ther*. 2010;48:571–579.
  68. Clarke HF, Walker SC, Dalley JW, Robbins TW, Roberts AC. Cognitive inflexibility after prefrontal serotonin depletion is behaviorally and neurochemically specific. *Cereb Cortex*. 2007;17:18–27.
  69. Chamberlain SR, Menzies L, Hampshire A, et al. Orbitofrontal dysfunction in patients with obsessive-compulsive disorder and their unaffected relatives. *Science*. 2008;321:421–422.
  70. Milella MS, Passarelli F, De Carolis L, et al. Opposite roles of dopamine and orexin in quinpirole-induced excessive drinking: a rat model of psychotic polydipsia. *Psychopharmacology (Berl)*. 2010;211:355–366.
  71. Schepisi C, De Carolis L, Nencini P. Effects of the 5HT<sub>2C</sub> antagonist SB242084 on the pramipexole-induced potentiation of water contrafreeloading, a putative animal model of compulsive behavior. *Psychopharmacology (Berl)*. 2013;227:55–66.
  72. De Carolis L, Schepisi C, Milella MS, Nencini P. Clomipramine, but not haloperidol or aripiprazole, inhibits quinpirole-induced water contrafreeloading, a putative animal model of compulsive behavior. *Psychopharmacology (Berl)*. 2011;218:749–759.
  73. Clarke HF, Dalley JW, Crofts HS, Robbins TW, Roberts AC. Cognitive inflexibility after prefrontal serotonin depletion. *Science*. 2004;304:878–880.
  74. Barlow RL, Alsiö J, Jupp B, et al. Markers of serotonergic function in the orbitofrontal cortex and dorsal raphe nucleus predict individual variation in spatial-discrimination serial reversal learning. *Neuropsychopharmacology*. 2015;40:1619–1630.
  75. Xiu Y, Kong XR, Zhang L, et al. White matter injuries induced by MK-801 in a mouse model of schizophrenia based on NMDA antagonism. *Anat Rec (Hoboken)*. 2014;297:1498–1507.
  76. Xiu Y, Kong XR, Zhang L, et al. The myelinated fiber loss in the corpus callosum of mouse model of schizophrenia induced by MK-801. *J Psychiatr Res*. 2015;63:132–140.
  77. Wischhof L, Irrsack E, Osorio C, Koch M. Prenatal LPS-exposure—a neurodevelopmental rat model of schizophrenia—differentially affects cognitive functions, myelination and parvalbumin expression in male and female offspring. *Prog Neuropsychopharmacol Biol Psychiatry*. 2015;57:17–30.
  78. Williams MR, Harb H, Pearce RK, Hirsch SR, Maier M. Oligodendrocyte density is changed in the basolateral amygdala in schizophrenia but not depression. *Schizophr Res*. 2013;147:402–403.
  79. Whitford TJ, Kubicki M, Schneiderman JS, et al. Corpus callosum abnormalities and their association with psychotic

- symptoms in patients with schizophrenia. *Biol Psychiatry*. 2010;68:70–77.
80. Lu LH, Zhou XJ, Keedy SK, Reilly JL, Sweeney JA. White matter microstructure in untreated first episode bipolar disorder with psychosis: comparison with schizophrenia. *Bipolar Disord*. 2011;13:604–613.
  81. Holleran L, Ahmed M, Anderson-Schmidt H, et al. Altered interhemispheric and temporal lobe white matter microstructural organization in severe chronic schizophrenia. *Neuropsychopharmacology*. 2014;39:944–954.
  82. Leroux E, Delcroix N, Dollfus S. Left-hemisphere lateralization for language and interhemispheric fiber tracking in patients with schizophrenia. *Schizophr Res*. 2015;165:30–37.
  83. Flores P, Sánchez-Kuhn A, Merchán A, Vilches O, García-Martín S, Moreno M. Schedule-induced polydipsia: searching for the endophenotype of compulsive behavior. *World J Neurosci*. 2014;4:253–260.
  84. Íbias J, Soria-Molinillo E, Kastanauskaite A, et al. Schedule-induced polydipsia is associated with increased spine density in dorsolateral striatum neurons. *Neuroscience*. 2015;300:238–245.