


Small but protective social capital against suicide ideation in poor communities

A community-based cross-sectional study

Minjae Choi, MPH^a , Myung Ki, MD, PhD^{a,b,*}, Paul S.F. Yip, PhD^c, Jungyoun Park, PhD^b, Areum Song, MS^a, Weon Young Lee, MD, PhD^d, Jong-Woo Paik, MD, PhD^e, Jiseun Lim, MD, PhD^f

Abstract

Coupled with the lowest level of social connectedness, South Korea has the highest suicide rate among the Organization for Economic Co-operation and Development countries. A possible link between community and suicide is social capital imprinted in social connectedness. This study explores whether social capital is protective against suicide ideation in relation to the poverty level of communities, and whether the associations are specific to certain elements of social capital.

A total of 908 participants were included to assess cross-sectional association of social capital at individual level with suicide ideation by comparing between poor (government-leased apartments) and non-poor communities (nongovernment-leased apartments). Logistic regression analyses were performed to examine various social capital dimensions in relation to suicide ideation.

Suicide ideation was far higher among those living in the poor communities (poor communities 12%; non poor communities 6.3%) and the level of social capital was lower in the poor communities. Nevertheless, the protective effect of social capital, in particular, the cognitive dimension against suicide ideation was demonstrated only in the poor communities (eg, odds ratio=0.27, 95% confidence interval: 0.12–0.58 for trust in the poor communities). Low income was significantly associated with suicide ideation only in the poor communities, but depression and resilience were associated with suicide ideation both in the poor and non-poor communities.

To increase the reliability of the results, established measures based on relevant literature were utilized, but measures on bridging social capital and social network might have relatively low reliability.

As to protection against suicide ideation, the extent of reliance on social capital was higher in poor communities than in non-poor communities, in particular, the cognitive dimension was likely to activate in this regard.

Abbreviations: SES = socioeconomic status, SI = synergy index.

Keywords: participation, resilience, social capital, socioeconomic status, suicidal ideation, trust

Editor: Ediriweera Desapriya.

Due to their containing information that could compromise the privacy of research participants, the data are not publicly available. However, the data supporting the findings of this current study are available on request from the corresponding author.

This study was supported by grants from the Korea Centers for Disease Control and Prevention [grant number: 2018P330300], Korea Health Industry Development Institute [grant number: HL19C0028] awarded to MK and by the Global PhD Fellowship Program through the National Research Foundation (NRF) of Korea funded by the Ministry of Education [grant number NRF-2018H1A2A1059973] awarded to MC. MK was also supported by the NRF [grant number: NRF-2019S1A5C2A03081040]. These funding bodies played no role of design, collection, analyses, and interpretation of the data.

The authors report no conflicts of interest.

Supplemental Digital Content is available for this article.

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

^a Department of Public Health, Korea University, ^b Department of Preventive Medicine, Korea University College of Medicine, Seoul, Republic of Korea, ^c Hong Kong Jockey Club Centre for Suicide Research and Prevention, The University of Hong Kong, Pokfulam, Hong Kong SAR, China, ^d Department of Preventive Medicine, Chung-Ang University College of Medicine, Dongjak-Gu, ^e Department of Psychiatry, College of Medicine, Kyung Hee University, Dongdaemun-gu, Seoul, ^f Department of Preventive Medicine, Eulji University, Daejeon Jung-gu, Daejeon, Republic of Korea.

* Correspondence: Myung Ki, Department Preventive Medicine, Korea University College of Medicine, 73 Incheon-ro, Seongbuk-gu, Seoul 136-705, Republic of Korea (e-mail: myungki@korea.ac.kr).

Copyright © 2020 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Choi M, Ki M, Yip PS, Park J, Song A, Lee WY, Paik JW, Lim J. Small but protective social capital against suicide ideation in poor communities: a community-based cross-sectional study. *Medicine* 2020;99:44(e22905).

Received: 6 April 2020 / Received in final form: 23 August 2020 / Accepted: 25 September 2020

<http://dx.doi.org/10.1097/MD.00000000000022905>

1. Introduction

South Korea has the highest suicide rate among the Organization for Economic Co-operation and Development countries.^[1] In particular, in 2018, the elderly population (≥ 65) leads the tragic epidemic with a prevalence of 48.6 per 100 thousand compared to 26.9 among other age groups (15–64 years). In conjunction with the growth of the elderly population in Korea, the far higher suicide rate among the elderly population poses an increasingly important social issue. Given the fact that Korea records the lowest level of social support among Organization for Economic Co-operation and Development countries in recent years^[2] and that social support has protective effects on suicide (ideation),^[3] the epidemic of suicide in Korea seems to be linked to its loose social connectedness.

Social capital, referring to the acquisition of resources embedded in social connectedness,^[4] has been recognized to influence suicide^[5,6] as well as a variety of other health outcomes such as mortality and mental health.^[7,8] In the absence of other forms of capital^[9–11] such as cultural and economic capital, social capital is recognized to have more pronounced protective effects on the health of the poor than the non-poor. However, some studies where social capital was mainly specified at the individual level showed that the protective effects of social capital on general health^[12] and suicide prevention^[6] may not be equal among subgroups, or even increase the risk of spreading undesirable consequences^[13] as adverse social interaction may bring negative impact on health.^[14] A recent review also reported that social capital functions either as a buffer or a dependency relationship against adverse socioeconomic circumstances in influencing health, although more evidence supported the former.^[15] Thus, social capital has both moderating and destructive potential under adverse socioeconomic circumstances, but uncertainty would depend on which elements of social capital are more significant in socioeconomically disadvantaged communities and how they are linked.

Social capital or social relations in the broad sense, is a multi-dimensional concept (eg, social support, network, trust, reciprocity, and participation), and the use of various measures contributes to the inconsistent demonstration of the effects of social capital. Some previous studies using a composite score such as the state social capital index,^[8] the summative social capital score,^[16] the social fragmented index,^[17] or the social embeddedness scale^[18] monitored the overall status of social capital, but this masked substantial variations across different items. Other studies focusing on selected dimensions of social capital such as trust^[19] or social support^[3] were limited to represent comprehensive dimensions of social capital.^[15] Similarly, most Korean studies on the association between social capital and suicide focused on 1 or 2 dimensions (eg, social support, social participation, and trust, and reported positive association with suicide (ideation)).^[20,21] However, studies including comprehensive dimensions of social capital showed that its association with mental health and suicide varies according to the types of social capital.^[22] In a cross-sectional ecological study,^[23] social trust, but not social participation or helpfulness, was linked with suicide rate. A study on an Australian rural population^[24] nonetheless showed that social support but not social network or a sense of community was associated with suicide ideation. A few Korean studies^[25,26] simultaneously included various dimensions of social capital, where, for example, trust, but not social participation, was associated with suicide ideation among the

elderly group.^[25] Thus, conceptual differentiation of social capital into several dimensions calls for assessing the association between social capital and suicide, but evidence remains unclear as to whether some aspects of social capital are more influential than others.

Further, social capital was differentiated into 2 aspects; bonding and bridging social capital.^[4,27] Bonding social capital is defined as horizontal connections among members of similar networks and bridging social capital as social connection between individuals in dissimilar groups. Some studies clarified that what ensures the beneficial effects of social capital for the disadvantaged population is bridging, but not bonding social capital.^[6,28] Thus, inclusion of both types of social capital would be necessary to better understand the potential mechanism of the interactions between social capital and adverse socioeconomic circumstances. In Korea, large apartment blocks (the typical form of residence in urban area) are developed separately for the poor and non-poor people. Large apartment blocks for the poor are mostly public-leased and are clearly demarcated from the outside, facing problems of stigmatization and social exclusion. These blocks are considerably self-contained and homogeneous, representing an independent community, isolated and dissimilar to the surrounding general residential area in terms of social identity and power relations. In the current study, both public-leased housing blocks and general apartment blocks were separately included to provide a unique opportunity for a natural experiment in the comparison of social capital between the poor and non-poor communities.

Based on the samples from the contrasting poor and non-poor communities, the current study aims to examine

- (1) whether the association of social capital with suicide ideation is protective, in particular, in relation to the poverty level of communities; and
- (2) whether the association is related to a specific dimension of social capital with simultaneous inclusion of various constructs of social capital (ie, social participation, trust, reciprocity, network, and bridging social capital).

2. Methods

2.1. Study population

A 2-stage convenience sampling was used to recruit the study participants: communities as a primary sampling unit and individuals as a secondary unit. Based on the information of public-leased apartments provided by the Seoul Housing & Communities Corporation, 2 largest public-leased apartment blocks in each district were selected from the 24 blocks in Dongdaemoon and 6 blocks in the Jungrang-gu district in Seoul. Non-poor communities were selected from the neighboring general apartment blocks in Dongdaemoon-gu, to enhance comparability of economic status between poor and non-poor apartment blocks, while minimizing differences in other regional characteristics. Samples of 607 (47%) participants were selected among the 1294 elderly population (≥ 60 years) in the poor communities, and another sample of 301 (7%) participants were selected among 4400 elderly population in the non-poor communities. This satisfied a required sample size,^[29] calculated based on difference in prevalence of elderly suicide ideation between poor community (12.14%) and non-poor (4.06%) reported in a prior Korean study.^[30] Then, any eligible

individuals were contacted mostly at home with prior announcements and flyers providing them with information on the survey. After excluding an individual with missing data ($n=1$), the final sample of 908 was included in the analyses. Ethical approval for data collection was obtained from the Institutional Research Board at Korea University (approval number: 1040548-KU-IRB-17-193-A-2).

2.2. Measure

For suicide ideation, the participants were asked to respond (yes or no) to a question on suicidal thoughts: “Have you ever felt like dying or killing yourself during the previous 12 months?”

Based on the appraisal of previous publications, relevant dimensions, and reliable measures of social capital were identified and generated (Supplementary Table 1, <http://links.lww.com/MD/F119>). Social network was assessed by totaling the responses to 3 questions on the quantity of contact per month with family and relatives, neighbors or friends.

Then, it was categorized into 2 groups (low and high) based on the median value of the summary score. Social trust was measured using the responses (very good, good, fair, bad, and very bad) to 2 questions (“In general, do you think that your neighbors can be trusted?” and “Do you think your community has a culture that neighbors help you with family events?”) and according to the summary score, the respondents were subdivided into 2 groups. Reciprocity was assessed using the responses on the Likert scale to the questions: “Among the people in each of the following 3 categories (family and relatives, neighbors and friends), how many of them will definitely help you upon your request?”^[31] The high and low reciprocity categories were defined based on a median value of summary score. Social participation was measured by asking whether the respondents regularly participate in five types of activities (religion, friendly meeting, senior citizen center, leisure, and voluntary organization). Those who responded positively to 1 or more of the 5 activities were classified as the high social participation group and others as the low social participation group. In this study, a single question on the general relationship outside a community was used to broadly imply bridging social capital: “Do you work or interact with other groups outside your apartment block?” This question was based on the Integrated Questionnaire for the Measurement of Social Capital developed by the World Bank (“Does this group work or interact with other groups with different/similar goals outside the village/neighborhood?”)^[32] with revision on the wording to consider the focus of the study unit (“you” instead of “group”) and the context (“an apartment block” instead of “village/neighborhood”). The 3 responses (“no,” “sometimes,” or “yes”) were dichotomized into no versus sometimes and yes. Resilience was measured using the responses to the questions on adaptation to change and on recovery after hardship; high and low resilience were classified based on the summary score.

The sociodemographic covariates considered in the analyses were age, gender, equalised household income, and marital status. Equalized household income was calculated by dividing the total household income by the square root of the number of household members based on the Luxembourg Income Study.^[33] Then, median equalized household income (1,125,000 Korean Won) was obtained from the distribution of the entire study population. To examine mental health status, the Geriatric Depression Scale was used where a score of 8+ is the cut-off for

depression.^[34] To assess their general health status, the respondents were asked to self-rate their health. All of these variables were dichotomized.

2.3. Statistical analyses

The prevalence of each of the components of social capital, demographic, socioeconomic, and health-related variables were calculated. To compare the associations between social capital and suicide ideation between the poor and non-poor communities, a bivariate distribution was examined, and covariate-adjusted odds ratios (ORs) using logistic regression models computed. All analyses were conducted separately for 2 contrasting communities.

Three models were used to assess the influence of covariates on the magnitude of the associations. Model 1 was unadjusted to account for basic differences across the 3 communities. In Model 2, income, along with age and gender, was adjusted to assess the influence of the socioeconomic factors. In Model 3, marital status and health-related factors (subjective health and depression) were adjusted in addition to those in Model 2. Social capital variables were entered in the 3 models separately to avoid multi-collinearity problem due to simultaneous modeling of highly inter-correlated variables.^[35] The age distributions of the 2 communities differed and, therefore, sample weights were developed for indirect age-standardization using the total study sample. To confirm the interaction effects between the social capital variables and socioeconomic circumstances on suicide ideation, the Synergy Index (“SI”) was calculated.^[36] All statistical analyses were conducted using SAS version 9.4.

3. Results

The general characteristics of individuals differed between the poor communities and the non-poor communities. The prevalence of suicide ideation was higher in the poor than the non-poor communities (12.0% vs 6.3%). The prevalence of social network and reciprocity in the non-poor communities were almost 2-fold higher than in the poor communities. In contrast, the prevalence of bridging social capital was higher in the poor communities. The other social capital variables showed no obvious pattern across the communities (Table 1).

The bivariate associations between the sociodemographic, health-related, and social-capital variables and suicide ideation in the poor and non-poor communities are shown in Table 2. In general, adverse sociodemographic and health-related factors (ie, without a spouse, poor self-rated health, and depression) were associated with suicide ideation. Similarly, except for social network and bridging social capital, a low level of social capital across measures was associated with a high level of suicide ideation, but this was mostly observed in the poor communities. The difference in the prevalence of suicide ideation between the high- and low-social capital groups was generally larger in the poor communities than in the non-poor communities (ie, 5.1% [high trust] and 14.9% [low trust] in the poor communities vs 3.3% and 7.1%, respectively, in the non-poor communities).

Multivariable associations between the measures of social capital and suicide ideation are shown in Table 3. The associations were only presented in poor communities; trust and reciprocity were significantly associated with suicide ideation

Table 1
General characteristics of participants in the poor and non-poor communities.

	Total	Poor communities	Non-poor communities
N (%)	908	607	301
Gender			
Male	288 (31.7)	187 (30.8)	101 (33.6)
Female	620 (68.3)	420 (69.2)	200 (66.4)
Age			
60–69	398 (43.8)	225 (37.1)	173 (57.5)
≥70	510 (56.2)	382 (62.9)	128 (42.5)
Marital status			
With spouse	641 (70.6)	407 (67.1)	236 (78.4)
Without spouse	267 (29.4)	200 (33.0)	65 (21.6)
Household income*			
Low	460 (50.7)	443 (73.0)	17 (5.7)
High	448 (49.3)	164 (27.0)	284 (94.3)
Self-rated health			
Good	260 (28.6)	135 (22.2)	125 (41.5)
Bad	648 (71.4)	427 (77.8)	176 (58.5)
Depression			
Yes	199 (21.9)	166 (27.3)	33 (11.0)
No	709 (78.1)	441 (72.7)	268 (89.0)
Resilience			
Low	410 (45.1)	289 (47.6)	121 (40.2)
High	498 (54.9)	381 (52.4)	180 (59.8)
Social network			
Low	472 (52.0)	389 (64.1)	83 (27.6)
High	436 (48.0)	218 (35.9)	218 (72.4)
Trust			
Low	671 (73.9)	430 (70.8)	241 (80.1)
High	237 (26.1)	177 (29.2)	60 (19.9)
Reciprocity			
Low	452 (49.8)	383 (63.1)	69 (22.9)
High	456 (50.2)	224 (36.9)	232 (77.1)
Social participation			
Low	110 (12.1)	73 (12.0)	37 (12.3)
High	798 (87.9)	534 (88.0)	264 (87.7)
Bridging social capital			
Low	281 (30.9)	158 (26.0)	123 (40.9)
High	627 (69.1)	449 (74.0)	178 (59.1)
Suicide ideation			
Yes	92 (10.1)	73 (12.0)	19 (6.3)
No	816 (89.9)	534 (88.0)	282 (93.7)

* Dichotomized based on median household equivalized income (1,125,000 Korean Won).

throughout all the 3 models (odds ratio=0.27; 95% confidence interval [CI]: 0.12–0.58 in trust and OR=0.24; 95% CI: 0.12–0.47 in reciprocity); associations of social participation and bridging social capital were present but attenuated in some models. Social network; however, was not associated with suicide ideation. Association of resilience with suicide ideation remained significant, but the association in the non-poor communities disappeared in Model 3. Low income was associated with suicide ideation in the poor communities, though again, it disappeared in Model 3. Depression was associated with suicide ideation in both the poor and non-poor communities but the magnitude of association was far larger in the non-poor communities. Interaction effects between social capital and socioeconomic status on suicide ideation were also identified (Fig. 1) in the poor communities (trust: SI=0.67, 95% CI: 0.47–0.96, reciprocity: SI=0.58, 95% CI: 0.41–0.79, resilience: SI=

0.66, 95% CI: 0.45–0.97 and social participation: SI=0.51, 95% CI: 0.40–0.65) (Supplementary Table 2, <http://links.lww.com/MD/F121>).

4. Discussion

The prevalence of suicide ideation was higher in the poor than in the non-poor communities. In general, the poor communities had lower levels of social capital but higher levels of trust and bridging social capital. Specific dimensions of social capital (eg, trust and reciprocity) were independently associated with suicide ideation but only reflected in the poor communities. Social network was not associated with suicide ideation. Resilience was consistently protective against suicide ideation in both the poor and non-poor communities. Depression appeared a stronger association with suicide ideation in the non-poor communities, while low income was associated to suicide ideation in the poor communities, though attenuated.

4.1. Methodological consideration

This study has several strengths. First, social capital was differentiated into several dimensions and this enabled examination of separate concepts of social capital on suicide ideation and simultaneous assessment of conceptual pairs (eg, bonding vs bridging and cognitive vs structural elements). Second, inclusion of the contrasting communities allowed exploration of the association between social capital and suicide ideation in relation to individual- and community-level poverty. This also facilitated a detailed examination of how social capital operated differently in the poor and non-poor communities. Nevertheless, due to the same reason that the present sample was selected from a few communities, the available data were limited to a small number of individuals in some subgroups (eg, poor people living in a non-poor communities).

This study also has several limitations. First, social capital is a broad concept, the results may rely on the number of social capital measures defined in a study.^[37] To increase the reliability of the results in the current study, established measures based on relevant literature were utilized, but some measures might have relatively low reliability. In particular, the bridging dimension of social capital was measured using a response to a single question in the relationship outside the apartment block. This may correspond to a central concept of bridging social capital, but the measure was narrowly defined to represent various bridging types on the relationship across divergent segments of society such as socioeconomic levels, religion, political orientation, age, and occupation.^[38] Likewise, social network was defined as the frequency of contact with friends, neighbors, and relatives but the quality of network was not inquired. Also, colleague networks, which is often regarded as a main form of network in Korea, were not included. Second, this study was based on a cross-sectional design, this limits the causal inference that refers to the direction between social capital and suicide ideation. Poor health reportedly increased the risk of low social capital,^[4] implying that the relationship between social capital and suicide ideation was bi-directional, and part of the association between social capital and suicidal ideation might be attributed to the reversed causal direction; that is, suicide ideation preceded and endorsed low levels of social capital. Therefore, a longitudinal approach is warranted to consider the situation when suicide ideation precedes low social capital. Third, social capital was specified

Table 2
Bivariate association between sociodemographic and social capital variables and suicide in the poor and non-poor communities.

	Poor communities				Non-poor communities			
	Total	Suicide ideation	No suicide ideation	Chi-square P-value	Total	Suicide ideation	No suicide ideation	Chi-square P-value
N (%)	607	73	534		301	19	282	
Gender								
Male	187 (30.8)	28 (15.0)	159 (85.0)	.14	101 (33.6)	5 (5.0)	96 (95.0)	.4899
Female	420 (69.2)	45 (10.7)	375 (89.3)		200 (66.4)	14 (7.0)	186 (93.0)	
Age								
60–69	225 (37.1)	36 (16.0)	189 (84.0)	.02	173 (57.5)	7 (4.1)	166 (95.9)	.0602
≥70	382 (62.9)	37 (9.7)	345 (90.3)		128 (42.5)	12 (9.4)	116 (90.6)	
Marital status								
With spouse	407 (67.1)	37 (9.1)	370 (90.9)	.002	236 (78.4)	9 (3.8)	227 (96.2)	<.001
Without spouse	200 (33.0)	36 (18.0)	164 (85.0)		65 (21.6)	15 (15.4)	55 (84.6)	
Household income*								
Low	443 (73.0)	63 (14.2)	380 (85.8)	.006	17 (5.7)	2 (11.8)	15 (88.2)	.3412
High	164 (27.0)	10 (6.1)	154 (93.9)		284 (94.3)	17 (6.0)	267 (94.0)	
Self-rated health								
Good	135 (22.2)	6 (4.4)	129 (95.6)	.002	125 (23.5)	2 (1.6)	123 (98.4)	.0046
Bad	427 (77.8)	67 (14.2)	405 (85.8)		176 (58.5)	17 (9.7)	159 (90.3)	
Depression								
Yes	166 (27.3)	41 (24.7)	125 (75.3)	<.001	33 (11.0)	11 (33.3)	22 (66.7)	<.001
No	441 (72.7)	32 (7.3)	409 (92.7)		268 (89.0)	8 (3.0)	260 (97.0)	
Resilience								
Low	289 (47.6)	57 (19.7)	232 (80.3)	<.001	121 (40.2)	13 (10.7)	108 (89.3)	.0095
High	381 (52.4)	16 (5.0)	302 (95.0)		180 (59.8)	6 (3.3)	174 (96.7)	
Social network								
Low	389 (64.1)	47 (12.1)	342 (88.9)	.96	83 (27.6)	7 (8.4)	76 (91.6)	.3504
High	218 (35.9)	26 (11.9)	192 (88.1)		218 (72.4)	12 (5.5)	206 (94.5)	
Trust								
Low	430 (70.8)	64 (14.9)	366 (85.1)	<.001	241 (80.1)	17 (7.1)	224 (92.9)	.2890
High	177 (29.2)	9 (5.1)	168 (94.9)		60 (19.9)	2 (3.3)	58 (96.7)	
Reciprocity								
Low	383 (63.1)	61 (15.9)	322 (84.1)	<.001	69 (22.9)	2 (2.9)	67 (97.1)	.1841
High	224 (36.9)	12 (5.4)	212 (94.6)		232 (77.1)	17 (7.3)	215 (92.7)	
Social participation								
Low	73 (12.0)	15 (15.9)	58 (84.1)	.017	37 (12.3)	4 (10.8)	33 (89.2)	.2296
High	534 (88.0)	58 (5.4)	476 (94.6)		264 (87.7)	15 (5.7)	249 (94.3)	
Bridging social capital								
Low	158 (26.0)	24 (15.2)	134 (84.8)	.1575	123 (40.9)	9 (7.3)	114 (92.7)	.5513
High	449 (74.0)	49 (10.9)	400 (89.1)		178 (59.1)	10 (5.6)	168 (94.4)	

* Dichotomized based on median household equivalized income (1,125,000 Korean Won).

as an individual attribute, though the features of social capital lie in the area-level as well.^[39] Despite the limited capacity of this type of approach, individual-level studies as the present, were able to highlight the associations between social capital, poverty, and suicide operating as individual characteristics.^[40] Fourth, another limitation concerns the unmeasured confounding factors such as anxiety and alcohol abuse. Alcohol drinking even at a moderate drinking level was reported to be a strong risk factor for suicide in Korea,^[41] therefore, a future study is required to assess the association between social capital and alcohol consumption levels in predicting suicides. Lastly, as there was restriction to obtaining the list of residents, the sampling was not conducted in a probabilistic or random manner. This approach may be subjected to a potential problem in representing a population. However, as the available samples from the poor and non-poor communities were relatively large (7%–50% of the total population) with low decline rates (3%–8%) to the survey request, the data can be considered to be generalizable.

4.2. Comparison with previous studies

In the current study, the level of social capital appeared to differ by dimensions between the poor and non-poor communities. However, those living in the poor communities are more often related to formal networks, and actual social capital is generally more restrictive in the poor communities. For example, the frequency of social participation was similar in the poor and non-poor communities, and the residents of the poor communities participated more frequently in activities in senior citizen centers but less frequently in leisure and voluntary activities (Supplementary Table 3, <http://links.lww.com/MD/F122>). Likewise, bridging social capital, which was higher among the residents of the poor communities may reflect aid-based relationships, as the senior citizen centers and welfare centers frequently provide basic services such as meals to the poor elderly persons. As the residents of the poor communities are more likely to rely on survival-oriented assistance, this may appear to increase their social capital.^[42] However, in contrast to previous findings, in the

Table 3
Multivariate association (OR [95% CI])[‡] between sociodemographic and social capital variables and suicide ideation in the poor and non-poor communities.

	Poor communities			Non-poor communities		
	Model 1 [§]	Model 2	Model 3	Model 1 [§]	Model 2	Model 3
Marital status						
Without spouse	1 (Reference)	1 (Reference)				
With living spouse	2.15 (1.32–3.51) [†]	2.45 (1.47–4.10) [†]	1.42 (0.81–2.51)	4.85 (1.96–11.97) [†]	4.34 (1.7–11.12) [†]	3.83 (1.16–9.88) [*]
Household Income [#]						
Median income or higher	1 (Reference)	1 (Reference)				
Lower than median income	2.42 (1.25–4.68) [†]	2.14 (1.08–4.23) [*]	1.77 (0.86–3.62)	1.74 (0.38–7.93)	1.32 (0.28–6.23)	0.29 (0.05–1.80)
Self-rated health						
Good	1 (Reference)	1 (Reference)				
Bad	3.89 (1.67–9.03) [†]	4.51 (1.90–10.70) [†]	5.38 (2.17–3.62) [†]	5.15 (1.39–19.09) [*]	4.55 (1.21–17.11) [*]	3.32 (0.77–14.24)
Depression						
No	1 (Reference)	1 (Reference)				
Yes	3.85 (2.34–6.31) [†]	3.84 (2.30–6.40) [†]	3.60 (2.07–6.26) [†]	18.36 (6.91–48.78) [†]	18.40 (6.52–51.93) [†]	11.50 (3.75–35.29) [†]
Resilience						
Low	1 (Reference)	1 (Reference)				
High	0.21 (0.12–0.37) [†]	0.25 (0.13–0.45) [†]	0.19 (0.10–0.35) [†]	0.27 (0.10–0.70) [†]	0.28 (0.10–0.73) [†]	0.47 (0.15–1.46)
Social network						
Low	1 (Reference)	1 (Reference)				
High	0.90 (0.54–1.50)	0.90 (0.53–1.51)	1.03 (0.59–1.78)	0.74 (0.29–1.89)	0.73 (0.28–1.90)	0.55 (0.18–1.69)
Trust						
Low	1 (Reference)	1 (Reference)				
High	0.28 (0.14–0.59) [†]	0.32 (0.15–0.67) [†]	0.27 (0.12–0.58) [†]	0.62 (0.16–2.31)	0.71 (0.18–2.73)	1.18 (0.26–5.41)
Reciprocity						
Low	1 (Reference)	1 (Reference)				
High	0.28 (0.15–0.54) [†]	0.29 (0.15–0.56) [†]	0.24 (0.12–0.47) [†]	3.48 (0.65–18.53)	3.29 (0.58–18.77)	5.96 (0.82–43.19)
Social participation						
Low	1 (Reference)	1 (Reference)				
High	0.46 (0.25–0.86) [*]	0.52 (0.28–0.98) [*]	0.56 (0.28–1.16)	0.50 (0.16–1.60)	0.42 (0.13–1.35)	0.76 (0.20–2.84)
Bridging social capital						
Low	1 (Reference)	1 (Reference)				
High	0.63 (0.38–1.05)	0.56 (0.33–0.96) [*]	0.80 (0.45–1.42)	0.77 (0.32–1.86)	0.85 (0.35–2.08)	1.09 (0.38–3.11)

CI = confidence interval, OR = odds ratio.

^{*} P-value < .05.

[†] P-value < .01.

[‡] All models were weighted to consider differences in age structure across 3 communities.

[§] Model 1: unadjusted.

^{||} Model 2: adjusted for age, gender, and household income.

^{||} Model 3: adjusted for age, gender, household income, marital status, self-rated health, and depression.

[#] Dichotomized based on median household equivalized income (1,125,000 Korean won).

current study, social trust was higher in the poor communities.^[43] Given that the present 2 questions of trust are primarily related to credit on the neighborhoods, the findings may reflect the

possibility that the affluent people are less concerned about their relationships with neighbors, which may reduce the level of trust among the residents of the non-poor communities.^[44]




Figure 1. SES-social capital interaction association on suicide ideation in the poor communities. ^{*}Significant interaction effect (P-value < .05) between social capital and SES on suicide ideation. SES = socioeconomic status.

The association of social capital with suicide ideation was demonstrated only in the poor communities, suggesting that the residents of those communities are more dependent on social capital. Previous studies on the impact of social capital on health support this notion that the impact of social capital was greater or only detected in deprived areas,^[45] ethnic minorities,^[12] and lower socioeconomic status (“SES”) groups.^[6,43] This may be because social capital compensates for insufficient material resources in poor communities^[15] and/or social relationships in poor communities are more cohesive and more likely to provide effective support necessary to maintain mental health.^[43]

An interaction effect between social capital and SES on suicide ideation was detected in the poor communities where the association between social capital and suicide ideation was strengthened among those with a low income.^[13,46] Also, some evidence was included by differentiating community level poverty, that the moderating role of social capital on the association between SES and suicide ideation was marked in the poor communities but minimal in the non-poor communities. This finding further supports that when other forms of capital are less available, social capital may be more important among those in poverty both at the individual- and community-levels.

Some dimensions of social capital, however, did not show protective associations with suicide ideation, while trust and reciprocity exerted a marked protective effect. This finding sheds some light on the distinction between the cognitive and structural dimensions of social capital. Cognitive social capital is typically defined as subjective perceptions of social relations such as trust, reciprocity, and norm, while structural social capital reflects directly observable measures such as social network and social participation.^[47,48] Similar to previous studies,^[22,48] and including a review,^[49] it is discovered that cognitive social capital is more beneficial in preventing suicide than structural social capital. Thus, the influence of social capital on suicide ideation in the poor communities may be more related to perceived rather than actual levels of social connectedness.

Resilience showed protective effects against suicide ideation both in the poor and non-poor communities. This is consistent with previous studies where regardless of the characteristics of the study population (eg, adolescents, the elderly, trauma patients, depressed persons, and rural residents),^[50–52] the associations of resilience on suicidality (eg, suicide ideation and suicide attempts) in Korea^[50,51] and elsewhere^[53,54] were observed. In these studies, the magnitude of the association of resilience has been particularly underscored, suggesting that psychological factors played important roles in preventing suicides in Korea. This is also consistent with the present finding that protection against suicide ideation is mediated primarily by the cognitive rather than the structural dimensions of social capital.

In the current study, low income was associated with suicide ideation, probably because low SES can lead to psychological distress and ultimately suicide ideation.^[55] Further, similar to previous studies where the association between low income and suicide is stronger in deprived than in non-deprived areas,^[56] low income was significantly associated with suicide ideation only in the poor communities, though it attenuated. Interestingly, depression was associated with suicide ideation in both communities. This is consistent with the well-known findings that suicidality could be attributable to being depressed, and in the current study, only a smaller magnitude of this association was shown in the poor communities.^[57] This may be because mental health issues alone are typically insufficient to induce

suicides. Other stressors, for example, debt and physical illnesses, are usually also present^[58]; these factors are more frequently encountered in the poor communities.^[57] This suggests that the relative importance between SES versus psychological factors of suicide risk differs between the poor and non-poor communities.

5. Conclusions

In the poor communities, social capital may be low in general, while it can be suicide protective among those with strong ties. This finding should be viewed as emphasizing the importance of both social capital and economic progress. The associations of social capital with suicide ideation were linked to specific dimensions and cognitive social capital such as trust and reciprocity, and likely to function in the poor communities. Further, social capital exerted a moderating role on the association between income and suicide ideation, in particular, in the poor communities.

Author contributions

Conceptualization: Minjae Choi, Myung Ki.

Formal analysis: Minjae Choi.

Methodology: Myung Ki.

Writing – original draft: Minjae Choi, Myung Ki.

Writing – review & editing: Paul Siu Fai Yip, Jungyoun Park, Aream Song, Weon Young Lee, Jong-Woo Paik, Jiseun Lim.

References

- [1] OECD. Health at a Glance 2017. Paris: OECD; 2017.
- [2] OECD. How's Life? 2017 Measuring Well-being. Paris: OECD; 2017.
- [3] Kleiman EM, Liu RT. Social support as a protective factor in suicide: findings from two nationally representative samples. *J Affect Disord* 2013;150:540–5.
- [4] Poortinga W. Social relations or social capital? Individual and community health effects of bonding social capital. *Soc Sci Med* 2006;63:255–70.
- [5] Congdon P. Latent variable model for suicide risk in relation to social capital and socio-economic status. *Soc Psychiatry Psychiatr Epidemiol* 2012;47:1205–19.
- [6] Fitzpatrick KM, Irwin J, Lagory M, et al. Just thinking about it: social capital and suicide ideation among homeless persons. *J Health Psychol* 2007;12:750–60.
- [7] Ehsan AM, De Silva MJ. Social capital and common mental disorder: a systematic review. *J Epidemiol Community Health* 2015;69:1021–8.
- [8] Smith ND, Kawachi I. State-level social capital and suicide mortality in the 50 U.S. states. *Soc Sci Med* 2014;120:269–77.
- [9] Mellor JM, Milyo J. State social capital and individual health status. *J Health Polit Policy Law* 2005;30:1101–30.
- [10] Perry M, Williams RL, Wallerstein N, et al. Social capital and health care experiences among low-income individuals. *Am J Public Health* 2008;98:330–6.
- [11] Story WT. Social capital and health in the least developed countries: a critical review of the literature and implications for a future research agenda. *Glob Public Health* 2013;8:983–99.
- [12] Baron-Epel O, Weinstein R, Haviv-Mesika A, et al. Individual-level analysis of social capital and health: a comparison of Arab and Jewish Israelis. *Soc Sci Med* 2008;66:900–10.
- [13] Moore S, Daniel M, Gauvin L, et al. Not all social capital is good capital. *Health Place* 2009;15:1071–7.
- [14] Villalonga-Olives E, Kawachi I. The dark side of social capital: a systematic review of the negative health effects of social capital. *Soc Sci Med* 2017;194:105–27.
- [15] Uphoff EP, Pickett KE, Cabieses B, et al. A systematic review of the relationships between social capital and socioeconomic inequalities in health: a contribution to understanding the psychosocial pathway of health inequalities. *Int J Equity Health* 2013;12:54.

- [16] McKenzie K, Whitley R, Weich S. Social capital and mental health. *Br J Psychiatry* 2002;181:280–3.
- [17] Congdon P. Suicide and parasuicide in London: a small-area study. *Urban Studies* 1996;33:137–58.
- [18] Kaslow NJ, Sherry A, Bethea K, et al. Social risk and protective factors for suicide attempts in low income African American men and women. *Suicide Life Threat Behav* 2005;35:400–12.
- [19] Kelly BD, Davoren M, Mhaolain AN, et al. Social capital and suicide in 11 European countries: an ecological analysis. *Soc Psychiatry Psychiatr Epidemiol* 2009;44:971–7.
- [20] Ra CK, Cho Y. Differentiated effects of social participation components on suicidal ideation across age groups in South Korea. *BMC Public Health* 2013;13:890.
- [21] Kwon H-J, Jeong J-U, Choi M. Social relationships and suicidal ideation among the elderly who live alone in republic of Korea: a logistic model. *Inquiry* 2018;55:0046958018774177.
- [22] Ferlander S, Stickle A, Kislitsyna O, et al. Social capital - a mixed blessing for women? A cross-sectional study of different forms of social relations and self-rated depression in Moscow. *BMC Psychol* 2016;4:37.
- [23] Okamoto M, Kawakami N, Kido Y, et al. Social capital and suicide: an ecological study in Tokyo. *Japan Environ Health Prev Med* 2013;18:306–12.
- [24] Handley TE, Inder KJ, Kelly BJ, et al. You've got to have friends: the predictive value of social integration and support in suicidal ideation among rural communities. *Soc Psychiatry Psychiatr Epidemiol* 2012;47:1281–90.
- [25] Ha M-O, Kim J-R, Jeong B, et al. Associations of social participation and trust with suicidal ideation and attempt in communities with high mortality. *J Agric Med and Community Health* 2013;38:116–29.
- [26] Shin S-J, Cho Y-T. Social capital and suicidal impulse. *Korean J Health Edu Promot* 2007;24:35–49.
- [27] Szreter S, Woolcock M. Health by association? Social capital, social theory, and the political economy of public health. *Int J Epidemiol* 2004;33:650–67.
- [28] Kushner HI, Sterk CE. The limits of social capital: Durkheim, suicide, and social cohesion. *Am J Public Health* 2005;95:1139–43.
- [29] Fleiss JL, Levin B, Paik MC. *Statistical Methods for Rates and Proportions*. New Jersey: John Wiley & Sons; 2013.
- [30] Lee S. A longitudinal study on predictors of suicide ideation in old people: using a panel logit model. *Health Soc Welfare Rev* 2017;6:191–229.
- [31] Wang P, Chen X, Gong J, et al. Reliability and validity of the personal social capital scale 16 and personal social capital scale 8: two short instruments for survey studies. *Soc Indic Res* 2014;119:1133–48.
- [32] Grootaert C, Narayan D, Jones VN, et al. *Measuring social capital: an integrated questionnaire*. Washington: The World Bank; 2004.
- [33] Atkinson AB, Rainwater L, Smeeding TM. *Income Distribution in OECD Countries: Evidence from the Luxembourg Income Study*. Paris: OECD; 1995.
- [34] Yesavage JA, Brink TL, Rose TL, et al. Development and validation of a geriatric depression screening scale: a preliminary report. *J Psychiatr Res* 1982;17:37–49.
- [35] Kiely JL, Sergievsky GH. Some conceptual problems in multivariable analyses of perinatal mortality. *Paediatr Perinat Epidemiol* 1991;5:243–57.
- [36] Lundberg M, Fredlund P, Hallqvist J, et al. A SAS program calculating three measures of interaction with confidence intervals. *Epidemiology* 1996;7:655–6.
- [37] Kawachi I. Commentary: social capital and health: making the connections one step at a time. *Int J Epidemiol* 2006;35:989–93.
- [38] Villalonga-Olives E, Adams I, Kawachi I. The development of a bridging social capital questionnaire for use in population health research. *SSM Popul Health* 2016;2:613–22.
- [39] Kawachi I, Kim D, Coutts A, et al. Commentary: Reconciling the three accounts of social capital. *Int J Epidemiol* 2004;33:682–90.
- [40] DeFilippis J. The myth of social capital in community development. *Hous Policy Debate* 2001;12:781–806.
- [41] Jung M. The relationship between alcohol abuse and suicide risk according to smoking status: a cross-sectional study. *J Affect Disord* 2019;244:164–70.
- [42] Saegert S, Thompson JP, Warren MR. *Social Capital and Poor Communities*. New York: Russell Sage Foundation; 2002.
- [43] Sun X, Rehnberg C, Meng Q. How are individual-level social capital and poverty associated with health equity? A study from two Chinese cities. *Int J Equity Health* 2009;8:2.
- [44] Caughy MO, O'Campo PJ, Muntaner C. When being alone might be better: neighborhood poverty, social capital, and child mental health. *Soc Sci Med* 2003;57:227–37.
- [45] Verhaeghe PP, Tampubolon G. Individual social capital, neighbourhood deprivation, and self-rated health in England. *Soc Sci Med* 2012;75:349–57.
- [46] Pearson JA, Geronimus AT. Race/ethnicity, socioeconomic characteristics, coethnic social ties, and health: evidence from the national Jewish population survey. *Am J Public Health* 2011;101:1314–21.
- [47] Murayama H, Wakui T, Arami R, et al. Contextual effect of different components of social capital on health in a suburban city of the greater Tokyo area: a multilevel analysis. *Soc Sci Med* 2012;75:2472–80.
- [48] Forsman AK, Nyqvist F, Wahlbeck K. Cognitive components of social capital and mental health status among older adults: a population-based cross-sectional study. *Scand J Public Health* 2011;39:757–65.
- [49] De Silva MJ, McKenzie K, Harpham T, et al. Social capital and mental illness: a systematic review. *J Epidemiol Community Health* 2005;59:619–27.
- [50] Ahn SJ, Park EH, Hong HJ, et al. The role of resilience as a protective factor for suicide in adolescent patients with depression. *Kor J Clin Psychol* 2013;32:351–66.
- [51] Kim S-H, Hwang K-R. The influence of social exclusion on suicidal impulse of senior citizens and the moderating effect of adjustment resilience. *J Korea Contents Assoc* 2016;16:263–73.
- [52] Yoon H-S, Yeom S-R. The mediating effects of resilience on depression and suicidal ideation of older adults in Korea. *Korean J Gerontol Soc Welfare* 2017;72:233–55.
- [53] Johnson J, Gooding PA, Wood AM, et al. Resilience as positive coping appraisals: testing the schematic appraisals model of suicide (SAMS). *Behav Res Ther* 2010;48:179–86.
- [54] Rutter PA, Freedenthal S, Osman A. Assessing protection from suicidal risk: psychometric properties of the suicide resilience inventory. *Death Stud* 2008;32:142–53.
- [55] Lorant V, Kunst AE, Huisman M, et al. Socio-economic inequalities in suicide: a European comparative study. *Br J Psychiatry* 2005;187:49–54.
- [56] Burrows S, Auger N, Gamache P, et al. Influence of social and material individual and area deprivation on suicide mortality among 2.7 million Canadians: a prospective study. *BMC Public Health* 2011;11:577.
- [57] Dupéré V, Leventhal T, Lacourse E. Neighborhood poverty and suicidal thoughts and attempts in late adolescence. *Psychol Med* 2009;39:1295–306.
- [58] Qin P, Agerbo E, Mortensen PB. Suicide risk in relation to socioeconomic, demographic, psychiatric, and familial factors: a national register-based study of all suicides in Denmark, 1981–1997. *Am J Psychiatry* 2003;160:765–72.