

# Impact of Antecedent Infections on the Antibodies against Gangliosides and Ganglioside Complexes in Guillain-Barré Syndrome: A Correlative Study

Debprasad Dutta<sup>#</sup>, Monojit Debnath<sup>#</sup>, Doniparthi V. Seshagiri<sup>1</sup>, Binu V. Sreekumaran Nair<sup>2</sup>, Sumit K. Das<sup>2</sup>, Rahul Wahatule<sup>1</sup>, Sanjib Sinha<sup>1</sup>, Vasanthapuram Ravi<sup>3</sup>, Arun B. Taly<sup>1</sup>, Madhu Nagappa<sup>1</sup>

Departments of Human Genetics, <sup>1</sup>Neurology, <sup>2</sup>Biostatistics and <sup>3</sup>Neurovirology, National Institute of Mental Health and Neurosciences (NIMHANS), Bengaluru, Karnataka, India

<sup>#</sup>These authors contributed equally

## Abstract

**Background and Aims:** Guillain-Barré Syndrome (GBS), an immune-mediated neuropathy, is characterized by antibodies against gangliosides/ganglioside complexes (GSCs) of peripheral nerves. Antecedent infections have been reported to induce antibodies that cross-react with the host gangliosides and thereby have a pivotal role in conferring an increased risk for developing GBS. Data pertaining to the impact of various antecedent infections, particularly those prevalent in tropical countries like India on the ganglioside/GSC antibodies is sparse. We aimed at exploring the association between six antecedent infections and the profile of ganglioside/GSC antibodies in GBS. **Methods:** Patients with GBS (n = 150) and healthy controls (n = 50) were examined for the serum profile of antibodies against GM1, GM2, GD1a, GD1b, GT1b, and GQ1b and their GSCs by ELISA. These antibodies were correlated with immunoreactivities against *Campylobacter jejuni*, Japanese encephalitis (JE), dengue, influenza, zika, and chikungunya infections. **Results:** The frequencies of antibodies against six single gangliosides ( $P < 0.001$ ) and their GSCs ( $P = 0.039$ ) were significantly higher in patients as compared to controls. Except for GT1b-antibody which was more frequent in axonal GBS, none of the other ganglioside/GSC antibodies correlated with the electrophysiological subtypes of GBS. Antecedent JE infection was significantly associated with increased frequency of antibodies against GD1a, GD1b, GT1b, and GQ1b. Antibodies against GSCs were not influenced by the antecedent infections. **Interpretation:** This study for the first time shows an association between antecedent JE infection and ganglioside antibodies in GBS. This finding reinforces the determining role of antecedent infections on ganglioside antibody responses and the subsequent immunological processes in GBS.

**Keywords:** Antecedent infections, autoantibodies, ganglioside complex, gangliosides, Guillain-Barré syndrome, Japanese encephalitis virus

## INTRODUCTION

Guillain-Barré Syndrome (GBS), an immune-mediated neuropathy, is the commonest cause of neuromuscular paralysis. Antecedent infection with *Campylobacter jejuni* has been identified as the predominant risk determinant of GBS. In addition, a number of other infectious pathogens, such as *Mycoplasma pneumoniae*, Epstein Barr Virus (EBV), Dengue virus, Chikungunya virus, *Hemophilus influenzae*, Cytomegalovirus (CMV), Zika virus, and the severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2) have been associated with increased risk of GBS.<sup>[1,2]</sup> However, there exist some variations in the patterns of association between these pathogens and the risk of developing GBS.<sup>[3-5]</sup> Contrary to the earlier studies, in our recent study, the Chikungunya virus was the most common infectious trigger of GBS, followed by *C. jejuni* infection.<sup>[6]</sup> It is noteworthy that not all individuals infected with these pathogens develop GBS. For example, only 1 in 1,000 individuals with *C. jejuni* infection develops GBS. This suggests that infectious triggers alone are not sufficient to drive the underlying pathogenetic processes in GBS.

The infectious pathogens potentially interact with the host immune cells and immune molecules and lead to the development of GBS.<sup>[7]</sup> One of the most widely recognized mechanisms through which *C. jejuni* causes GBS is ‘molecular mimicry’, i.e., cross-reaction between antibodies raised against pathogens and the gangliosides of peripheral nerves.<sup>[8,9]</sup> Antibodies against gangliosides and ganglioside complexes (GSCs) have been

**Address for correspondence:** Dr. Madhu Nagappa, Additional Professor, Department of Neurology, National Institute of Mental Health & Neurosciences (NIMHANS), Hosur Road, Bengaluru - 560 029, Karnataka, India.  
E-mail: madhu\_nagappa@yahoo.co.in

Submitted: 03-Feb-2022 Revised: 10-Apr-2022 Accepted: 21-Apr-2022  
Published: 14-Jun-2022

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

DOI: 10.4103/aian.aian\_121\_22

reported in GBS patients across various populations and they have been suggested to be the major drivers of the pathogenic processes as well as the severity of GBS.<sup>[10-12]</sup> However, the repertoire of autoantibodies targeting the peripheral nerves is rather heterogeneous in GBS.<sup>[13]</sup> Notably, the patterns of distribution of the ganglioside- and GSC- antibodies are not uniform across different populations. The precise factors that contribute to the variations in the frequencies of these antibodies in GBS are not well understood.

There is a growing recognition that the type of these antibodies, as well as the magnitude of their production in GBS, may depend on the burden of infectious pathogens in a population. Further, specific genes in these pathogens may also play a role in determining the induction of various ganglioside antibodies.<sup>[14]</sup> A positive correlation between preceding infections and the profile of ganglioside antibodies has been reported by previous studies in the French, Japanese, and Chinese populations.<sup>[15-17]</sup> These studies focussed mainly on the association between *C. jejuni*, and to some extent on *M. pneumoniae* and CMV, and the induction of ganglioside antibodies. However, these studies are limited and they do not provide adequate insights into the causal relationship between the spectrum of preceding infections and antibodies against the individual gangliosides/GSCs in GBS. As such the associations between all the major risk pathogens that are prevalent across various geographical territories and the ganglioside/GSC antibodies have not been tested. In tropical countries, arboviral infections such as chikungunya, dengue, Japanese Encephalitis (JE), etc. are more prevalent and they have also been linked to the risk of developing GBS.<sup>[18,19]</sup> We have reported the association between JE, dengue, and chikungunya virus infections and the risk of GBS in the Indian population.<sup>[6]</sup> However, the impact of such preceding arboviral infections on the ganglioside- and GSC-antibody profile is not known. To address these knowledge gaps, this study was aimed at exploring the association between six infections and antibodies against gangliosides and GSCs in patients with GBS.

## SUBJECTS AND METHODS

### Study participants

The present study was conducted on patients with GBS admitted to the emergency services of a single neurology unit of the National Institute of Mental Health & Neurosciences (NIMHANS), Bangalore, India. Adults (age  $\geq 18$  years) fulfilling the National Institute of Neurological Disorders and Stroke (NINDS) diagnostic criteria were enrolled for the study.<sup>[20]</sup> Patients who received treatment with intravenous immunoglobulin and/or underwent plasmapheresis prior to the study entry were excluded. Nerve conduction studies were carried out in all patients using standard protocols, and subtyping was done based on the criteria recommended by Rajabally *et al.*<sup>[21]</sup> Healthy community controls were recruited from the blood donation camps organized by the Department of Transfusion Medicine and Haematology, NIMHANS. The patients and controls were matched for age, gender, and ethnicity. The

study was approved by the Institute Ethics Committee [No. NIMH/DO/Ethics Sub-committee (BS&NS) 5<sup>th</sup> Meeting/2017, dated 13.6.2017]. All the study participants provided written informed consent prior to their participation in the study.

### Collection of blood samples

Ten ml. of peripheral blood was drawn from the median cubital vein under aseptic conditions, of which 5 ml. into sterile Becton Dickinson (BD<sup>®</sup>) serum vacutainers and the remaining 5 ml. into EDTA vacutainers. The serum and the EDTA tubes were centrifuged at 3000 rpm for 12 minutes to separate the serum and plasma, respectively. The serum and plasma samples were aliquoted and stored at  $-80^{\circ}\text{C}$  until the immunoassays were performed and they underwent only one freeze and thaw cycle.

### Profiling of antibodies against gangliosides and ganglioside complexes

A manual and validated Enzyme-Linked Immune Sorbent Assay (ELISA) was employed to determine antibodies against single gangliosides and GSCs in the sera of patients ( $n = 150$ ) and controls ( $n = 50$ ). The selection of ganglioside antigens was based on (i) homology in molecular architecture with the lipo-polysaccharides (LPS) in *C. jejuni*, and (ii) their abundance in peripheral nerves. Assessment of antibodies against six gangliosides (GM1, GM2, GD1a, GD1b, GT1b, GQ1b) and 15 GSCs (GM1 + GM2, GM1 + GD1a, GM1 + GD1b, GM1 + GT1b, GM1 + GQ1b, GM2 + GD1a, GM2 + GD1b, GM2 + GT1b, GM2 + GQ1b, GD1a + GD1b, GD1a + GT1b, GD1a + GQ1b, GD1b + GT1b, GD1b + GQ1b, and GT1b + GQ1b) were performed. The detection of anti-ganglioside and anti-GSC antibodies was accomplished by horseradish peroxidase (HRP) labelled anti-human antibody, which was visualized by a color-shifting substrate reagent and was read spectrophotometrically. A commercial ganglioside autoantibody detection kit (Bühlmann Laboratories AG, Schönenbuch, Switzerland) was utilized for standardizing the manually developed assay. Positive controls from the kit were used for assay validation and quality assurance.

To assay antibodies against single gangliosides, each individual ganglioside was dissolved in ethanol ( $0.2 \mu\text{g}/50 \mu\text{l}$ ) and was added to the wells of the ELISA plate, while for antibodies against GSCs, two ( $0.1 \mu\text{g}$  each) gangliosides were mixed in a microwell and left for approximately 30 minutes. The plates were kept at  $37^{\circ}\text{C}$  for several minutes for drying and complete evaporation of ethanol. Thereafter,  $50 \mu\text{l}$  of the blocking solution [1% Bovine Serum Albumin (BSA) in Phosphate-Buffered Saline (PBS)] was added to each well and was allowed to stand for 30 minutes at room temperature. The blocking solution was removed from the microwells, and the serum sample diluted (1:40) with 1% BSA in PBS was added to each well ( $50 \mu\text{l}/\text{well}$ ) and was left to stand for 90 minutes at room temperature. The plate was washed three times with  $300 \mu\text{l}$  of 0.1% BSA in PBS. Following this, HRP-conjugated secondary antibody diluted with 1% BSA in PBS was added to the wells ( $50 \mu\text{l}/\text{well}$ ) and the plate was left for 90 minutes

at room temperature. The plate was washed again with 0.1% BSA in PBS three times. Subsequently, 100  $\mu$ L of OPD substrate solution (ortho-phenylenediamine dihydrochloride dissolved in 0.1M citrate-phosphate buffer) was added and the plate was left at room temperature for 2 minutes. The color reaction was stopped by the addition of 50  $\mu$ L of 8N H<sub>2</sub>SO<sub>4</sub>. The optical densities (ODs) of the reactions were read with an ELISA plate reader at 490 nm, and the OD values were corrected by subtracting the OD of a well that was not coated with gangliosides (blank control) to obtain the ganglioside autoantibody reactivity.

The OD value 0.1 was used for defining the threshold level of seroreactivity. An OD  $\geq$ 0.1 was considered seropositive for ganglioside antibodies and the OD  $<$ 0.1 was considered seronegative. Seroreactivity to GSCs (e.g., GSC X + Y comprising gangliosides X and Y) was considered as 'anti-X + Y autoantibody positive' when the OD of anti-X + Y autoantibody was higher by 0.2 than the OD of anti-X or anti-Y autoantibody. When a serum sample showed both anti-X and anti-Y autoantibody reactivities, the serum was considered anti-X + Y autoantibody-positive only when the OD value of the anti-X + Y autoantibody was higher compared to the sum of the anti-X and anti-Y autoantibodies.

### Detection of antecedent infection

The IgM antibody capture (MAC) micro-ELISA kit was used to detect *C. jejuni* antibodies in the sera (MyBioSource, San Diego, California, USA). Serum IgM antibodies to JE virus, dengue virus, and chikungunya virus were detected using the ELISA kits manufactured by the National Institute of Virology (NIV, Pune, India) and these findings have been recently published.<sup>[6]</sup> Besides this, in the current study, we examined seroreactivities to influenza and zika viruses in the patients with GBS (n = 150) and control subjects (n = 150). For the detection of the influenza virus, throat/nasal swabs samples were collected from the study participants. The QIAamp Viral RNA Mini Kit (Qiagen) was used for the extraction and purification of viral RNA from throat/nasal swabs. Molecular detection of influenza virus RNA was carried out by real time RT-PCR using a Center for Disease Control and Prevention (CDC, USA) standardized protocol. For the detection of the Zika virus, total RNA was extracted from the plasma samples using QIAamp® Viral RNA Mini kit. A real-time PCR assay standardized by the CDC, USA, was used for the qualitative detection of zika virus.

### Statistical analysis

Statistical analyses were performed using SPSS-27 for Windows (SPSS Inc., Chicago, Illinois, USA).  $P <$ 0.05 was considered to be statistically significant. Gaussian distribution checkpoint was verified by the Shapiro-Wilk test to confirm the normality of the variables. The profile of antibodies against the single gangliosides and GSCs was compared between the patient and control groups using the Chi-square test. Further, study participants were stratified into two groups based on the presence or absence of ganglioside/GSC

antibodies. The associations of ganglioside/GSC antibodies with electrophysiological subtypes as well as immunoreactivity to infectious pathogens were tested using the Chi-square or Fisher's exact test. Benjamini-Hochberg correction at  $\alpha\%$  was applied to control the false discovery rate (FDR).

## RESULTS

### Clinical and demographic profile of the study participants


The cohort comprised 97 men (64.7%) and 53 women (35.3%) with GBS. In the control group, there were 30 men (60.0%) and 20 women (40.0%). The median age at the time of study entry was 37 years (IQR = 27 to 47 years) and 36.5 years (IQR = 30 to 43 years) in the patient and control groups respectively. Thus, the patient and control groups were matched for age and also for gender ( $P = 0.55$  and  $0.87$ , respectively). The median duration of GBS was 6 days (IQR = 4 to 10 days). Antecedent infections reported by patients included fever (n = 18, 12%), acute gastroenteritis (n = 18, 12%), and respiratory infection (n = 9, 6%). The Hughes disability scale (HDS) score at the time of study entry was 1, 2, 3, 4, and 5 in 1 (0.7%), 14 (9.3%), 44 (29.3%), 90 (60.0%) and 1 (0.7%) patient, respectively. Twelve patients (8%) eventually developed respiratory muscle weakness and required mechanical ventilation. Based on the criteria of Rajabally *et al.*,<sup>[21]</sup> there were 67 (44.7%), 43 (28.7%), 33 (22.0%), 5 (3.3%), and 2 (1.3%) patients with axonal, primary demyelinating, equivocal, inexcitable and normal electrophysiology, respectively.

### Autoantibodies against single gangliosides

The frequency of autoantibodies against all the studied gangliosides was significantly higher among patients with GBS than in the controls ( $P <$  0.001). GM1 autoantibody was the most common (80%), whilst GQ1b autoantibody was the least common (53.3%) among patients with GBS [Table 1].

### Antibodies against GSCs

In the present cohort, 43 patients (28.7%) had autoantibody positivity for any one of the tested GSCs. Autoantibodies against GSCs consisting of GM1 as one of the components were the most common [Figure 1]. None of the control subjects had autoantibodies against GSCs.



**Figure 1:** Autoantibodies against various ganglioside complexes (GSCs) in patients with Guillain-Barré syndrome

### Correlation between ganglioside/GSC antibodies and electrophysiological subtypes

The IgM autoantibodies against single gangliosides were compared between the two major electrophysiological subtypes of GBS viz. demyelinating (n = 43) and axonal (n = 67) [Table 2]. A significantly higher frequency of GT1b autoantibody was noted in the axonal (n = 54, 80.6%) as compared to demyelinating (n = 27, 62.8%) subtype of GBS ( $P = 0.039$ ). However, there were no differences in the frequencies of GSC autoantibodies between the axonal and demyelinating subtypes of GBS in the present cohort.

### Correlation between ganglioside/GSC autoantibodies and infection immunoreactivities

The data pertaining to the antecedent infections were taken from our recently published article.<sup>[6]</sup> Zika virus RNA was detected neither among patients nor in healthy controls. Two patients and none of the healthy controls tested positive for influenza virus RNA. The association of IgM immunoreactivity against the six tested pathogens namely *C. jejuni*, JE virus, dengue virus, chikungunya virus, influenza virus, and zika virus and antibodies against gangliosides and GSCs were examined. Except for preceding JE virus infection, none of the other pathogens showed a statistically significant association with antibodies against the single gangliosides. GBS patients with evidence of preceding JE virus infection exhibited significantly higher frequency of autoantibodies against GD1a ( $P = 0.001$ ), GD1b ( $P < 0.001$ ), GT1b ( $P = 0.008$ ), and GQ1b ( $P = 0.008$ ) gangliosides [Table 3]. Notably, none of the preceding infections showed any association with antibodies against GSCs.

## DISCUSSION

It has long been understood that GBS is a post-infection autoimmune disease of the peripheral nervous system. Nevertheless, the immunological process underlying GBS pathogenesis still is an enigma in a substantial number of patients. Antecedent infections and ‘molecular mimicry’ between antibodies against infectious pathogens and the host gangliosides are recognized as the key underlying mechanism of GBS. Efforts have been made to delineate the interactions between antecedent infections and antibodies against gangliosides, but the previous studies were limited only to a few pathogens. Besides, there exists a lack of clear understanding regarding the association between antecedent infections and antibodies against GSCs. The current study is the first of its kind from India to report the association between antecedent infections and antibodies against gangliosides as well as GSCs.

In the present study, the prevalence of antibodies against the six tested gangliosides was found to be in the order of GM1>GM2>GT1b>GD1b>GD1a>GQ1b. GM1 autoantibody was the commonest and was observed in majority (80%) of the patients. Similar to the present study, GM1 antibody was reported to be common in GBS patients in several other

**Table 1: Profile of antibodies against single gangliosides in Guillain-Barré syndrome**

Ganglioside autoantibodies	GBS (n=150)	Controls (n=50)	Statistics ( $\chi^2$ )	P
GM1 autoantibody	120 (80.0)	3 (6.0)	86.72	<0.001
GM2 autoantibody	117 (78.0)	2 (4.0)	85.21	<0.001
GD1a autoantibody	105 (70.0)	0	73.68	<0.001
GD1b autoantibody	107 (71.3)	1 (2.0)	72.57	<0.001
GT1b autoantibody	110 (73.3)	2 (4.0)	73.16	<0.001
GQ1b autoantibody	80 (53.3)	1 (2.0)	41.00	<0.001
Any ganglioside autoantibody	142 (94.7)	8 (16.0)	123.76	<0.001

‘GBS’: Guillain-Barré syndrome \*Numbers in parentheses represent percentages

**Table 2: Comparison of ganglioside antibody profile between demyelinating and axonal subtypes of Guillain-Barré syndrome**

Ganglioside autoantibodies	Demyelinating GBS (n=43)	Axonal GBS (n=67)	Statistics ( $\chi^2$ )	P
GM1	32 (74.4)	57 (85.1)	1.92	0.165
GM2	32 (74.4)	54 (80.6)	0.586	0.444
GD1a	28 (65.1)	50 (74.6)	1.14	0.284
GD1b	31 (72.1)	48 (71.6)	0.006	0.959
GT1b	27 (62.8)	54 (80.6)	4.27	0.039
GQ1b	27 (62.8)	33 (49.2)	1.93	0.164
Positivity for any ganglioside autoantibody	42 (97.7)	62 (92.5)	1.34	0.247

‘GBS’: Guillain-Barré syndrome. \*Numbers in parentheses represent percentages

**Table 3: Correlation of Japanese encephalitis IgM immunoreactivity and ganglioside antibodies in Guillain-Barré syndrome**

Ganglioside Autoantibodies	JE IgM immunoreactivity		Statistics ( $\chi^2$ )	P
	Positive (n=60)	Negative (n=90)		
GD1a	51 (85.0)	54 (60.0)	10.71	0.001
GD1b	55 (91.7)	52 (57.8)	20.21	<0.001
GT1b	51 (85.0)	59 (65.6)	6.69	0.008
GQ1b	40 (66.7)	40 (44.4)	7.14	0.008

‘JE’: Japanese Encephalitis. \*Numbers in parentheses represent percentages

populations, including Korean,<sup>[22]</sup> Spanish,<sup>[13]</sup> and Chinese.<sup>[23]</sup> It is important to note that the presence of GM1 antibody has been associated with the severity and prognosis of GBS.<sup>[24,25]</sup> Autoantibody against GM1 from patients with GBS was observed to impede voltage-gated calcium channels ( $Ca^{+2}$ ).<sup>[26]</sup> This leads to neuromuscular weakness due to  $Ca^{+2}$  channel dysfunction in the motor nerve-endings in patients with GBS. It is interesting to note that 16% of the healthy controls in the present study had antibodies to at least one of the gangliosides. Previous studies have reported the presence of antibodies

against gangliosides in up to 15% of healthy individuals, who were considered to be 'healthy controls', but their prevalence in the general population is not known.<sup>[11,27]</sup>

In the present study, GSC autoantibodies were detected in 28.7% of GBS patients (43/150), but in none of the control subjects. Autoantibody against the GM1 + GM2 complex was the most common anti-GSC antibody (n = 12, 8%) among patients. Similarly, in an earlier study from Italy, GSC autoantibodies were reported in 27% (17/63) of patients with GBS.<sup>[28]</sup> However, in an Italian GBS cohort, the most frequent anti-GSC antibody was against the GD1a + GD1b complex.<sup>[28]</sup> In a UK cohort of GBS, the frequency of anti-GSC antibodies was reported to be 21.7% (39/180).<sup>[29]</sup> Contrary to these studies, the prevalence of anti-GSC antibodies was found to be only 17% (39/234) in a Japanese cohort.<sup>[30]</sup> Thus, the distribution pattern of GSC autoantibodies varies across different ethnic groups. There exists a dearth of understanding on the individual as well as ethnic differences in anti-ganglioside and anti-GSC antibodies. It may be hypothesized that microbial exposures, being apparently unique to each individual and also to each ethnic group, may have some influence on the individual or ethnic differences in the autoantibody profiles. Besides, the profile of different ganglioside autoantibodies in GBS was reported to vary depending upon the technique used.<sup>[29-32]</sup> Various studies have employed different types of immunoassays in profiling antiganglioside autoantibodies. Therefore, more precise information is required to explain the methodical and ethnic differences in anti-ganglioside and anti-GSC antibodies in GBS patients.

The antibodies against the individual gangliosides/GSCs were reported to be associated with the subtypes of GBS. In the present study, only the GT1b autoantibody was more frequent in the axonal subtype. The earlier studies demonstrated an association between the axonal form of GBS and GM1 and GD1a antibodies.<sup>[16,25]</sup> In a large multi-centric study, GSC antibodies were associated with the axonal form of GBS.<sup>[33]</sup> However, in our study anti-GSC antibodies were not found to be associated with the electrophysiological subtypes of GBS.

The most salient finding in the current study was the association of antecedent JE virus infection with antibodies against GD1a, GD1b, GT1b, and GQ1b gangliosides in GBS patients. This is the first study showing an association between antecedent JE infection and ganglioside antibodies in GBS patients. Most of the previous studies focused on the association of *C. jejuni*, *M. pneumoniae*, CMV, and Epstein-Barr Virus. The current study focused on the association between the arboviral infections that are endemic to India and ganglioside antibodies. Of the studied pathogens, an association was observed between antecedent JE infection and ganglioside antibodies. It is noteworthy that no association between antecedent *C. jejuni* infection and antibodies against gangliosides was observed in the current study. Several studies reported an association between antecedent *C. jejuni* infection and ganglioside antibodies. Elevated titers of antibodies against GM1,

GD1a, GD1b, and GQ1b were reported in GBS patients with antecedent *C. jejuni* infection in Japan.<sup>[15]</sup> Antibodies against GM1 and GD1a were reported to be associated with GBS developing after *C. jejuni* infection in the French population.<sup>[16]</sup> Other studies demonstrated an association between *C. jejuni* infection and GM1 antibodies.<sup>[34-36]</sup> Interestingly, in a previous study, antibodies to *C. jejuni* were reported more frequently in GBS than controls (17.2% vs 7%) and antibodies against gangliosides such as GM1 and GD1b were present in 20% of patients with *C. jejuni* antibodies, while 9.6% of patients without *C. jejuni* antibodies also had anti-GM1 or anti-GD1b antibodies.<sup>[37]</sup> Thus, the findings on the association between antecedent *C. jejuni* infection and gangliosides antibodies are not consistent across studies. In addition, GM2 antibody was reported to be associated with CMV-associated GBS and GM1 antibody with *M. pneumoniae*-associated GBS.<sup>[16]</sup> Notably, in the current study, antecedent infections were not found to be associated with any of the anti-GSC antibodies. In contrast, in an earlier cohort of GBS, an association between antecedent gastrointestinal infection with *C. jejuni* and antibodies against GSC was noted.<sup>[30]</sup> Studies on the association between antecedent infections and GSC-antibodies are albeit limited.

## CONCLUSION

Infectious pathogens are the major risk determinants of GBS. The spectrum of microbial risk determinants of GBS is expanding, the two recently added viruses such as Zika and SARS-CoV-2 are examples of such an expansion. The functional interactions between infectious triggers and gangliosides/GSCs influence the risk, severity as well as prognosis of GBS. Given the determining role of pathogens on the risk of developing GBS, it is essential to identify their interacting immune partners and the subsequent pathophysiological trajectories. GT1b antibody was more frequent in the axonal variant of GBS in the current study. The association of antecedent JE infection with GD1a, GD1b, GT1b, and GQ1b antibodies in the present cohort of GBS provides additional insights into the role of antecedent infections in the immunobiology of GBS. This adds to the existing knowledge that besides *C. jejuni*, CMV, and *M. pneumoniae*, other pathogens also have the potential to influence the production of ganglioside antibodies. This may imply that the profile of infections antedating the onset of GBS in the tropics may differ from those in temperate regions. Further research is warranted in ethnically diverse populations with a larger number of gangliosides and GSC antigens and a wider spectrum of antecedent pathogens to obtain better insights into the functional interactions between infections and antibody responses.

## Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form the patient(s) has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

## Financial support and sponsorship

The study was funded by the Science and Engineering Research Board (SERB), Government of India (Grant No.: YSS/2015/000620). The funder was not involved in the study design, collection, analysis and interpretation of data, writing of the report, or submitting the article for publication.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Leonhard SE, Mandarakas MR, Gondim F, Bateman K, Ferreira M, Cornblath DR, *et al.* Diagnosis and management of Guillain-Barré syndrome in ten steps. *Nature reviews. Neurology* 2019;15:671-83.
- Toscano G, Palmerini F, Ravaglia S, Ruiz L, Invernizzi P, Cuzzoni MG, *et al.* Guillain-Barré syndrome associated with SARS-CoV-2. *N Engl J Med* 2020;382:2574-6. <https://doi.org/10.1056/NEJMc2009191>.
- Jacobs BC, Rothbarth PH, van der Meché FG, Herbrink P, Schmitz PI, de Klerk MA, *et al.* The spectrum of antecedent infections in Guillain-Barré syndrome: A case-control study. *Neurology* 1998;51:1110-5.
- Hao Y, Wang W, Jacobs BC, Qiao B, Chen M, Liu D, *et al.* Antecedent infections in Guillain-Barré syndrome: A single-center, prospective study. *Ann Clin Transl Neurol* 2019;6:2510-7.
- Sinha S, Prasad KN, Jain D, Pandey CM, Jha S, Pradhan S. Preceding infections and anti-ganglioside antibodies in patients with Guillain-Barré syndrome: A single centre prospective case-control study. *Clin Microbiol Infect* 2007;13:334-7.
- Dutta D, Debnath M, Nagappa M, Das SK, Wahatule R, Sinha S, *et al.* Antecedent infections in Guillain-Barré syndrome patients from south India. *J Peripher Nerv Syst* 2021;26:298-306.
- Hadden RD, Karch H, Hartung HP, Zielasek J, Weissbrich B, Schubert J, *et al.* Preceding infections, immune factors, and outcome in Guillain-Barré syndrome. *Neurology* 2001;56:758-65.
- Yu RK, Usuki S, Ariga T. Ganglioside molecular mimicry and its pathological roles in Guillain-Barré syndrome and related diseases. *Infect Immun* 2006;74:6517-27.
- Yuki N, Susuki K, Koga M, Nishimoto Y, Odaka M, Hirata K, *et al.* Carbohydrate mimicry between human ganglioside GM1 and *Campylobacter jejuni* lipooligosaccharide causes Guillain-Barre syndrome. *Proc Natl Acad Sci U S A* 2004;101:11404-9.
- Press R, Matá, S., Lolli F, Zhu J, Andersson T, Link H. Temporal profile of anti-ganglioside antibodies and their relation to clinical parameters and treatment in Guillain-Barré syndrome. *J Neurol Sci* 2001;190:41-7.
- Cutillo G, Saariaho AH, Meri S. Physiology of gangliosides and the role of anti-ganglioside antibodies in human diseases. *Cell Mol Immunol* 2020;17:313-22.
- Kaida K, Ariga T, Yu RK. Antiganglioside antibodies and their pathophysiological effects on Guillain-Barré syndrome and related disorders--a review. *Glycobiology* 2009;19:676-92.
- Lleixà C, Martín-Aguilar L, Pascual-Goñi E, Franco T, Caballero M, de Luna N, *et al.* Autoantibody screening in Guillain-Barré syndrome. *J Neuroinflammation* 2021;18:251. [10.1186/s12974-021-02301-0](https://doi.org/10.1186/s12974-021-02301-0).
- Godschalk PC, Heikema AP, Gilbert M, Komagamine T, Ang CW, Glerum J, *et al.* The crucial role of *Campylobacter jejuni* genes in anti-ganglioside antibody induction in Guillain-Barre syndrome. *J Clin Invest* 2004;114:1659-65.
- Hao Q, Saida T, Kuroki S, Nishimura M, Nukina M, Obayashi H, *et al.* Antibodies to gangliosides and galactocerebroside in patients with Guillain-Barré syndrome with preceding *Campylobacter jejuni* and other identified infections. *J Neuroimmunol* 1998;81:116-26.
- Caudie C, Quittard Pinon A, Taravel D, Sivadon-Tardy V, Orlikowski D, Rozenberg F, *et al.* Preceding infections and anti-ganglioside antibody profiles assessed by a dot immunoassay in 306 French Guillain-Barré syndrome patients. *J Neurol* 2011;258:1958-64.
- Wang L, Shao C, Yang C, Kang X, Zhang G. Association of anti-gangliosides antibodies and anti-CMV antibodies in Guillain-Barré syndrome. *Brain Behavior* 2017;7:e00690. [doi: 10.1002/brb3.690](https://doi.org/10.1002/brb3.690).
- Agarwal A, Vibha D, Srivastava AK, Shukla G, Prasad K. Guillain-Barre syndrome complicating chikungunya virus infection. *J Neurovirol* 2017;23:504-7.
- Leonhard SE, Tan CY, van der Eijk AA, Reisin RR, Franken SC, Huizinga R, *et al.* Antecedent infections in Guillain-Barré syndrome in endemic areas of arbovirus transmission: A multinational case-control study. *J Peripher Nerv Syst* 2021;26:449-60.
- Asbury AK, Cornblath DR. Assessment of current diagnostic criteria for Guillain-Barré syndrome. *Ann Neurol* 1990;27 Suppl, S21-4.
- Rajabally YA, Durand MC, Mitchell J, Orlikowski D, Nicolas G. Electrophysiological diagnosis of Guillain-Barré syndrome subtype: Could a single study suffice? *J Neurol Neurosurg Psychiatry* 1990;86:115-9.
- Kim JK, Bae JS, Kim DS, Kusunoki S, Kim JE, Kim JS, *et al.* Prevalence of anti-ganglioside antibodies and their clinical correlates with Guillain-Barré syndrome in Korea: A nationwide multicenter study. *J Clin Neurol (Seoul, Korea)* 2014;10:94-100.
- Zhu J, Zhang Y, Li R, Lin Y, Fu Y, Yan Y, *et al.* Anti-ganglioside Antibodies in Guillain-Barre syndrome: A novel immunoblotting-panel assay. *Front Neurol* 2021;12:760889. [doi: 10.3389/fneur.2021.760889](https://doi.org/10.3389/fneur.2021.760889).
- Lardone RD, Yuki N, Odaka M, Daniotti JL, Irazoqui FJ, Nores GA. Anti-GM1 IgG antibodies in Guillain-Barré syndrome: Fine specificity is associated with disease severity. *J Neurol Neurosurg Psychiatry* 2021;81:629-33.
- Kuwabara S, Yuki N, Koga M, Hattori T, Matsuura D, Miyake M, *et al.* IgG anti-GM1 antibody is associated with reversible conduction failure and axonal degeneration in Guillain-Barré syndrome. *Ann Neurol* 1998;44:202-8.
- Nakatani Y, Hotta S, Utsunomiya I, Tanaka K, Hoshi K, Ariga T, *et al.* Cav2.1 voltage-dependent Ca<sup>2+</sup> channel current is inhibited by serum from select patients with Guillain-Barré syndrome. *Neurochem Res* 2009;34:149-57.
- Kollewe K, Wurster U, Sinzenich T, Körner S, Dengler R, Mohammadi B, *et al.* Anti-ganglioside antibodies in amyotrophic lateral sclerosis revisited. *PLoS One* 2015;10:e0125339. [doi: 10.1371/journal.pone.0125339](https://doi.org/10.1371/journal.pone.0125339).
- Notturmo F, Luciani M, Caporale CM, Ciarelli A, Uncini A. Antibodies to ganglioside complexes in Guillain-Barré syndrome: Clinical correlates, fine specificity and complement activation. *Int J Immunopathol Pharmacol* 2015;22:437-45.
- Rinaldi S, Brennan KM, Kalna G, Walgaard C, van Doorn P, Jacobs BC, *et al.* Antibodies to heteromeric glycolipid complexes in Guillain-Barré syndrome. *PLoS One* 2013;8:e82337. [doi: 10.1371/journal.pone.0082337](https://doi.org/10.1371/journal.pone.0082337).
- Kaida K, Morita D, Kanzaki M, Kamakura K, Motoyoshi K, Hirakawa M, *et al.* Anti-ganglioside complex antibodies associated with severe disability in GBS. *J Neuroimmunol* 2007;182:212-8.
- Irie S, Saito T, Kanazawa N, Ogino M, Ogino Y, Sakai F. Detection of serum anti-ganglioside antibodies by latex agglutination assay in Guillain-Barré syndrome: Comparison with ELISA. *Intern Med* 2003;42:490-5.
- Bonyadi MR, Barzegar M, Badalzadeh R, Hashemilar M. Comparison of immunoblotting and ELISA for detection of anti-ganglioside antibodies in children with Guillain-Barre syndrome. *Iran J Immunol* 2010;7:117-23.
- Shahrizaaila N, Kokubun N, Sawai S, Umaphathi T, Chan YC, Kuwabara S, *et al.* Antibodies to single glycolipids and glycolipid complexes in Guillain-Barré syndrome subtypes. *Neurology* 2014;83:118-24.
- Rees JH, Gregson NA, Hughes RA. Anti-ganglioside GM1 antibodies in Guillain-Barré syndrome and their relationship to *Campylobacter jejuni* infection. *Ann Neurol* 1995;38:809-16.
- Jacobs BC, van Doorn PA, Schmitz PI, Tio-Gillen AP, Herbrink P, Visser LH, *et al.* *Campylobacter jejuni* infections and anti-GM1 antibodies in Guillain-Barré syndrome. *Ann Neurol* 1996;40:181-7.
- Tang J, Yuan J, Hao H. GM1 antibody in Guillain-Barre syndrome after *Campylobacter jejuni* infection. *Chin Med J* 1995;108:262-4.
- Vriesendorp FJ, Mishu B, Blaser MJ, Koski CL. Serum antibodies to GM1, GD1b, peripheral nerve myelin, and *Campylobacter jejuni* in patients with Guillain-Barré syndrome and controls: Correlation and prognosis. *Ann Neurol* 1993;34:130-5.