



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Society of Cardiovascular Computed Tomography guidance for use of cardiac computed tomography amidst the COVID-19 pandemic Endorsed by the American College of Cardiology

Andrew D. Choi<sup>\*</sup>, Suhny Abbara, Kelley R. Branch, Gudrun M. Feuchtner, Brian Ghoshhajra, Gianluca Pontone, Koen Nieman, Todd C. Villines, Michelle C. Williams, Ron Blankstein<sup>\*\*</sup>

*Division of Cardiology & Department of Radiology, The George Washington University School of Medicine, Washington, DC, United States*

*Division of Cardiothoracic Imaging, Department of Radiology, University of Texas Southwestern Medical Center, Dallas, TX, United States*

*Division of Cardiology, Department of Medicine, University of Washington Medical Center, Seattle, WA, United States*

*Department of Radiology, Innsbruck Medical University, Vienna, Austria*

*Division of Cardiovascular Imaging, Massachusetts General Hospital, Boston, MA, United States*

*Department of Cardiovascular Medicine and Radiology, Stanford University, Stanford, CA, United States*

*Department of Cardiovascular Imaging, Centro Cardiologico Monzino IRCCS, University of Milan, Italy*

*Division of Cardiovascular Medicine, University of Virginia Health System, Charlottesville, VA, United States*

*University of Edinburgh/British Heart Foundation Centre for Cardiovascular Science, Edinburgh, United Kingdom*

*Cardiovascular Imaging Program, Departments of Medicine (Cardiovascular Division) and Radiology, Brigham and Women's Hospital, Boston, MA, United States*

### ARTICLE INFO

#### Keywords:

COVID-19

Coronavirus

Social distancing

Cardiovascular Computed Tomography

### ABSTRACT

The world is currently suffering through a pandemic outbreak of severe respiratory syndrome coronavirus 2 (SARS-CoV-2) known as Coronavirus Disease 2019 (COVID-19). The United States (US) Centers for Disease Control and Prevention (CDC) currently advises medical facilities to “reschedule non-urgent outpatient visits as necessary”. The European Centre for Disease Prevention and Control, the United Kingdom National Health Service and several other international agencies covering Asia, North America and most regions of the world have recommended similar “social distancing” measures. The Society of Cardiovascular Computed Tomography (SCCT) offers guidance for cardiac CT (CCT) practitioners to help implement these international recommendations in order to decrease the risk of COVID-19 transmission in their facilities while deciding on the timing of outpatient and inpatient CCT exams. This document also emphasizes SCCT’s commitment to the health and well-being of CCT technologists, imagers, trainees, and research community, as well as the patients served by CCT.

### 1. Introduction

The world is currently suffering through a pandemic outbreak of severe respiratory syndrome coronavirus 2 (SARS-CoV-2) known as Coronavirus Disease 2019 (COVID-19). The United States (US) Centers for Disease Control and Prevention (CDC) currently advises medical facilities to “reschedule non-urgent outpatient visits as necessary”.<sup>1</sup> The European Centre for Disease Prevention and Control, the United Kingdom National Health Service and several other international agencies covering Asia, North America and most regions of the world have recommended similar “social distancing” measures.<sup>2,3</sup>

The Society of Cardiovascular Computed Tomography (SCCT) offers

guidance that fully support and extend these international recommendations specifically for cardiac CT (CCT) practitioners to decrease risk of COVID-19 transmission in their facilities while allowing for optimal timing considerations for effective utilization of CCT to improve cardiovascular health outcomes. While many institutions will have their own guidelines for clinicians and imagers to follow, these recommendations are meant to help CCT labs which are interested in developing or refining such policies. It is important to emphasize the SCCT’s commitment to the health and well-being of CCT technologists, imagers, trainees, and research community, as well as the patients served by CCT by preventing the spread of disease.

As this represents initial guidance for a rapidly evolving pandemic,

<sup>\*</sup> Corresponding author.

<sup>\*\*</sup> Corresponding author.

E-mail addresses: [adchoi@mfa.gwu.edu](mailto:adchoi@mfa.gwu.edu) (A.D. Choi), [rblankstein@bwh.harvard.edu](mailto:rblankstein@bwh.harvard.edu) (R. Blankstein).

**Table 1**

Guiding points to consider when deciding on the role and timing of CCT.

- The delivery of CCT services should be performed in a manner which will be safe to technologists and imagers, as well as patients.
- Consider deferring CCT exams which can be safely postponed in order to minimize risk of exposure to patients and staff.
- CCT may be preferred to transesophageal echocardiography (TEE) in order to rule-out left atrial appendage and intracardiac thrombus prior to cardioversion in order to reduce coughing and aerosolization related to TEE.
- The ability of CCT to decisively exclude coronary disease or high risk anatomy may prevent the need for inpatient admissions and resource use.
- Consider that elderly patients, those with co-morbidities, and those who may be immunosuppressed are at greater risk of morbidity/mortality from COVID-19, and the benefit and risk of cardiac CT should be evaluated on a case by case basis.
- In patients under investigation (PUI) and with confirmed COVID-19, the benefit of CCT in most clinical scenarios will likely be lower than the risk of exposure and infection to healthcare personnel. These cases should be considered on a case-by-case basis.

the SCCT advises that CCT practitioners work closely with their referring physicians to determine the appropriateness and timing of each individual study on a case by case basis, while also considering the local epidemiology of COVID-19 and local institutional guidelines for practice.

## 2. Basic concepts

- Social distancing — keeping at least six feet (1.8 m) between individuals in waiting rooms and work spaces as much as feasible.
- Encourage sick employees to stay home. Personnel who develop respiratory symptoms (e.g., cough, shortness of breath) or unexplained fever should be instructed not to report to work.
- Ensure that your sick leave policies are flexible and consistent with public health guidance and that employees are aware of these policies. Make contingency plans for increased absenteeism
- Screen patients and visitors for symptoms of acute respiratory illness (e.g., fever, cough, difficulty breathing) or gastrointestinal symptoms and coronavirus exposure in the last 2 weeks before entering one's healthcare facility.<sup>4</sup>
- Ensure technologist and CCT imager hand hygiene best practices. If soap and water are not readily available, use of a hand sanitizer that contains at least 60% alcohol.
- Consider standard droplet precautions for patients and healthcare personnel as per institutional infection control protocols.
- Increase scheduling intervals or appointment times to allow adequate time to clean equipment as needed.
- Leverage telemedicine technologies and isolated workstations to allow for reading and interpretation, that allow for social distancing to limit staff exposure, when possible.
- Assign a team member to monitor and incorporate regular updates from the CDC and appropriate regional jurisdictions.

## 3. Patients under investigation (PUI) and confirmed COVID-19

In patients under investigation (PUI) and with confirmed COVID-19, the benefit of CCT scanning in most clinical scenarios **will likely be lower than the risk of COVID-19 exposure and infection** to healthcare personnel. These cases should be considered on a case-by-case basis.

**For these PUI and confirmed COVID-19 patients in which CCT scanning is determined to be necessary, the following issues**

**should be considered:**

- **Ensure proper use of personal protection equipment (PPE).** Healthcare personnel including technologists, radiologists and cardiac imagers who come in close contact with confirmed or suspected COVID-19 should wear the appropriate personal protective equipment.<sup>5,6</sup> Patients should wear a surgical mask during imaging to ensure standard droplet precautions.
- **Appropriate environmental cleaning and decontamination** of rooms by thorough cleaning of the surfaces by a staff member with appropriate PPE as per CDC and local institutional guidelines for airborne viral diseases.<sup>7</sup>

## 4. Cardiac CT indications and timing

To advise practitioners of cardiovascular CT on how to implement the CDC recommendation of rescheduling non-urgent visits as necessary and international guidelines on social distancing, the SCCT offers the following guiding points (Table 1) and suggestions for CCT timing based on various indications (Table 2). As this is not an exhaustive list, the SCCT advises CCT practitioners to work with referring physicians on a case by case basis.

## 5. Incidental pulmonary findings in patients at risk of COVID-19 exposure

COVID-19 is a viral pneumonia, with a spectrum of findings ranging from normal lungs to acute respiratory distress syndrome. Typical chest CT findings in known cases are described elsewhere.<sup>8,9</sup> If typical or atypical pulmonary findings are encountered, consultation with a radiologist with thoracic expertise is encouraged, and appropriate documentation and timely communication of these findings is important, especially in cases not known or suspected to have the disease.

## 6. Conclusion

As this situation is shifting rapidly, the information contained within this document is likely to evolve. The SCCT will maintain an updated version of this statement as more information becomes available on the Society's website at <https://scct.org/page/COVID-19>. The SCCT advises that members keep informed regarding future updates from the medical and radiological communities on protecting patients, staff, trainees and providers from COVID-19 while deciding on the optimal timing of outpatient and inpatient CCT exams.

## Other Resources

American College of Cardiology COVID-19 bulletin: <https://www.acc.org/latest-in-cardiology/features/accs-coronavirus-disease-2019-covid-19-hub>

American College of Radiology recommendations for the use of chest radiography and computed Tomography (CT) for suspected COVID-19 infection, <https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Recommendations-for-Chest-Radiography-and-CT-for-Suspected-COVID19-Infection>

British Society of thoracic imaging: COVID-19 resources, <https://www.bsti.org.uk/covid-19-resources/>

Centers for disease control and prevention steps for healthcare facilities, <https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/steps-to-prepare.html>

Coronavirus (COVID-19): UK government response, <https://www.gov.uk/government/topical-events/coronavirus-covid-19-uk-government-response>

**Table 2**  
Timing considerations for common indications for CCT amidst COVID-19.

	Elective Indications (May be rescheduled > 8 weeks)	Semi-Urgent Indications (Consider scanning within 4-8 weeks)	Urgent Indications (Consider scanning within hours to < 2-4 weeks)
<b>CAD</b>	<ul style="list-style-type: none"> <li>Asymptomatic coronary artery calcium imaging</li> </ul>		<ul style="list-style-type: none"> <li>Acute chest pain when sufficient clinical suspicion for CAD</li> </ul>
	<ul style="list-style-type: none"> <li>Stable chest pain without high suspicion for CAD</li> </ul>		<ul style="list-style-type: none"> <li>Stable chest pain at high risk for events, or when there is concern for possible high-risk coronary anatomy</li> </ul>
<b>SHD</b>	<ul style="list-style-type: none"> <li>Stable structural heart patients (eg TAVR, TMVR, LAA closure in conjunction with Heart Team)*</li> </ul>		<ul style="list-style-type: none"> <li>Patient requiring urgent structural intervention (eg, TAVR, TMVR, LAA closure)</li> </ul>
<b>A-FIB</b>	<ul style="list-style-type: none"> <li>Pulmonary vein assessment for A-Fib Ablation planning*</li> </ul>	<ul style="list-style-type: none"> <li>Evaluation of left atrial appendage in <u>chronic</u> atrial arrhythmia prior to restoration of sinus rhythm</li> </ul>	<ul style="list-style-type: none"> <li>Evaluation of left atrial appendage in <u>acute</u> atrial arrhythmia prior to restoration of sinus rhythm<sup>+</sup></li> </ul>
<b>Heart Failure</b>	<ul style="list-style-type: none"> <li>Stable cardiomyopathy patients</li> </ul>		<ul style="list-style-type: none"> <li>Acute inpatient cardiomyopathy in low to intermediate pretest probability of CAD, only if CCT would change management</li> <li>Evaluation of LVAD dysfunction</li> </ul>
<b>Valvular</b>	<ul style="list-style-type: none"> <li>Evaluation for aortic stenosis severity</li> </ul>	<ul style="list-style-type: none"> <li>Sub-acute to chronic prosthetic valve dysfunction</li> </ul>	<ul style="list-style-type: none"> <li>Acute symptomatic prosthetic heart valve dysfunction, endocarditis, perivalvular extension of endocarditis or possible valve abscess</li> </ul>
<b>Masses/Congenital</b>	<ul style="list-style-type: none"> <li>Cardiac masses, which are suspected to be benign or unlikely to plan biopsy or surgery</li> </ul>		<ul style="list-style-type: none"> <li>New cardiac masses which are suspected to be malignant, if necessary to plan biopsy or surgery</li> </ul>
	<ul style="list-style-type: none"> <li>Elective evaluation of congenital anatomy</li> </ul>		<ul style="list-style-type: none"> <li>Rule-out left ventricular thrombus following equivocal echocardiography when alternative diagnostic tests (e.g. MRI) are not feasible</li> </ul>

\*Especially in institutions that will delay such elective cases

<sup>+</sup>When cardioversion is deemed necessary

CAD = coronary artery disease; SHD = structural heart disease; A-Fib = atrial fibrillation; LAA = left atrial appendage; TAVR = transcatheter aortic valve replacement/implantation; TMVR = transcatheter mitral valve replacement/implantation; LVAD = left ventricular assist device

European Centre for disease prevention and control: COVID-19, <https://www.ecdc.europa.eu/en/novel-coronavirus-china>  
 Radiology department preparedness for COVID-19: radiology scientific expert panel, <https://pubs.rsna.org/doi/10.1148/radiol.2020200988>  
 Society of Thoracic Radiology/American Society of Emergency Radiology COVID-19 position statement, <https://thoracicrad.org/>

**References**

- Centers for disease control and prevention: steps healthcare facilities can take now to prepare for coronavirus disease 2019 (COVID-19). Available from <https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/steps-to-prepare.html>.
- European Centre for disease prevention and control: COVID-19. Available from <https://www.ecdc.europa.eu/en/novel-coronavirus-china>.
- NHS England: coronavirus guidance for clinicians. Available from <https://www.england.nhs.uk/coronavirus/>.
- Centers for disease control and prevention: what healthcare personnel should know about caring for patients with confirmed or possible COVID-19 infection. Available from <https://www.cdc.gov/coronavirus/2019-ncov/hcp/caring-for-patients.html>.
- Centers for disease control and prevention: infection control. Available from <https://www.cdc.gov/coronavirus/2019-ncov/infection-control/index.html>.
- Centers for disease control and prevention: interim infection prevention and control recommendations for patients with suspected or confirmed coronavirus disease 2019 (COVID-19) in healthcare settings. Available from <https://www.cdc.gov/coronavirus/2019-ncov/infection-control/control-recommendations.html>.
- Centers for disease control and prevention: environmental infection control guidelines. Available from <https://www.cdc.gov/infectioncontrol/guidelines/environmental/background/air.html#c2c>.
- Mossa-Basha M, Meltzer CC, Kim DC, Tuite MJ, Kolli KP, Tan BS. Radiology department preparedness for COVID-19: radiology scientific expert panel. *Radiology*. 2020:200988.
- British Society of thoracic imaging: COVID-19 Resources. Available from <https://www.bsti.org.uk/covid-19-resources/>.

**SCCT COVID-19 Guidance Statement (Endorsed by the ACC) - Relationships With Industry**

Committee Member	Employment	Consultant	Speakers Bureau	Ownership/ Partnership/ Principal	Personal Research	Employment or Salary Support	Institutional, Organizational or Other Financial Benefit	Expert Witness
Andrew D. Choi, Co-Chair	The George Washington University School of Medicine, US	None	None	None	None	None	None	None
Ron Blankstein, Co-Chair	Brigham and Women's Hospital and Harvard Medical School, US	None	None	None	None	None	Research grant Amgen Inc., Astellas Inc	None
Suhny Abbara	University of Texas Southwestern Medical Center, US	None	None	None	None	None	Elsevier (Royalties)	None
Kelley R. Branch	University of Washington Medical Center, US	None	None	None	None	None	None	None
Gudrun M. Feuchtner	Innsbruck Medical University, Austria	None	None	None	None	None	None	None
Brian Ghoshhajra	Massachusetts General Hospital, US	Siemens Healthineers	None	None	None	None	None	None
Koen Nieman	Stanford University, US	None	None	None	None	None	Unrestricted research support from Siemens Healthineers, Bayer Healthcare and Heartflow, Inc.	None
Gianluca Pontone	University of Milan, Italy	None	GE Healthcare, Heartflow, Inc., Bracco	None	GE Healthcare, Heartflow, Inc.	None	None	None
Todd C. Villines	University of Virginia, US	None	None	None	None	None	None	None
Michelle C. Williams	University of Edinburgh/ British Heart Foundation, UK	None	None	None	None	None	None	None