

A Case of Acute Pancreatitis developing after Extracorporeal Shock Wave Lithotripsy

¹Vedat Goral, ²Erkan Sahin, ³Murat Arslan

ABSTRACT

Extracorporeal shock wave lithotripsy (ESWL) is a standard treatment method used for the treatment of renal calculi and upper ureteral calculi. Acute pancreatitis is a serious condition which develops due to multiple etiologic factors and is characterized by autodigestion of the pancreas. A case of acute pancreatitis which developed following ESWL performed for right renal calculi treatment is presented here.

Keywords: ESWL, Complication, Acute pancreatitis.

How to cite this article: Goral V, Sahin E, Arslan M. A Case of Acute Pancreatitis developing after Extracorporeal Shock Wave Lithotripsy. *Euroasian J Hepato-Gastroenterol* 2015;5(1):52-54.

Source of support: Nil

Conflict of interest: None

Copyright and License information: Copyright © 2015; Jaypee Brothers Medical Publishers (P) Ltd. This work is licensed under a Creative Commons Attribution 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/3.0/>

Received Date: 07-07-2014

Accepted Date: 08-09-2014

INTRODUCTION

A 41-year-old female patient had received extracorporeal shock wave lithotripsy (ESWL) at the Urology Department for calculi in the right kidney in our hospital. The ESWL procedure involved 3,000 shots at each session with ELMED (Orlando, FL, USA) lithotripter, starting from 6 voltage and increasing up to 18 to 20 voltage depending on the hardness of the calculus. The patient developed acute pancreatitis after the third session. Upon completion of the third ESWL session, the patient presented to the emergency department with abdominal pain localized in the epigastrium, nausea and vomiting. Acute pancreatitis was diagnosed with the examinations performed, and the patient was, therefore, hospitalized and started conservative treatment. When the patient first presented, she had severe pain and tenderness localized in the epigastrium. Measurements at presentation: amylase: 5178 IU/L (N: 25-125 U/L), lipase: 1030 U/L (N: 10-140 U/L), ALT: 74 IU/ml, AST: 132 IU/ml, WBC: 10300/mm³, CRP: 2.7 mgr/dl, and RANSON score was 1 at admission. Patient's fasting glucose, urea, creatinine, sodium, potassium and calcium values were within normal ranges. In addition, there were 7 mm calculi at the

right kidney (mid and inferior calyx) and a 7 mm calculus at the right proximal ureter (Fig. 1). Abdominal computed tomography (CT) revealed acute pancreatitis diagnosis on the basis of volume increases in the pancreas head, corpus and body, contamination of the peripancreatic fatty lanes and minimal peripancreatic fluid (Fig. 2) and Balthazar score was C. Gallbladder and intra/extrahepatic bile ducts were normal. Results of MRCP X-ray were normal. Other causes which may cause acute pancreatitis (e.g. gallstones, alcohol, hypercalcemia, hyperparathyroidism, medications, infections, hypertriglyceridemia) were excluded. Conservative treatment for acute pancreatitis was given at the hospital. Oral intake was stopped except for fluid intake. Follow-up examinations 6 days later demonstrated overall improvement and also improvements in laboratory parameters. The calculus was found to be eliminated at the third session of ESWL.

DISCUSSION

Acute pancreatitis results from the activation of pancreatic enzymes within the pancreas and autodigestion of the pancreatic tissue.¹ Although, its etiology differs among countries, gallbladder small stones and alcohol

¹Department of Gastroenterology, Izmir University School of Medicine, Izmir, Turkey

²Department of Radiology, Izmir University School of Medicine, Izmir, Turkey

³Department of Urology, Izmir University School of Medicine, Izmir, Turkey

Address reprint requests to: Vedat Goral, Professor, Department of Gastroenterology, Izmir University School of Medicine, Izmir, Turkey, Phone: 0090 232 3995050, Fax: 0090 232 3995070, e-mail: vegoral@hotmail.com




Fig. 1: Right renal stone



Fig. 2: Acute pancreatitis at upper abdominal CT (Balthazar score C)

are mainly responsible for about 70 to 80% of the cases. Alcohol ranks first among etiologic factors in western countries while biliary factors are the leading causes in our country. Other causes include abdominal trauma, hypertriglyceridemia, pancreas or ampulla Vater tumor, drugs, infectious causes, endoscopic retrograde cholangiopancreatography (ERCP), autoimmune causes and post-surgical conditions.¹ Acute pancreatitis has two distinct courses, i.e. mild disease and severe disease. Of the acute pancreatitis cases, 70 to 80% is mild and 20 to 30% is severe. A wide range of clinical manifestations, from mild interstitial edema to severe hemorrhagic gangrene and necrosis may be manifested.

Extracorporeal shock wave lithotripsy is a non-invasive standard method with recent successful use in the treatment of renal calculi.^{2,3} Its complications are mostly transient and mild. There have been occasional reports of potentially life-threatening subcapsular hemorrhage and perirenal hematoma, urosepsis, venous thrombosis, biliary obstruction, intestinal perforation, lung damage, aortic aneurysm rupture, intracranial hemorrhage, pneumonia, cardiac arrhythmia and acute pancreatitis.⁴⁻¹⁵ The incidence of these major complications is less than 1%. The incidence of acute pancreatitis developing post-ESWL has been reported as 0.4 to 0.7% in the literature. Our patient's medical history did not include gallbladder stones, alcohol or drug use, metabolic causes (e.g. hypercalcemia, hyperparathyroidism), infectious agents (e.g. CMV, EBV, HBV) or any other etiologic factors. Extracorporeal shock wave lithotripsy was considered as etiologic factor for acute pancreatitis in this case report. Few isolated reports of acute pancreatitis following ESWL exist in literature, but the exact mechanism of pancreatitis is not known.⁴⁻¹⁵ Possibly as a result of widening of the focus during ESWL, a part of

the pancreas was exposed to extracorporeal shock waves resulting in acute cellular damage (trauma effect) leading to small hematoma, microvascular damage and finally to acute pancreatitis.^{5,7,9,10}

CONCLUSION

In cases of sudden abdominal pain, nausea and vomiting following ESWL, consideration should be given to acute pancreatitis, and ESWL should be discontinued immediately to avoid pancreatic damage in these patients.

REFERENCES

1. Sah RP, Dawra RK, Saluja AK. New insights into the pathogenesis of pancreatitis. *Curr Opin Gastroenterol* 2013; 29(5):523-530.
2. Zhong P, Preminger GM. Physics of shock-wave lithotripsy. In: Coe FL, Favus MJ, Pak CY, Parks JH, Preminger GM, editors. *Kidney Stones: Medical and Surgical Management*. Philadelphia: Lippincott-Raven Publishers; 1996. p. 529-548.
3. Rassweiler JJ, Köhrmann KU, Seemann O, Tschada R, Alken PM. Clinical comparison of ESWL. In: Coe FL, Favus MJ, Pak CY, Parks JH, Preminger GM, editors. *Kidney stones: medical and surgical management*. Philadelphia: Lippincott-Raven Publishers; 1996. p. 571-605.
4. Weng CH, Ho PY, Tsai CC, Hsu JM, Chen M, Lin WR. Severe acute pancreatitis with abscess after extracorporeal shock wave lithotripsy: a rare complication. *Urolithiasis* 2013 Apr; 41(2):133-136.
5. Karakayali F, Sevmiş S, Ayvaz I, Tekin I, Boyvat G. Acute necrotizing pancreatitis as a rare complication of extracorporeal shock wave lithotripsy. *Int J Urol* 2006 May; 13(5):613-615.
6. Hassan I, Zietlow SP. Acute pancreatitis after extracorporeal shock wave lithotripsy for a renal calculus. *Urology* 2002 Dec; 60(6):1111.
7. Abe H, Nisimura T, Osawa S, Miura T, Oka F. Acute pancreatitis caused by extracorporeal shock wave lithotripsy for bilateral renal pelvic calculi. *Int J Urol* 2000 Feb;7(2):65-68.

8. Mylarappa P, Javali T, Prathvi, Ramesh D. Acute pancreatitis and development of pancreatic pseudo cyst after extracorporeal shock wave lithotripsy to a left renal calculus: A rare case with review of literature. *Indian J Urol* 2014 Jan;30(1): 110-112.
9. Conlin MJ. Complications of extracorporeal shock wave lithotripsy. In: Taneja SS, Smith RB, Ehrlich RM, editors. *Complications of Urologic Surgery*. Philadelphia: WB Saunders; 2001. p. 155-164.
10. Lazarides MK, Drista H, Arvanitis DP, Dayantas JN. Aortic aneurysm rupture after extracorporeal shock wave lithotripsy. *Surgery* 1997 Jul;122(1):112-113.
11. Geh JL, Curley P, Mayfield MP, Price JJ. Small bowel perforation after extracorporeal shock wave lithotripsy. *Br J Urol* 1997 Apr;79(4):648-649.
12. Hung SY, Chen HM, Jan YY, Chen MF. Common bile duct and pancreatic injury after extracorporeal shock wave lithotripsy for renal stone. *Hepatogastroenterology* 2000 Jul-Aug;47(34):1162-1163.
13. Ell C, Kerzel W, Heyder N, et al. Tissue reactions under piezoelectric shockwave application for the fragmentation of biliary calculi. *Gut* 1989 May;30(5):680-685.
14. Gülüm M, Yeni E, Savas M, Çelik H, Çiftçi H. Do the applied shock wave numbers and the application side affect pancreatic exocrine functions in patients who have undergone extracorporeal shock wave lithotripsy for upper urinary tract calculi? *Turk J Med Sci* 2012;42(4):591-596.
15. Florio G, Cicia S, Del Papa M, Carnì D. Pancreatic pseudocyst caused by extracorporeal shock wave lithotripsy for renal pelvic calculi. *Chir Ital* 2003 Mar-Apr;55(2):279-282.