

RESEARCH

Open Access

Thermal limits of two biting midges, *Culicoides imicola* Kieffer and *C. bolitinos* Meiswinkel (Diptera: Ceratopogonidae)

F Arné Verhoef^{1*}, Gert J Venter² and Christopher W Weldon¹

Abstract

Background: *Culicoides imicola* Kieffer and *Culicoides bolitinos* Meiswinkel (Diptera: Ceratopogonidae) are both of veterinary importance, being vectors of Schmallenberg, bluetongue and African horse sickness (AHS) viruses. Within South Africa, these *Culicoides* species show a marked difference in their abundances according to altitude, with *C. imicola* highly abundant in lower altitudes, but being replaced as the dominant species by *C. bolitinos* in cooler, high-altitude regions.

Methods: The thermal physiology of field collected adults of each species was determined to evaluate whether it could account for differences in their distribution and abundance. Critical thermal maxima (CT_{max}) and minima (CT_{min}), as well as upper and lower lethal temperatures (ULT and LLT) were assessed after acclimation temperatures of 19°C, 24°C and 29°C. Critical thermal limits were determined using an ecologically relevant rate of temperature change of 0.06°C.min⁻¹.

Results: Significant differences in CT_{min} and CT_{max} were found between acclimation temperatures for *C. imicola* and *C. bolitinos*. In *C. bolitinos*, the LLT of individuals acclimated at 24°C was significantly improved ($LLT_{50} = -6.01^\circ\text{C}$) compared with those acclimated at the other temperatures ($LLT_{50} = -4^\circ\text{C}$). Acclimation had a weak (difference in LLT_{50} of only 1°C) but significant effect on the LLT of *C. imicola*. When CT_{min} , CT_{max} , LLT and ULT were superimposed on daily maximum and minimum temperature records from locations where each tested *Culicoides* species is dominant, it was found that temperatures frequently declined below the CT_{min} and LLT of *C. imicola* at the location where *C. bolitinos* was dominant.

Conclusions: The distribution and abundance of *C. imicola* is likely directly constrained by their relatively poor tolerance of lower temperatures. Results for *C. bolitinos* suggest that the adult phase is hardy, and it is hypothesised that the thermal biology of other life stages could determine their range.

Keywords: Bluetongue, African horse sickness, Orbiviruses, Phenotypic plasticity, Thermal biology, Tolerance limits

Background

Biting midges in the genus *Culicoides* (Diptera: Ceratopogonidae) are incredibly abundant, small flies (ranging from 1–4 mm). Female biting midges feed on vertebrate blood, and in so doing, act as vectors of up to 66 viruses that affect animals including humans [1,2]. At least three of these viruses, namely bluetongue (BTV), epizootic haemorrhagic disease, and African horse sickness (AHSV) viruses,

all belonging within the genus *Orbivirus* (Reoviridae), cause diseases of such international significance in livestock that they have been classified as notifiable to the World Organisation for Animal Health (OIE). The rapid spread of BTV in Europe, outbreaks of AHSV in Spain since 1987 [3], and recent outbreaks of a novel orthobunyavirus, Schmallenberg virus [4,5], has demonstrated the devastating effect of these viruses on naïve livestock populations. Due to the emergence and over-wintering potential of these viruses in Europe, whereby novel European *Culicoides* species have been incriminated as the vector [6], there is an increased interest in the vector competence

* Correspondence: arne.verhoef@zoology.up.ac.za

¹Flies of Economic Significance Research Group, Department of Zoology and Entomology, University of Pretoria, Private Bag X20, Hatfield 0028, South Africa

Full list of author information is available at the end of the article

and ecology of *Culicoides* species [7]. Similarly, in sub-Saharan Africa where AHS is endemic and widespread, long term economic losses have resulted from export embargoes on horses as well as direct loss of livestock [8]. The population dynamics of *Culicoides* species are markedly influenced by environmental conditions [9]. On account of these factors, there is a need to understand the biotic and abiotic drivers of *Culicoides* distribution and abundance.

Two *Culicoides* species have been implicated in the transmission of BTV and AHSV in southern Africa, namely *C. imicola* and *C. bolitinos*. *Culicoides (Avaritia) imicola* Kieffer is one of the most widely distributed members of *Culicoides* in the world, extending over the African continent into southern Europe and eastwards to south China [10,11]. Based on its wide geographical distribution and host preference for larger mammals, as indicated by the high numbers collected near livestock, *C. imicola* is considered the principal vector of AHSV, equine encephalitis virus and BTV in South Africa [12,13] and BTV in southern Europe [14,15]. Often found in great densities, with single trap captures able to reach 10^6 individuals [16], they may make up 99% of all *Culicoides* species collected near livestock in areas where they are abundant [10,17]. The larvae develop in moist soils rich in organic matter, often in full sun. In years with above-average rainfall, a 200-fold increase in abundance may be observed [16] due to increased availability of breeding sites. In contrast, *C. bolitinos* Meiswinkel occurs in low numbers across sub-Saharan Africa, but can become the dominant species in cooler, high altitude regions [16,18] such as the Free State province of South Africa, and Lesotho. In South Africa, *Culicoides bolitinos* was also shown to be abundant in the winter rainfall region of the Western Cape province [19,20], and the dominant *Culicoides* species, in the absence of *C. imicola*, in the sandy dunefields adjoining Port Elizabeth in the Eastern Cape Province [21]. The absence of *C. imicola* at Port Elizabeth in light trap collections made at Struisbaai and Alexander Bay on the southern and western coastline was attributed to the sandiness of the soil [22]. Described in 1989 [11], *C. bolitinos* was identified as a potential vector of AHSV as recently as 1998 [16]. The larval cycle is completed in bovid dung (buffalo, domestic cattle and blue wildebeest) [11]. The abundance of *C. bolitinos* is thus affected by the availability of suitable dung, with the decomposition of dung potentially acting to incubate *C. bolitinos* larvae.

Despite the role of *Culicoides* species as potential vectors and their increased relevance to epidemiological research [23] little is known about their basic physiology. This is most likely due to their small size, nocturnal behaviour as well as ease of injury during handling, which has discouraged or hampered experimental manipulation [10]. As noted above, *C. imicola* and *C. bolitinos* show

marked differences in abundance, seemingly according to regional climate. It is likely, therefore, that thermal physiology may be a determining factor in their distribution and abundance. Most of the physiological processes of insects are temperature-dependent [24] and thermal limits are significant for fitness because they determine the ability of an organism to remain active and survive during extreme conditions [25]. Extending from this tenet, thermal limits form a fundamental part of mechanistic niche modelling [26] that aid in the generation of distribution maps and risk management strategies. These distribution models are increasingly important with a changing climate, in order to assist in predicting future outbreaks or range expansions or to assess local changes in dominance of species [27]. Some work has been done on the role of temperature on the geographical distribution of *Culicoides* species [28], including *C. imicola* [29-32], mainly through the use of climate envelope models. The effect of temperature on BTV replication within both *C. imicola* and *C. bolitinos* has also been investigated [33,34]. However, the tolerance of these species to temperature extremes has never been established, and spatial models are coarse [23]. Knowledge of the thermal biology of these species may aid in understanding their population dynamics, particularly their distribution and abundance, and may even provide insight into the epidemiology of the arboviruses they transmit.

This study determined the thermal limits of adult *C. imicola* and *C. bolitinos*. Phenotypic plasticity in measured thermal tolerance traits was also determined. Results were related to the temperature conditions of locations within the known geographical distribution of these two livestock associated species.

Methods

Collections and study animals

Rearing *Culicoides* in the laboratory is extremely difficult [10], and due to unsuccessful rearing attempts during pilot studies, only the thermal limits of adult, field collected individuals were investigated in this study. All midges were collected using Onderstepoort light traps during March, April and October, 2013. Trap catches of both species are highly female biased, so all individuals tested using the methods outlined below were female. All *C. imicola* were trapped at ARC-Onderstepoort Veterinary Institute, Gauteng, South Africa (ARC-OVI; 25° 39' 2.58" S 28°11' 8.58" E; 1219 m a.s.l.). The Onderstepoort area is relatively frost-free with only a few days where the minimum temperature will fall below 0°C. *Culicoides bolitinos* were trapped on Koeberg farm near the Clarens Valley, Free State, South Africa (28° 0' 0" S, 28° 31' 0.12" E, 1631 m a.s.l.). Clarens regularly experiences minimum daily temperatures below freezing, with occasional snow in winter. Both species were transported in cool boxes to the Hatfield Campus of the University of Pretoria where they were

housed as described by Venter et al. [35] and fed a 10% sucrose solution (in distilled water) from saturated cotton wool. Adult *C. imicola* and *C. bolitinos* were kept at acclimation temperatures of 19°C, 24°C and 29°C with a 12 L:12D photoperiod for three days before being subjected to thermal tolerance assays. In most cases, insects held at constant temperatures for a period of three days will exhibit phenotypic changes in thermal tolerance traits [36].

Temperature data from meteorological stations close to ARC-OVI (Pretoria-Arcadia, 30164, -25° 44' 18.8514" S 28° 12' 26.3874" E, 1400 m a.s.l) and Clarens (Golden Gate-Clarens, 30647, -28° 30' 13.716" S 28° 35' 1.68" E, 1849 m a.s.l) were obtained for the period 1st January 2011 until 31st October 2013 from the ARC-Institute for Soil, Climate and Water (Arcadia, South Africa). These data were used to determine the relevance of the thermal limits of both species.

Critical thermal limits

A dynamic, ramping temperature protocol was used to determine the critical thermal minimum (CT_{min}) and maximum (CT_{max}) for each species after having been exposed to the three acclimation temperatures. It has been reported that critical thermal limits vary with methodological context [37], so an ecologically relevant rate of change was determined and used. Little of the microhabitat preference of adult *C. imicola* is known, except their heliophilic and exophilic nature [31]. Consequently, a known larval habitat and adult catch site was chosen. The appropriate rate of temperature change was determined using iButtons with a 0.5°C resolution housed in waterproof silicone enclosures (SL50-ACC06, Maxim, Dallas, TX, USA). The iButtons were buried just beneath the soil surface in two waterlogged areas, in full sun, at ARC-OVI. Emergence traps indicated this area to be a habitat for *C. imicola*. The mean daily temperature change of this microhabitat was found to be 0.06°Cmin⁻¹.

To determine critical thermal limits, individual midges were placed in an aluminium 8-chamber stage, made for use under a dissecting microscope (Additional file 1) and connected to a programmable circulating refrigerated waterbath (CC-K25, Huber Kältemaschinenbau, Offenburg, Germany). A 1:1 blend of propylene glycol and water was pumped through silicone tubing from the waterbath to the stage, through which a number of channels had been drilled to permit flow of liquid and the exchange of heat. Biting midges were placed in individual 0.2 mL plastic microcentrifuge tubes (430-11, WhiteSci, Bellville, South Africa) and subjected to a heating or cooling rate of 0.06°C.min⁻¹ after a ten minute equilibration period at the start temperature of 24°C. Sample size for each treatment was $n = 14$ (two trials, each with $n = 7$). CT_{min} was defined as the point where muscle contraction

ceased (i.e. cold stupor). CT_{max} was initially defined as the onset of muscle spasms [36]. However, no discernible difference between erratic movements due to discomfort or muscle spasms could be seen, and thus CT_{max} was redefined as the heat stupor point, when muscle contractions ceased. In both cases, midges were gently prodded with a horse hair attached to a wooden toothpick to ascertain an inability to move. A type-T thermocouple connected to an eight-channel temperature data logger (TC-08, Pico Technology, St. Neots, UK) was inserted into an empty tube to record temperature in the remaining chamber of the stage.

For each species, data were checked for equality of variances (Levene's test) and inspected for normality. In all cases, the data met these assumptions. The effect of acclimation on both CT_{min} and CT_{max} were assessed by subjecting data to a one way analysis of variance. Post-hoc multiple comparisons were performed using Fisher's LSD tests. Analyses were implemented using Statistica v. 7 (Statsoft, Tulsa, Oklahoma, USA).

Upper and lower lethal temperatures

A static temperature, plunging protocol was adopted to determine lethal temperatures. Ten to 20 individuals from each species and acclimation temperature were placed in 2 ml centrifuge tubes, with 5 replicates per test temperature. The centrifuge tubes were sealed before being placed in a resealable plastic bag that was immersed in a programmable circulating refrigerated waterbath. The fluid (1:1 blend of propylene glycol and water) in the waterbath was held at a set test temperature. The midges were exposed to the test temperature for 2 hours, after which they were placed in a 70 mm Petri dish lined with a moistened 2 cm² filter paper, and given 6 hours to recover. Survival was noted after the recovery period. Temperatures tested were: -6°C, -3°C and 0°C for lower lethal temperatures (LLT), and 36°C, 39°C and 42°C for upper lethal temperatures (ULT). Pilot studies indicated no temperature-related mortality above (for LLT) or below (for ULT) these temperatures and they were omitted. Complete mortality for *C. bolitinos* acclimated at 19 and 29°C was recorded at a test temperature of -6°C, and so tests at lower temperatures were performed on these flies. *Culicoides bolitinos* acclimated at 24°C survived below -6°C (100% mortality was recorded at -8°C) but this data was not included in analysis of LLT.

Generalised linear models with binomial distribution and logit link function were used to test the effect of test temperature, acclimation and the interaction between test temperature and acclimation on survival. Probit regression analyses were used to determine upper and lower median lethal temperatures (ULT₅₀ and LLT₅₀). All lethal temperature data analyses were performed in SPSS Statistics 21 (IBM Corporation, Armonk, New York, USA).

Results

Critical thermal limits

Critical thermal limits for *C. imicola* responded significantly to acclimation (CT_{min} : $F_{2,39} = 23.553$; $p < 0.0001$; CT_{max} : $F_{2,39} = 22.2$; $p < 0.0001$) with an increase of 2.10°C in CT_{min} and 1.63°C for CT_{max} between acclimation temperatures of 19 and 29°C (Figure 1). Post hoc comparisons indicated that the CT_{min} of *C. imicola* from each acclimation temperature was significantly different from all other acclimation temperatures. For CT_{max} , *C. imicola* acclimated at 19°C and 24°C did not differ significantly from each other but both differed significantly from the individuals acclimated at 29°C.

For *C. bolitinos*, variation in critical thermal limits in response to acclimation was less pronounced than in *C. imicola* (Figure 1; CT_{min} : 1.37°C; CT_{max} : 1.30°C), but these differences were statistically significant (CT_{min} : $F_{2,38} = 4.64907$; $p = 0.0156$; CT_{max} : $F_{2,39} = 11.3$; $p = 0.0001$). Fisher's LSD tests revealed that the CT_{min} between 19°C- and 24°C-acclimated individuals did not differ significantly, but both differed significantly from *C. bolitinos* acclimated at 29°C. For CT_{max} the 24°C treatment was lowest and differed significantly from the 19°C- and 29°C-acclimated individuals, which did not differ significantly from each other.

Lethal temperatures

The interaction of LLT test temperatures and acclimation temperature had a significant effect on survival of both species (Table 1, Figure 2 A and B). *Culicoides bolitinos* acclimated to 24°C had the lowest LLT_{50} (-6.01°C), whereas those acclimated at 29°C had the highest LLT_{50} (-3.98°C). Parameter estimates relative to an acclimation temperature of 29°C were inspected and indicated that survival of *C. bolitinos* was significantly higher ($p < 0.001$) when acclimated at 24°C. LLT of those acclimated at 19°C and 29°C were not significantly different from each other. There was no significant effect of acclimation on the LLT of *C. imicola*. The LLT_{50} between acclimation treatments varied by 1°C, with the lowest estimated for 19°C (-4.39°C).

Interactions between acclimation and temperature were significant for the ULT of both species (Table 1) Upper lethal temperatures were generally similar for both species, (mean ULT_{50} : *C. imicola* = 40.05°C; *C. bolitinos* = 40.39°C). No significant difference relative to 29°C was noted for 19°C-acclimated *C. bolitinos*, though individuals acclimated to 24°C did differ significantly from 29°C, with 24°C-acclimated *C. bolitinos* having the highest ULT_{50} (41.59°C). For *C. imicola*, parameter estimates relative to 29°C indicated a significant




Table 1 Effects of temperature and acclimation on the thermal limits of *Culicoides bolitinos* and *C. imicola*

Species	Trait	Effect	Wald χ^2	d.f.	p
<i>C. bolitinos</i>	ULT	Intercept	276.072	1	<0.0001
		Acclimation	341.425	2	<0.0001
		Temperature	5.826	1	0.016
		Acclimation × Temperature	521.357	5	<0.0001
	LLT	Intercept	59.813	1	<0.0001
		Acclimation	37.235	2	<0.0001
		Temperature	128.514	2	<0.0001
		Acclimation × Temperature	107.761	4	<0.0001
<i>C. imicola</i>	ULT	Intercept	4.1	1	0.043
		Acclimation	29.793	2	<0.0001
		Temperature	63.892	2	<0.0001
		Acclimation × Temperature	67.629	4	<0.0001
	LLT	Intercept	167.244	1	<0.0001
		Acclimation	6.978	2	0.031
		Temperature	0.993	1	0.319
		Acclimation × Temperature	425.118	4	<0.0001

difference between the 19°C and 29°C acclimated flies but none between 24°C and 29°C. *Culicoides imicola* showed near-total mortality when exposed to 42°C, with only 5.3% of the 29°C treatment surviving, and 0% of the other treatments. *Culicoides bolitinos* showed similar mortality at 42°C, with 0% survival of 19°C and 29°C. The 24°C-acclimated *C. bolitinos*, however, had a survivorship of 26.13%, with complete mortality documented only when subjected to 44°C (Figure 2).


Discussion

Temperature is a determining factor in insect life history, and thermal tolerance plays a key role in determining insect distribution [24,38,39]. Critical thermal limits are valuable for determining the relative temperature tolerances of insects acclimated to different temperatures, or originating from different latitudinal and altitudinal ranges [40]. Typically critical thermal minima are more variable in response to acclimation than critical thermal maxima, both inter- and intra-specifically [41,42]. Interspecific differences in our study agree with this notion, with the CT_{max} of *C. imicola* and *C. bolitinos* being similar (41.87°C SE ± 0.976, and 41.56°C SE ± 0.828 respectively), whilst the CT_{min} of *C. imicola* (1.61°C SE ± 1.175) is nearly a degree higher than that of *C. bolitinos* (0.65°C SE ± 1.117). For the *Culicoides* species tested in this study, expected intra-specific variation was found only for *C. imicola*, with acclimation at temperatures from 19 to 29°C leading to a change in CT_{min} of 2.10°C, but only 1.63°C for CT_{max} . In the case of *C. bolitinos*, while acclimation had a statistically significant effect on both CT_{min} and CT_{max} , the change in CT_{min} (1.37°C)

was similar to that of CT_{max} (1.30°C). For *C. imicola*, CT_{min} changed in a stepwise manner, with approximately 1°C difference between acclimation temperatures. *Culicoides bolitinos* showed little variation between the two lower acclimation temperatures (<0.5°C), but the 29°C-acclimated midges showed a significant increase in CT_{min} , possibly because 24°C still falls within the optimal performance breadth for *C. bolitinos*, whereas 29°C might be considered an 'extreme' temperature that reduces performance (see Deere & Chown [43]).

Little difference in the upper lethal temperatures was found between *C. imicola* and *C. bolitinos*, highlighting limited variability in ULT. Similar to critical temperatures, it has previously been found that variability in lethal limits is greater at a family or generic level, with limited variability between species [44]. Upper thermal limits are also less pliable than lower limits, among species, populations, and over time (i.e. plasticity responses) [26,42]. Our study corroborates this, with little difference in survival of *C. imicola* and *C. bolitinos* at high temperatures. However, our study also points to a dissimilar response of different species to experimental conditions, and possibly environmental variation, as suggested by Chown et al. [37]. *Culicoides imicola* had a clear response to acclimation, with prior exposure to high temperatures improving heat tolerance in a stepwise manner. This was not the case with *C. bolitinos* however, as individuals acclimated to 24°C outperformed both those acclimated at 19°C and 29°C. The relatively poor performance of *C. bolitinos* when exposed to high temperatures after acclimation at 29°C contrasts to the idea of the beneficial acclimation hypothesis, and may indicate a deleterious acclimation response. The deleterious acclimation hypothesis predicts substantial negative performance implications for individuals held at extreme temperatures, both relative to those at median temperatures and those that typically do not experience such extremes in the field [43]. Indeed, although not formally tested, *C. bolitinos* acclimated at 29°C exhibited higher mortality in holding cages, with no individuals surviving past a week in the laboratory, whereas conspecifics acclimated at 19°C and 24°C were capable of surviving in the laboratory for at least two weeks. Determining the degree to which stress proteins and other thermal protectants are up-regulated would provide clarification as to whether a temperature of 29°C is indeed detrimental [40], as would longevity and other performance assays.

Acclimation had no significant effect on the LLT of *C. imicola*, with LLT_{50} values similar for all acclimation temperatures (c. -4°C), and total mortality at -6°C. In contrast, acclimation had a significant effect on *C. bolitinos*, with the LLT_{50} for the 24°C-acclimated individuals reaching -6.01°C, but no significant difference between 19°C and 29°C was found. The deleterious acclimation hypothesis might once again explain the observed pattern, as



would a mismatch between optimum temperatures and realised habitat. Another possibility may be the uncontrolled age and physiological state of the field-collected insects. To obtain sufficient numbers for all experiments it was necessary to collect additional individuals from the field by trapping on several occasions. Senescence has been suggested to reduce stress tolerance in insects [41]. This is supported by Luckinbill [44], who found longevity selection improved low temperature tolerance in *Drosophila melanogaster* [41]. Adult *C. bolitinos*, however, have an average life expectancy of approximately 20 days, but may

survive up to 90 days [15]. Similarly, *C. imicola* adults can survive for more than 15 days and *C. pycnostictus* for 54 days at temperatures of -1.5°C [17]. Given that our collection methods were consistent and that the period of time that elapsed before testing was an equivalent proportion of the life expectancy of both species, we believe that our data are a true representation of the ability of *C. imicola* and *C. bolitinos* to withstand temperature extremes.

This study indicates that lower temperature limits may play a role in constraining the distribution of *C. imicola*. Modelling attempts often rely on a mean temperature

threshold of 12.5°C to predict presence or absence of *C. imicola* [29,45] as established by Purse et al. [46]. Peters et al. [45] on the other hand, found absence indicated by a mean temperature of 14.4°C and presence by a mean of 15.7°C. For the two years of weather data we analysed, Clarens had a mean annual temperature of 14.51°C, two degrees above the suggested thermal intercept suggested by Purse, but very close to that of Peters and colleagues. Furthermore, the suggested mean minimum temperature thresholds of Peters et al. (Absence: 9.4°C; Presence: 10.0°C) coincide strongly with the mean minimum temperatures of our study sites (Clarens: 8.67°C; Onderstepoort: 10.74°C). However, we propose that mean temperatures are only an indication of potential distribution, and it is extreme minimum temperatures reached that determine population persistence of *C. imicola*. The lowest temperature recorded for Clarens in 2012 was -8°C (Figure 3), with temperatures regularly below 0°C and often dropping below the LLT of adult *C. imicola* in winter. In tandem with a relatively high CT_{min} , it seems likely that the thermal environment in the areas surrounding Clarens may prevent adult *C. imicola* from surviving or establishing in any great abundance, concurring with the negative correlation found between *C. imicola* abundance and low winter temperatures [47]. In addition, it is highly likely that the low winter temperatures

of Clarens and similar areas affect larval development and survival because the lower developmental threshold of larvae is ~11.75°C (calculated from [10]). Indeed, Baylis et al. [30] suggested that minimum temperatures determine the ability of the *C. imicola* to successfully overwinter as larvae (for one or two months [47]), as the larvae inhabit the O horizon and are easily affected by ambient temperature in the open habitats they prefer [31]. Furthermore, Legg et al. [48] suggested that median lower developmental thresholds are useful for determining phenology (and thus also distribution potential).

In the case of *C. bolitinos* it appears that the thermal limits of adults are not related to their distribution because their upper temperature tolerance is similar to that of *C. imicola* and they are tolerant of lower temperatures. It may be that the thermal limits of other life stages, especially the larvae, play an important role in limiting the abundance and distribution of *C. bolitinos*. In warmer environments where they are less abundant, the exothermic decomposition of dung could lead to dung temperatures too high for larvae to survive. Evidence for this scenario is currently not available, as when we placed iButtons in fresh dung, mean dung temperatures were isothermic with moderate ambient temperature (c. 14°C), making inference difficult. Non-lethal thermal stress effects could also explain




Figure 3 Daily temperature range for Onderstepoort and Clarens from the end of winter 2011, to end of winter 2013 (1 August 2011 – 31 July 2013). Horizontal lines represent thermal limits of *C. imicola* and *C. bolitinos*: ULT_{50} and CT_{max} of *C. bolitinos* and ULT_{50} of *C. imicola* (red); ULT_{50} of *C. imicola* (orange); CT_{min} of *C. imicola* (purple); LLT_{50} of *C. imicola* (magenta); CT_{min} of *C. bolitinos* (blue); LLT_{50} of *C. bolitinos* (light blue).

their limited numbers in warmer regions. *Scatophaga stercoraria*, another dipteran with coprophagous larvae, displays temperature-mediated variation in egg size as a maternal physiological response [49]. In this study, Blanckenhorn [49] found reduced egg-size and reduced survivorship of offspring even at intermediate temperatures of 24°C. Similar stress effects could suppress populations of *C. bolitinos* in hotter environments. Modelling done on an allopatric, identical sister species *C. brevitarsis* in Australia found that mean maximum temperatures could predict BTV infection (and thus *C. brevitarsis* distribution) with an accuracy of ~75% [50]. Considering the notable similarity of *C. bolitinos* and *C. brevitarsis* in morphology, habitat and habit, it is possible that it could be similarly employed.

Conclusion

It appears that although factors such as precipitation and availability of breeding sites may affect *C. imicola* distribution and abundance, thermal limits play an important role in determining abundance and distribution of this species. Temperatures lower than the CT_{min} of *C. imicola* may prohibit its dominance in high-altitude areas or its expansion into more temperate climes. The presence of *C. imicola* in Clarens, even though at low abundance (<1% of all *Culicoides* spp. captured) [16], indicates that some individuals of this species may utilise protected microclimates to escape severe cold, similar to some European species that overwinter in stables [51]. Another possibility may be that *C. imicola* reinvades these areas annually, as *Culicoides* species have been known to travel long distances by wind-mediated dispersal [52]. The results for *C. bolitinos* are more complex, as adults of this species show a relative wide thermal performance breadth in comparison to *C. imicola*. Further research on the thermal biology of juvenile life stages is required to explain the more temperate distribution of *C. bolitinos*. Due to difficulty in rearing *Culicoides* in the laboratory, this resolution may prove to be logistically challenging. This study shows that thermal limits and tolerance varies between closely related species and that this may play a regulatory role in the presence and abundance of species. A better understanding of thermal phenomena may contribute to the creation of more realistic risk maps and the elucidation of the influence of climate change on the presence and expansion of viral diseases.

Additional file

Additional file 1: Technical drawing of water-jacketed eight-chamber stage, made for use under microscope. Each chamber can accommodate a 0.2 ml micro-centrifuge tube.

Abbreviations

AHSV: African horse sickness virus; BTV: Blue tongue virus; CT_{min} : Critical thermal minimum; CT_{max} : Critical thermal maximum; LLT: Lower lethal temperature; ULT: Upper lethal temperature.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

FAV and CWW conceived and designed the experiment. Experiments and data analyses were performed by FAV. GJV provided access to light traps and provided guidance on *Culicoides* identification and maintenance. All authors contributed to preparation of the manuscript, and read and approved the final manuscript.

Acknowledgements

The water-jacketed eight-chamber stage was kindly designed and built by Wietsche Penny and Schalk Els, Department of Mechanical Engineering, University of Pretoria. Weather data were kindly supplied by the ARC-Institute for Soil, Climate and Water. Light traps and other collection and rearing materials were made available to us by the ARC-Onderstepoort Veterinary Institute. We are grateful to the owners and staff of Koeberg Farm for providing access to and use of their facilities. Solomon Boikanyo and Steven Jones provided invaluable field assistance.

Author details

¹Flies of Economic Significance Research Group, Department of Zoology and Entomology, University of Pretoria, Private Bag X20, Hatfield 0028, South Africa. ²ARC-Onderstepoort Veterinary Institute, Private Bag X5, Onderstepoort 0110, South Africa.

Received: 24 June 2014 Accepted: 8 August 2014

Published: 20 August 2014

References

1. Borkent A: **The Biting Midges, the Ceratopogonidae (Diptera)**. In *Biology of Disease Vectors*. 2nd edition. Edited by Marquardt WC. San Diego: Elsevier Academic Press; 2005.
2. Meiswinkel R, Venter G, Nevill E, Coetzer J, Tustin R: *Vectors: Culicoides spp.*, Infectious Diseases of Livestock, Volume One. 2nd edition. Cape Town: Oxford University Press; 2004:93–136.
3. Rodriguez M, Hooghuis H, Castano M: **African horse sickness in Spain**. *Vet Microbiol* 1992, **33**(1):129–142.
4. Hoffmann B, Scheuch M, Höper D, Jungblut R, Holsteg M, Schirmeier H, Eschbaumer M, Goller KV, Wernike K, Fischer M: **Novel orthobunyavirus in Cattle, Europe, 2011**. *Emerg Infect Dis* 2012, **18**(3):469–472.
5. Van den Brom R, Lutikholt S, Lievaart-Peterson K, Peperkamp N, Mars M, van der Poel W, Vellema P: **Epizootie uitbraak van congenital afwijkingen bij Schmallenbergvirus**. *Tijdschr Diergeneeskd* 2012, **137**:106–111.
6. de Vos C, Hoek C, Nodelijk G: **Risk of introducing African horse sickness virus into the Netherlands by international equine movements**. *Prev Vet Med* 2012, **106**(2):108–122.
7. Purse BV, Mellor PS, Rogers DJ, Samuel AR, Mertens PP, Baylis M: **Climate change and the recent emergence of bluetongue in Europe**. *Nat Rev Microbiol* 2005, **3**(2):171–181.
8. Sinclair M, Buhrmann G, Gummow B: **An epidemiological investigation of the African horse sickness outbreak in the Western Cape Province of South Africa in 2004 and its relevance to the current equine export protocol**. *J S Afr Vet Assoc* 2006, **77**(4):191–196.
9. Tilibaşa E, Dărăbuş G: **Preliminary studies on dynamics of Culicoides spp. in western Romania in conjunction with some environmental factors**. *Parasite Vector* 2014, **7**(Suppl 1):O7.
10. Veronesi E, Venter G, Labuschagne K, Mellor P, Carpenter S: **Life-history parameters of culicoides (avaritia) imicola kieffer in the laboratory at different rearing temperatures**. *Vet Parasitol* 2009, **163**(4):370–373.
11. Meiswinkel R: **Afrotropical Culicoides: a redescription of C.(avaritia) imicola kieffer, 1913 (diptera: ceratopogonidae) with description of the closely allied C.(A.) bolitinos sp. nov. reared from the dung of the African buffalo, blue wildebeest and cattle in South Africa**. *Onderstepoort J Vet* 1989, **56**(1):23–39.
12. Nevill E, Venter G, Edwardes M: **Potential Culicoides vectors of livestock orbiviruses in South Africa**. In *Bluetongue, African horse sickness, and related orbiviruses: Proceedings of the Second International Symposium: 1992*. CRC Press, Inc; 1992:306–314.
13. Paweska J, Venter G: **Vector competence of Culicoides species and the seroprevalence of homologous neutralizing antibody in horses for six**

- serotypes of equine encephalosis virus (EEV) in South Africa. *Med Vet Entomol* 2004, **18**(4):398–407.
14. Mellor P, Walton T, Osburn B: **Culicoides as Potential Orbivirus Vectors in Europe.** In *Bluetongue, African horse sickness, and related orbiviruses: Proceedings of the Second International Symposium: 1992*. CRC Press, Inc; 1992:278–283.
 15. Mellor P, Boorman J, Baylis M: **Culicoides biting midges: their role as arbovirus vectors.** *Annu Rev Entomol* 2000, **45**(1):307–340.
 16. Meiswinkel R, Paweska J: **Evidence for a new field Culicoides vector of African horse sickness in South Africa.** *Prev Vet Med* 2003, **60**(3):243–253.
 17. Nevill E: **Cattle and Culicoides biting midges as possible overwintering hosts of bluetongue virus.** *Onderstepoort J Vet* 1971, **38**(2):65.
 18. Venter G, Meiswinkel R: **The Virtual Absence of Culicoides Imicola (Diptera: Ceratopogonidae) in a Light-Trap Survey of the Colder, High-Lying Area of the Eastern Orange Free State, South Africa, and Implications for the Transmission of Arboviruses.** *Onderstepoort J Vet* 1994, **61**(4):327–340.
 19. Nevill E, Venter G, Edwardes M, Pajor I, Meiswinkel R, Van Gas J: **Culicoides species associated with livestock in the Stellenbosch area of the Western Cape Province, Republic of South Africa (Diptera: Ceratopogonidae).** *Onderstepoort J Vet* 1988, **55**(2):101–106.
 20. Venter G, Nevill E, Van der Linde T: **Geographical distribution and relative abundance of stock-associated culicoides species (diptera: ceratopogonidae) in southern Africa in relation to their potential as viral vectors.** *Onderstepoort J Vet* 1996, **63**(1):25.
 21. Meiswinkel R: **Discovery of a culicoides imicola-free zone in South Africa: preliminary notes and potential significance.** *Onderstepoort J Vet* 1997, **64**:43.
 22. Meiswinkel R, Labuschagne K, Baylis M, Mellor P: **Multiple vectors and their differing ecologies: observations on two bluetongue and African horse sickness vector culicoides species in South Africa.** *Veter Ital* 2004, **40**(3):297.
 23. Kirkeby C, Bødker R, Stockmarr A, Lind P: **Spatial abundance and clustering of culicoides (diptera: ceratopogonidae) on a local scale.** *Parasite Vector* 2013, **6**(43):1–14.
 24. Chown SL, Nicolson S: *Insect Physiological Ecology: Mechanisms and Patterns*. Oxford: Oxford University Press; 2004.
 25. Terblanche JS, Marais E, Chown SL: **Stage-related variation in rapid cold hardening as a test of the environmental predictability hypothesis.** *J Insect Physiol* 2007, **53**(5):455–462.
 26. Lyons CL, Coetzee M, Terblanche JS, Chown SL: **Thermal limits of wild and laboratory strains of two African malaria vector species, anopheles arabiensis and anopheles funestus.** *Malar J* 2012, **11**(226):10.1186.
 27. Kearney M, Porter W: **Mechanistic niche modelling: combining physiological and spatial data to predict species' ranges.** *Ecol Lett* 2009, **12**(4):334–350.
 28. Racz V, Venter G, Griot C, Stärk K: **Estimating the temporal and spatial risk of bluetongue related to the incursion of infected vectors into Switzerland.** *BMC Vet Res* 2008, **4**(1):42.
 29. Wittmann E, Mellor P, Baylis M: **Using climate data to map the potential distribution of culicoides imicola (diptera: ceratopogonidae) in Europe.** *Rev Sci Tech OIE* 2001, **20**(3):731–736.
 30. Baylis M, Meiswinkel R, Venter G: **A preliminary attempt to use climate data and satellite imagery to model the abundance and distribution of culicoides imicola (diptera: ceratopogonidae) in southern Africa.** *J S Afr Vet Assoc* 1999, **70**(2):80–89. (2), pp. 80–89.
 31. Conte A, Goffredo M, Ippoliti C, Meiswinkel R: **Influence of biotic and abiotic factors on the distribution and abundance of culicoides imicola and the obsoletus complex in Italy.** *Vet Parasitol* 2007, **150**(4):333–344.
 32. Brugger K, Rubel F: **Characterizing the species composition of European culicoides vectors by means of the köppen-geiger climate classification.** *Parasite Vector* 2013, **6**:333.
 33. Paweska J, Venter G, Mellor P: **Vector competence of South African culicoides species for bluetongue virus serotype 1 (BTV-1) with special reference to the effect of temperature on the rate of virus replication in C. Imicola and C. Bolitinos.** *Med Vet Entomol* 2002, **16**(1):10–21.
 34. Carpenter S, Wilson A, Barber J, Veronesi E, Mellor P, Venter G, Gubbins S: **Temperature dependence of the extrinsic incubation period of orbiviruses in culicoides biting midges.** *PLoS One* 2011, **6**(11):e27987.
 35. Venter G, Paweska J, Van Dijk A, Mellor P, Tabachnick W: **Vector competence of culicoides bolitinos and C. Imicola for South African bluetongue virus serotypes 1, 3 and 4.** *Med Vet Entomol* 1998, **12**(4):378–385.
 36. Terblanche JS, Clusella-Trullas S, Deere JA, Chown SL: **Thermal tolerance in a south-east African population of the tsetse fly glossina pallidipes (diptera, glossinidae): implications for forecasting climate change impacts.** *J Insect Physiol* 2008, **54**:114–127.
 37. Chown SL, Jumbam KR, Sørensen JG, Terblanche JS: **Phenotypic variance, plasticity and heritability estimates of critical thermal limits depend on methodological context.** *Funct Ecol* 2009, **23**(1):133–140.
 38. Addo-Bediako A, Chown S, Gaston K: **Thermal tolerance, climatic variability and latitude.** *P R Soc London* 2000, **267**(1445):739–745.
 39. Mitchell KA, Hoffmann AA: **Thermal ramping rate influences evolutionary potential and species differences for upper thermal limits in Drosophila.** *Funct Ecol* 2010, **24**(3):694–700.
 40. Bowler K, Terblanche JS: **Insect thermal tolerance: what is the role of ontogeny, ageing and senescence?** *Biol Rev* 2008, **83**(3):339–355.
 41. Goto S: **Expression of drosophila homologue of senescence marker protein-30 during cold acclimation.** *J Insect Physiol* 2000, **46**(7):1111–1120.
 42. Weldon C, Terblanche J, Chown S: **Time-course for attainment and reversal of acclimation to constant temperature in two ceratitis species.** *J Therm Biol* 2011, **36**:479–485.
 43. Deere JA, Chown SL: **Testing the beneficial acclimation hypothesis and its alternatives for locomotor performance.** *Am Nat* 2006, **168**(5):630–644.
 44. Luckinbill L: **Selection for longevity confers resistance to low-temperature stress in drosophila melanogaster.** *J Gerontol A-Biol* 1998, **53**(2):B147–B153.
 45. Peters J, De Baets B, Calvete C, Lucientes J, De Clercq E, Ducheyne E, Verhoest N: **Absence reduction in entomological surveillance data to improve niche-based distribution models for Culicoides imicola.** *Prev Vet Med* 2011, **100**(1):15–28.
 46. Purse BV, McCormick BJ, Mellor PS, Baylis M, Boorman J, Borrás D, Burgu I, Capela R, Caracappa S, Collantes F: **Incriminating bluetongue virus vectors with climate envelope models.** *J App Ecol* 2007, **44**(6):1231–1242.
 47. De Liberato C, Purse BV, Goffredo M, Scholl F, Scaramozzino P: **Geographical and seasonal distribution of the bluetongue virus vector, Culicoides imicola, in central Italy.** *Med Vet Entomol* 2003, **17**(4):388–394.
 48. Legg D, Van Vleet S, Lloyd J: **Simulated predictions of insect phenological events made by using mean and median functional lower developmental thresholds.** *J Econ Entomol* 2000, **93**(3):658–661.
 49. Blanckenhorn WU: **Temperature effects on egg size and their fitness consequences in the yellow dung fly Scathophaga stercoraria.** *Evol Ecol* 2000, **14**(7):627–643.
 50. Ward M: **The use of discriminant analysis in predicting the distribution of bluetongue virus in Queensland, Australia.** *Vet Res Comm* 1994, **18**(1):63–72.
 51. Mehlhorn H, Walldorf V, Klimpel S, Schmahl G, Al-Quraishy S, Walldorf U, Mehlhorn B, Bätza H-J: **Entomological survey on vectors of Bluetongue virus in Northrhine-Westfalia (Germany) during 2007 and 2008.** *Parasitol Res* 2009, **105**(2):321–329.
 52. Bishop A, Barchia I, Spohr L: **Models for the dispersal in Australia of the arbovirus vector, Culicoides brevitarsis Kieffer (Diptera: Ceratopogonidae).** *Prev Vet Med* 2000, **47**(4):243–254.

doi:10.1186/1756-3305-7-384

Cite this article as: Verhoef et al.: Thermal limits of two biting midges, *Culicoides imicola* Kieffer and *C. bolitinos* Meiswinkel (Diptera: Ceratopogonidae). *Parasites & Vectors* 2014 **7**:384.