

## Article

# Excess Folic Acid Supplementation before and during Pregnancy and Lactation Alters Behaviors and Brain Gene Expression in Female Mouse Offspring

Xingyue Yang <sup>1,2,†</sup>, Wenyan Sun <sup>3,†</sup>, Qian Wu <sup>1</sup>, Hongyan Lin <sup>2</sup>, Zhixing Lu <sup>3</sup>, Xin Shen <sup>1</sup>, Yongqi Chen <sup>3</sup>, Yan Zhou <sup>3</sup>, Li Huang <sup>1</sup>, Feng Wu <sup>2</sup>, Fei Liu <sup>4,\*</sup> and Dandan Chu <sup>1,\*</sup> 

- <sup>1</sup> Key Laboratory of Neuroregeneration of Jiangsu and Ministry of Education, NMPA Key Laboratory for Research and Evaluation of Tissue Engineering Technology Products, Co-Innovation Center of Neuroregeneration, Nantong University, Nantong 226001, China; yangxingyue618@163.com (X.Y.); wq15706295797@163.com (Q.W.); sx827477223@outlook.com (X.S.); Huangli@ntu.edu.cn (L.H.)
- <sup>2</sup> Department of Pharmacology, School of Pharmacy, Nantong University, Nantong 226001, China; LHY031583@163.com (H.L.); wf619@ntu.edu.cn (F.W.)
- <sup>3</sup> Department of Biochemistry and Molecular Biology, School of Medicine, Nantong University, Nantong 226001, China; sunjiajun51601@163.com (W.S.); Tsukiato0412@gmail.com (Z.L.); cyq1530852896@163.com (Y.C.); ntzhouy@ntu.edu.cn (Y.Z.)
- <sup>4</sup> Department of Neurochemistry, Inge Grundke-Iqbal Research Floor, New York State Institute for Basic Research in Developmental Disabilities, Staten Island, NY 10314, USA
- \* Correspondence: feiliu63@hotmail.com (F.L.); chudd@ntu.edu.cn (D.C.)
- † These authors contributed equally to this work.



**Citation:** Yang, X.; Sun, W.; Wu, Q.; Lin, H.; Lu, Z.; Shen, X.; Chen, Y.; Zhou, Y.; Huang, L.; Wu, F.; et al. Excess Folic Acid Supplementation before and during Pregnancy and Lactation Alters Behaviors and Brain Gene Expression in Female Mouse Offspring. *Nutrients* **2022**, *14*, 66. <https://doi.org/10.3390/nu14010066>

Academic Editor: Marilyn Cornelis

Received: 17 November 2021

Accepted: 21 December 2021

Published: 24 December 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Use of folic acid (FA) during early pregnancy protects against birth defects. However, excess FA has shown gender-specific neurodevelopmental toxicity. Previously, we fed the mice with 2.5 times the recommended amount of FA one week prior to mating and during the pregnancy and lactation periods, and detected the activated expression of *Fos* and related genes in the brains of weaning male offspring, as well as behavioral abnormalities in the adults. Here, we studied whether female offspring were affected by the same dosage of FA. An open field test, three-chamber social approach and social novelty test, an elevated plus-maze, rotarod test and the Morris water maze task were used to evaluate their behaviors. RNA sequencing was performed to identify differentially expressed genes in the brains. Quantitative real time-PCR (qRT-PCR) and Western blots were applied to verify the changes in gene expression. We found increased anxiety and impaired exploratory behavior, motor coordination and spatial memory in FA-exposed females. The brain transcriptome revealed 36 up-regulated and 79 down-regulated genes in their brains at weaning. The increase of *Tlr1*; *Sult1a1*; *Tph2*; *Acacb*; *Etnppl*; *Angptl4* and *Apold1*, as well as a decrease of *Ppara* mRNA were confirmed by qRT-PCR. Among these genes; the mRNA levels of *Etnppl*; *Angptl4* and *Apold1* were increased in the both FA-exposed female and male brains. The elevation of *Sult1a1* protein was confirmed by Western blots. Our data suggest that excess FA alters brain gene expression and behaviors in female offspring, of which certain genes show apparent gender specificity.

**Keywords:** folic acid; pregnancy; lactation; behaviors; gene expression; female

## 1. Introduction

The water-soluble micronutrient folate (vitamin B-9) participates in multiple biological processes in mammals, such as DNA and protein synthesis, as well as nucleic acid and protein methylation [1]. As the fully oxidized synthetic form of natural folate, folic acid (FA) is catalyzed into 5-methyltetrahydrofolate (5-MTHF) in the liver, and finally produces the universal methyl donor S-adenosyl methionine (SAM), utilized in methylation reactions [1]. Appropriate FA supplement (0.4 mg per day) is widely accepted to prevent against neural tube defects (NTD) and other birth defects [2]. Moreover, FA fortification has been introduced in the United States, Canada and several other countries to diminish the number of

NTD [3]. Due to vitamin supplements and mandatory fortification, high concentrations of folate have been detected in maternal serum as well as in breast milk [4–7]. Whether excess FA supplementation during pregnancy and lactation affects the neurodevelopment of offspring has become an important public health issue.

The negative impact of excessive maternal FA intake on infant neurodevelopment has been reported by several groups. Periconceptional use of FA exceeding the maximum tolerated dose ( $\geq 1$  mg/day) was associated with decreased cognitive development in children [8]. FA supplementation during gestation is considered to be related to an increased risk for autism [9]. Extremely high concentrations of maternal plasma FA ( $>60.3$  nmol/L) elevated the risk of autism spectrum disorder by 2.5 times [10,11]. In rodents, female mice supplemented with high dose of FA (40 mg/kg) before and during pregnancy led to embryonic delay and neural tube defects of the offspring [12]. Maternal over-supplementation with FA resulted in behavioral abnormalities in the offspring, such as anxiety-like behavior, hyperactivity, increased ultrasonic vocalizations [13] and seizure susceptibility [14], as well as impaired reversal learning [15]. The gene expression and DNA methylation profiling were altered in the brains of FA-exposed offspring [13,16–18]. Sex-specific influences of excess FA on behavioral defects of the offspring have been reported. Pregnant mice fed with 4 mg/kg diet of FA induced hyperactivity of male but not female offspring [13]. Maternal over-supplementation with FA was associated with attentional dysfunction in children, particularly in boys [19]. However, the roles of excess FA intake during the perigestational and lactational period on the brain transcriptome of weaning female mice and their behaviors after adulthood are not fully understood.

Previously, we fed the mice with approximately 2.5 times the recommended amount of FA one week prior to mating and during pregnancy and the lactation period, and detected behavioral defects in adult male offspring, including increased anxiety, impairments in social preference, and motor and spatial learning [18]. Transcriptome sequencing of the brains revealed that *Fos* and associated genes were upregulated in weaning male offspring. Here we applied the same dosage of FA to the mice and examined whether excess FA altered behaviors and gene expression in the brains of female offspring, and whether the changes are different from those in males. Behavior test showed that early exposure to excessive FA resulted in long-term alterations of behaviors in adult female offspring, such as increased anxiety-like behavior, decreased exploratory activities, motor coordination and spatial memory. RNA sequencing was used to identify differentially expressed genes (DEGs) in the brains of FA-exposed female mice at weaning. Quantitative real time-PCR (qRT-PCR) revealed the increased transcription of *Tlr1*, *Sult1a1*, *Tph2*, *Acacb*, *Etnppl*, *Angptl4* and *Apold1*, and decreased transcription of *Ppara* in FA-exposed female brains. Western blots confirmed the promoted expression of *Sult1a1* protein in FA-exposed brains. These results indicated that excess FA supplementation in pregnancy and lactation altered brain transcriptome in weaning female mice and changes adult behaviors, and that the effects were different from those of males.

## 2. Materials and Methods

### 2.1. FA Supplementation and Animals

The animal experiments were performed according to the National Institutes of Health Guidelines for animal research. ICR mice were purchased from Shanghai, China and maintained in the laboratory animal center of Nantong University. Mice were housed at 3–5 per cage on a 12 h light/12 h dark cycle with free access to food and water. The mice were supplied normal chow diet, containing the routinely recommended supplement of 2 mg FA/kg diet (Xietong Pharmaceutical Biotechnology, Nanjing, China) [20]. Formula of the diet is supplied in Supplementary Table S1. At 8 weeks of age, 4 males and 8 females were randomly divided into two groups and supplied with no (control) or 3.75 mg/L FA (Sigma-Aldrich, St. Louis, MO, USA) in their drinking water for one week [18,21,22]. Then, the mice were allowed to mate naturally by introducing 2 females in the cage of 1 male. FA was administrated throughout the mating, pregnancy and lactation period. A breeding

mouse consumes about 5 g in their diet per day [23], which provides 10 µg FA. An adult mouse consumes an average of 4–8 mL of water daily [24], that is, at least 15 µg of FA from the drinking water. Therefore, the total amount of FA taken by the mice from food and water is at least 2.5 times ( $2.5 \times$  FA) the normal dietary intake [18]. This dosage of FA was used because the total intake of folate from FA fortification and vitamin supplement exceeded twice the recommended dietary allowance (RDA) in the North American populations [25] and the serum levels of total folate increased approximately 2.5-fold after fortification in the U.S. population [6].

After weaning (postnatal day 21, P21d), the pups were supplied with standard chow and water. For RNA sequencing, qRT-PCR and Western blots, the pups were sacrificed at weaning. The cerebrums were quickly dissected, incubated in phosphate-buffered saline (PBS) on ice for 5 min [26], quickly frozen in liquid nitrogen and then stored at  $-80\text{ }^{\circ}\text{C}$  before use. The body weight was recorded at 21 days and 1, 2, 3, 4 and 5 months of age (Con,  $n = 12$ ;  $2.5 \times$  FA,  $n = 9$ ).

## 2.2. Behavioral Analysis

All behavioral experiments were performed between 8 am and 5 pm. Female offspring were collected from 3–4 litters per group (2–5 females/litter). Two-month-old offspring were subjected to the behavioral tests in the following order: open field, three-chamber social approach and social novelty test, elevated plus-maze, rotarod test and Morris water maze task (Con,  $n = 12$ ;  $2.5 \times$  FA,  $n = 9$ ). Except for the rotarod, the behaviors were recorded and analyzed by a video-tracking system (ANYmaze, Stoelting Co., Wood Dale, IL, USA). The apparatuses were wiped with water first, then with 75% ethanol, and were allowed to dry between mice.

### 2.2.1. Open Field Test

The mice were placed in a 48 cm L  $\times$  48 cm W  $\times$  40 cm H black plastic arena (Xinruan, Shanghai, China) and tracked every 2.5 min for 4 consecutive sessions, with a total time of 10 min. A 15  $\times$  15 cm area in the center of the open field was designated as the central zone. The total distance, immobile time, mean speed, time in the central zone, entries into the central zone, and mean distance from the central zone of the mice were recorded.

### 2.2.2. Three-Chamber Social Approach and Social Novelty Test

Sociability and social novelty of the mice were evaluated in a three-chambered box [27] (Ugo Basile, Gemonio, Italy), with each chamber being 20 cm W  $\times$  40 cm L  $\times$  22 cm H. The three chambers were separated by transparent Plexiglas walls with an 8 cm H  $\times$  5 cm W door in the middle. As described [18], the subject mouse was first introduced into the central chamber (Central) and allowed to habituate to the apparatus freely for 5 min. Then, a 2 month-old stranger female (Social) was settled in a wire cage and put in the left chamber. In the right chamber, an identical empty wire cage was placed as the inanimate target (Inanimate). The subject was allowed to explore the whole apparatus for 10 min. Time spent in each chamber was recorded to evaluate the sociability of the mice. In the sociability phase, another 2 month-old stranger female was enclosed in the cage in the right chamber (Novel). The subject was then allowed to explore the whole apparatus for another 10 min. The time spent exploring each chamber was calculated to evaluate the preference for social novelty.

### 2.2.3. Elevated Plus-Maze

The elevated plus-maze apparatus was described previously [18]. The subject was introduced in the central square facing a closed arm, and allowed to explore the elevated plus-maze for 5 min. The time of mice exploring the closed or open arms were recorded to access anxiety-like behavior.

#### 2.2.4. Rotarod

The female offspring were accessed on the rotarod (Ugo Basile, Gemonio, Italy) 3 trails per day for 5 consecutive days. Rotarod steadily accelerated from 4 revolutions per minute (rpm) to 40 rpm in 2 min. One trial lasted for a maximum of 500 s. Between each trial, the mice were allowed to rest for 30 min.

#### 2.2.5. Morris Water Maze Task

Morris water maze tasks were performed as previously described [18]. Briefly, the mice were tested in a black circular pool with a diameter of 180 cm and a height of 60 cm that was theoretically divided into four equal quadrants. The 13 cm diameter escape platform, which was submersed 1 cm below the water surface, was put in the center of one quadrant. In each training trial, the mice explored freely in the maze for up to 90 s. Those who did not reach the escape platform within 90 s were gently led to it by the experimenter. After each trial, the mouse was left on the escape platform for another 20 s. The training was conducted 4 times a day for 4 consecutive days. Between each trial, the interval was more than 30 min. Twenty-four hours after the last training, a probe trial was conducted for 90 s without the escape platform. The swimming speed, the first latency to the platform (target) zone, and the distance traveled until the first entry into the target zone were measured to evaluate spatial memory.

#### 2.3. Transcriptome Sequencing

The brain transcriptome was sequenced by Vazyme Biotech Co., Ltd. (Nanjing, China) and described previously [18]. In brief, the brains of 21-day old female mice ( $n = 4$  per group) were collected from 2 litters (2 mice per litter) with different parents. Total RNA was isolated from the cerebrum using Trizol (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions and cleaned up by RNeasy spin columns (Qiagen, Valencia, CA, USA). The concentration and purity of isolated RNA were determined by a Qubit<sup>®</sup> RNA Assay Kit (Life Technologies, Carlsbad, CA, USA) and NanoDrop<sup>®</sup> spectrophotometers (Thermo Fisher Scientific, Waltham, MA, USA), respectively. The integrity of isolated RNA was determined by a LabChip GX system (Caliper, Newton, MA, USA). The quality score of RNA ranges between 7.6 and 8.4. The sequencing libraries were prepared using VAHTS mRNAseq v2 Library Prep Kit for Illumina R (Vazyme, Nanjing, China) following the manufacturer's recommendations. The sequencing was performed on an Illumina HiSeq<sup>™</sup> X Ten platform (Illumina, San Diego, CA, USA) with 150 bp paired-end reads. Low quality reads and those containing adapter or ploy-N were removed from the raw reads. Then the clean reads were aligned to the reference mouse genome (mm10) using TopHat (version 2.1.1) and assembled using the reference-based assembler Cufflinks (v2.2.1). The fragments per kilobase of transcript sequence per millions base pairs, mapped (FPKM) for coding genes, was calculated by Cuffdiff (v1.3.0). Comparing FPKM values of the control and  $2.5 \times$  FA mice brains, transcripts with a fold change  $> 2$  and an adjusted  $p < 0.05$  were considered as significantly differentially expressed. Gene ontology (GO) enrichment analysis of DEGs was conducted with GO::TermFinder. Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses for the DEGs were performed using the Database for Annotation, Visualization, and Integrated Discovery (DAVID) bioinformatic resources (<https://david.ncifcrf.gov/>, accessed on 25 August 2017).

#### 2.4. qRT-PCR

Cerebral RNA ( $n = 3$ ) was extracted from 2 litters with different parents. HiScript II Q RT SuperMix for qPCR (with gDNA wiper) (Vazyme, Nanjing, China) was used to reverse transcribed equal amount of RNA of each sample to complementary DNA. Real-time PCR was conducted with AceQ qPCR SYBR<sup>®</sup> Green Master Mix (Vazyme, Nanjing, China) on a LightCycler<sup>®</sup> 480 (Roche Diagnostics, Basel, Switzerland).

The amplifying conditions for cDNA were as follows: denaturation at 95 °C for 10 min, 45 cycles of 95 °C for 10 s and 60 °C for 30 s, followed by 95 °C for 10 s, 65 °C for 60 s, 97 °C for 1 s and 37 °C for 30 s. Quantitation of the relative levels of mRNA were carried out by the comparative  $2^{-\Delta\Delta C_t}$  method and normalized to Ribosomal Protein S18 (RPS18) levels (B661301, Sangon Biotech, Shanghai, China). Primer sequences are listed in Table 1.

**Table 1.** Sequences of qRT-PCR primers.

Gene	Forward Sequence	Reverse Sequence	Amplicon Length (bp)
<i>Tlr1</i>	GAGGCATGAAGAGAGCGGAA	TAGGGGTGTCCACAATTGCC	292
<i>Sult1a1</i>	GCTAGATAAGTGTGGCCGGG	TTCGGGCAACGTAGATCACC	194
<i>Tph2</i>	CAATCGAGTTCCGGCCTTTC	CTGCGTGTAGGGGTTGAAGT	275
<i>Acacb</i>	CCGCTCAAGATCGAGGAGTC	ATGCAGGCTACCTTGCTTGT	255
<i>Ppara</i>	TGTGAACTGACGTTTGTGGC	CCACAGAGCGCTAAGCTGT	70
<i>Etnppl</i>	CGGTCATGTGCGAGCTCTAT	GTCGTGGAGGAAGCGAGAAT	263
<i>Angptl4</i>	GGGGACCTTAACTGTGCCAA	CTGCAGAGGATAGTAGCGGC	165
<i>Apold1</i>	CCGTCCTGAAGCCAAGATT	AGAAAAACAACGCTGCGTCC	168

### 2.5. Western Blot

Mice cerebrums ( $n = 6$ ) from 2 litters (3 females per litter) with different parents were homogenized in a 9-fold volume of ice-cold brain lysis buffer (50 mM Tris-HCl, pH 7.4, 2.0 mM EDTA, 8.5% sucrose and 10 mM  $\beta$ -mercaptoethanol) containing a protease inhibitor cocktail (Roche, Basel, Switzerland). Protein concentration of the samples was measured by a Pierce™ 660 nm protein assay kit (Thermo Fisher Scientific). The same levels of protein in each sample were loaded and separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). The protein was then transferred onto polyvinylidene difluoride (PVDF) membrane. The membrane was blocked with 5% skim milk in Tris-buffered saline (TBS, 150 mM NaCl, 50 mM Tris-HCl, pH 7.4) for 30 min and incubated with rabbit anti-Sult1a1 antibody (1:500, bs-6283R, Bioss Antibodies, Beijing, China) overnight at room temperature. Then the membrane was rinsed three times with TBST (TBS containing 0.05% Tween-20) and incubated with peroxidase affinitive goat anti-rabbit IgG (111-035-144) (Jackson ImmunoResearch, West Grove, PA, USA) for 2 hr at room temperature, and then developed with an Enhanced chemiluminescence (ECL) kit (Thermo Fisher Scientific) and exposed to X-ray film (Carestream Health, White City, OR, Canada). The intensity of bands was quantified using Multi Gauge software (Fuji Film, Tokyo, Japan).


### 2.6. Statistical Analysis

Data were presented as mean  $\pm$  standard error of the mean (SEM). The statistical significance was analyzed using GraphPad Prism 8 (GraphPad Software Inc., San Diego, CA, USA) by a two-way analysis of variance (ANOVA) followed by a post hoc analysis using the Bonferroni's multiple comparison test or the unpaired two-tailed Student's  $t$  test.  $p < 0.05$  was considered statistically significant.

## 3. Results

### 3.1. Excess FA Supplementation throughout Pregnancy and Lactation Modifies Behaviors in the Adult Female Offspring

The mice were fed with drinking water dissolved with 3.75 mg/L FA (about 2.5 times the dietary requirement) from one week before mating, and throughout the mating, gestation and lactation periods (Figure 1A). The female offspring were weaned to normal drinking water at postnatal day 21 (P21d). The body weight of female offspring ( $2.5 \times$  FA hereafter) was tracked for 5 months, and there was no difference when compared with the control (Figure 1B, FA  $\rightarrow F_{1,114} = 1.418$ ,  $p = 0.2362$ ; Day  $\rightarrow F_{5,114} = 138.9$ ,  $p < 0.0001$ ; Day  $\times$  FA  $\rightarrow F_{5,114} = 1.058$ ,  $p = 0.3873$ ), suggesting that early-life FA supplementation did not affect the physical growth of the female offspring.




**Figure 1.** Effect of excess FA intake on body weight and the performance in open field test of the female offspring. (A) Schematic of the study design. (B) Control (Con) and 2.5 × FA female mice were weighed from postnatal day 21 to 150. (C–H) In the open field test, the distance traveled by the female offspring in the arena (C), the immobile time (D), the mean speed (E), the time spent in the central square (F), the numbers of entries into the central square (G) and the mean distance from the boundary of the central zone (H) were measured. Con,  $n = 12$ ; 2.5 × FA,  $n = 9$ . \*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$ .

Two-month old 2.5 × FA female mice were subjected to several behavioral tests. First, their locomotor/exploratory activities and anxiety-like behavior were assessed by the open field test. Although the mean speed of 2.5 × FA mice showed no difference from that of the control (Figure 1E,  $p = 0.1465$ ), the overall distance that 2.5 × FA mice traveled in the open field decreased (Figure 1C, Duration  $F_{3,76} = 14.50$ ,  $p < 0.0001$ ; FA  $F_{1,76} = 6.618$ ,  $p = 0.0120$ ; Duration × FA  $F_{3,76} = 0.7745$ ,  $p = 0.5118$ ), and their immobile time increased significantly (Figure 1D, Duration  $F_{3,76} = 37.44$ ,  $p < 0.0001$ ; FA  $F_{1,76} = 21.02$ ,  $p < 0.0001$ ; Duration × FA  $F_{3,76} = 2.121$ ,  $p = 0.1045$ ). The 2.5 × FA mice spent less time in the central zone of the open field (Figure 1F, Duration  $F_{3,76} = 5.928$ ,  $p = 0.0011$ ; FA  $F_{1,76} = 12.23$ ,  $p = 0.0008$ ; Duration × FA  $F_{3,76} = 0.6138$ ,  $p = 0.6081$ ) and entered the central zone less than the control (Figure 1G, Duration  $F_{3,76} = 5.179$ ,  $p = 0.0026$ ; FA  $F_{1,76} = 11.52$ ,  $p = 0.0011$ ; Duration × FA  $F_{3,76} = 0.8008$ ,  $p = 0.4972$ ). Moreover, 2.5 × FA mice were farther away from the boundary of the central zone than the control mice (Figure 1H,  $p = 0.0263$ ). These results showed that excess FA

supplementation impaired exploratory activity and increased anxiety-like behavior of the adult female offspring.


Previously, we reported that excess FA impaired social preference of the adult male offspring in the three-chamber sociability test; that is,  $2.5 \times$  FA males spent less time in the chamber with a social target (a stranger mouse) [18]. Here, we performed the three-chamber social approach and social novelty test on  $2.5 \times$  FA female mice and found no obvious abnormality (Figure 2A,B). When compared with the control,  $2.5 \times$  FA mice spent a similar time in the chamber containing the social target or an inanimate target during the sociability phase (Figure 2A, FA  $F_{1,19} = 1.349$ ,  $p = 0.2598$ ; Chamber  $F_{2,38} = 5.138$ ,  $p = 0.0106$ ; Chamber  $\times$  FA  $F_{2,38} = 0.7914$ ,  $p = 0.4606$ ), and in the chamber with a familiar mouse or a novel target (a new unfamiliar mouse) during the social novelty phase (Figure 2B, FA  $F_{1,19} = 0.0488$ ,  $p = 0.8275$ ; Chamber:  $F_{2,38} = 3.498$ ,  $p = 0.0403$ ; Chamber  $\times$  FA  $F_{2,38} = 0.0569$ ,  $p = 0.9448$ ), indicating that FA did not affect the social performance of the female offspring.



**Figure 2.**  $2.5 \times$  FA did not affect the sociability and anxiety-like behavior of the female offspring. (A,B) In the three-chamber social approach and social novelty test, the time of mice spent in different chambers during the sociability phase (A) or the social novelty phase (B) were recorded. (C,D) In the elevated plus maze, the time of mice spent in the closed or open arms (C) and the number of entries (D) into the closed or open arms were recorded. Con,  $n = 12$ ;  $2.5 \times$  FA,  $n = 9$ .

Next, the mice were assessed for anxiety-like behavior using the elevated plus-maze. The residence time and number of entries of  $2.5 \times$  FA mice in the open arms or in the closed arms was not different from that of the control mice (Figure 2C, FA  $F_{1,38} = 0.1213$ ,  $p = 0.7295$ ; Arm  $F_{1,38} = 33.44$ ,  $p < 0.0001$ ; FA  $\times$  Arm  $F_{1,38} = 0.1059$ ,  $p = 0.7466$ ; Figure 2D, FA  $F_{1,38} = 0.6638$ ,  $p = 0.4203$ ; Arm  $F_{1,38} = 11.86$ ,  $p = 0.0014$ ; FA  $\times$  Arm  $F_{1,38} = 0.0258$ ,  $p = 0.8733$ ), suggesting that  $2.5 \times$  FA supplementation did not induce anxiety in the female offspring in the elevated plus-maze.

Motor learning and coordination of the mice were analyzed by an accelerating rotarod. Although both groups showed an increasing latency to fall from the rotarod with the increase of trials,  $2.5 \times$  FA mice displayed less improvement in their performance (Figure 3A, FA  $F_{1, 285} = 5.679$ ,  $p = 0.0178$ ; Trial  $F_{14, 285} = 5.575$ ,  $p < 0.0001$ ; Trial  $\times$  FA  $F_{14, 285} = 1.121$ ,  $p = 0.3386$ ), implying that excess FA impaired the motor coordination of the female offspring.



**Figure 3.** The motor learning and spatial memory abilities were impaired in  $2.5 \times$  FA mice. (A) Motor learning and coordination of the mice was assessed by rotarod. (B–E) Spatial learning and memory of  $2.5 \times$  FA mice were assessed by Morris water maze test. (B) During the training days, the time it took the mice to reach the escape platform (latency) was recorded. Twenty-four hours after the last training day, the probe trials were performed. Swimming speed of the mice (C), the first latency (D) and the travel distance to reach the escape platform (target) (E) were recorded. Con,  $n = 12$ ;  $2.5 \times$  FA,  $n = 9$ . \*,  $p < 0.05$ ; \*\*,  $p < 0.01$ .


We used the Morris water maze to assess the spatial learning and memory of  $2.5 \times$  FA female mice. During the training days,  $2.5 \times$  FA and control mice spent similar time before finding the escape platform (Figure 3B, FA  $F_{1, 76} = 0.0268$ ,  $p = 0.8703$ ; Day  $F_{3, 76} = 8.384$ ,  $p < 0.0001$ ; Day  $\times$  FA  $F_{3, 76} = 0.4246$ ,  $p = 0.7359$ ). In the probe trial, no significant difference was found in the swimming speed between  $2.5 \times$  FA and control mice (Figure 3C,  $p = 0.0652$ ). However,  $2.5 \times$  FA mice took more time (Figure 3D,  $p = 0.0306$ ) and traveled a longer distance (Figure 3E,  $p = 0.0052$ ) to reach the target zone, displaying deficiency of spatial memory.

Taken together, these data revealed that excess FA supplementation during pregnancy and lactation period increased anxiety-like behavior and reduced the exploration, motor coordination and spatial memory of the adult female offspring, without affecting their general growth, athletic ability, sociability and preference for social novelty.



### 3.2. Excess FA Alters Brain Transcriptome of the Female Offspring at Weaning

We previously identified 176 DEGs in the brains of FA-exposed male mice at P21d using RNA sequencing (103 up-regulated and 73 down-regulated) [18]. In the present study, 2.5 × FA female brains were also collected at P21d and subjected to RNA sequencing. Analysis of the brain transcriptome revealed 115 DEGs in 2.5 × FA mice when compared to the control female brains (36 up-regulated genes and 79 down-regulated genes, Figure 4A,B, Supplementary Table S2). The number of DEGs induced by excess FA exposure in female brain is less than two-thirds of that in male brain, implying that the female brain transcriptome was less sensitive to the same dosage of FA. Thirty nine genes were differentially expressed in both male and female 2.5 × FA brains (16 up-regulated and 23 down-regulated, Figure 4A, Supplementary Table S3). The differential changes of the brain transcriptome might explain distinct behavioral abnormalities of female and male mice upon early-life FA exposure.




**Figure 4.** The transcriptome was altered in the brains of 2.5 × FA mice at P21d. RNA extracted from the cerebrum of control and 2.5 × FA mice ( $n = 4$ ) was analyzed by transcriptome sequencing. (A) Venn diagram indicates the number of DEGs that are unique in each comparison or shared (blue) between groups. (B) The number of up-regulated (orange) or down-regulated (blue) genes in 2.5 × FA male or female mice brain.

### 3.3. Analysis of Pathway and Gene Ontology Enrichment of the DEGs in 2.5 × FA Brains

The KEGG (Kyoto Encyclopedia of genes and Genomes) analysis of the 115 DEGs in the brains of female 2.5 × FA mice displayed five major enrichment pathways involving alcoholism (Pathway ID: ko05034), amphetamine addiction (ko05031), cocaine addiction (ko05030), histidine metabolism (ko00340) and tyrosine metabolism (ko00350) (Figure 5A). Among these pathways, alcoholism, amphetamine addiction and cocaine addiction pathways were also enriched in 2.5 × FA male brains (Table 2).

**Table 2.** Top 5 statistics of KEGG pathway enrichment for DEGs in the brains of 2.5 × FA female and male mice.

	Female	Male
1	Alcoholism	Amphetamine addiction
2	Amphetamine addiction	Dopaminergic synapse
3	Cocaine addiction	Cocaine addiction
4	Histidine metabolism	Alcoholism
5	Tyrosine metabolism	Neuroactive ligand–receptor interaction



**Figure 5.** KEGG and gene ontology (GO) functional enrichment analysis of DEGs in 2.5 × FA female brains. **(A)** Top 5 statistics of KEGG pathway enrichment for the DEGs in 2.5 × FA female brains. **(B)** Top 3 statistics of GO term enrichment for the DEGs in 2.5 × FA female brains.


The DEGs were then categorized according to the gene ontology (GO) analysis. For 2.5 × FA females, the top significantly enriched GO terms in the category of biological process, cell component and molecular function were regulation of RNA metabolic process (GO ID: 0051252), cellular component (GO: 0005575) and nucleic acid binding transcription factor activity (GO: 0001071), respectively (Figure 5B). However, the GO enrichment of DEGs in 2.5 × FA female and male brains showed considerable differences (Table 3).

**Table 3.** Top 5 statistics of GO enrichment for DEGs in the brains of 2.5 × FA female and male mice.

Biological Process	Female	Male
1	Regulation of RNA metabolic process	Cell-cell signaling
2	Regulation of transcription, DNA-templated	Muscle tissue development
3	Regulation of nucleic acid-templated transcription	Feeding behavior
4	Phenol-containing compound metabolic process	Positive regulation of amine transport
5	Phenol-containing compound biosynthetic process	Positive regulation of anion transport
Cellular component	Female	Male
1	Cellular_component	Plasma membrane part
2	Cytosol	Cell projection
3	Neuron projection	Neuron projection
4	Cytoplasmic membrane-bounded vesicle	Neuron part
5	Synapse part	Integral component of plasma membrane
Molecular function	Female	Male
1	Sequence-specific DNA binding transcription factor activity	Receptor binding
2	Nucleic acid binding transcription factor activity	Enzyme inhibitor activity
3	Organic acid binding	Hormone activity
4	Carboxylic acid binding	Drug binding
5	Amino acid binding	Neuropeptide hormone activity


### 3.4. The Expression of Genes Enriched in the GO Category of Cellular Component Were Changed in 2.5 × FA Brains at Weaning

Since most DEGs were enriched in the GO category of cellular component (Figure 5B), we further validated the transcription levels of these genes by qPCR and found that the mRNA levels of *Tlr1* (*toll like receptor 1*) ( $p = 0.0138$ ), *Sult1a1* (*sulfotransferase family 1A, phenol-preferring, member 1*) ( $p = 0.0052$ ), *Tph2* (*tryptophan hydroxylase 2*) ( $p = 0.0311$ ), *Acacb* (*acetyl-CoA carboxylase beta*) ( $p = 0.0246$ ), *Etnppl* (*ethanolamine-phosphate phospho-lyase*) ( $p = 0.0017$ ), *Angptl4* (*angiopoietin like 4*) ( $p < 0.0001$ ) and *Apold1* (*apolipoprotein L domain containing 1*) ( $p = 0.0039$ ) were significantly increased, while the expression of *Ppara* (*peroxisome proliferator activated receptor alpha*) was down-regulated ( $p = 0.0370$ ) in 2.5 × FA brains (Figure 6).



**Figure 6.** DEGs enriched in the GO category of cellular component were validated by qRT-PCR. The relative mRNA expression level of *Tlr1*, *Sult1a1*, *Tph2*, *Acacb*, *Ppara*, *Etnppl*, *Angptl4* and *Apold1* genes in control and 2.5 × FA brains ( $n = 3$ ) were determined by qRT-PCR. The levels of ribosomal protein S18 (RPS18) mRNA were used for normalization. \*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$ .

The protein expression of 2.5 × FA brains was then analyzed using Western blots. The protein levels of *Sult1a1* in 2.5 × FA brains increased by more than 1.5-fold (Figure 7,  $p = 0.0003$ ). These data revealed that excess FA supplement altered the brain transcriptome in female offspring at weaning.



**Figure 7.** The protein levels of *Sult1a1* were elevated in 2.5 × FA brains at P21d. (A) The brains of control and 2.5 × FA mice were lysed and analyzed by Western blots, comparing anti-*Sult1a1* levels ( $n = 6$ ). GAPDH was used as a loading control. (B) The protein levels of *Sult1a1* were calculated by normalizing to GAPDH. \*\*\*,  $p < 0.001$ .

#### 4. Discussion

Moderate FA intake from food fortification and vitamin supplementation has been accepted to protect against neural tube defects and other birth defects [2]. However, there is also evidence indicating that excess FA supplementation has side effects on neurodevelopment [19,28–31]. We have reported that excess FA supplement before and throughout the pregnancy and lactation periods leads to long-term weight gain and abnormal behaviors in adult male offspring, such as deficits in social preference, anxiety-related behavior and abnormal motor and spatial learning abilities [18]. Here, we examined the influence of the same dosage of FA on the behaviors of female offspring and found distinct responses (summarized in Table 4). Unlike the results in males, 2.5 × FA neither affected the general growth nor caused defects in social behavior and performance in the elevated plus-maze of the adult female offspring (Figures 1B and 2). In the accelerating rotarod test, the motor learning of males has been shown to lag behind from the fourth trial [18], while the difference between 2.5 × FA and control females is far less dramatic (Figure 3A). Nevertheless, the anxiety-like behavior and decreased exploration in open field (Figure 1C–H), and the impaired spatial memory in the water maze (Figure 3C–E) were specifically identified in females but not in males. These results showed that excessive FA during pregnancy and lactation had adverse effects on the behaviors of adult female offspring, but the specific effects were different from those in males.

**Table 4.** Summary of body weight changes and behavioral results of 2.5 × FA female and male mice.

	2.5 × FA vs. Control	
	Female	Male [18]
Body weight	NS <sup>a</sup>	Increased
Open field	Decreased exploration Increased anxiety	NS <sup>a</sup>
Three-chamber sociability	NS <sup>a</sup>	Decreased sociability
Three-chamber social novelty	NS <sup>a</sup>	NS <sup>a</sup>
Elevated plus-maze	NS <sup>a</sup>	Increased anxiety
Accelerating rotarod	Impaired motor learning	Impaired motor learning
Morris water maze	Impaired spatial memory	Delay in spatial learning

<sup>a</sup> No significant difference was identified.

As the fully oxidized form of folate, FA is reduced to tetrahydrofolate by the two-step catalysis of dihydrofolate reductase (DHFR) *in vivo*, and then to 5-MTHF through methylenetetrahydrofolate reductase (MTHFR). The biologically active 5-MTHF transfers a methyl group to homocysteine (Hcy) to form methionine. The latter is further converted to SAM, the universal methyl group donor. After participating in various methylation reactions, SAM is converted to S-adenosyl-homocysteine (SAH) [32]. Excess FA intake leads to the high levels of unmetabolized FA and total folate in maternal and fetal circulation [33] as well as in breast milk [4]. Furthermore, FA supplementation has been shown to elevate the SAM/SAH ratio in human neuroblastoma SH-SY5Y cells [34] and human plasma [35]. At present, we cannot distinguish whether the abnormal behaviors and brain gene expression of the mice offspring are caused by unmetabolized FA or excessive methyl donors, or both. If the levels of unmetabolized FA, reduced folates, SAM, SAH and the SAM/SAH ratio could be measured in the blood and brains of 2.5 × FA mice, it may help to better understand this issue.

The gender-specific influences of excess FA on behavioral abnormalities have been reported by other groups. Pregnant mice fed with 4 mg/kg diet of FA induced hyperactivity of male but not female offspring [13]. Maternal over-supplementation with FA has been associated with attentional dysfunction in children, particularly in boys [19]. The potential mechanism for the different behavioral responses between genders might be the sexual variation in FA metabolism. For example, the expression of MTHFR, a key enzyme that generates 5-MTHF for Hcy remethylation in the folate and Hcy metabolic pathways, was

higher in male liver than in the female liver. 10 mg FA/kg diet (5 times the recommended amount) increased the ratio of phosphorylated MTHFR and decreased glycerophosphocholine and sphingomyelin in the brains of male but not female offspring [29]. Whether the expression and activity of one-carbon metabolism enzymes, such as DHFR, MTHFR, serine hydroxymethyltransferase (SHMT) and methionine synthetase (MS), are different between males and females in the liver and/or the brain at various embryonic and postnatal ages, and how the enzymes respond to excess FA intake, are worthy of further investigation.

Excess maternal FA has been shown to modulate gene expression levels and DNA methylation profiling in the cerebellum, depending on the sex of mouse offspring [13,17,36]. In the present study, the transcriptome of FA-exposed mouse brain also exhibited gender-related differences (Tables 2 and 3). The male and female differences of brain transcriptome alternations were mainly concentrated in two areas: gene number and interrupted processes. Firstly, when compared with males, the total number of DEGs in the brain of weaning  $2.5 \times$  FA female mice decreased by one-third, and the number of up-regulated genes decreased by two-thirds (Figure 4A,B). Secondly, the DEGs in males were mostly enriched in the GO biological process category-“Cell-cell signaling”; while in females, the largest number of genes were concentrated in the category “regulation of RNA metabolic process” (Figure 5B, Table 3), which contains various zinc finger proteins. Unlike in males, there were no significant changes in the expression of *Fos* and associated genes [18] in  $2.5 \times$  FA female brains. Instead, several genes in the GO cell component category-“cellular component” were dysregulated in female brains. In general, the differences in the number and categories of DEGs caused by  $2.5 \times$  FA in female and male brains may contribute to distinct behavioral responses between genders. Whether gender and/or gene polymorphism-induced differences in FA metabolism [37,38] also lead to variant neurodevelopmental responses upon excess early-life FA exposure in humans is worth further investigation.

Transcriptome analysis of the whole brain in this study provided a general view of gene expression changes caused by excess FA. Nevertheless, it is also necessary to focus on specific brain regions that are relevant to FA-induced behavioral abnormalities. The performance of mice in the open field is mainly related to brain regions such as the amygdala [39], hypothalamus [40] and prefrontal cortex [41,42]. The cerebellum plays a role in sensorimotor functions, balance control as well as spatial memory and other cognitive functions [43]. The hippocampus functions in spatial navigation, the forming, storing and retrieval of episodic memory and the processing of many other types of memory [44]. Further studies on the gene expression in these specific brain regions could provide more mechanisms for the changes in mouse behavior caused by excess FA.

The transcription of *Sult1a1*, *Tlr1*, *Tph2*, *Ppara* and *Acacb* gene were exclusively increased in  $2.5 \times$  FA female brains. We detected the elevated expression of the *Sult1a1* protein, but no significant changes in *Tlr1* and *Tph2* proteins (data not shown), which may be due to the limited sensitivity of Western blots and/or the spatially restricted expression of these proteins. *Sult1a1* catalyzes the sulfonation of a wide range of endogenous metabolites, drugs and xenobiotics, such as neurotransmitters, iodothyronines, catechins, estrogens and hydroxylated aromatic amines [45,46]. It is highly expressed in liver, kidney and heart of C57BL/6 mice and the expression in females is significantly higher than that in males [47]. *Sult1a1* expression is also detected in wild-type mice brains in both neurons and glial cells [48] without apparent sex differences [47]. *Sult1a1* has been reported to regulate neurotransmitter metabolism through sulfonating monoamines (e.g., dopamine) in vivo [49–51]. *Tph2* is the rate-limiting enzyme of 5-hydroxytryptamine synthesis in the brain. In most psychiatric diseases, the neurotransmitter serotonin shows an increase in the brain and is considered to be a major therapeutic target, implying that activation of TPH2 would be associated with the pathogenesis of human psychiatric disorders [52]. Accordingly, we speculate that excess FA-induced overexpression of *Sult1a1* and/or *Tph2* might disturb neurotransmitter metabolism, and therefore contribute to neurodevelopmental disorders.

Toll like receptors (TLRs), which localize on the cell surface or in intracellular vesicular compartments, play an essential role in innate immune responses against pathogens. TLR1 forms heterodimers with TLR2 and is activated in response to bacterial infections, during embryonic brain development or in neuropathogenesis [53–55]. Postnatal activation of TLR1/2 but not TLR2/6 heterodimer impairs the performance of adult mice in the Morris water maze, suggesting that early activation of TLR1/2 confers long-term consequences on hippocampus-dependent spatial learning and memory [54]. PPAR $\alpha$  is a metabolic regulator involved in lipid metabolism. Knockout of *Ppara* leads to schizophrenia-like phenotypes, including repetitive behavior and lowered prepulse inhibition, and impaired synaptogenesis in the cortex of mice [56]. Activation of TLR1/2 by the agonist peptidoglycan suppressed mRNA, protein and activity of PPAR $\alpha$  [57]. Whether excess early-life FA exposure induced by the behavioral deficiencies in adult female mice relies on the Tlr1-dependent inhibition of PPAR $\alpha$  remains to be further elucidated. Besides, it has been reported that PPAR $\alpha$  mRNA levels are significantly decreased in the liver of rat pups fed vitamin B12-deficient but not FA-deficient diets [58]. Here, we showed the expression of *Ppara* was down-regulated in  $2.5 \times$  FA brains. Thus, it is worth exploring whether the expression of *Ppara* in the liver of  $2.5 \times$  FA female changes, whether the change is consistent with that found in the brain and the potential mechanisms.

Among the DEGs induced by excess FA intake, the mRNA levels of *Etnnpl*, *Angptl4* and *Apold1* were significantly decreased in both male and female  $2.5 \times$  FA mice brains (Figure 6, Supplementary Table S3) [18]. *Etnnpl*, a gene specifically expressed in astrocytes [59], is involved in the catabolism of phosphoethanolamine, which is implicated in membrane synthesis [60]. *Apold1* is mainly expressed in endothelial cells [61] and functions in lipid binding, transportation and localization [62]. *Angptl4* is usually highly expressed in adipose tissue and is known as a fasting-induced inhibitor of lipoprotein lipase [63]. In the brain, *Angptl4* is secreted by astrocytes under hypoxia and induces brain endothelial cell migration and promotes angiogenesis [64]. These studies suggest that  $2.5 \times$  FA might affect lipid metabolism and angiogenesis in the brains of both female and male offspring.

## 5. Conclusions

In conclusion, we found that  $2.5 \times$  FA supplementation before and during pregnancy and lactation altered the brain transcriptome of weaning female offspring and impaired the exploratory behavior, motor coordination and spatial memory of female offspring in adulthood. Transcriptome sequencing indicated that excess early-life FA exposure may influence neurodevelopment in multiple aspects, such as neurotransmitter metabolism, immune/inflammation, lipid metabolism and angiogenesis, and involve a variety of cell types including neuron, glial cell and endothelial cell. Therefore, how excess FA mediates the differential gene expression in the brain, how FA-triggered alterations of infant gene expression contribute to long-term influences on the behaviors of adult offspring and whether excess FA supplementation throughout pregnancy and lactation has similar effects on human behaviors deserve more intensive investigation.

**Supplementary Materials:** The following are available online at <https://www.mdpi.com/article/10.3390/nu14010066/s1>, Table S1: Diet formula, Table S2: Differential expressed genes in  $2.5 \times$  FA compared to control female mice brains at P21d, Table S3: Genes differentially expressed in both  $2.5 \times$  FA female and male brains.

**Author Contributions:** Conceptualization, F.L.; methodology, F.W. and D.C.; software, F.W. and D.C.; validation, X.Y., W.S., F.W., F.L. and D.C.; formal analysis, X.Y., W.S. and Z.L.; investigation, Q.W. and H.L.; resources, Q.W. and X.S.; data curation, D.C.; writing—original draft preparation, D.C.; writing—review and editing, X.Y., W.S., Y.C., F.W. and F.L.; visualization, D.C.; supervision, F.W., Y.Z. and L.H.; project administration, F.W. and D.C.; funding acquisition, F.L., D.C., Y.Z. and L.H. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was supported by the Neural Regeneration Co-innovation Center of Jiangsu Province, Nantong University, the National Natural Science Foundation of China (Grants 81300978) and the Innovation and Entrepreneurship Training Program for College Students in Jiangsu Province (202110304100Y).

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Laboratory Animal Ethics Committee of Nantong University (protocol code S20200408-016, 16 March 2020).

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The data presented in this study are available on request from the corresponding author. The data are not publicly available due to confidentiality reasons.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Naninck, E.F.G.; Stijger, P.C.; Brouwer-Brolsma, E.M. The Importance of Maternal Folate Status for Brain Development and Function of Offspring. *Adv. Nutr.* **2019**, *10*, 502–519. [[CrossRef](#)] [[PubMed](#)]
2. Viswanathan, M.; Treiman, K.A.; Kish-Doto, J.; Middleton, J.C.; Coker-Schwimmer, E.J.; Nicholson, W.K. Folic Acid Supplementation for the Prevention of Neural Tube Defects: An Updated Evidence Report and Systematic Review for the US Preventive Services Task Force. *JAMA* **2017**, *317*, 190–203. [[CrossRef](#)] [[PubMed](#)]
3. Crider, K.S.; Bailey, L.B.; Berry, R.J. Folic Acid Food Fortification—Its History, Effect, Concerns, and Future Directions. *Nutrients* **2011**, *3*, 370–384. [[CrossRef](#)] [[PubMed](#)]
4. Page, R.; Robichaud, A.; Arbuckle, T.E.; Fraser, W.D.; MacFarlane, A.J. Total folate and unmetabolized folic acid in the breast milk of a cross-section of Canadian women. *Am. J. Clin. Nutr.* **2017**, *105*, 1101–1109. [[CrossRef](#)]
5. Cheng, T.L.; Mistry, K.B.; Wang, G.; Zuckerman, B.; Wang, X. Folate Nutrition Status in Mothers of the Boston Birth Cohort, Sample of a US Urban Low-Income Population. *Am. J. Public Health* **2018**, *108*, 799–807. [[CrossRef](#)]
6. Pfeiffer, C.M.; Hughes, J.P.; Lacher, D.A.; Bailey, R.L.; Berry, R.J.; Zhang, M.; Yetley, E.A.; Rader, J.I.; Sempos, C.T.; Johnson, C.L. Estimation of trends in serum and RBC folate in the U.S. population from pre- to postfortification using assay-adjusted data from the NHANES 1988–2010. *J. Nutr.* **2012**, *142*, 886–893. [[CrossRef](#)] [[PubMed](#)]
7. Murray, L.K.; Smith, M.J.; Jadavji, N.M. Maternal oversupplementation with folic acid and its impact on neurodevelopment of offspring. *Nutr. Rev.* **2018**, *76*, 708–721. [[CrossRef](#)] [[PubMed](#)]
8. Valera-Gran, D.; Navarrete-Munoz, E.M.; Garcia de la Hera, M.; Fernandez-Somoano, A.; Tardon, A.; Ibarluzea, J.; Balluerka, N.; Murcia, M.; Gonzalez-Safont, L.; Romaguera, D.; et al. Effect of maternal high dosages of folic acid supplements on neurocognitive development in children at 4–5 y of age: The prospective birth cohort Infancia y Medio Ambiente (INMA) study. *Am. J. Clin. Nutr.* **2017**, *106*, 878–887. [[CrossRef](#)] [[PubMed](#)]
9. Desoto, M.C.; Hitlan, R.T. Synthetic folic acid supplementation during pregnancy may increase the risk of developing autism. *J. Pediatr. Biochem.* **2012**, *2*, 251–261.
10. Raghavan, R.; Riley, A.W.; Volk, H.; Caruso, D.; Hironaka, L.; Sices, L.; Hong, X.; Wang, G.; Ji, Y.; Brucato, M.; et al. Maternal Multivitamin Intake, Plasma Folate and Vitamin B12 Levels and Autism Spectrum Disorder Risk in Offspring. *Paediatr. Perinat. Epidemiol.* **2018**, *32*, 100–111. [[CrossRef](#)]
11. Egorova, O.; Myte, R.; Schneede, J.; Hagglof, B.; Bolte, S.; Domellof, E.; Ivars A'roch, B.; Elgh, F.; Ueland, P.M.; Silfverdal, S.A. Maternal blood folate status during early pregnancy and occurrence of autism spectrum disorder in offspring: A study of 62 serum biomarkers. *Mol. Autism* **2020**, *11*, 7. [[CrossRef](#)] [[PubMed](#)]
12. Pickell, L.; Brown, K.; Li, D.; Wang, X.L.; Deng, L.; Wu, Q.; Selhub, J.; Luo, L.; Jerome-Majewska, L.; Rozen, R. High intake of folic acid disrupts embryonic development in mice. *Birth Defects Res. Part. A Clin. Mol. Teratol.* **2011**, *91*, 8–19. [[CrossRef](#)] [[PubMed](#)]
13. Barua, S.; Chadman, K.K.; Kuizon, S.; Buenaventura, D.; Stapley, N.W.; Ruocco, F.; Begum, U.; Guariglia, S.R.; Brown, W.T.; Junaid, M.A. Increasing maternal or post-weaning folic acid alters gene expression and moderately changes behavior in the offspring. *PLoS ONE* **2014**, *9*, e101674. [[CrossRef](#)]
14. Giroto, F.; Scott, L.; Avchalumov, Y.; Harris, J.; Iannattone, S.; Drummondmain, C.; Tobias, R.; Belloespinosa, L.; Rho, J.M.; Davidsen, J. High dose folic acid supplementation of rats alters synaptic transmission and seizure susceptibility in offspring. *Sci. Rep.* **2013**, *3*, 1465. [[CrossRef](#)] [[PubMed](#)]
15. Henzel, K.S.; Ryan, D.P.; Schroder, S.; Weiergraber, M.; Ehninger, D. High-dose maternal folic acid supplementation before conception impairs reversal learning in offspring mice. *Sci. Rep.* **2017**, *7*, 3098. [[CrossRef](#)]
16. Huot, P.S.; Ly, A.; Szeto, I.M.; Reza-Lopez, S.A.; Cho, D.; Kim, Y.I.; Anderson, G.H. Maternal and postweaning folic acid supplementation interact to influence body weight, insulin resistance, and food intake regulatory gene expression in rat offspring in a sex-specific manner. *Appl. Physiol. Nutr. Metab.* **2016**, *41*, 411–420. [[CrossRef](#)] [[PubMed](#)]
17. Barua, S.; Kuizon, S.; Brown, W.T.; Junaid, M.A. DNA Methylation Profiling at Single-Base Resolution Reveals Gestational Folic Acid Supplementation Influences the Epigenome of Mouse Offspring Cerebellum. *Front. Neurosci.* **2016**, *10*, 168. [[CrossRef](#)]

18. Chu, D.; Li, L.; Jiang, Y.; Tan, J.; Ji, J.; Zhang, Y.; Jin, N.; Liu, F. Excess Folic Acid Supplementation Before and During Pregnancy and Lactation Activates Fos Gene Expression and Alters Behaviors in Male Mouse Offspring. *Front. Neurosci.* **2019**, *13*, 313. [[CrossRef](#)] [[PubMed](#)]
19. Compan Gabucio, L.M.; Garcia de la Hera, M.; Torres Collado, L.; Fernandez-Somoano, A.; Tardon, A.; Guxens, M.; Vrijheid, M.; Rebagliato, M.; Murcia, M.; Ibarluzea, J.; et al. The Use of Lower or Higher Than Recommended Doses of Folic Acid Supplements during Pregnancy Is Associated with Child Attentional Dysfunction at 4–5 Years of Age in the INMA Project. *Nutrients* **2021**, *13*, 327. [[CrossRef](#)]
20. Reeves, P.G. Components of the AIN-93 diets as improvements in the AIN-76A diet. *J. Nutr.* **1997**, *127* (Suppl. S5), 838S–841S. [[CrossRef](#)] [[PubMed](#)]
21. Mochizuki, S.; Takano, M.; Sugano, N.; Ohtsu, M.; Tsunoda, K.; Koshi, R.; Yoshinuma, N. The effect of B vitamin supplementation on wound healing in type 2 diabetic mice. *J. Clin. Biochem. Nutr.* **2016**, *58*, 64–68. [[CrossRef](#)] [[PubMed](#)]
22. Qipshidze, N.; Metreveli, N.; Lominadze, D.; Tyagi, S.C. Folic acid improves acetylcholine-induced vasoconstriction of coronary vessels isolated from hyperhomocysteinemic mice: An implication to coronary vasospasm. *J. Cell. Physiol.* **2011**, *226*, 2712–2720. [[CrossRef](#)]
23. Huerkamp, M.J.; Dowdy, M.R. Diet replenishment for ad-libitum-fed mice housed in social groups is compatible with shelf life. *J. Am. Assoc. Lab. Anim. Sci.* **2008**, *47*, 47–50.
24. Bachmanov, A.A.; Reed, D.R.; Beauchamp, G.K.; Tordoff, M.G. Food intake, water intake, and drinking spout side preference of 28 mouse strains. *Behav. Genet.* **2002**, *32*, 435–443. [[CrossRef](#)]
25. Kim, Y.I. Folic acid supplementation and cancer risk: Point. *Cancer Epidemiol. Biomark. Prev.* **2008**, *17*, 2220–2225. [[CrossRef](#)]
26. Wang, Y.; Zhang, Y.; Hu, W.; Xie, S.; Gong, C.X.; Iqbal, K.; Liu, F. Rapid alteration of protein phosphorylation during postmortem: Implication in the study of protein phosphorylation. *Sci. Rep.* **2015**, *5*, 15709. [[CrossRef](#)]
27. Moy, S.S.; Nadler, J.J.; Perez, A.; Barbaro, R.P.; Johns, J.M.; Magnuson, T.R.; Piven, J.; Crawley, J.N. Sociability and preference for social novelty in five inbred strains: An approach to assess autistic-like behavior in mice. *Genes Brain Behav.* **2004**, *3*, 287–302. [[CrossRef](#)] [[PubMed](#)]
28. Harlan De Crescenzo, A.; Panoutsopoulos, A.A.; Tat, L.; Schaaf, Z.; Racherla, S.; Henderson, L.; Leung, K.Y.; Greene, N.D.E.; Green, R.; Zarbalis, K.S. Deficient or Excess Folic Acid Supply During Pregnancy Alter Cortical Neurodevelopment in Mouse Offspring. *Cereb. Cortex* **2021**, *31*, 635–649. [[CrossRef](#)]
29. Cosin-Tomas, M.; Luan, Y.; Leclerc, D.; Malysheva, O.V.; Lauzon, N.; Bahous, R.H.; Christensen, K.E.; Caudill, M.A.; Rozen, R. Moderate Folic Acid Supplementation in Pregnant Mice Results in Behavioral Alterations in Offspring with Sex-Specific Changes in Methyl Metabolism. *Nutrients* **2020**, *12*, 1716. [[CrossRef](#)]
30. Wiens, D.; DeSoto, M.C. Is High Folic Acid Intake a Risk Factor for Autism?—A Review. *Brain Sci.* **2017**, *7*, 149. [[CrossRef](#)]
31. Wiens, D.; DeWitt, A.; Kosar, M.; Underriner, C.; Finsand, M.; Freese, M. Influence of Folic Acid on Neural Connectivity during Dorsal Root Ganglion Neurogenesis. *Cells Tissues Organs* **2016**, *201*, 342–353. [[CrossRef](#)]
32. Schaevitz, L.; Berger-Sweeney, J.; Ricceri, L. One-carbon metabolism in neurodevelopmental disorders: Using broad-based nutraceuticals to treat cognitive deficits in complex spectrum disorders. *Neurosci. Biobehav. Rev.* **2014**, *46*, 270–284. [[PubMed](#)]
33. Plumptre, L.; Masih, S.P.; Ly, A.; Aufreiter, S.; Sohn, K.J.; Croxford, R.; Lausman, A.Y.; Berger, H.; O'Connor, D.L.; Kim, Y.I. High concentrations of folate and unmetabolized folic acid in a cohort of pregnant Canadian women and umbilical cord blood. *Am. J. Clin. Nutr.* **2015**, *102*, 848–857. [[CrossRef](#)] [[PubMed](#)]
34. Li, W.; Jiang, M.; Xiao, Y.; Zhang, X.; Cui, S.; Huang, G. Folic acid inhibits tau phosphorylation through regulation of PP2A methylation in SH-SY5Y cells. *J. Nutr. Health Aging* **2015**, *19*, 123–129. [[CrossRef](#)] [[PubMed](#)]
35. Ma, F.; Li, Q.; Zhou, X.; Zhao, A.; Song, A.; Li, W.; Liu, H.; Xu, W.; Huang, G. Effects of folic acid supplementation on cognitive function and Aβeta-related biomarkers in mild cognitive impairment: A randomized controlled trial. *Eur. J. Nutr.* **2019**, *58*, 345–356. [[CrossRef](#)] [[PubMed](#)]
36. Barua, S.; Kuizon, S.; Chadman, K.K.; Brown, W.T.; Junaid, M.A. Microarray analysis reveals higher gestational folic Acid alters expression of genes in the cerebellum of mice offspring—a pilot study. *Brain Sci.* **2015**, *5*, 14–31. [[CrossRef](#)]
37. Bailey, S.W.; Ayling, J.E. The extremely slow and variable activity of dihydrofolate reductase in human liver and its implications for high folic acid intake. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 15424–15429.
38. Ismail, S.; Senna, A.A.; Behiry, E.G.; Ashaat, E.A.; Zaki, M.S.; Ashaat, N.A.; Salah, D.M. Study of C677T variant of methylene tetrahydrofolate reductase gene in autistic spectrum disorder Egyptian children. *Am. J. Med. Genet. Part. B Neuropsychiatr. Genet. Off. Publ. Int. Soc. Psychiatr. Genet.* **2019**, *180*, 305–309. [[CrossRef](#)]
39. Sun, Y.; Qian, L.; Xu, L.; Hunt, S.; Sah, P. Somatostatin neurons in the central amygdala mediate anxiety by disinhibition of the central subnucleus extended amygdala. *Mol. Psychiatry* **2020**, 1–12. [[CrossRef](#)]
40. Zhang, R.; Asai, M.; Mahoney, C.E.; Joachim, M.; Shen, Y.; Gunner, G.; Majzoub, J.A. Loss of hypothalamic corticotropin-releasing hormone markedly reduces anxiety behaviors in mice. *Mol. Psychiatry* **2017**, *22*, 733–744. [[CrossRef](#)]
41. Reznikov, R.; Bambico, F.R.; Diwan, M.; Raymond, R.J.; Nashed, M.G.; Nobrega, J.N.; Hamani, C. Prefrontal Cortex Deep Brain Stimulation Improves Fear and Anxiety-Like Behavior and Reduces Basolateral Amygdala Activity in a Preclinical Model of Posttraumatic Stress Disorder. *Neuropsychopharmacology* **2018**, *43*, 1099–1106. [[CrossRef](#)] [[PubMed](#)]



42. Zhang, H.T.; Huang, Y.; Masood, A.; Stolinski, L.R.; Li, Y.; Zhang, L.; Dlaboga, D.; Jin, S.L.; Conti, M.; O'Donnell, J.M. Anxiogenic-like behavioral phenotype of mice deficient in phosphodiesterase 4B (PDE4B). *Neuropsychopharmacology* **2008**, *33*, 1611–1623. [[CrossRef](#)] [[PubMed](#)]
43. Hatten, M.E.; Roussel, M.F. Development and cancer of the cerebellum. *Trends Neurosci.* **2011**, *34*, 134–142. [[CrossRef](#)]
44. Fjell, A.M.; Mcevoy, L.; Holland, D.; Dale, A.M.; Walhovd, K.B. What is normal in normal aging? Effects of aging, amyloid and Alzheimer's disease on the cerebral cortex and the hippocampus. *Prog. Neurobiol.* **2014**, *117*, 20–40. [[CrossRef](#)]
45. Riches, Z.; Bloomer, J.C.; Coughtrie, M.W. Comparison of 2-aminophenol and 4-nitrophenol as in vitro probe substrates for the major human hepatic sulfotransferase, SULT1A1, demonstrates improved selectivity with 2-aminophenol. *Biochem. Pharm.* **2007**, *74*, 352–358. [[CrossRef](#)]
46. Cook, I.; Wang, T.; Girvin, M.; Leyh, T.S. The structure of the catechin-binding site of human sulfotransferase 1A1. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 14312–14317. [[CrossRef](#)] [[PubMed](#)]
47. Alnouti, Y.; Klaassen, C.D. Tissue distribution and ontogeny of sulfotransferase enzymes in mice. *Toxicol. Sci.* **2006**, *93*, 242–255. [[CrossRef](#)] [[PubMed](#)]
48. Salman, E.D.; Kadlubar, S.A.; Falany, C.N. Expression and localization of cytosolic sulfotransferase (SULT) 1A1 and SULT1A3 in normal human brain. *Drug Metab. Dispos.* **2009**, *37*, 706–709. [[CrossRef](#)]
49. Tsapakis, E.M.; Fernandes, C.; Moran-Gates, T.; Basu, A.; Sugden, K.; Aitchison, K.J.; Tarazi, F.I. Effects of antidepressant drug exposure on gene expression in the developing cerebral cortex. *Synapse* **2014**, *68*, 209–220. [[CrossRef](#)]
50. Sidharthan, N.P.; Minchin, R.F.; Butcher, N.J. Cytosolic sulfotransferase 1A3 is induced by dopamine and protects neuronal cells from dopamine toxicity: Role of D1 receptor-N-methyl-D-aspartate receptor coupling. *J. Biol. Chem.* **2013**, *288*, 34364–34374. [[CrossRef](#)] [[PubMed](#)]
51. Cook, I.; Wang, T.; Leyh, T.S. Tetrahydrobiopterin regulates monoamine neurotransmitter sulfonation. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, E5317–E5324. [[CrossRef](#)]
52. Bader, M. Inhibition of serotonin synthesis: A novel therapeutic paradigm. *Pharm. Ther.* **2020**, *205*, 107423. [[CrossRef](#)] [[PubMed](#)]
53. Wang, F.X.; Liu, S.Y.; Zheng, X.; Chen, X.; Lu, L.X.; Chen, B.; Xiong, X.Y.; Shu, H.F.; Yang, Q.W.; Yang, H. TLR1 expression in mouse brain was increased in a KA-induced seizure model. *Inflamm. Res.* **2015**, *64*, 487–495. [[CrossRef](#)]
54. Madar, R.; Rotter, A.; Waldman Ben-Asher, H.; Mughal, M.R.; Arumugam, T.V.; Wood, W.H., 3rd; Becker, K.G.; Mattson, M.P.; Okun, E. Postnatal TLR2 activation impairs learning and memory in adulthood. *Brain Behav. Immun.* **2015**, *48*, 301–312. [[CrossRef](#)]
55. Daniele, S.G.; Beraud, D.; Davenport, C.; Cheng, K.; Yin, H.; Maguire-Zeiss, K.A. Activation of MyD88-dependent TLR1/2 signaling by misfolded alpha-synuclein, a protein linked to neurodegenerative disorders. *Sci. Signal.* **2015**, *8*, ra45. [[CrossRef](#)]
56. Wada, Y.; Maekawa, M.; Ohnishi, T.; Balan, S.; Matsuoka, S.; Iwamoto, K.; Iwayama, Y.; Ohba, H.; Watanabe, A.; Hisano, Y.; et al. Peroxisome proliferator-activated receptor alpha as a novel therapeutic target for schizophrenia. *EBioMedicine* **2020**, *62*, 103130. [[CrossRef](#)]
57. Chistyakov, D.V.; Aleshin, S.E.; Astakhova, A.A.; Sergeeva, M.G.; Reiser, G. Regulation of peroxisome proliferator-activated receptors (PPAR) alpha and -gamma of rat brain astrocytes in the course of activation by toll-like receptor agonists. *J. Neurochem.* **2015**, *134*, 113–124. [[CrossRef](#)]
58. Meher, A.; Joshi, A.; Joshi, S. Differential regulation of hepatic transcription factors in the Wistar rat offspring born to dams fed folic acid, vitamin B12 deficient diets and supplemented with omega-3 fatty acids. *PLoS ONE* **2014**, *9*, e90209. [[CrossRef](#)] [[PubMed](#)]
59. White, C.J.; Ellis, J.M.; Wolfgang, M.J. The role of ethanolamine phosphate phospholyase in regulation of astrocyte lipid homeostasis. *J. Biol. Chem.* **2021**, *297*, 100830. [[CrossRef](#)] [[PubMed](#)]
60. Leventoux, N.; Augustus, M.; Azar, S.; Riquier, S.; Villemain, J.P.; Guelfi, S.; Falha, L.; Bauchet, L.; Goze, C.; Ritchie, W.; et al. Transformation Foci in IDH1-mutated Gliomas Show STAT3 Phosphorylation and Downregulate the Metabolic Enzyme ETNPPL, a Negative Regulator of Glioma Growth. *Sci. Rep.* **2020**, *10*, 5504. [[CrossRef](#)]
61. Diaz-Canestro, C.; Bonetti, N.R.; Wust, P.; Nageswaran, V.; Liberale, L.; Beer, J.H.; Montecucco, F.; Luscher, T.F.; Bohacek, J.; Camici, G.G. Apold1 deficiency associates with increased arterial thrombosis in vivo. *Eur. J. Clin. Investig.* **2020**, *50*, e13191. [[CrossRef](#)] [[PubMed](#)]
62. Sosa-Madrid, B.S.; Hernandez, P.; Blasco, A.; Haley, C.S.; Fontanesi, L.; Santacreu, M.A.; Pena, R.N.; Navarro, P.; Ibanez-Escriche, N. Genomic regions influencing intramuscular fat in divergently selected rabbit lines. *Anim. Genet.* **2020**, *51*, 58–69. [[CrossRef](#)] [[PubMed](#)]
63. Cushing, E.M.; Chi, X.; Sylvers, K.L.; Shetty, S.K.; Potthoff, M.J.; Davies, B.S.J. Angiotensin-like 4 directs uptake of dietary fat away from adipose during fasting. *Mol. Metab.* **2017**, *6*, 809–818. [[CrossRef](#)] [[PubMed](#)]
64. Chakraborty, A.; Kamermans, A.; van Het Hof, B.; Castricum, K.; Aanhane, E.; van Horssen, J.; Thijssen, V.L.; Scheltens, P.; Teunissen, C.E.; Fontijn, R.D.; et al. Angiotensin like-4 as a novel vascular mediator in capillary cerebral amyloid angiopathy. *Brain* **2018**, *141*, 3377–3388. [[CrossRef](#)]