



EDITORIAL

Trauma occurs in social contexts

This paper is part of the Special Issue: Trauma occurs in social contexts. More papers from this issue can be found at www.ejpt.net

ilnius, Lithuania's capital, was the beautiful location of the 14th Conference of the European Society of Traumatic Stress studies (ESTSS) from June 10 to 13, 2015. The main theme of this two-yearly conference was "Trauma in changing societies: social contexts and clinical practice." The topic of the conference is timely, unfortunately. The current crisis in Syria changed (and is still changing) societies, especially in the Middle East. Over half of the population of Syria has left their homes, of which over 4 million have fled to other countries for an unknown future (UNHCR, 2016; see also Hall & Olff, 2016; Purgato & Olff, 2015). Other recent events such as recent terrorist attacks in Europe have also had an impact on the social contexts in affected countries. Such circumstances are likely to have implications for clinical practice in psychotrauma. The idea that social context in which trauma occurs strongly affects the mental health impact of trauma has been acknowledged within ESTSS (Ajduković, 2013). The lack of social support after trauma is one of the most consistent predictors of PTSD (e.g., Ozer et al., 2003); the neurobiology underlying the beneficial effects of social support receives increasing attention (Koch et al., 2016a, 2016b; Olff, 2012; Olff et al., 2014) and research shows how the opposite, that is, social exclusion, may explain PTSD symptoms through tonic immobility (Mooren & Van Minnen, 2014). But social support is a complex construct and differences exist among countries in the perception of received and desired support (Zepinic, Bogic, & Priebe, 2012). This special issue presents the papers based on the keynote presentations and invited panel discussions during the meeting, and the majority of these papers are related to this overarching conference theme.

The first lines of the popular Lithuanian song "Trys milijonai" sound "Perhaps we fought for too long—and in too many wars. Perhaps too long we have been chanting Glory." They refer to the many regional and world-wide conflicts in which Lithuania has been involved during the past century. The first paper of this special issue was written by Evaldas Kazlauskas and Pauline Želvienè from Vilnius University in Lithuania (Kazlauskas & Želvienè, 2016) and presents an overview of the emerging number of studies from the Baltic countries (Lithuania, Estonia, Latvia) on the psychological consequences of these events. Studies are presented on the effects of adverse events resulting from Russian occupation for 50 years until 1990, and on the effects of two major disasters that involved the Baltic populations (the Chernobyl Nuclear Power Station accident in 1986 and the sinking of the MS Estonia in 1994).

Stuart Turner (2015) focuses on the recent increase in refugees seeking asylum in Europe due to conflicts in Syria and Afghanistan, and implications that this "refugee crisis" may have on policy of refugee admissions and mental health care. Turner (2015) emphasises the importance of supporting economic growth and human rights in low- and middle-income countries, and of recognising the weaknesses of the policies and decisionmaking processes concerning asylum procedures for refugees.

Maercker and Hecker (2016) describe a social interpersonal framework model of posttraumatic stress disorder (PTSD), in which the role of affective reactions in response to others, relationships with close ones, and the broader culture and society such as the cultural values held on posttraumatic stress and resilience are outlined. Maercker and Hecker (2016) also provide implications for psychological trauma interventions addressing social affects, relationships, or communities.

The paper "Social attachments and traumatic stress" by Richard Bryant (2016) provides an overview of his exciting keynote lecture that—similar to the previous keynote—challenged the traditional individual focus on trauma and recovery in favour of a focus that includes also the larger social context in which trauma occurs. Bryant (2016) presents both experimental studies and large-scale social network analysis studies on social connections and attachments in relation to the recovery after traumatic experiences. Based on this review, Bryant (2016) argues that an individual's attachment style (either secure or avoidant) is likely to interact with the response to receiving social support, a notion that should be tested in further empirical studies.

European Journal of Psychotraumatology 2016. © 2016 Marit Sijbrandij and Miranda Olff. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), allowing third parties to copy and redistribute the material in any medium or format, and to remix, transform, and build upon the material, for any purpose, even commercially, under the condition that appropriate credit is given, that a link to the license is provided, and that you indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use. Citation: European Journal of Psychotraumatology 2016, **7**: 31389 - http://dx.doi.org/10.3402/ejpt.v7.31389 (page number not for citation purpose) In addition, three papers integrated invited panel discussions that were held at the conference. First, Magruder, Kassam-Adams, Thoresen, and Olff (2016) discuss and illustrate how a public health approach to trauma may inform research and prevention efforts. This is illustrated by an example of the response to the 2011 Utøya attacks in Norway (see also Thoresen, Aakvaag, Wentzel-Larsen, Dyb, & Hjemdal, 2012) by showing the policy responses and intervention efforts on the individual, community, and the societal levels.

Kazlauskas et al. (2016b) describe the current situation with respect to the availability of different treatment approaches for trauma-related symptoms in seven European countries: Georgia, Germany, Lithuania, the Netherlands, Poland, Switzerland, and Turkey. The countries appear to be very different in terms of awareness of health problems related to trauma, and whether evidence-based trauma treatments are available.

Finally, Thomaes et al. (2016) from the ESTSS Taskforce on Neurobiology provided a general overview of the main research challenges that trauma researchers in the field of neurobiology are facing. They provide suggestions towards enhancing the methodology in future studies of biological traces underlying the aetiology of PTSD, and pharmacological strategies to prevent or treat PTSD. This paper is also a call for more standardisation and greater European and international collaboration in this area.

With this special issue that presents highlights of the biennial psychotrauma conference in Europe with contributions from experts inside and outside Europe, we hope to boost future research on the role of the broader social context in which trauma occurs. This offers many possibilities for novel research topics and methods. To name a few, big data studies may examine the use of social media, and specific characteristics of communication between individuals early after mass trauma by tracking smartphone data. Studies may also apply frameworks and variables from related disciplines such as sociology or economics, for instance, human value orientations (cf. Maercker & Hecker, 2016) to study the development of trauma-related psychological distress within larger communities or to compare countries. We may need more research into social circumstances for PTSD treatment, like this study showing that EMDR may be effective in reducing PTSD and depression symptoms among Syrian refugees located in a refugee camp (Acarturk et al., 2015). An exciting new research area opens up, with many possibilities for collaboration between researchers in the field of trauma and beyond.

Marit Sijbrandij

Department of Clinical Psychology The Netherlands and EMGO Institute for Health and Care Research VU University Amsterdam NL-1081 HV Amsterdam, The Netherlands Email: e.m.sijbrandij@vu.nl

Miranda Olff

Department of Psychiatry Academic Medical Center University of Amsterdam Amsterdam, The Netherlands Arq Psychotrauma Expert Group Diemen, The Netherlands

References

- Acarturk, C., Konuk, E., Cetinkaya, M., Senay, I., Sijbrandij, M., Cuijpers, P., & Aker, T. (2016). EMDR for Syrian refugees with posttraumatic stress disorder symptoms: results of a pilot randomized controlled trial. *European Journal of Psychotraumatology*, 6, 27414, doi: http://dx.doi.org/10.3402/ejpt.v6.27414
- Ajduković, D. (2013). Introducing the notion of social context of collective trauma to ESTSS. *European Journal of Psychotraumatology*, 4, 21258, doi: http://dx.doi.org/10.3402/ejpt.v4i0.21258
- Bryant, R.A. (2016). Social attachments and traumatic stress. *European Journal of Psychotraumatology*, 7, 29065, doi: http:// dx.doi.org/10.3402/ejpt.v7.29065
- Hall, B.J., & Olff, M. (2016). Global mental health: Trauma and adversity among populations in transition. *European Journal of Psychotraumatology*, 7, 31140, doi: http://dx.doi.org/10.3402/ ejpt.v7.31140
- Kazlauskas, E., Javakhishvilli, J., Meewisse, M., Merecz-Kot, D., Sar, V., Schäfer, I., ... Gersons, B.P.R. (2016). Trauma treatment across Europe: Where do we stand now from a perspective of seven countries? *European Journal of Psychotraumatology*, 7, 29450, doi: http://dx.doi.org/10.3402/ejpt.v7.29450
- Kazlauskas, E., & Zelvienè, P. (2016). Trauma research in the Baltic countries: From political oppression to recovery. *European Journal of Psychotraumatology*, 7, 29295, doi: http://dx.doi.org/ 10.3402/ejpt.v7.29295
- Koch, S.B., Van Zuiden, M., Nawijn, L., Frijling, J.L., Veltman, D.J., & Olff, M. (2016a). Intranasal oxytocin administration dampens amygdala reactivity towards emotional faces in male and female PTSD patients. *Neuropsychopharmacology*. doi: http://dx.doi.org/10.1038/npp.2015.299
- Koch, S.B., Van Zuiden, M., Nawijn, L., Frijling, J.L., Veltman, D.J., & Olff, M. (2016b). Intranasal oxytocin normalizes amygdala functional connectivity in post-traumatic stress disorder. *Neuropsychopharmacology*. doi: http://dx.doi.org/ 10.1038/npp.2016.1
- Maercker, A., & Hecker, T. (2016). Broadening perspectives on trauma and recovery: A socio-interpersonal view of PTSD. *European Journal of Psychotraumatology*, 7, 29303, doi: http:// dx.doi.org/10.3402/ejpt.v7.29303
- Magruder, K.M., Kassam-Adams, N., Thoresen, S., & Olff, M. (2016). Prevention and public health approaches to trauma and traumatic stress: State of the art and next step. *European Journal of Psychotraumatology*, 7, 29715, doi: http://dx.doi.org/ 10.3402/ejpt.v7.29715
- Mooren, N., & Van Minnen, A. (2014). Feeling psychologically restrained: The effect of social exclusion on tonic immobility. *European Journal of Psychotraumatology*, 5, 22928, doi: http:// dx.doi.org/10.3402/ejpt.v5.22928
- Olff, M. (2012). Bonding after trauma: On the role of social support and the oxytocin system in traumatic stress. *European Journal*

of Psychotraumatology, 3, 18597, doi: http://dx.doi.org/10.3402/ ejpt.v3i0.18597

- Olff, M., Koch, S., Nawijn, L., Frijling, J., Van Zuiden, M., & Veltman, D. (2014). Social support, oxytocin and PTSD. *European Journal of Psychotraumatology*, 5, 26513, doi: http://dx.doi.org/10.3402/ejpt.v5.26513
- Ozer, E.J., Best, S.R., Lipsey, T.L., & Weiss, D.S. (2003). Predictors of posttraumatic stress disorder and symptoms in adults: A meta-analysis. *Psychological Bulletin*, 129(1), 52–73.
- Purgato, M., & Olff, M. (2015). Global mental health and trauma: The current evidence and the long road ahead. *European Journal of Psychotraumatology*, 6, 30120, doi: http://dx.doi.org/ 10.3402/ejpt.v6.30120
- Thomaes, K., De Kloet, C.S., Wilker, S., El-Hage, W., Schäfer, I., Kleim, B., ... Van Zuiden, M. (2016). Investigating biological traces of traumatic stress in changing societies: Challenges and directions from the ESTSS Taskforce on Neurobiology. *European Journal of Psychotraumatology*, 7, 29453, doi: http:// dx.doi.org/10.3402/ejpt.v7.29453
- Thoresen, S., Aakvaag, H.F., Wentzel-Larsen, T., Dyb, G., & Hjemdal, O.K. (2012). The day Norway cried: Proximity and distress in Norwegian citizens following the 22nd July 2011 terrorist attacks in Oslo and on Utøya Island. *European Journal of Psychotraumatology*, 3, 19709, doi: http://dx.doi. org/10.3402/ejpt.v3i0.19709
- Turner, S. (2015). Refugee blues: A UK and European perspective. European Journal of Psychotraumatology, 6, 29328, doi: http:// dx.doi.org/10.3402/ejpt.v6.29328
- UNHCR. (2016). Supporting Syria and the region: London conference 2016. First thematic pledging session (humanitarian). Remarks by Filippo Grandi United Nations High Commissioner for Refugees. London. Retrieved February 22, 2016, from http://www.unhcr.org/56b32a6d9.html
- Zepinic, V., Bogic, M., & Priebe, S. (2012). Refugees' views of the effectiveness of support provided by their host countries. *European Journal of Psychotraumatology*, 3, 8447, doi: http:// dx.doi.org/10.3402/ejpt.v3i0.8447