

Poster presentation

Open Access

## P04-36. HIV-1-specific antibody dependant cellular cytotoxicity (ADCC) mediated by primary NK cells

M Delaporte\*, V Holl, T Decoville, J Penichon and C Moog

Address: UMR INSERM/UMS, Strasbourg, France

\* Corresponding author

from AIDS Vaccine 2009  
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, 6(Suppl 3):P64 doi:10.1186/1742-4690-6-S3-P64

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P64>

© 2009 Delaporte et al; licensee BioMed Central Ltd.

### Background

Natural killer (NK) cells play an important role in antiviral immune responses; they kill HIV-1-infected cells either by direct lysis or through an antibody-dependent cellular cytotoxicity (ADCC) mechanism. HIV-1-specific ADCC were mainly detected using the CEM-NK target cell line. The aim of our study was to analyze the ADCC activity of various HIV-1-specific antibodies with an assay involving primary activated NK cells and autologous infected CD4 T-lymphocytes.

### Methods

NK cells were purified from PBMC by magnetic bead selection. Autologous lymphocytes were stimulated by PHA for 3 days before being infected with different R5 HIV-1 strains for 3 additional days. NK cells previously activated with IL-2 or IL-15 were added to infected lymphocytes for 4 hours in the presence of different concentrations of anti-HIV-1-specific antibodies. The percentages of HIV-1-infected CD4 T-lymphocytes were measured by the detection of intracellular viral p24 antigen using flow cytometry. The immuno-phenotype of NK cells and the expression of CD107a (marker of NK cell degranulation) were determined in parallel.

### Results

Our results show that in the absence of anti-HIV-1-specific antibodies, activated NK cells reduce by up to 50% the number of infected CD4 T-lymphocytes by direct lysis. In the presence of antibodies, the percentage of HIV-1-infected CD4 T-lymphocytes was further reduced and the percentage of NK cell degranulation was increased indi-

cating HIV-1-specific ADCC activity. This activity was compared to other antibody inhibitory activities i.e. HIV-1 neutralization and Fcγ receptors-mediated HIV-1 inhibition. Moreover, a correlation between the HIV-1-specific ADCC and the immuno-phenotype of NK cells was analyzed.

### Conclusion

These results demonstrate that HIV-1-specific antibodies can inhibit HIV-1 replication *in vitro* by different mechanisms, including ADCC. If ADCC participates in HIV-1 protection, antibodies displaying ADCC should be induced by vaccination together with a stimulation of an innate NK immune response.