

Lipocalin 2 (LCN2) Knockdown Regulates Treg/Th17 Balance to Improve Asthma in Mice

Le Xu, Minkang Shi 

Department of Pediatrics, Liyang People's Hospital, Liyang, JiangSu, 213300, People's Republic of China

Correspondence: Minkang Shi, Department of Pediatrics, Liyang People's Hospital, No. 70 Jianshe West Road, Liyang, JiangSu, 213300, People's Republic of China, Email smk198403231411@163.com

Purpose: Asthma substantially affects the quality of life and health of children. Lipocalin 2 (LCN2) is an immune-related protein, which is predicted to be highly expressed in asthma. Here, we investigated the role of LCN2 in ovalbumin (OVA)-induced asthma mouse model.

Methods: We knocked down LCN2 in an asthma mouse model and performed histopathological analysis using hematoxylin and eosin (H&E) staining assay. Differentiated cells were assessed using Diff-Quick staining assay. We investigated the regulatory T (Treg) cell/T helper 17 (Th17) cell balance using flow cytometry and enzyme-linked immunosorbent assay (ELISA). Inflammatory factors were measured using quantitative real-time reverse transcription PCR (qRT-PCR). The involved pathways were assessed using Western blotting.

Results: LCN2 was upregulated in patients with asthma. OVA promoted pathological deterioration in the lungs, increased IgE levels in the plasma, and elevated the number of differentiated inflammatory cells, whereas LCN2 knockdown abrogated the OVA-induced effects. Additionally, the Treg/Th17 imbalance and increased inflammatory cytokine levels were improved by LCN2 knockdown in OVA-treated mice. Moreover, LCN2 knockdown reversed the activation of the janus kinase (JNK) pathway.

Conclusion: LCN2 knockdown improved the Treg/Th17 balance, alleviated inflammation, and inactivated the JNK pathway in OVA-induced asthma mouse model, suggesting that LCN2 may be a novel therapeutic target for asthma in children.

Keywords: innate immunity, T lymphocyte differentiation, peripheral blood, JNK pathway, inflammation

Introduction

Asthma is a chronic respiratory disease characterized by airway inflammation and remodeling.¹ Patients with asthma usually present with cough, shortness of breath, and chest tightness. Asthma affects more than 300 million people, especially children; its prevalence in children below the age of 18 years is approximately 8%.^{2,3} Severe asthma causes poor outcomes in children, such as drug side effects, poor quality of life, and death.⁴ Because asthma is a heterogeneous condition with different phenotypes, it cannot be completely cured despite the availability of several biologics that target type 2 inflammation.⁵ Thus, the pathogenesis of asthma should be elucidated.

CD4⁺ T cells participate in the progression of asthma and determine the phenotype of asthma, ie, eosinophilic or neutrophilic.⁶ Naïve CD4⁺ T cells differentiate into regulatory T (Treg) cells and T helper 17 (Th17) cells.^{7,8} Th17 cells induce inflammation, leading to autoimmune and inflammatory disorders, whereas the Treg cells act antagonistically and maintain immune homeostasis and prevent diseases.⁹ Moreover, previous studies have reported Treg/Th17 imbalance during asthma. Therefore, reducing Treg/Th17 imbalance could be a potential strategy to attenuate asthma.^{10,11}

Lipocalin 2 (LCN2), is associated with innate immunity. It functions as an iron regulator under inflammatory conditions.¹² In normal tissues, LCN2 protects against bacterial infection and oxidative stress damage. Moreover, LCN2 is expressed in several types of cells, such as macrophages, neutrophils, granulocytes, and epithelial cells.¹³ LCN2 expression is high in CD4⁺ T cells, and it plays a crucial role in type 1 helper T (Th1) cell differentiation.¹⁴ Previous studies have reported that LCN2 is involved in the pathological progression of many diseases, such as

malignancy, cardiovascular disease, and inflammation.^{13,15,16} Dysregulation of LCN2 has been observed in the serum and plasma of patients with asthma, indicating that LCN2 is associated with the development of asthma.^{17,18} However, the underlying mechanisms of LCN2 are not yet understood.

Therefore, in the present study, we investigated the role of LCN2 in the pathogenesis of asthma. We evaluated the effect of LCN2 knockdown on the histopathology, inflammatory cytokine levels, and Treg/Th17 balance on an asthma mouse model. The findings of this study suggest that LCN2 is a potential therapeutic target for asthma in children.

Materials and Methods

Microarray Analysis

The GSE31773 dataset was downloaded from the Gene Expression Omnibus (GEO) database (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE31773>). Abnormally expressed genes were analyzed using the R programming language. Differentially expressed genes were defined based on the following criteria: $|\log_2(\text{fold change})| > 1$ and P value < 0.05 .

Clinical Samples

This study was approved by the Ethics Committee of Liyang People's Hospital (No. [2022]02-283-01). Thirty-three children (age: 3–12 years) diagnosed with asthma participated in this study. Patients with the following conditions were excluded from the study: other respiratory diseases, lung disease, allergic disease, history of asthma, or those taking drugs. Moreover, 33 age-matched healthy children were included in the control group. Fasting peripheral blood samples were collected from all participants. Written informed consent was obtained from the legal guardians of all participants. Blood samples were collected according to the Declaration of Helsinki.

Animal Study

The animal experiments were approved by the Ethics Committee of Liyang People's Hospital (No. [2022]02-283-01) and were performed in accordance with the Guide for the Care and Use of Laboratory Animals (National Institutes of Health). Male BALB/c mice (4-week-old; Shanghai Laboratory Animal Centre, Shanghai, China) were housed in specific pathogen-free (SPF) environment, with 12-h light/12-h dark cycle, $23 \pm 2^\circ\text{C}$, 50–60% humidity, and free water and food. The mice were randomly divided into the following four groups, with six mice per group: control, ovalbumin (OVA), OVA + sh-NC, and OVA + sh-LCN2. To establish the asthma model, OVA (25 μg ; Sigma-Aldrich, St. Louis, MO, USA) and aluminum hydroxide (1 mg; Sigma-Aldrich) were dissolved in 200 μL of normal saline and intraperitoneally injected into the mice from OVA, OVA + sh-NC, and OVA + sh-LCN2 groups on days 0, 7, and 14. On day 15, the mice were injected with 5% OVA in the airway for 30 min every day using an ultrasonic sprayer until day 22. The control mice were sensitized using normal saline as per the abovementioned dosage of OVA. Lentiviruses carrying shRNA (sh)-negative control (NC) and sh-LCN2 (GenePharma, Shanghai, China) were intravenously injected into the mice from groups OVA + sh-NC, and OVA + sh-LCN2 before model establishment. All mice were sacrificed 23 d after peripheral blood collection. Bronchoalveolar lavage fluid (BALF) and lung tissue samples were obtained from all mice.

BALF Collection

After sacrificing the mice, the trachea was intubated using polyethylene tubing. An incision was made at the midline of the neck to expose the trachea. Tracheostomy was performed using a 14-gauge needle. Cold sterile phosphate-buffered saline (PBS; 0.5 mL) was added through the needle to the right lung of each mouse, and the lung was washed twice. The liquid obtained via bronchoalveolar lavage or BALF was collected and centrifuged at $500 \times g$ for 10 min. The collected BALF was stored at -20°C until further use.

Hematoxylin and Eosin (H&E) Staining Analysis

The left lung tissue samples were fixed in 10% formalin for histopathological testing and cut into paraffin sections. After dewaxing and rehydration, the sections were stained using hematoxylin (Beyotime, Shanghai, China) for 5 min and eosin (Beyotime) for 3 min. The cells were visualized under a microscope (Olympus, Tokyo, Japan).

Inflammation score was quantified as the following criteria¹⁹: 0 point (no inflammatory cells), 1 point (inflammatory cells were observed occasionally), 2 points (1–3 layers of inflammatory cells around the bronchi), 3 points (4–5 layers of inflammatory cells around the bronchi or vessels), and 4 points (more than 5 layers of inflammatory cells around the bronchi or vessels).

Diff-Quick Staining Assay

The differentiated cells in the BALF were evaluated using a Diff-Quick stain kit (Solarbio, Beijing, China). Briefly, BALF was prepared as a BALF smear, which was incubated with Diff-Quick I and Diff-Quick II for 10 and 15s, respectively. The differentiated cells were observed under a microscope, and their number was counted. The total number of inflammatory cells was counted using a hemocytometer. The percentage of neutrophils, macrophages, eosinophils, and lymphocytes were calculated by the number of differentiated cells/total inflammatory cells *100%.

Flow Cytometry

For flow cytometry, peripheral blood mononuclear cells (PBMCs) were isolated from peripheral blood using Percoll solution (Sigma-Aldrich). After centrifugation at $800 \times g$ for 30 min and washing with PBS, the PBMCs were maintained in Dulbecco's Modified Eagle Medium (DMEM; Hyclone, South Logan, UT, USA) supplemented with 10% fetal bovine serum (FBS; Hyclone) at 37°C with 5% CO₂.

To investigate Th17 cells, the PBMCs were stimulated with 25 ng/mL PMA, 1 mg/mL ionomycin, and 1 µg/mL BFA at 37°C for 5 h. Next, the samples were incubated with a fluorescein isothiocyanate (FITC)-conjugated anti-CD4 antibody at 4°C for 45 min, followed by incubation with a phycoerythrin (PE)-conjugated anti-interleukin-17 (IL-17) antibody for 30 min in the dark. To investigate Treg cells, CD4⁺ T cells were isolated using a CD4⁺ T-cell isolation kit (Novo Biotechnology, Beijing, China). The samples were incubated with an FITC-conjugated anti-CD25 antibody at 4°C for 45 min, followed by incubation with a PE-conjugated anti-forkhead box p3 (Foxp3) antibody for 30 min in the dark. Finally, all samples were analyzed using a flow cytometer. The antibodies were purchased from BioLegend (San Diego, CA, USA).

Enzyme-Linked Immunosorbent Assay (ELISA)

The levels of tumor growth factor beta (TGF-β), interleukin-10 (IL-10), IL-17, and interleukin-23 (IL-23) in the serum and the levels of immunoglobulin E (IgE) in the plasma of mice were measured using specific mouse ELISA kits for each molecule, according to the manufacturer's protocol. All ELISA kits were purchased from Thermo Fisher Scientific (Waltham, MA, USA).

Quantitative Reverse Transcription PCR (qRT-PCR)

Total RNA was isolated from the peripheral blood and lung tissue samples using TRIzol reagent (Invitrogen, Carlsbad, CA, USA). RNA (1 µg) was reverse-transcribed to cDNA using PrimeScript RT Master Mix. qPCR was conducted using a real-time PCR master mix (SYBR Green). mRNA expression was calculated using the $2^{-\Delta\Delta CT}$ method and normalized to that of GAPDH. The primer sequences are shown in Table 1.

Western Blotting

Proteins were isolated from lung tissue samples of the mice using RIPA buffer (Beyotime). After the measurement of protein concentration using a BCA kit (Beyotime), equal amount of protein was added to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) for separation, followed by the transfer of the proteins to polyvinylidene fluoride (PVDF) membranes (Beyotime). The membranes were blocked with 5% not-fat milk at 22°C for 1 h, incubated with primary antibodies at 4°C overnight, and incubated with a secondary antibody at 22°C for 2 h. Each protein band was visualized using BeyoECL plus (Beyotime). The antibodies used are listed in Table 2.

Table 1 The Primer Sequences Used in qRT-PCR

Name	Forward (3'-5')	Reverse (3'-5')
LCN2	AGACAAAGACCCGCAAAAAG	TGGCAACCTGGAACAAAAG
IL-5	GAAAGAGACCTTGACACA	GAACTCTTGAGGTAATCCAGG
IL-13	ATGAGTCTGCAGTATCCCG	CCGTGGCAGACAGGAGTGTT
IL-33	GGTGTTACTGAGTTATATGAG	GGAGCTCCACAGAGTGTCCTTG
GAPDH	AGCCACATCGCTCAGACAC	AATGAAGGGGTCATTGATGG

Table 2 The Antibodies Used in Western Blot

Name	Catalog. No	Dilution
PI3K	ab191606	1:1000
p-PI3K	ab182651	1:1000
AKT	ab8805	1:500
p-AKT	ab38449	1:1000
DVL2	ab228804	1:1000
AXIN1	ab133221	1:1000
p-GSK3 β	ab75814	1/10,000
GSK3 β	ab32391	1/5000
β -catenin	ab32572	1/5000
p-IKK α	ab38515	1:1000
IKK α	ab32041	1:1000
p-IKB	ab194519	1:1000
IKB	ab124957	1:2000
p-p65	ab76302	1:1000
P65	ab16502	1:2000
p-JNK	ab124956	1:1000
JNK	ab179461	1:1000
GAPDH	ab9485	1:2500
Goat anti-rabbit IgG & HRP	ab6721	1:10,000

Statistical Analysis

The data are shown as the mean \pm standard deviation (SD) and were analyzed using GraphPad Prism 8.0 software. For comparison of multiple groups, one-way analysis of variance was used, and the comparison of two groups, Student's *t*-test was used. Statistical significance was defined as $P < 0.05$.

Results

LCN2 Was Highly Expressed in Patients with Asthma

To identify the genes involved in the pathogenesis of asthma, we investigated the differentially expressed genes using microarray analysis; several genes were upregulated or downregulated. *LCN2* was upregulated in patients with severe asthma compared with that in healthy donors (Figure 1A and B). Moreover, *LCN2* levels in the peripheral blood samples of patients with asthma were higher than those in the healthy controls (Figure 1C). These results demonstrated that *LCN2* expression increased in patients with asthma.

Knockdown of *LCN2* Improved the Pathological Changes of Asthma

To investigate the role of *LCN2* in the development of asthma, lentiviruses carrying sh-NC or sh-*LCN2* were injected into the mice and *LCN2* expression was assessed. The results revealed that *LCN2* expression in the peripheral blood samples of the mice decreased following sh-*LCN2* injection compared with that in the sh-NC group (Figure 2A). The plasma IgE levels increased after OVA treatment compared with those of the control mice; however, decrease in *LCN2*




Figure 1 Lipocalin 2 (LCN2) is highly expressed in patients with asthma. **(A)** Differentially expressed genes predicted via the analysis of dataset GSE31773 are presented in the heatmap. Blue: low expression; red: high expression. **(B)** Differentially expressed genes predicted via the analysis of dataset GSE31773 are presented in the bubble diagram. Green: downregulated genes; red: upregulated genes. **(C)** LCN2 levels in the peripheral blood samples of children with asthma ($n=33$) and healthy controls ($n=33$). Data were analyzed using Student's *t*-test. $***P<0.001$.




Figure 2 Knockdown of lipocalin 2 (LCN2) improves the pathological changes that occur due to asthma. **(A)** LCN2 levels in the peripheral blood samples of mice were measured using quantitative reverse transcription PCR (qRT-PCR) after sh-NC and sh-LCN2 were intravenously injected into the mice. **(B)** Plasma IgE levels in the mice of each group. **(C)** Histopathology of lung tissue samples was assessed using hematoxylin and eosin (H&E) staining. (magnification: 100 \times , scale bar: 200 μ m; magnification: 400 \times , scale bar: 50 μ m). **(D)** Inflammation score; $n=6$. Data were analyzed using Student's *t*-test in **(A)** and using one-way ANOVA in **(B–D)**. $***P<0.001$. $####P<0.001$.

levels reduced IgE levels in the OVA-administered groups (Figure 2B). H&E staining revealed that OVA caused lung wall thickening, inflammatory cell infiltration, and alveolar endothelial cell shedding. Nevertheless, LCN2 silencing reversed the lung damage induced by OVA (Figure 2C). Thus, LCN2 knockdown decreased plasma IgE levels and progressive pathology in asthmatic mice.

LCN2 Knockdown Regulated Differentiated Inflammatory Cells

Next, we stained the mice BALF samples using Giemsa, and counted the number of differentiated cells according to cell size. We observed that LCN2 knockdown reduced the percentage of OVA-induced neutrophils, macrophages, and eosinophils (Figure 3A–C); in contrast, OVA decreased the number of lymphocytes, whereas LCN2 knockdown reversed this effect (Figure 3D). Thus, LCN2 knockdown reduced the number of neutrophils, macrophages, and eosinophils, and increased the number of lymphocytes in OVA-treated mice.

LCN2 Knockdown Promoted Treg/Th17 Balance and Regulated Cytokine Levels

To evaluate the Treg/Th17 balance, we performed flow cytometry and measured the levels of Treg and Th17 cells. The results elucidated that OVA increased the number of Th17 cells and decreased that of Treg cells, whereas LCN2




Figure 3 Lipocalin 2 (LCN2) knockdown inhibited differentiation of inflammatory cells. **(A)** Neutrophils, **(B)** macrophages, **(C)** eosinophils, and **(D)** lymphocytes in bronchoalveolar lavage fluid (BALF) were assessed; n=6. Statistical analysis was performed using one-way ANOVA. ***P<0.001. ###P<0.001.

knockdown counteracted this OVA-induced effect (Figure 4A–D). Thus, OVA reduced the Treg/Th17 cell ratio, which increased after LCN2 knockdown (Figure 4E). Serum TGF- β and IL-10 levels reduced in OVA-treated mice, and sh-LCN2 reversed this effect (Figure 4F and G). Moreover, OVA elevated serum IL-17 and IL-23 levels, whereas LCN2 knockdown reversed this effect (Figure 4H and I). The levels of the inflammation-related factors IL-5, IL-13, and IL-33 increased in the lung tissue and BALF samples of OVA-treated mice, and LCN2 knockdown decreased their levels (Figure 4J–O). Thus, LCN2 knockdown improved the Treg/Th17 imbalance in the serum samples of OVA-treated mice and inhibited inflammation in their lung tissue and BALF samples.

Knockdown of LCN2 Inactivated the Janus Kinase (JNK) Pathway

To evaluate the signaling pathways involved in the progression of asthma, we performed Western blotting and investigated several pathway-related factors. The results revealed that LCN2 knockdown did not affect the protein levels of PI3K/AKT, Wnt/ β -catenin, and NF- κ B pathway-related factors in the lung tissue samples of the mice (Figure 5A–C). Additionally, LCN2 knockdown downregulated p-JNK, but not JNK (Figure 5D). OVA upregulated p-JNK, whereas LCN2 lockdown abrogated this effect (Figure 5E). Thus, we demonstrated that LCN2 knockdown inhibited the activation of the JNK pathway in OVA-treated mice.

Discussion

Asthma is a common chronic disease affecting both adults and children. The pathogenesis of asthma is complex, which could be due to the interactions between various genes and the environment.²⁰ LCN2 is an important regulator of the immune response, which is upregulated in asthma. Moreover, because LCN2 is a biomarker of early intervention in asthma,¹⁹ we investigated the role of LCN2 in asthma. IgE is a pro-allergic regulator, and IgE is commonly used for allergic asthma treatment.²¹ In this study, we predicted differentially expressed genes in asthma and normal controls via microarray analysis. LCN2 was predicted to be upregulated in patients with asthma, which was confirmed via the analysis of clinical samples. Next, we elucidated that LCN2 knockdown inhibited IgE levels in the plasma of OVA-treated mice, suggesting that LCN2 reduces allergic reactions.

One of the hallmarks of asthma is airway inflammation; thus, inflammatory cells mediate asthma development. For example, eosinophil count increases in type 2 asthma; macrophages release allergenic cytokines; and neutrophils are associated with non-allergic asthma. The infiltration by these inflammatory cells causes the narrowing of the airway.^{22,23} Herein, we found that LCN2 knockdown reduced the counts of neutrophils, macrophages, and eosinophils and increased those of lymphocytes in OVA-treated mice, suggesting that downregulation of LCN2 improved the inflammation caused due to asthma.

Th17 cells promote inflammation, whereas Treg cells prevent inflammation. Therefore, the balance between Treg and Th17 cells is important for normal physiological processes, and an imbalance between them has been proven to be closely associated with inflammation.²⁴ Moreover, the Treg/Th17 balance regulates the development of neutrophilic, eosinophilic, or allergic asthma.^{10,11,25} Thus, the Treg/Th17 balance plays an important regulatory in the pathogenesis of




Figure 4 Lipocalin 2 (LCN2) promotes the Treg/Th17 balance and regulates cytokines levels. (A) Foxp3, which regulates regulatory T (Treg) cells, was investigated using flow cytometry and quantified in (B). (C) Interleukin-17 (IL-17), the regulator of T helper 17 (Th17) cells, was detected using flow cytometry and quantified in (D). (E) Treg/Th17 cell-ratio. The levels of (F) tumor growth factor beta (TGF- β), (G) interleukin-10 (IL-10), (H) IL-17, and (I) interleukin-23 (IL-23) were detected in the serum of mice from all groups using enzyme-linked immunosorbent assay (ELISA). The levels of (J) interleukin-5 (IL-5), (K) interleukin-13 (IL-13), and (L) interleukin-33 (IL-33) were examined in the lung tissue samples of mice using quantitative reverse transcription PCR (qRT-PCR). (M) IL-5, (N) IL-13, and (O) IL-33 expression in bronchoalveolar lavage fluid (BALF) of mice was measured using qRT-PCR; n=6. Statistical analysis was performed using one-way ANOVA. ***P<0.001. ###P<0.001.




Figure 5 Knockdown of lipocalin 2 (LCN2) inactivates the janus kinase (JNK) pathway. Effects of LCN2 knockdown on the (A) PI3K/AKT pathway, (B) Wnt/ β -catenin pathway, (C) NF- κ B pathway, and (D) JNK pathway-related factors in lung tissue samples were evaluated using Western blotting. (E) Protein levels of p-JNK and JNK were detected in the lung tissue samples of mice from all groups using Western blotting; n=6.

asthma. Foxp3, TGF- β , and IL-10 are regulators of Treg cells, and IL-17 and IL-23 are regulators of Th17 cells.²⁶ In the present study, OVA decreased Treg cell count and reduced the levels of TGF- β and IL-10. Additionally, OVA increased Th17 cell count and IL-17 and IL-23 levels in the mice serum samples. LCN2 knockdown abrogated the OVA-induced Treg/Th17 imbalance. Moreover, we investigated the inflammatory factors in the lung tissue samples of mice. IL-5, IL-13, and IL-33 are type 2 cytokines that increase eosinophilia and mucus in the airway and promote IgE production.^{27,28} We found that the LCN2 knockdown decreased the OVA-induced elevation in IL-5, IL-13, and IL-33 levels in the lung tissue samples of mice. Thus, these findings suggest that LCN2 knockdown inhibits OVA-induced Treg/Th17 imbalance and inflammation.

The JNK pathway is involved in biological processes, including inflammation, cell differentiation, proliferation, and apoptosis.²⁹ It is activated under various stress conditions.³⁰ In asthma, the JNK pathway regulates airway inflammation.^{31,32} Moreover, LCN2 mediates the JNK pathway in several diseases, such as cancer and acute liver injury.^{33,34} However, the effect of LCN2 on the JNK pathway in asthma is unclear. We elucidated that LCN2 knockdown only inhibited the activation of the JNK pathway and not that of the PI3K/AKT-,catenin, and NF- κ B pathways. These data suggested that the JNK pathway participates in the progression of LCN2-mediated asthma.

Conclusions

Therefore, this is the first study to reveal that LCN2 knockdown inhibits inflammation, Treg/Th17 imbalance, and JNK pathway activation in OVA-induced asthma mouse model, suggesting that LCN2 silencing could relieve asthma, and LCN2 is a potential new target for asthma treatment.

Abbreviations

BALF, bronchoalveolar lavage fluid; DMEM, Dulbecco's Modified Eagle Medium; ELISA, enzyme-linked immunosorbent assay; FITC, fluorescein isothiocyanate; Foxp3, forkhead box p3; GEO, Gene Expression Omnibus; IgE, immunoglobulin E; IL-10, interleukin-10; IL-17, interleukin-17; IL-23, interleukin-23; LCN2, lipocalin 2; MSFI, migration stimulating factor inhibitor; NGAL, neutrophil gelatinase-associated lipocalin; OVA, ovalbumin; PBS, phosphate-buffered saline; PBMCs, peripheral blood mononuclear cells; PE, phycoerythrin; PVDF, polyvinylidene fluoride; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis; SPF, specific pathogen-free; TGF- β , tumor growth factor beta.

Data Sharing Statement

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics Approval and Informed Consent

The study was approved by the Ethics Committee of LiYang People's Hospital (No. [2022]02-283-01). Informed consent was obtained from all individual participants included in the study. All animal experiments should comply with the ARRIVE guidelines.

Funding

There is no funding to report.

Disclosure

The authors declare there is no conflicts of interest in this work.

References

1. Chung KF, Dixey P, Abubakar-Waziri H, et al. Characteristics, phenotypes, mechanisms and management of severe asthma. *Chin Med J (Engl)*. 2022;135(10):1141–1155. doi:10.1097/CM9.0000000000001990
2. Zahran HS, Bailey CM, Damon SA, Garbe PL, Breyse PN. Vital Signs: asthma in Children - United States, 2001-2016. *MMWR Morb Mortal Wkly Rep*. 2018;67(5):149–155. doi:10.15585/mmwr.mm6705e1
3. GBD 2015 Chronic Respiratory Disease Collaborators. Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Respir Med*. 2017;5(9):691–706. doi:10.1016/S2213-2600(17)30293-X
4. Haktanir Abul M, Phipatanakul W. Severe asthma in children: evaluation and management. *Allergol Int*. 2019;68(2):150–157. doi:10.1016/j.alit.2018.11.007
5. Pijnenburg MW, Fleming L. Advances in understanding and reducing the burden of severe asthma in children. *Lancet Respir Med*. 2020;8(10):1032–1044. doi:10.1016/S2213-2600(20)30399-4
6. Jeong J, Lee HK. The Role of CD4+ T Cells and Microbiota in the Pathogenesis of Asthma. *Int J Mol Sci*. 2021;22(21):11822. doi:10.3390/ijms222111822
7. Lee GR. The Balance of Th17 versus Treg Cells in Autoimmunity. *Int J Mol Sci*. 2018;19(3):730. doi:10.3390/ijms19030730
8. Yan JB, Luo MM, Chen ZY, He BH. The Function and Role of the Th17/Treg Cell Balance in Inflammatory Bowel Disease. *J Immunol Res*. 2020;2020:8813558. doi:10.1155/2020/8813558
9. Cai F, Jin S, Chen G. The Effect of Lipid Metabolism on CD4+ T Cells. *Mediators Inflamm*. 2021;2021:6634532. doi:10.1155/2021/6634532
10. Zhou Y, Hu L, Zhang H, et al. Guominkang formula alleviate inflammation in eosinophilic asthma by regulating immune balance of Th1/2 and Treg/Th17 cells. *Front Pharmacol*. 2022;13:978421. doi:10.3389/fphar.2022.978421
11. Wei C, Huang L, Zheng Y, Cai X. Selective activation of cannabinoid receptor 2 regulates Treg/Th17 balance to ameliorate neutrophilic asthma in mice. *Ann Transl Med*. 2021;9(12):1015. doi:10.21037/atm-21-2778
12. Xiao X, Yeoh BS, Vijay-Kumar M. Lipocalin 2: an Emerging Player in Iron Homeostasis and Inflammation. *Annu Rev Nutr*. 2017;37:103–130. doi:10.1146/annurev-nutr-071816-064559
13. Xu WX, Zhang J, Hua YT, Yang SJ, Wang DD, Tang JH. An Integrative Pan-Cancer Analysis Revealing LCN2 as an Oncogenic Immune Protein in Tumor Microenvironment. *Front Oncol*. 2020;10:605097. doi:10.3389/fonc.2020.605097
14. Chen W, Li W, Zhang Z, et al. Lipocalin-2 Exacerbates Lupus Nephritis by Promoting Th1 Cell Differentiation. *J Am Soc Nephrol*. 2020;31(10):2263–2277. doi:10.1681/ASN.2019090937
15. Jang HM, Lee JY, An HS, et al. LCN2 deficiency ameliorates doxorubicin-induced cardiomyopathy in mice. *Biochem Biophys Res Commun*. 2022;588:8–14. doi:10.1016/j.bbrc.2021.12.048
16. Ma J, Chen J, Xue K, et al. LCN2 Mediates Skin Inflammation in Psoriasis through the SREBP2–NLRC4 Axis. *J Invest Dermatol*. 2022;142(8):2194–2204.e11. doi:10.1016/j.jid.2022.01.012

17. Hong L, Herjan T, Bulek K, et al. Mechanisms of Corticosteroid Resistance in Type 17 Asthma. *J Immunol.* 2022;209(10):1860–1869. doi:10.4049/jimmunol.2200288
18. Kawagoe J, Kono Y, Togashi Y, et al. Serum Neutrophil Gelatinase-associated Lipocalin (NGAL) Is Elevated in Patients with Asthma and Airway Obstruction. *Curr Med Sci.* 2021;41(2):323–328. doi:10.1007/s11596-021-2350-1
19. Louten J, Mattson JD, Malinao MC, et al. Biomarkers of disease and treatment in murine and cynomolgus models of chronic asthma. *Biomark Insights.* 2012;7:87–104. doi:10.4137/BMI.S9776
20. Papi A, Brightling C, Pedersen SE, Reddel HK. Asthma. *Lancet.* 2018;391(10122):783–800. doi:10.1016/S0140-6736(17)33311-1
21. Bush RK. The use of anti-IgE in the treatment of allergic asthma. *Med Clin North Am.* 2002;86(5):1113–1129. doi:10.1016/s0025-7125(02)00036-6
22. Boonpiyathad T, Sözener ZC, Satitsuksanoa P, Akdis CA. Immunologic mechanisms in asthma. *Semin Immunol.* 2019;46:101333. doi:10.1016/j.smim.2019.101333
23. Nakagome K, Nagata M. Involvement and Possible Role of Eosinophils in Asthma Exacerbation. *Front Immunol.* 2018;9:2220. doi:10.3389/fimmu.2018.02220
24. Hanidziar D, Koulmanda M. Inflammation and the balance of Treg and Th17 cells in transplant rejection and tolerance. *Curr Opin Organ Transplant.* 2010;15(4):411–415. doi:10.1097/MOT.0b013e32833b7929
25. Wang L, Wan H, Tang W, et al. Critical roles of adenosine A2A receptor in regulating the balance of Treg/Th17 cells in allergic asthma. *Clin Respir J.* 2018;12(1):149–157. doi:10.1111/crj.12503
26. Talaat RM, Mohamed SF, Bassyouni IH, Raouf AA. Th1/Th2/Th17/Treg cytokine imbalance in systemic lupus erythematosus (SLE) patients: correlation with disease activity. *Cytokine.* 2015;72(2):146–153. doi:10.1016/j.cyto.2014.12.027
27. Lambrecht BN, Hammad H, Fahy JV. The Cytokines of Asthma. *Immunity.* 2019;50(4):975–991. doi:10.1016/j.immuni.2019.03.018
28. Drake LY, Kita H. IL-33: biological properties, functions, and roles in airway disease. *Immunol Rev.* 2017;278(1):173–184. doi:10.1111/imr.12552
29. Kumar A, Singh UK, Kini SG, et al. JNK pathway signaling: a novel and smarter therapeutic targets for various biological diseases. *Future Med Chem.* 2015;7(15):2065–2086. doi:10.4155/fmc.15.132
30. Cui J, Zhang M, Zhang YQ, Xu ZH. JNK pathway: diseases and therapeutic potential. *Acta Pharmacol Sin.* 2007;28(5):601–608. doi:10.1111/j.1745-7254.2007.00579.x
31. Huang G, Su J, Zhao W, et al. JNK modulates RAGE/ β -catenin signaling and is essential for allergic airway inflammation in asthma. *Toxicol Lett.* 2021;336:57–67. doi:10.1016/j.toxlet.2020.10.002
32. Jin X, Jin W, Li G, Zheng J. Zeaxanthin attenuates OVA-induced allergic asthma in mice by regulating the p38 MAPK/ β -catenin signaling pathway. *Allergol Immunopathol.* 2022;50(5):75–83. doi:10.15586/aei.v50i5.664
33. Schröder SK, Asimakopoulou A, Tillmann S, Koschmieder S, Weiskirchen R. TNF- α controls Lipocalin-2 expression in PC-3 prostate cancer cells. *Cytokine.* 2020;135:155214. doi:10.1016/j.cyto.2020.155214
34. Borkham-Kamphorst E, van de Leur E, Zimmermann HW, et al. Protective effects of lipocalin-2 (LCN2) in acute liver injury suggest a novel function in liver homeostasis. *Biochim Biophys Acta.* 2013;1832(5):660–673. doi:10.1016/j.bbdis.2013.01.014

Journal of Asthma and Allergy

Dovepress

Publish your work in this journal

The Journal of Asthma and Allergy is an international, peer-reviewed open-access journal publishing original research, reports, editorials and commentaries on the following topics: Asthma; Pulmonary physiology; Asthma related clinical health; Clinical immunology and the immunological basis of disease; Pharmacological interventions and new therapies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-asthma-and-allergy-journal>