

POSTER PRESENTATION

Open Access

# Acyclic nucleoside phosphonates containing a second phosphonate group are potent inhibitors of the 6-oxopurine phosphoribosyltransferases and have antimalarial activity

Dianne Keough<sup>1</sup>, Petr Špaček<sup>2</sup>, Dana Hocková<sup>2</sup>, Tomáš Tichý<sup>2</sup>, Silvie Vrbková<sup>2</sup>, Lenka Slavětínská<sup>2</sup>, Zlatko Janeba<sup>2</sup>, Lieve Naesens<sup>3</sup>, Michael Edstein<sup>4</sup>, Marina Chavchich<sup>4</sup>, Tzu Wang<sup>1\*</sup>, John de Jersey<sup>1</sup>, Luke Guddat<sup>1</sup>

From Challenges in malaria research: Core science and innovation  
Oxford, UK. 22-24 September 2014

## Background

The 6-oxopurine phosphoribosyltransferases have been suggested to be a target for the discovery of new antimalarial drugs. This is because protozoan parasites rely solely on the salvage of purines from their host to make the nucleotides needed for RNA and DNA synthesis and lack the de novo pathway. Acyclic nucleoside phosphonates (ANPs) that contain a 6-oxopurine base are good inhibitors of the *Plasmodium falciparum* (Pf) and *Plasmodium vivax* (Pv) 6-oxopurine phosphoribosyltransferases (PRTs) [1]. Chemical modifications based on the crystal structure of 2-(phosphonoethoxy) ethylguanine (PEEG) in complex with human HGPRT have led to the design of new ANPs [2]. These novel compounds contain a second phosphonate group attached to the ANP scaffold [3].

## Results

{[(2-[(Guanine-9Hyl)methyl]propane-1,3-diyl)bis(oxy)]bis(methylene)} diphosphonic acid exhibited a  $K_i$  value of 30 nM for human HGPRT and 70 nM for Pf HGPRT. The crystal structure of this compound in complex with human HGPRT shows that it fills or partially fills three critical locations in the active site: the binding sites of the purine base, the 5'-phosphate group, and pyrophosphate [3]. This is the first HG(X) PRT inhibitor that has been able to achieve this result. Pro-drugs have been synthesized resulting in  $IC_{50}$  values as low as 3.8  $\mu$ M for Pf grown in cell

culture, which is up to 25-fold lower compared to the parent compounds [3].

## Conclusion

The crystal structure of {[(2-[(Guanine-9Hyl)methyl]propane-1,3-diyl)bis(oxy)]bis(methylene)} diphosphonic acid in complex with human HGPRT provides a template for chemical modifications to increase both potency and selectivity for the parasite enzymes.

## Acknowledgements

This work was financially supported by Australia National Health and Medical Research Council (Grants 569703 and 1030353), the Grant Agency of Czech Republic (Grant P207/11/0108) and Gilead Sciences (Foster city, CA).

## Authors' details

<sup>1</sup>School of Chemistry and Molecular Biosciences, University of Queensland, Brisbane, Queensland, Australia. <sup>2</sup>Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, Prague, Czech Republic. <sup>3</sup>Rega Institute for Medical Research, Katholieke Universiteit Leuven, Leuven, Belgium. <sup>4</sup>Australia Army Malaria Institute, Brisbane, Queensland, Australia.

Published: 22 September 2014

## References

1. Hockova D, Holy A, Masojdkova M, Keough DT, de Jersey J, Guddat LW: Synthesis of branched 9-[2-(2-phosphonoethoxy)ethyl]purines as a new class of acyclic nucleoside phosphonates which inhibit *Plasmodium falciparum* hypoxanthine-guanine-xanthine phosphoribosyltransferase. *Bioorg Med Chem* 2009, **17**:6218-6232.
2. Keough DT, Hockova D, Holy A, Naesens LM, Skinner-Adams TS, Jersey J, Guddat LW: Inhibition of hypoxanthine-guanine phosphoribosyltransferase by acyclic nucleoside phosphonates: a new class of antimalarial therapeutics. *J Med Chem* 2009, **52**:4391-4399.
3. Keough DT, Spacek P, Hockova D, et al: Acyclic nucleoside phosphonates containing a second phosphonate group are potent inhibitors of

<sup>1</sup>School of Chemistry and Molecular Biosciences, University of Queensland, Brisbane, Queensland, Australia

Full list of author information is available at the end of the article

6-oxopurine phosphoribosyltransferases and have antimalarial activity. *J Med Chem* 2013, **56**:2513-2526.

doi:10.1186/1475-2875-13-S1-P91

**Cite this article as:** Keough *et al.*: Acyclic nucleoside phosphonates containing a second phosphonate group are potent inhibitors of the 6-oxopurine phosphoribosyltransferases and have antimalarial activity. *Malaria Journal* 2014 **13**(Suppl 1):P91.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

