

## Focus: Induced Pluripotency &amp; Cellular Reprogramming

# Reprogramming of cell fate: epigenetic memory and the erasure of memories past

 Buhe Nashun<sup>†</sup>, Peter WS Hill<sup>†</sup> & Petra Hajkova<sup>\*</sup>

## Abstract

Cell identity is a reflection of a cell type-specific gene expression profile, and consequently, cell type-specific transcription factor networks are considered to be at the heart of a given cellular phenotype. Although generally stable, cell identity can be reprogrammed *in vitro* by forced changes to the transcriptional network, the most dramatic example of which was shown by the induction of pluripotency in somatic cells by the ectopic expression of defined transcription factors alone. Although changes to cell fate can be achieved in this way, the efficiency of such conversion remains very low, in large part due to specific chromatin signatures constituting an epigenetic barrier to the transcription factor-mediated reprogramming processes. Here we discuss the two-way relationship between transcription factor binding and chromatin structure during cell fate reprogramming. We additionally explore the potential roles and mechanisms by which histone variants, chromatin remodelling enzymes, and histone and DNA modifications contribute to the stability of cell identity and/or provide a permissive environment for cell fate change during cellular reprogramming.

**Keywords** cell fate; chromatin; induced pluripotent stem cells; reprogramming; transcription factors

DOI 10.15252/embj.201490649 | Received 25 November 2014 | Revised 26 February 2015 | Accepted 18 March 2015 | Published online 27 March 2015

The EMBO Journal (2015) 34: 1296–1308

## Introduction

During the differentiation process, the developmental capacity of totipotent cells in the early embryo is progressively lost as these undertake cell fate decisions. This process is driven by the expression of cross-antagonistic transcription factors (TF) promoting development towards one cell fate while repressing an alternative differentiation path (Graf & Enver, 2009). Cell fate decisions are fortified by progressive acquisition of complex layers of epigenetic modifications at both the DNA and chromatin level (Goldberg *et al*, 2007; Xie *et al*, 2013; Ho *et al*, 2014). While cell identity is undeniably dictated by the expression profile guided by cell type-specific TFs (Davidson & Erwin, 2006), the robustness of the acquired

transcriptional state is additionally crucially dependent on the configuration of the chromatin context in which these TFs operate (Voss & Hager, 2014). As the key epigenetic modifications acquired during developmental progression are stable and inherited through subsequent cell divisions, an ‘epigenetic memory’ is established that underlies the phenotypic stability of the differentiated cell state (Zhu *et al*, 2013; Jost, 2014; Shipony *et al*, 2014).

Although generally stable *in vivo*, cell fate decisions can be manipulated and even reversed, *in vitro*. The experimental demonstration that every cell of an organism contains the complete genetic information, and that the acquired somatic state can be reversed by exposing the somatic nucleus to the oocyte environment (Gurdon *et al*, 1958; Gurdon, 1960, 1962), set off a search for mechanisms implicated in the erasure of epigenetic memory and the re-establishment of pluri- or totipotency. It has subsequently been shown that cell identity is also amenable to reprogramming using cell fusion (Miller & Ruddle, 1976) and by overexpression of master regulator TFs (Davis *et al*, 1987). Ultimately, reprogramming of somatic cells back to pluripotency was achieved by the ectopic expression of (only) four TFs (Takahashi & Yamanaka, 2006).

In agreement with the role of TFs and gene regulatory networks in defining cell identity, reprogramming of cell fate requires extinction of the existing transcriptional programme followed by the establishment and stabilisation of the transcriptional network specific to the cell type of interest. It has, however, become increasingly obvious that the successful reprogramming process entails, and in fact requires, complete erasure of the existing somatic epigenetic memory followed by the establishment of a new cell type-specific epigenetic signature. Thus, although changes to cell identity can be achieved by ectopic expression of key TFs alone, the efficiency of conversion remains painfully low, with existing chromatin modifications constituting a well-described barrier to the reprogramming process (Mikkelsen *et al*, 2008; Pasque *et al*, 2012; Chen *et al*, 2013b; Gaspar-Maia *et al*, 2013; Sridharan *et al*, 2013).

Here we summarise the current knowledge regarding the complex relationship between chromatin structure and reprogramming of cell fate. We additionally consider whether epigenetic changes are secondary to the newly established transcriptional networks, or whether establishing a permissive chromatin template is a necessary—or potentially even sufficient—step for cell reprogramming to occur.

Medical Research Council Clinical Sciences Centre, Faculty of Medicine, Imperial College London, London, UK

<sup>\*</sup>Corresponding author. Tel: +44 20 83838264; E-mail: petra.hajkova@csc.mrc.ac.uk

<sup>†</sup>These authors contributed equally to this work

## Transcription factors and chromatin structure: a two-way relationship

In a model whereby TF cross-antagonism is the central mechanism by which cell fate is determined, cell fate transitions, such as those observed during de-differentiation and trans-differentiation events, are possible through the ectopic expression of the required cell type instructive TFs (Graf & Enver, 2009). The most extreme and best studied example of this is the direct reprogramming of somatic cells to induced pluripotent stem cells (iPSCs) through the ectopic expression of the pluripotency-associated TFs: Oct4, Sox2, Klf4, and Myc (OSKM) (Takahashi & Yamanaka, 2006). The expression of these TFs destabilises the transcriptional network of differentiated somatic cells and induces the expression of the embryonic stem (ES) cell transcriptional network that eventually leads to the establishment of an ES-like phenotype (Adachi & Scholer, 2012; Niwa, 2014).

In addition to changing the transcriptional network, overexpression of the OSKM transcription factors during iPSC reprogramming has been shown to induce large-scale chromatin changes that ultimately lead to the establishment of a chromatin template highly similar to that of ES cells (Orkin & Hochedlinger, 2011; Liang *et al*, 2012; Apostolou & Hochedlinger, 2013). Of note, the establishment of this chromatin template appears to be finely regulated by OSKM expression levels: sustained high transgene levels appear to hinder the proper establishment of specific (bivalent) chromatin marks during the later stages of iPSC induction, while establishment of the normal ESC-like epigenetic signature can be achieved upon lowering/attenuating expression of the four transgenes at an intermediate point during the induction process (Hussein *et al*, 2014; Tonge *et al*, 2014).

In general, TFs (including OSK) are known to reshape the chromatin landscape in the regions where they bind, both by enabling the binding of other TFs and through direct recruitment of various histone modifiers (Mal & Harter, 2003; Ancelin *et al*, 2006; Magnani *et al*, 2011; Zaret & Carroll, 2011; Soufi *et al*, 2012; Drouin, 2014; Sherwood *et al*, 2014) (Fig 1A). Moreover, the binding of TFs is known to induce locus-specific DNA demethylation (Stadler *et al*, 2011; Feldmann *et al*, 2013). In accordance with these observations, large-scale chromatin changes associated with iPSC reprogramming may be a secondary phenomenon that follows destabilisation of the somatic transcriptional network and establishment of the new pluripotency network. The observed chromatin changes would thus not themselves be directly implicated in the reprogramming process, but rather would reflect successful establishment of the pluripotent state they are associated with.

Contrary to this view, accumulating evidence points towards an important role for chromatin in early stages of reprogramming. It has been shown that the initial engagement of OSK factors during iPSC reprogramming is hindered by repressive histone modifications (Soufi *et al*, 2012), and the failure to successfully establish new gene regulatory networks in trans-differentiation experiments clearly correlates with the presence of closed inaccessible chromatin in the original somatic cell type (Cahan *et al*, 2014; Morris *et al*, 2014). Additionally, both repressive H3K9me2/3 histone methylation and the presence of 5mC have been documented to act as a barrier to the reprogramming process (Mikkelsen *et al*, 2008; Lister *et al*, 2011; Chen *et al*, 2013b; Sridharan *et al*, 2013). Considering these observations, efficient reprogramming appears to require an optimal chromatin configuration that not only allows for fast and

effective engagement of the introduced TFs, but additionally promotes the exchange of chromosomal components, thus enabling fast and efficient erasure of pre-existing DNA and histone modifications (Fig 1B).


## Reprogramming requires opening of the compacted somatic chromatin template

Developmental progression from a totipotent to a differentiated cell is a gradual process accompanied by deposition of repressive histone marks and by increasing chromatin compaction (Gifford *et al*, 2013; Xie *et al*, 2013; Zhu *et al*, 2013). Successful iPSC reprogramming thus requires removal of the somatic repressive chromatin to allow for conversion to a highly dynamic pluripotent chromatin state that is largely devoid of heterochromatin (Meshorer *et al*, 2006). In agreement with this, accumulating evidence suggests that chromatin remodelling complexes and selective deposition or eviction of certain histone variants play important roles in the acquisition and subsequent maintenance of the permissive pluripotent chromatin state (Fig 2 and Table 1).

### Chromatin remodelling factors

Multiple chromatin remodelling factors have been shown to regulate both ES cell identity and somatic cell reprogramming by their chromatin shaping activities. Of the SWI/SNF family of chromatin remodelling factors, esBAF (Brm/Brg-associated factor in ES cells) and Ino80 (inositol requiring 80) have been shown to be important both for the maintenance of ES self-renewal and pluripotency, and also for iPSC reprogramming (Ho *et al*, 2009b, 2011; Wang *et al*, 2014b). In ES cells, esBAF, as well as Ino80, co-localise genome-wide with the pluripotency factors (Ho *et al*, 2009a; Singhal *et al*, 2010; Wang *et al*, 2014b). The activity of these remodelling complexes leads to the generation of open chromatin structure and is thought to promote binding and transcriptional activity of the OSKM factors during reprogramming (Singhal *et al*, 2010; Wang *et al*, 2014b). In a similar manner, the CHD (chromodomain helicase DNA binding) family remodelling factor, Chd1, is also required to maintain open chromatin in ES cells and has been shown to be important for ES cell self-renewal and pluripotency (Gaspar-Maia *et al*, 2009). Down-regulation of Chd1 leads to accumulation of heterochromatin and significantly reduces reprogramming efficiency (Gaspar-Maia *et al*, 2009). These results thus collectively indicate that the potential to open chromatin, or to maintain a less compacted chromatin state, is a prerequisite for the acquisition of pluripotency.

Contrary to the remodelling complexes implicated in the generation of open chromatin structure discussed above, the NuRD (nucleosome remodelling deacetylase) complex contains histone deacetylase activity implicated in gene repression. In the absence of Mbd3, one of the core subunits of the complex, embryonic stem cells exhibit LIF-independent self-renewal capacity associated with elevated expression of pluripotency-related genes (Kaji *et al*, 2006; Reynolds *et al*, 2012). Upon differentiation, Mbd3-null ES cells fail to fully repress genes that are expressed in pre-implantation embryos, which in turn leads to deficiency in lineage commitment (Kaji *et al*, 2006). Interestingly, Mbd3 depletion dramatically increases reprogramming efficiency and results in deterministic and synchronised iPSC reprogramming (Rais *et al*, 2013), even in the absence of c-Myc




**Figure 1. Relationship between transcription factors and chromatin configuration during cell reprogramming.**

(A) Pioneer transcription factors (TFs) are known to reshape the chromatin landscape in the regions where they bind, both by enabling the binding of other TFs and through direct recruitment of various histone modifiers. In addition, the binding of both pioneer and non-pioneer TFs is known to induce locus-specific DNA demethylation. (B) Closed inaccessible chromatin in the original somatic cell type, marked by repressive histone modifications and DNA methylation, hinders the initial engagement of reprogramming-associated TFs. In turn, the activity of chromatin-modifying enzymes results in a permissive chromatin configuration that allows for fast and effective engagement of the introduced TFs, enabling efficient reprogramming.

or Sox2 (Luo *et al*, 2013). It has been suggested that Mbd3/NuRD is recruited through direct interaction with OSKM transcription factors to downstream OSKM target genes and counteracts their reactivation during iPS induction. In the absence of Mbd3, this inhibitory effect is relieved, favouring re-activation of pluripotency genes and leading to improved reprogramming efficiency (Rais *et al*,

2013). However, another recent study reported that Mbd3/NuRD is required for efficient iPS generation from neural stem cells (NSC), pre-iPS cells and epiblast-derived stem cells (EpiSCs) (dos Santos *et al*, 2014). Although overexpression of Mbd3/NuRD does not have any positive or negative effect on iPSC induction efficiency, combined overexpression with Nanog improves both reprogramming



**Figure 2. Chromatin components and modifiers affecting reprogramming efficiency.**

Reprogramming requires the establishment of permissive chromatin and is associated with chromatin opening and changes to histone and DNA modifications. Multiple factors have been implicated in these processes: marked in green and red are factors whose presence/activity is associated with increased and decreased reprogramming efficiency, respectively; marked in purple are those factors whose presence/activity has been shown to both increase and decrease reprogramming efficiency in a context-dependent manner; factors whose influence on reprogramming requires further investigation are marked by (?).

kinetics and efficiency, which is in stark contrast with previous reports showing that overexpression of Mbd3 inhibits iPSC induction (Luo *et al*, 2013; Rais *et al*, 2013). The reported difference may be due to different induction methods and culture conditions used in these studies; however, further investigation is required to clarify the exact role of Mbd3/NuRD in iPSC generation.

**Histone chaperones and variants**

In support of the idea of open chromatin structure promoting reprogramming, overexpression of the histone chaperone Asf1a favours the maintenance of ES cell pluripotency and enhances iPSC induction efficiency from human adult dermal fibroblasts (hADF). Asf1 (anti-silencing factor 1A) non-selectively binds to an H3-H4 heterodimer and facilitates its import from the cytoplasm into the nucleus thus directly regulating the availability of H3-H4 dimer for turnover by the canonical histone H3.1/2-chaperone Caf-1 or by the

H3.3-chaperone Hira (Burgess & Zhang, 2013). Asf1a is also essential for acetylation of newly synthesised H3 at lysine 56 (H3K56ac) (Burgess & Zhang, 2013; Gonzalez-Munoz *et al*, 2014), and it has been suggested that Asf1a regulates the expression of core pluripotency genes during reprogramming by increasing global H3K56 acetylation levels (Gonzalez-Munoz *et al*, 2014).

The incorporation of various histone variants into nucleosomes has a marked impact on local chromatin structure and dynamics. In the context of iPSC reprogramming, combined over-expression of the histone variants TH2A and TH2B, which are normally enriched in the oocyte and early embryo (Montellier *et al*, 2013; Shinagawa *et al*, 2014), has been shown to enhance the efficiency of iPSC generation ninefold. This effect is further enhanced by additional over-expression of the phosphorylation-mimic form of nucleoplasmin (P-Npm), a factor implicated in chromatin remodelling and zygotic gene activation following fertilisation (Shinagawa *et al*, 2014).

**Table 1. The roles of chromatin modifiers during somatic cell reprogramming.**

Category	Chromatin modifiers	Roles in reprogramming	References
Histone modifications	H3K4me2/3	Marks promoters and enhancers of pluripotency- or differentiation-associated genes during initial steps of reprogramming	Ang et al (2011); Koche et al (2011)
	H3K9me2/3	Marks broad heterochromatin regions refractory to initial OSKM binding; acts as an epigenetic barrier towards reprogramming	Soufi et al (2012); Chen et al (2013b); Sridharan et al (2013); Matoba et al (2014)
	H3K27me3	Represses pluripotency-associated genes in somatic cells and differentiation-associated genes in iPSCs	Mansour et al (2012)
	H3K36me2/3	Marks promoter regions of early responsive (MET) genes and represses their activation	Liang et al (2012)
	H3K79me2	Marks transcriptionally active genes; acts as a barrier for efficient repression of lineage-specific genes	Onder et al (2012)
Heterochromatin proteins	HP-1 $\gamma$	Impedes reprogramming by repressing Nanog reactivation	Sridharan et al (2013)
Histone modifiers	Wdr5	Enhances reprogramming by physically interacting with Oct4 and maintaining H3K4me3 on pluripotency-associated gene promoters	Ang et al (2011)
	SUV39H1/2	Enhances reprogramming by facilitating Oct4/Sox2 binding through H3K9me3 demethylation	Onder et al (2012)
	G9a	Inhibition or down-regulation of G9a enhances reprogramming by regulating global H3K9me2/3 levels	Ma et al (2008); Shi et al (2008); Chen et al (2013b); Sridharan et al (2013)
	Setdb1	(?) Down-regulation enhances reprogramming by facilitating H3K9me3 status at core pluripotency genes in one study while opposite effect was observed in another study	Onder et al (2012); Chen et al (2013b)
	Ehmt1	(?) Down-regulation enhances reprogramming by regulating global H3K9me2/3 levels in one study but opposite effect was observed in another study	Onder et al (2012); Sridharan et al (2013)
	PRC1 (Ring1, Bmi1)	Down-regulation of Ring1 or Bmi1 reduces reprogramming efficiency, while overexpression of Bmi1 enhances reprogramming efficiency by regulating H3K27me3 levels	Pereira et al (2010); Moon et al (2011); Onder et al (2012)
	PRC2 (Ezh2, Suz12, Eed)	Down-regulation of Ezh2, Suz12, or Eed reduces reprogramming efficiency, while overexpression of Ezh2 enhances reprogramming efficiency by maintaining H3K27me3 at lineage-specific genes	Pereira et al (2010); Buganim et al (2012); Onder et al (2012); Fragola et al (2013)
	Utx	Physically interacts with OSK; facilitates iPS formation by H3K27me3 demethylation at pluripotency-associated genes	Mansour et al (2012)
	Jmjd3 (Kdm6b)	Depletion increases iPS generation efficiency while overexpression inhibits reprogramming through up-regulating <i>Ink4a/Arf</i> locus expression by H3K27me3 demethylation; also promotes degradation of PHF20 independent of its demethylase activity	Zhao et al (2013)
	Jhdm1a/b (Kdm2a/b)	(?) Down-regulation reduces reprogramming efficiency, while overexpression enhances reprogramming by activating early responsive (MET) genes and the expression of microRNA cluster 302/367	Wang et al (2011); Liang et al (2012)
Chromatin remodellers	Dot1L	Down-regulation enhances reprogramming by promoting the silencing of lineage-specific genes through loss of H3K79me2	Onder et al (2012)
	MBD3/NuRD	Down-regulation enhances reprogramming by facilitating the reactivation of downstream OSKM target genes in one study, while opposite effect was observed in another study	Rais et al (2013); dos Santos et al (2014)
	Ino80	Down-regulation leads to more closed chromatin structure near pluripotency gene promoters and reduces reprogramming efficiency	Wang et al (2014b)
	Chd1	Down-regulation leads to accumulation of heterochromatin and reduces reprogramming efficiency	Gaspar-Maia et al (2009)
	BAF (Brg1, Baf155)	Brg1 and Baf155 synergistically increase reprogramming efficiency by enhancing Oct4 binding and facilitating de-methylation of <i>Oct4</i> and <i>Nanog</i> promoters	Singhal et al (2010)



Table 1 (continued)

Category	Chromatin modifiers	Roles in reprogramming	References
Histone variants	H1foo	Overexpression maintains the pluripotency gene expression and maintains global low methylation status	Hayakawa <i>et al</i> (2012)
	H2A.X	Down-regulation of H2A.X completely inhibits iPS generation	Wu <i>et al</i> (2014)
	H3.3	H3.3 counteracts H1 binding, and down-regulation of H3.3 in oocyte leads to compromised somatic cell reprogramming	Braunschweig <i>et al</i> (2009); Wen <i>et al</i> (2014)
	macroH2A	Co-occupies pluripotency genes with H3K27me3 and acts as an epigenetic barrier to induced pluripotency. Down-regulation significantly enhances iPS generation	Pasque <i>et al</i> (2012); Barrero <i>et al</i> (2013); Gaspar-Maia <i>et al</i> (2013)
	TH2A/B	Co-overexpression enhances reprogramming by inducing an open chromatin structure	Shinagawa <i>et al</i> (2014)
Histone chaperones	ASF1A	Overexpression enhances reprogramming by increasing global H3K56ac levels in the presence of GDF9 in culture medium	Gonzalez-Munoz <i>et al</i> (2014)
DNA modifiers	Dnmt1	Inhibiting activity by small molecules or knockdown significantly increases reprogramming efficiency	Mikkelsen <i>et al</i> (2008)
	TET1/2	Physically interacts and acts in synergy with Nanog. Oxidises 5mC in <i>Oct4</i> regulatory elements, although the importance of this is unclear; induces TDG-mediated demethylation at the <i>mir200</i> cluster, which is necessary for MET during fibroblast reprogramming	Doerge <i>et al</i> (2012); Costa <i>et al</i> , (2013); Gao <i>et al</i> (2013); Hu <i>et al</i> (2014)
	PARP1	Functions in the regulation of 5mC; promotes Oct4 accessibility to <i>Nanog</i> and <i>Esrrb</i> loci	Doerge <i>et al</i> (2012)
	Dnmt3a/b	Dispensable for nuclear reprogramming of somatic cells to pluripotent state	Pawlak and Jaenisch (2011)

Increased DNase I sensitivity upon forced expression of TH2A and TH2B and the synergistic effect of P-Npm suggests that the enhancement of somatic cell reprogramming occurs through the induction of an open chromatin structure (Shinagawa *et al*, 2014).

Similarly, histone variant H3.3 counteracts linker histone H1-mediated chromatin compaction, keeping diverse genomic sites in an open chromatin conformation (Braunschweig *et al*, 2009). H3.3 incorporation into donor nuclei is required for successful somatic cell nuclear transfer (SCNT) (Nashun *et al*, 2011; Jullien *et al*, 2012; Wen *et al*, 2014), and down-regulation of histone H3.3 in mouse oocytes leads to compromised reprogramming efficiency (Wen *et al*, 2014). This appears to parallel the *in vivo* situation, where, following fertilisation, the selective incorporation of H3.3 into the paternal genome by the H3.3-specific histone chaperone Hira is essential for its de-condensation (Inoue & Zhang, 2014; Lin *et al*, 2014), and loss of H3.3 leads to over-condensation during early embryonic development (Lin *et al*, 2013). While Hira-mediated H3.3 deposition is required for proper establishment of H3K27me3 at the promoters of developmentally regulated genes in embryonic stem cells, depletion of H3.3 or Hira has only minor transcriptional effects (Banaszynski *et al*, 2013). The role of these factors during induction of pluripotency remains largely unknown.

In comparison with TH2A, TH2B, and H3.3, macroH2A, with its unique macro-domain, is associated with a repressive chromatin state. In agreement with the open chromatin structure found in pluripotent cells, the pluripotent state is associated with low macroH2A levels that increase following cell differentiation (Creppe *et al*, 2012). MacroH2A is abundant in differentiated somatic cells, but disassociates immediately from somatic donor chromosomes during SCNT (Chang *et al*, 2010). Recent studies indicated that macroH2A acts as an epigenetic barrier to induced pluripotency: the absence of this particular histone variant enhances iPSC reprogramming up to 25-fold (Pasque *et al*, 2012), while its overexpression

prevents efficient reprogramming of epiblast stem cells to naïve pluripotency (Pasque *et al*, 2012; Barrero *et al*, 2013). It has additionally been shown that macroH2A and H3K27me3 co-occupy the regulatory regions of pluripotency genes in somatic cells (Barrero *et al*, 2013; Gaspar-Maia *et al*, 2013). Although iPSCs induced in the absence of this histone variant are able to differentiate, they retain the ability to return to a stem cell-like state (Gaspar-Maia *et al*, 2013) likely due to the incomplete inactivation of pluripotent genes during differentiation (Creppe *et al*, 2012).

Recent reports have shed new light on a possible role of another H2A histone variant in the reprogramming process. Ectopic expression of reprogramming factors increases the level of phosphorylated histone H2A.X, and high basal levels of  $\gamma$ -H2A.X have been observed in both iPSCs and ESCs, decreasing upon differentiation (Banath *et al*, 2009; Turinetto *et al*, 2012). Depletion of H2A.X reduces the efficiency of iPSC derivation (Wu *et al*, 2014) and compromises self-renewal activity in ES cells (Turinetto *et al*, 2012). Although typically associated with the DNA damage response, high  $\gamma$ -H2A.X levels do not correlate with elevated levels of DNA damage response proteins (Turinetto *et al*, 2012). Thus, while these recent findings suggest that they play an important role during reprogramming, the exact mechanism by which H2A.X or its phosphorylated form ( $\gamma$ -H2A.X) contribute to self-renewal and iPSC reprogramming requires further investigation.

### Changes in histone post-translational modification linked to the reprogramming process

#### *Early iPSC reprogramming is marked by rapid acquisition of active post-translational histone modifications*

Rapid genome-wide changes of H3K4me2 distribution are one of the earliest events observed in the initial phase of reprogramming

(Koche *et al.*, 2011). H3K4me2 peaks exhibit dramatic changes at promoter and enhancer regions of more than a thousand genes, including both pluripotency-related and developmentally regulated loci. As positive H3K4me2 changes are observed on both pluripotent and developmentally regulated genes (including those expressed in MEFs), the observed initial histone modification changes thus likely predominantly reflect chromatin accessibility. Interestingly (and in line with above), H3K4me2 is targeted to the pluripotency-associated genes before their transcriptional activation. Wdr5, the key component of Set/MLL histone methyltransferase complex responsible for H3K4 methylation, has been shown to directly interact with Oct4 (Ang *et al.*, 2011) and promoters gaining H3K4me2 are significantly enriched for targets of Oct4 and Sox2 (Koche *et al.*, 2011). This interaction thus possibly explains the rapid acquisition of H3K4 methylation early during iPSC reprogramming at loci bound by ectopic Oct4. Consistently, Wdr5 is required not only for ES cell self-renewal but also for efficient reprogramming of somatic cells to pluripotency (Ang *et al.*, 2011).

#### *Erasure and remodelling of repressive histone modifications*

Although the initial observed epigenetic changes during the reprogramming process are connected with the acquisition of transcriptionally permissive histone marks (see above), the cumulative evidence suggests that it is the erasure and remodelling of repressive histone modifications that constitute the true barrier to the reprogramming process.

**H3K9me2/3** In stark contrast to H3K4me3-containing regions, broad chromatin domains enriched for repressive H3K9me3 are refractory to initial OSKM binding (Soufi *et al.*, 2012). Reduction of H3K9me3 levels through down-regulation of methyltransferases Suv39H1&2 enhances Oct4 and Sox2 binding at these regions and increases reprogramming efficiency (Onder *et al.*, 2012; Soufi *et al.*, 2012). Consequently, H3K9me3-marked broad heterochromatin regions are considered as an epigenetic barrier during somatic cell reprogramming (Soufi *et al.*, 2012; Chen *et al.*, 2013b; Sridharan *et al.*, 2013). In support of this, a recent publication has also documented an inhibitory role for H3K9me3 during reprogramming by SCNT (Matoba *et al.*, 2014). It should be however noted that the role of H3K9me3 in iPSC generation is context dependent, as the downregulation of Setdb1, another H3K9me3 methyltransferase, has been reported to both facilitate and impede reprogramming (Onder *et al.*, 2012; Chen *et al.*, 2013b). In this context, it has been argued that H3K9me3 is important for silencing of lineage-specific genes; consistently, Setdb1 has been shown to repress trophoblast-specific genes in ES cells (Yeap *et al.*, 2009; Yuan *et al.*, 2009).

Next to H3K9me3, reduction of H3K9me2 through knockdown or inhibition of G9a methyltransferase also favours somatic cell reprogramming both in transcription factor- and in cell fusion-based reprogramming systems (Ma *et al.*, 2008; Shi *et al.*, 2008; Chen *et al.*, 2013b; Sridharan *et al.*, 2013). Contrary to G9a, the role of Ehmt1/Glp (a binding partner of G9a) during iPSC generation remains controversial (Onder *et al.*, 2012; Sridharan *et al.*, 2013).

**H3K27me3 and PRC2** Large regions of metazoan chromatin containing developmentally regulated genes are silenced by H3K27me3 catalysed by polycomb repressive complex-2 (PRC2). In agreement

with the necessity to remove somatic heterochromatin patterns, loss of H3K27me3 is observed during the earliest stages of reprogramming yielding a transient open/primed chromatin state (Koche *et al.*, 2011; Hussein *et al.*, 2014; Tonge *et al.*, 2014). The removal of this repressive histone mark is likely mediated by Utx, a JmjC-domain-containing enzyme that specifically mediates H3K27me2/3 demethylation (Klose *et al.*, 2006). The importance of Utx in iPSC reprogramming is highlighted by greatly reduced reprogramming efficiency and aberrant global H3K27me3 and H3K4me3 epigenetic profiles in the iPSC cells generated from Utx-depleted mouse embryonic fibroblasts (Mansour *et al.*, 2012). Furthermore, complete absence of Utx completely abolishes the ability of somatic cells to be reprogrammed back to the ground state of pluripotency (Mansour *et al.*, 2012). While Utx physically interacts with OSK reprogramming factors and removes the repressive mark from pluripotency-promoting genes such as *Fgf4*, *Sall4*, and *Sall1*, Utx overexpression does not increase the efficiency of iPSC formation, suggesting that it does not represent a rate-limiting factor in the process (Mansour *et al.*, 2012). In contrast to Utx, Jmjd3 (Kdm6b), another histone H3K27me3 demethylase, negatively regulates somatic cell reprogramming, highlighting the locus specificity and partially non-overlapping functions of these enzymes. Depletion of Jmjd3 is thought to reduce cell senescence by inhibiting Ink4a/Arf expression through maintenance of H3K27me3 levels at its promoter, enhancing both the kinetics and efficiency of reprogramming (Zhao *et al.*, 2013).

Although the somatic pattern of H3K27me3 needs to be erased during the iPSC reprogramming, global loss of H3K27me3 through down-regulation of Eed (resulting in loss of all PRC2 complexes) leads to a severe decline in the efficiency of iPSC reprogramming (Fragola *et al.*, 2013); thus, silencing through H3K27me3 appears indispensable for the establishment of iPSCs (Fragola *et al.*, 2013). Consistent with this idea, overexpression of the PRC2 catalytic subunit Ezh2 enhances reprogramming efficiency (Buganim *et al.*, 2012); down-regulation of other PRC2 complex components (Suz12 and Eed) significantly hinders iPSC generation (Onder *et al.*, 2012); and additional subunits of PRC2 in mouse ES cells (Jarid2, Mtf2 and esPRC2p48) act synergistically to enhance OSK (Oct4/Sox2/Klf4)-mediated mouse embryonic fibroblast reprogramming (Zhang *et al.*, 2011). Moreover, components of the PRC1 complex, Ring1a and Bmi1, are also required for efficient reprogramming, with the combined overexpression of Bmi1 and Oct4 sufficient to induce iPSCs from mouse fibroblasts (Moon *et al.*, 2011). Finally, and of note, PRC1 subunit Ring1b and PRC2 subunit Ezh2 are also required for ES cells to efficiently reprogramme somatic cells in cell fusion-based systems (Pereira *et al.*, 2010).

#### *Epigenetic changes linked to the memory of active transcriptional state*

One of the key steps during the reprogramming process is the extinction of the initial somatic transcriptional programme. Although the expression of somatic genes is typically down-regulated early during reprogramming (Brambrink *et al.*, 2008; Stadtfeld *et al.*, 2008; Polo *et al.*, 2012), stalled reprogramming intermediates often show incomplete silencing of the somatic programme suggesting that the maintenance of the original gene expression profile constitutes one of the hurdles in the reprogramming process. From the chromatin point of view, actively transcribed genes are characterised by

Set2-mediated H3K36me2/3 and Dot11-mediated H3K79me2 histone modification marks present in gene bodies (Nguyen & Zhang, 2011; Venkatesh *et al*, 2012). Down-regulation of either of these modifications by knockdown of the relevant histone methyltransferase (Onder *et al*, 2012) or by an overexpression of the relevant histone demethylase (Wang *et al*, 2011; Liang *et al*, 2012) prior to the iPSC reprogramming significantly enhances the reprogramming process. On the molecular level, removal of these histone marks leads to the efficient down-regulation of the original somatic transcription profile, thus promoting cell fate change.

### DNA modifications: 5-methylcytosine, 5-hydroxymethylcytosine and higher oxidative products during reprogramming

#### *Faithful reprogramming requires establishment of the pluripotent methylome*

In contrast to the high levels of DNA methylation consistently observed in somatic cells, DNA methylation levels are low in the naïve pluripotent cells both *in vivo* (Mayer *et al*, 2000; Oswald *et al*, 2000; Smith *et al*, 2012, 2014; Guo *et al*, 2014; Wang *et al*, 2014c) and *in vitro* (Ficz *et al*, 2013; Habibi *et al*, 2013; Leitch *et al*, 2013; Takashima *et al*, 2014). Considering this, it has been suggested that global DNA demethylation is a conserved and required feature of reprogramming events (Hill *et al*, 2014).

The functional relationship between faithful transcriptome and methylome reprogramming in iPSC and SCNT experimental systems has been recently shown by Mitalipov and colleagues (Ma *et al*, 2014). Using genetically matched starting somatic cells, the authors of this study used whole-genome bisulphite sequencing and RNA sequencing to extensively compare the DNA methylomes and transcriptomes of iPSC lines, ES cell lines generated through SCNT, and ES cell lines generated through traditional *in vitro* fertilisation (IVF) (Ma *et al*, 2014). They observed that both the DNA methylome and transcriptome of SCNT-derived ES cell lines, where the somatic nucleus was exposed to the cytoplasm of the host oocyte, were highly similar to that of ES cell lines derived through IVF. In contrast, iPSC reprogramming, involving only the ectopic expression of reprogramming TFs, generated cell lines with both significant differences in gene expression and high numbers of aberrantly methylated regions (Ma *et al*, 2014). The study revealed a strong correlation between incomplete reactivation of gene expression during reprogramming and high promoter methylation in iPSCs (Ma *et al*, 2014), suggesting that incomplete demethylation during iPSC reprogramming may be responsible for the observed incomplete transcriptional reprogramming.

#### *DNA methyltransferase activity inhibits reprogramming efficiency*

Studies investigating the direct relationship between DNA methylation and reprogramming efficiency have revealed that inhibition of global DNA methyltransferase activity through addition of 5-azadC to growth medium or targeted knockdown of the maintenance DNA methyltransferase Dnmt1 greatly increases reprogramming efficiency (Mikkelsen *et al*, 2008) (Fig 2 and Table 1). In contrast, the reprogramming potential of somatic cells depleted of the *de novo* methyltransferases Dnmt3a and Dnmt3b appeared largely unaffected (Pawlak & Jaenisch, 2011). These observations

suggest that, while maintenance of the somatic methylome is a barrier that must be overcome, *de novo* deposition of methylation is not a requirement for successful iPSC reprogramming. In fact, the observation that the large majority of differentially methylated regions (DMRs) between iPSCs and IVF ES cells do not overlap DMRs between donor somatic cells and IVF ES cells suggest that *de novo* DNA methylation may potentially contribute to the aberrant transcriptional profiles observed in iPSCs (Ma *et al*, 2014). Combined, these observations suggest that faithful reprogramming of the methylome may be a rate-limiting step to successful cell reprogramming. Consistent with such a model, deposition of H3K4me2 in the earliest stages of iPSC reprogramming only occurs at promoters that are already hypomethylated in the somatic nucleus, while acquisition of H3K4me2 at hypermethylated somatic promoters appears to be an event restricted to late stages of iPSC reprogramming, presumably following DNA demethylation at these regions (Koche *et al*, 2011).

#### *Context-specific requirement for Tet enzymes and oxidation of 5-methylcytosine oxidation in reprogramming*

The recently discovered Tet family of oxygenases (Tet1-3), which catalyse the oxidation of 5-methylcytosine (5-mC) to 5-hydroxymethylcytosine (5-hmC), 5-formylcytosine (5-fC), and 5-carboxycytosine (5-caC) through iterative rounds of oxidation (Tahiliani *et al*, 2009; He *et al*, 2011; Ito *et al*, 2011), have been implicated in reprogramming processes *in vivo* and *in vitro*, as we have recently reviewed (Hill *et al*, 2014).

In the context of iPSC generation, Tet proteins were originally identified as key mediators of reprogramming, as depletion of Tet1 and Tet2 resulted in significantly reduced efficiency of iPSC colony formation (Doegel *et al*, 2012; Chen *et al*, 2013a; Costa *et al*, 2013; Gao *et al*, 2013; Hu *et al*, 2014). While it was originally suggested that Tet1-mediated 5-hmC formation was required for Oct4 reactivation through demethylation of Oct4 regulatory elements (Gao *et al*, 2013), later studies were unable to reproduce these results (Hu *et al*, 2014). Further investigation revealed that Tet proteins are only necessary for somatic cells to undergo the mesenchymal-to-epithelial transition (MET) during iPSC reprogramming (Hu *et al*, 2014). Tet1-3 triple knockout somatic cells of epithelial morphology, and fibroblasts acutely depleted of all three Tet proteins only following MET, could both be efficiently reprogrammed to iPSCs (Hu *et al*, 2014). Further characterisation of the involvement of Tet proteins during MET revealed that Tet2 mediates the oxidation of 5mC at the MET-regulating *mir200* microRNA cluster, resulting in DNA demethylation and expression of the relevant microRNAs (Hu *et al*, 2014).

By comparison, the role of Tet proteins and 5-hmC in other *in vitro* reprogramming systems is less well characterised. A requirement for Tet2 has been described for reactivation of the somatic pluripotency-associated genes *Oct4*, *Nanog*, and *Cripto* during cell fusion experiments, although the mechanism by which this is achieved, and whether this is dependent on 5-hmC formation, is still unclear (Piccolo *et al*, 2013). Similarly, oocyte-derived Tet3 has been implicated in demethylation and reactivation of the somatic *Oct4* promoter in SCNT experiments (Gu *et al*, 2011). More experimental work remains to be done to understand the relative importance of Tet proteins and 5-mC oxidation for both cell fusion and SCNT reprogramming systems.



### DNA replication and cell division: a window of permissive chromatin?


Considering the stability of heterochromatin and its restrictive role in the reprogramming process, it is important to consider that cells undergo dynamic cell cycle-associated chromatin changes with the existing chromatin structure disrupted by passage of the replication fork during S phase (MacAlpine & Almouzni, 2013). In view of this, studies investigating the effect of cell cycle and cell division on reprogramming can provide additional functional insights into the role of chromatin structure in the reprogramming process. Using cells in distinct stages of the cell cycle, Boiani and colleagues and Fisher and colleagues definitively identified DNA synthesis in the somatic nucleus as an essential requirement for reprogramming in both SCNT (Wang *et al*, 2014a) and cell fusion (Tsubouchi *et al*, 2013) experimental systems. Consistently, early analysis of iPSC reprogramming mechanisms revealed that increased cell division rates achieved through down-regulation of the p53/p21 pathway or over-expression of Lin28 markedly accelerated reprogramming (Hanna *et al*, 2009), suggesting that increased frequency of cell cycling is associated with accelerated iPSC reprogramming.

It remains to be fully understood why DNA synthesis is a prerequisite for reprogramming by cell fusion or SCNT (Tsubouchi *et al*, 2013; Wang *et al*, 2014a), or why accelerated cell division decreases the latency time of iPSC reprogramming (Hanna *et al*, 2009). One potential hypothesis considers the complex nature of chromatin replication during S phase. As mentioned above, an immediate consequence of DNA replication (and consequently cell division) is the disruption of the existing chromatin structure by passage of the replication fork (Alabert & Groth, 2012; MacAlpine & Almouzni, 2013). For faithful re-establishment of the parental epigenome, a link must exist between the DNA replication fork and the factors that propagate DNA modifications, histone modifications, the correct incorporation of histone variants and other non-histone chromatin proteins (Alabert & Groth, 2012; MacAlpine & Almouzni, 2013). Although the mechanisms for maintenance of DNA methylation patterns are relatively well understood, the abundance of

histone modifications seems to fluctuate with progression through the cell cycle (Bonenfant *et al*, 2007). It is thus conceivable that chromatin changes associated with S phase can provide a window of opportunity for the ectopic TFs to bind their response elements. Additionally, over the course of a number of cell divisions, minor stochastic disruptions to the epigenetic inheritance could result in additional loss of epigenetic memory. In the context of differentiated cell states with robust transcriptional networks, minor disruptions would not likely result in overt phenotypic changes. However, upon exposure of the somatic nucleus to the pluripotent TFs (either through ectopic expression of the iPSC reprogramming factors, cell fusion, or SCNT), errors in the maintenance of epigenetic information and aberrant DNA accessibility may facilitate the recruitment of pioneer factors to regions normally recalcitrant to their binding (Soufi *et al*, 2012), or, more generally, of non-pioneer pluripotency-associated TFs to their DNA targets (Sherwood *et al*, 2014).

### Conclusions: is a permissive chromatin template sufficient for reprogramming in the absence of ectopic expression of instructive transcription factors?

The model whereby the presence of cell type-specific TFs is the central mechanism by which cell fate is determined, and chromatin structure simply regulates the probability that TFs bind their genomic targets, suggests that reprogramming to pluripotency can only be induced when the somatic nucleus is exposed to the pluripotent TFs (through ectopic iPSC factor expression, cell fusion, or SCNT). According to this model, simple disruption of the underlying chromatin structure would be insufficient to drive reprogramming alone. Remarkably, however, it has recently been shown that full reprogramming of somatic cells can be achieved in the absence of forced TF overexpression through chemical manipulation of signalling pathways and epigenetic modifiers alone (Hou *et al*, 2013). It should be noted that reprogramming in the absence of instructive TFs is likely only possible when cells are being reprogrammed back to pluripotency, as trans-differentiation would ultimately depend on



**Figure 3. Is transition through a state characterised by open, dynamic chromatin a pre-requisite for all cell fate transitions?**

(A) Reprogramming to pluripotency appears to require increased chromatin plasticity. (B, C) Possible relationship between chromatin dynamics and trans-differentiation: (B) trans-differentiation via an upstream progenitor may be connected with a transient increase in chromatin permissiveness, and/or (C) direct trans-differentiation between two somatic states without transition through an intermediary state characterised by more plastic chromatin may be possible, although this has yet to be experimentally validated.

the presence of lineage specifying TFs. Additionally, the presence of specific culture conditions (e.g. agonists or antagonists of specific signalling pathways) potentially compensates in part for the absence of ectopic OSKM expression by providing a selective 'environment' during the reprogramming process. Nevertheless, with these caveats in mind, the ability to reprogramme somatic cells in the absence of instructive TFs clearly shows that it is possible to induce cell fate reversal by synergistically destabilising the chromatin template and the existing transcriptional network (using inhibitors of signalling pathways). Although most of our current understanding regarding chromatin dynamics during reprogramming stems from reprogramming back to pluripotency using the iPSC reprogramming system, the reached conclusions seem relevant also for trans-differentiation experiments, where manipulation of chromatin accessibility/dynamics might be an important factor to consider next to the establishment of the relevant gene regulatory network. In this context, trans-differentiation studies have documented that reprogramming somatic cells directly to a cell of another somatic fate involves de-differentiation and passage through a less differentiated progenitor state (Xie *et al*, 2004) (Fig 3). While this progenitor state is likely to be characterised by a more plastic chromatin configuration, whether transitioning through a more open dynamic chromatin state during the intermediary steps of trans-differentiation is an absolute prerequisite for cell fate change, or whether this appears to be a unique requirement for reprogramming to pluripotency, will have to be addressed in future work (Fig 3).

### Acknowledgements

We would like to apologise to all colleagues whose work could not be cited due to the space constraints. We are grateful to all members of Hajkova lab for stimulating discussions. The work in Hajkova lab is funded by the MRC programme grant (MC\_US\_A652\_5PY70) to P.H. and by EpigeneSys FP7 EU network. P.H. is a member of the EMBO Young Investigator programme. B.N. is a recipient of the Marie Curie International Incoming Fellowship.

### Conflict of interest

The authors declare that they have no conflict of interest.

### References

- Adachi K, Scholer HR (2012) Directing reprogramming to pluripotency by transcription factors. *Curr Opin Genet Dev* 22: 416–422
- Alabert C, Groth A (2012) Chromatin replication and epigenome maintenance. *Nat Rev Mol Cell Biol* 13: 153–167
- Ancelin K, Lange UC, Hajkova P, Schneider R, Bannister AJ, Kouzarides T, Surani MA (2006) Blimp1 associates with Prmt5 and directs histone arginine methylation in mouse germ cells. *Nat Cell Biol* 8: 623–630
- Ang YS, Tsai SY, Lee DF, Monk J, Su J, Ratnakumar K, Ding J, Ge Y, Darr H, Chang B, Wang J, Rendl M, Bernstein E, Schaniel C, Lemischka IR (2011) Wdr5 mediates self-renewal and reprogramming via the embryonic stem cell core transcriptional network. *Cell* 145: 183–197
- Apostolou E, Hochedlinger K (2013) Chromatin dynamics during cellular reprogramming. *Nature* 502: 462–471
- Banaszynski LA, Wen D, Dewell S, Whitcomb SJ, Lin M, Diaz N, Elsasser SJ, Chapgier A, Goldberg AD, Canaan E, Rafii S, Zheng D, Allis CD (2013) Hira-dependent histone H3.3 deposition facilitates PRC2 recruitment at developmental loci in ES cells. *Cell* 155: 107–120
- Banath JP, Banuelos CA, Klokov D, MacPhail SM, Lansdorp PM, Olive PL (2009) Explanation for excessive DNA single-strand breaks and endogenous repair foci in pluripotent mouse embryonic stem cells. *Exp Cell Res* 315: 1505–1520
- Barrero MJ, Sese B, Kuebler B, Bilic J, Boue S, Marti M, Izpisua Belmonte JC (2013) Macrohistone variants preserve cell identity by preventing the gain of H3K4me2 during reprogramming to pluripotency. *Cell Rep* 3: 1005–1011
- Bonenfant D, Towbin H, Coulot M, Schindler P, Mueller DR, van Oostrum J (2007) Analysis of dynamic changes in post-translational modifications of human histones during cell cycle by mass spectrometry. *Mol Cell Proteomics* 6: 1917–1932
- Brambrink T, Foreman R, Welstead GG, Lengner CJ, Wernig M, Suh H, Jaenisch R (2008) Sequential expression of pluripotency markers during direct reprogramming of mouse somatic cells. *Cell Stem Cell* 2: 151–159
- Braunschweig U, Hogan GJ, Pagie L, van Steensel B (2009) Histone H1 binding is inhibited by histone variant H3.3. *EMBO J* 28: 3635–3645
- Buganim Y, Faddah DA, Cheng AW, Itskovich E, Markoulaki S, Ganz K, Klemm SL, van Oudenaarden A, Jaenisch R (2012) Single-cell expression analyses during cellular reprogramming reveal an early stochastic and a late hierarchic phase. *Cell* 150: 1209–1222
- Burgess RJ, Zhang Z (2013) Histone chaperones in nucleosome assembly and human disease. *Nat Struct Mol Biol* 20: 14–22
- Cahan P, Li H, Morris SA, Lummertz da Rocha E, Daley GQ, Collins JJ (2014) Cell Net: network biology applied to stem cell engineering. *Cell* 158: 903–915
- Chang CC, Gao S, Sung LY, Corry GN, Ma Y, Nagy ZP, Tian XC, Rasmussen TP (2010) Rapid elimination of the histone variant MacroH2A from somatic cell heterochromatin after nuclear transfer. *Cell Reprogram* 12: 43–53
- Chen J, Guo L, Zhang L, Wu H, Yang J, Liu H, Wang X, Hu X, Gu T, Zhou Z, Liu J, Liu J, Wu H, Mao SQ, Mo K, Li Y, Lai K, Qi J, Yao H, Pan G *et al* (2013a) Vitamin C modulates TET1 function during somatic cell reprogramming. *Nat Genet* 45: 1504–1509
- Chen J, Liu H, Liu J, Qi J, Wei B, Yang J, Liang H, Chen Y, Chen J, Wu Y, Guo L, Zhu J, Zhao X, Peng T, Zhang Y, Chen S, Li X, Li D, Wang T, Pei D (2013b) H3K9 methylation is a barrier during somatic cell reprogramming into iPSCs. *Nat Genet* 45: 34–42
- Costa Y, Ding J, Theunissen TW, Faiola F, Hore TA, Shliaha PV, Fidalgo M, Saunders A, Lawrence M, Dietmann S, Das S, Levasseur DN, Li Z, Xu M, Reik W, Silva JC, Wang J (2013) NANOG-dependent function of TET1 and TET2 in establishment of pluripotency. *Nature* 495: 370–374
- Creppe C, Janich P, Cantarino N, Noguera M, Valero V, Musulen E, Douet J, Posavec M, Martin-Caballero J, Sumoy L, Di Croce L, Benitah SA, Buschbeck M (2012) MacroH2A1 regulates the balance between self-renewal and differentiation commitment in embryonic and adult stem cells. *Mol Cell Biol* 32: 1442–1452
- Davidson EH, Erwin DH (2006) Gene regulatory networks and the evolution of animal body plans. *Science* 311: 796–800
- Davis RL, Weintraub H, Lassar AB (1987) Expression of a single transfected cDNA converts fibroblasts to myoblasts. *Cell* 51: 987–1000
- Doegge CA, Inoue K, Yamashita T, Rhee DB, Travis S, Fujita R, Guarnieri P, Bhagat G, Vanti WB, Shih A, Levine RL, Nik S, Chen EI, Abeliovich A (2012) Early-stage epigenetic modification during somatic cell reprogramming by Parp1 and Tet2. *Nature* 488: 652–655
- Drouin J (2014) Minireview: pioneer transcription factors in cell fate specification. *Mol Endocrinol* 28: 989–998
- Feldmann A, Ivanek R, Murr R, Gaidatzis D, Burger L, Schubeler D (2013) Transcription factor occupancy can mediate active turnover of DNA methylation at regulatory regions. *PLoS Genet* 9: e1003994
- Ficz G, Hore TA, Santos F, Lee HJ, Dean W, Arand J, Krueger F, Oxley D, Paul YL, Walter J, Cook SJ, Andrews S, Branco MR, Reik W (2013) FGF signaling

- inhibition in ESCs drives rapid genome-wide demethylation to the epigenetic ground state of pluripotency. *Cell Stem Cell* 13: 351–359
- Fragola G, Germain PL, Laise P, Cuomo A, Blasimme A, Gross F, Signaroldi E, Bucci G, Sommer C, Pruneri G, Mazzarol G, Bonaldi T, Mostoslavsky G, Casola S, Testa G (2013) Cell reprogramming requires silencing of a core subset of polycomb targets. *PLoS Genet* 9: e1003292
- Gao YW, Chen JY, Li K, Wu T, Huang B, Liu WQ, Kou XC, Zhang Y, Huang H, Jiang YH, Yao C, Liu XL, Lu ZW, Xu ZJ, Kang L, Chen J, Wang HL, Cai T, Gao SR (2013) Replacement of Oct4 by Tet1 during iPSC induction reveals an important role of DNA methylation and hydroxymethylation in reprogramming. *Cell Stem Cell* 12: 453–469
- Gaspar-Maia A, Alajem A, Polesso F, Sridharan R, Mason MJ, Heidersbach A, Ramalho-Santos J, McManus MT, Plath K, Meshorer E, Ramalho-Santos M (2009) Chd1 regulates open chromatin and pluripotency of embryonic stem cells. *Nature* 460: 863–868
- Gaspar-Maia A, Qadeer ZA, Hasson D, Ratnakumar K, Leu NA, Leroy G, Liu S, Costanzi C, Valle-Garcia D, Schaniel C, Lemischka I, Garcia B, Pehrson JR, Bernstein E (2013) MacroH2A histone variants act as a barrier upon reprogramming towards pluripotency. *Nat Commun* 4: 1565
- Gifford CA, Ziller MJ, Gu HC, Trapnell C, Donaghey J, Tsankov A, Shalek AK, Kelley DR, Shishkin AA, Issner R, Zhang XL, Coyne M, Fostel JL, Holmes L, Meldrim J, Guttman M, Epstein C, Park H, Kohlbacher O, Rinn J et al (2013) Transcriptional and epigenetic dynamics during specification of human embryonic stem cells. *Cell* 153: 1149–1163
- Goldberg AD, Allis CD, Bernstein E (2007) Epigenetics: a landscape takes shape. *Cell* 128: 635–638
- Gonzalez-Munoz E, Arboleda-Estudillo Y, Otu HH, Cibelli JB (2014) Cell reprogramming. Histone chaperone ASF1A is required for maintenance of pluripotency and cellular reprogramming. *Science* 345: 822–825
- Graf T, Enver T (2009) Forcing cells to change lineages. *Nature* 462: 587–594
- Gu TP, Guo F, Yang H, Wu HP, Xu GF, Liu W, Xie ZG, Shi LY, He XY, Jin SG, Iqbal K, Shi YJG, Deng ZX, Szabo PE, Pfeifer GP, Li JS, Xu GL (2011) The role of Tet3 DNA dioxygenase in epigenetic reprogramming by oocytes. *Nature* 477: 606–610
- Guo H, Zhu P, Yan L, Li R, Hu B, Lian Y, Yan J, Ren X, Lin S, Li J, Jin X, Shi X, Liu P, Wang X, Wang W, Wei Y, Li X, Guo F, Wu X, Fan X et al (2014) The DNA methylation landscape of human early embryos. *Nature* 511: 606–610
- Gurdon JB, Elsdale TR, Fischberg M (1958) Sexually mature individuals of *Xenopus laevis* from the transplantation of single somatic nuclei. *Nature* 182: 64–65
- Gurdon JB (1960) The developmental capacity of nuclei taken from differentiating endoderm cells of *Xenopus laevis*. *J Embryol Exp Morphol* 8: 505–526
- Gurdon JB (1962) The developmental capacity of nuclei taken from intestinal epithelium cells of feeding tadpoles. *J Embryol Exp Morphol* 10: 622–640
- Habibi E, Brinkman AB, Arand J, Kroeze LI, Kerstens HH, Matarese F, Lepikhov K, Gut M, Brun-Heath I, Hubner NC, Benedetti R, Altucci L, Jansen JH, Walter J, Gut IG, Marks H, Stunnenberg HG (2013) Whole-genome bisulfite sequencing of two distinct interconvertible DNA methylomes of mouse embryonic stem cells. *Cell Stem Cell* 13: 360–369
- Hanna J, Saha K, Pando B, van Zon J, Lengner CJ, Creighton MP, van Oudenaarden A, Jaenisch R (2009) Direct cell reprogramming is a stochastic process amenable to acceleration. *Nature* 462: 595–601
- Hayakawa K, Ohgane J, Tanaka S, Yagi S, Shiota K (2012) Oocyte-specific linker histone H1foo is an epigenomic modulator that decondenses chromatin and impairs pluripotency. *Epigenetics* 7: 1029–1036
- He YF, Li BZ, Li Z, Liu P, Wang Y, Tang QY, Ding JP, Jia YY, Chen ZC, Li L, Sun Y, Li XX, Dai Q, Song CX, Zhang KL, He C, Xu GL (2011) Tet-Mediated Formation of 5-Carboxylcytosine and Its Excision by TDG in Mammalian DNA. *Science* 333: 1303–1307
- Hill PW, Amouroux R, Hajkova P (2014) DNA demethylation, Tet proteins and 5-hydroxymethylcytosine in epigenetic reprogramming: an emerging complex story. *Genomics* 104: 324–333
- Ho L, Jothi R, Ronan JL, Cui K, Zhao K, Crabtree GR (2009a) An embryonic stem cell chromatin remodeling complex, esBAF, is an essential component of the core pluripotency transcriptional network. *Proc Natl Acad Sci USA* 106: 5187–5191
- Ho L, Ronan JL, Wu J, Staahl BT, Chen L, Kuo A, Lessard J, Nesvizhskii AI, Ranish J, Crabtree GR (2009b) An embryonic stem cell chromatin remodeling complex, esBAF, is essential for embryonic stem cell self-renewal and pluripotency. *Proc Natl Acad Sci USA* 106: 5181–5186
- Ho L, Miller EL, Ronan JL, Ho WQ, Jothi R, Crabtree GR (2011) esBAF facilitates pluripotency by conditioning the genome for LIF/STAT3 signalling and by regulating polycomb function. *Nat Cell Biol* 13: 903–913
- Ho JW, Jung YL, Liu T, Alver BH, Lee S, Ikegami K, Sohn KA, Minoda A, Tolstorukov MY, Appert A, Parker SC, Gu T, Kundaje A, Riddle NC, Bishop E, Egelhofer TA, Hu SS, Alekseyenko AA, Rechtsteiner A, Asker D et al (2014) Comparative analysis of metazoan chromatin organization. *Nature* 512: 449–452
- Hou P, Li Y, Zhang X, Liu C, Guan J, Li H, Zhao T, Ye J, Yang W, Liu K, Ge J, Xu J, Zhang Q, Zhao Y, Deng H (2013) Pluripotent stem cells induced from mouse somatic cells by small-molecule compounds. *Science* 341: 651–654
- Hu X, Zhang L, Mao SQ, Li Z, Chen JK, Zhang RR, Wu HP, Gao J, Guo F, Liu W, Xu GF, Dai HQ, Shi YG, Li XL, Hu BQ, Tang FC, Pei DQ, Xu GL (2014) Tet and TDG mediate DNA demethylation essential for mesenchymal-to-epithelial transition in somatic cell reprogramming. *Cell Stem Cell* 14: 512–522
- Hussein SM, Puri MC, Tonge PD, Benevento M, Corso AJ, Clancy JL, Mosbergen R, Li M, Lee DS, Cloonan N, Wood DL, Munoz J, Middleton R, Korn O, Patel HR, White CA, Shin JY, Gauthier ME, Le Cao KA, Kim JI et al (2014) Genome-wide characterization of the routes to pluripotency. *Nature* 516: 198–206
- Inoue A, Zhang Y (2014) Nucleosome assembly is required for nuclear pore complex assembly in mouse zygotes. *Nat Struct Mol Biol* 21: 609–616
- Ito S, Shen L, Dai Q, Wu SC, Collins LB, Swenberg JA, He C, Zhang Y (2011) Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-carboxylcytosine. *Science* 333: 1300–1303
- Jost D (2014) Bifurcation in epigenetics: implications in development, proliferation, and diseases. *Phys Rev E Stat Nonlin Soft Matter Phys* 89: 010701
- Jullien J, Astrand C, Szenker E, Garrett N, Almouzni G, Gurdon JB (2012) HIRA dependent H3.3 deposition is required for transcriptional reprogramming following nuclear transfer to *Xenopus* oocytes. *Epigenet Chromatin* 5: 17–26
- Kaji K, Caballero IM, MacLeod R, Nichols J, Wilson VA, Hendrich B (2006) The NuRD component Mbd3 is required for pluripotency of embryonic stem cells. *Nat Cell Biol* 8: 285–292
- Klose RJ, Kallin EM, Zhang Y (2006) JmjC-domain-containing proteins and histone demethylation. *Nat Rev Genet* 7: 715–727
- Koche RP, Smith ZD, Adli M, Gu H, Ku M, Gnirke A, Bernstein BE, Meissner A (2011) Reprogramming factor expression initiates widespread targeted chromatin remodeling. *Cell Stem Cell* 8: 96–105
- Leitch HG, McEwen KR, Turp A, Encheva V, Carroll T, Grabole N, Mansfield W, Nashun B, Knezovich JG, Smith A, Surani MA, Hajkova P (2013) Naive pluripotency is associated with global DNA hypomethylation. *Nat Struct Mol Biol* 20: 311–316
- Liang G, He J, Zhang Y (2012) Kdm2b promotes induced pluripotent stem cell generation by facilitating gene activation early in reprogramming. *Nat Cell Biol* 14: 457–466

- Lin CJ, Conti M, Ramalho-Santos M (2013) Histone variant H3.3 maintains a decondensed chromatin state essential for mouse preimplantation development. *Development* 140: 3624–3634
- Lin CJ, Koh FM, Wong P, Conti M, Ramalho-Santos M (2014) Hira-mediated H3.3 incorporation is required for DNA replication and ribosomal RNA transcription in the mouse zygote. *Dev Cell* 30: 268–279
- Lister R, Pelizzola M, Kida YS, Hawkins RD, Nery JR, Hon G, Antosiewicz-Bourget J, O'Malley R, Castanon R, Klugman S, Downes M, Yu R, Stewart R, Ren B, Thomson JA, Evans RM, Ecker JR (2011) Hotspots of aberrant epigenomic reprogramming in human induced pluripotent stem cells. *Nature* 471: 68–73
- Luo M, Ling T, Xie W, Sun H, Zhou Y, Zhu Q, Shen M, Zong L, Lyu G, Zhao Y, Ye T, Gu J, Tao W, Lu Z, Grummt I (2013) NuRD blocks reprogramming of mouse somatic cells into pluripotent stem cells. *Stem Cells* 31: 1278–1286
- Ma DK, Chiang CH, Ponnusamy K, Ming GL, Song H (2008) G9a and Jhdm2a regulate embryonic stem cell fusion-induced reprogramming of adult neural stem cells. *Stem Cells* 26: 2131–2141
- Ma H, Morey R, O'Neil RC, He Y, Daughtry B, Schultz MD, Hariharan M, Nery JR, Castanon R, Sabatini K, Thiagarajan RD, Tachibana M, Kang E, Tippner-Hedges R, Ahmed R, Gutierrez NM, Van Dyken C, Polat A, Sugawara A, Sparman M et al (2014) Abnormalities in human pluripotent cells due to reprogramming mechanisms. *Nature* 511: 177–183
- MacAlpine DM, Almouzni G (2013) Chromatin and DNA replication. *Cold Spring Harb Perspect Biol* 5: a010207
- Magnani L, Eeckhoutte J, Lupien M (2011) Pioneer factors: directing transcriptional regulators within the chromatin environment. *Trends Genet* 27: 465–474
- Mal A, Harter ML (2003) MyoD is functionally linked to the silencing of a muscle-specific regulatory gene prior to skeletal myogenesis. *Proc Natl Acad Sci USA* 100: 1735–1739
- Mansour AA, Gafni O, Weinberger L, Zviran A, Ayyash M, Rais Y, Krupalnik V, Zerbib M, Amann-Zalcenstein D, Maza I, Geula S, Viukov S, Holtzman L, Pribluda A, Canaani E, Horn-Saban S, Amit I, Novershtern N, Hanna JH (2012) The H3K27 demethylase Utx regulates somatic and germ cell epigenetic reprogramming. *Nature* 488: 409–413
- Matoba S, Liu Y, Lu F, Iwabuchi Kumiko A, Shen L, Inoue A, Zhang Y (2014) Embryonic Development following Somatic Cell Nuclear Transfer Impeded by Persisting Histone Methylation. *Cell* 159: 884–895
- Mayer W, Niveleau A, Walter J, Fundele R, Haaf T (2000) Demethylation of the zygotic paternal genome. *Nature* 403: 501–502
- Meshorer E, Yellajoshula D, George E, Scambler PJ, Brown DT, Misteli T (2006) Hyperdynamic plasticity of chromatin proteins in pluripotent embryonic stem cells. *Dev Cell* 10: 105–116
- Mikkelsen TS, Hanna J, Zhang X, Ku M, Wernig M, Schorderet P, Bernstein BE, Jaenisch R, Lander ES, Meissner A (2008) Dissecting direct reprogramming through integrative genomic analysis. *Nature* 454: 49–55
- Miller RA, Ruddle FH (1976) Pluripotent teratocarcinoma-thymus somatic cell hybrids. *Cell* 9: 45–55
- Montellier E, Boussovar F, Rousseaux S, Zhang K, Buchou T, Fenaille F, Shiota H, Debernardi A, Hery P, Curtet S, Jamshidikia M, Barral S, Holota H, Bergon A, Lopez F, Guardiola P, Pernet K, Imbert J, Petosa C, Tan M et al (2013) Chromatin-to-nucleoprotamine transition is controlled by the histone H2B variant TH2B. *Genes Dev* 27: 1680–1692
- Moon JH, Heo JS, Kim JS, Jun EK, Lee JH, Kim A, Kim J, Whang KY, Kang YK, Yeo S, Lim HJ, Han DW, Kim DW, Oh S, Yoon BS, Scholer HR, You S (2011) Reprogramming fibroblasts into induced pluripotent stem cells with Bmi1. *Cell Res* 21: 1305–1315
- Morris SA, Cahan P, Li H, Zhao AM, San Roman AK, Shivdasani RA, Collins JJ, Daley GQ (2014) Dissecting engineered cell types and enhancing cell fate conversion via Cell Net. *Cell* 158: 889–902
- Nashun B, Akiyama T, Suzuki MG, Aoki F (2011) Dramatic replacement of histone variants during genome remodeling in nuclear-transferred embryos. *Epigenetics* 6: 1489–1497
- Nguyen AT, Zhang Y (2011) The diverse functions of Dot1 and H3K79 methylation. *Genes Dev* 25: 1345–1358
- Niwa H (2014) The pluripotency transcription factor network at work in reprogramming. *Curr Opin Genet Dev* 28C: 25–31
- Onder TT, Kara N, Cherry A, Sinha AU, Zhu N, Bernt KM, Cahan P, Marcarci BO, Unteraehrer J, Gupta PB, Lander ES, Armstrong SA, Daley GQ (2012) Chromatin-modifying enzymes as modulators of reprogramming. *Nature* 483: 598–602
- Orkin SH, Hochedlinger K (2011) Chromatin connections to pluripotency and cellular reprogramming. *Cell* 145: 835–850
- Oswald J, Engemann S, Lane N, Mayer W, Olek A, Fundele R, Dean W, Reik W, Walter J (2000) Active demethylation of the paternal genome in the mouse zygote. *Curr Biol* 10: 475–478
- Pasque V, Radzishuskaya A, Gillich A, Halley-Stott RP, Panamarova M, Zernicka-Goetz M, Surani MA, Silva JC (2012) Histone variant macroH2A marks embryonic differentiation in vivo and acts as an epigenetic barrier to induced pluripotency. *J Cell Sci* 125: 6094–6104
- Pawlak M, Jaenisch R (2011) De novo DNA methylation by Dnmt3a and Dnmt3b is dispensable for nuclear reprogramming of somatic cells to a pluripotent state. *Genes Dev* 25: 1035–1040
- Pereira CF, Piccolo FM, Tsubouchi T, Sauer S, Ryan NK, Bruno L, Landeira D, Santos J, Banito A, Gil J, Koseki H, Merkenschlager M, Fisher AG (2010) ESCs require PRC2 to direct the successful reprogramming of differentiated cells toward pluripotency. *Cell Stem Cell* 6: 547–556
- Piccolo FM, Bagci H, Brown KE, Landeira D, Soza-Ried J, Feytout A, Mooijman D, Hajkova P, Leitch HG, Tada T, Kriaucionis S, Dawlaty MM, Jaenisch R, Merkenschlager M, Fisher AG (2013) Different roles for Tet1 and Tet2 proteins in reprogramming-mediated erasure of imprints induced by EGC fusion. *Mol Cell* 49: 1023–1033
- Polo JM, Anderssen E, Walsh RM, Schwarz BA, Nefzger CM, Lim SM, Borkent M, Apostolou E, Alaei S, Cloutier J, Bar-Nur O, Cheloufi S, Stadtfeld M, Figueroa ME, Robinton D, Natesan S, Melnick A, Zhu JF, Ramaswamy S, Hochedlinger K (2012) A Molecular roadmap of reprogramming somatic cells into iPS cells. *Cell* 151: 1617–1632
- Rais Y, Zviran A, Geula S, Gafni O, Chomsky E, Viukov S, Mansour AA, Caspi I, Krupalnik V, Zerbib M, Maza I, Mor N, Baran D, Weinberger L, Jaitin DA, Lara-Astiaso D, Blecher-Gonen R, Shipony Z, Mukamel Z, Hagai T et al (2013) Deterministic direct reprogramming of somatic cells to pluripotency. *Nature* 502: 65–70
- Reynolds N, Latos P, Hynes-Allen A, Loos R, Leaford D, O'Shaughnessy A, Mosaku O, Signolet J, Brennecke P, Kalkan T, Costello I, Humphreys P, Mansfield W, Nakagawa K, Strouboulis J, Behrens A, Bertone P, Hendrich B (2012) NuRD suppresses pluripotency gene expression to promote transcriptional heterogeneity and lineage commitment. *Cell Stem Cell* 10: 583–594
- dos Santos RL, Tosti L, Radzishuskaya A, Caballero IM, Kaji K, Hendrich B, Silva JC (2014) MBD3/NuRD facilitates induction of pluripotency in a context-dependent manner. *Cell Stem Cell* 15: 102–110
- Sherwood RI, Hashimoto T, O'Donnell CW, Lewis S, Barkal AA, van Hoff JP, Karun V, Jaakkola T, Gifford DK (2014) Discovery of directional and nondirectional pioneer transcription factors by modeling DNase profile magnitude and shape. *Nat Biotechnol* 32: 171–178



- Shi Y, Do JT, Despons C, Hahm HS, Scholer HR, Ding S (2008) A combined chemical and genetic approach for the generation of induced pluripotent stem cells. *Cell Stem Cell* 2: 525–528
- Shinagawa T, Takagi T, Tsukamoto D, Tomaru C, Huynh LM, Sivaraman P, Kumarevel T, Inoue K, Nakato R, Katou Y, Sado T, Takahashi S, Ogura A, Shirahige K, Ishii S (2014) Histone variants enriched in oocytes enhance reprogramming to induced pluripotent stem cells. *Cell Stem Cell* 14: 217–227
- Shipony Z, Mukamel Z, Cohen NM, Landan G, Chomsky E, Zelig SR, Fried YC, Ainbinder E, Friedman N, Tanay A (2014) Dynamic and static maintenance of epigenetic memory in pluripotent and somatic cells. *Nature* 513: 115
- Singhal N, Graumann J, Wu G, Arauzo-Bravo MJ, Han DW, Greber B, Gentile L, Mann M, Scholer HR (2010) Chromatin-Remodeling Components of the BAF Complex Facilitate Reprogramming. *Cell* 141: 943–955
- Smith ZD, Chan MM, Mikkelsen TS, Gu H, Gnirke A, Regev A, Meissner A (2012) A unique regulatory phase of DNA methylation in the early mammalian embryo. *Nature* 484: 339–344
- Smith ZD, Chan MM, Humm KC, Karnik R, Mekhoubad S, Regev A, Eggan K, Meissner A (2014) DNA methylation dynamics of the human preimplantation embryo. *Nature* 511: 611–615
- Soufi A, Donahue G, Zaret KS (2012) Facilitators and impediments of the pluripotency reprogramming factors' initial engagement with the genome. *Cell* 151: 994–1004
- Sridharan R, Gonzales-Cope M, Chronis C, Bonora G, McKee R, Huang C, Patel S, Lopez D, Mishra N, Pellegrini M, Carey M, Garcia BA, Plath K (2013) Proteomic and genomic approaches reveal critical functions of H3K9 methylation and heterochromatin protein-1gamma in reprogramming to pluripotency. *Nat Cell Biol* 15: 872–882
- Stadler MB, Murr R, Burger L, Ivanek R, Lienert F, Scholer A, Wirbelauer C, Oakeley EJ, Gaidatzis D, Tiwari VK, Schubeler D (2011) DNA-binding factors shape the mouse methylome at distal regulatory regions. *Nature* 480: 490–495
- Stadtfeld M, Maherali N, Breault DT, Hochedlinger K (2008) Defining molecular cornerstones during fibroblast to iPS cell reprogramming in mouse. *Cell Stem Cell* 2: 230–240
- Tahiliani M, Koh KP, Shen YH, Pastor WA, Bandukwala H, Brudno Y, Agarwal S, Iyer LM, Liu DR, Aravind L, Rao A (2009) Conversion of 5-Methylcytosine to 5-Hydroxymethylcytosine in Mammalian DNA by MLL Partner TET1. *Science* 324: 930–935
- Takahashi K, Yamanaka S (2006) Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 126: 663–676
- Takahashi Y, Guo G, Loos R, Nichols J, Ficiz G, Krueger F, Oxley D, Santos F, Clarke J, Mansfield W, Reik W, Bertone P, Smith A (2014) Resetting transcription factor control circuitry toward ground-state pluripotency in human. *Cell* 158: 1254–1269
- Tongue PD, Corso AJ, Monetti C, Hussein SM, Puri MC, Michael IP, Li M, Lee DS, Mar JC, Cloonan N, Wood DL, Gauthier ME, Korn O, Clancy JL, Preiss T, Grimmond SM, Shin JY, Seo JS, Wells CA, Rogers IM et al (2014) Divergent reprogramming routes lead to alternative stem-cell states. *Nature* 516: 192–197
- Tsubouchi T, Soza-Ried J, Brown K, Piccolo FM, Cantone I, Landeira D, Bagci H, Hochegger H, Merckenschlager M, Fisher AG (2013) DNA synthesis is required for reprogramming mediated by stem cell fusion. *Cell* 152: 873–883
- Turinetto V, Orlando L, Sanchez-Ripoll Y, Kumpfmüller B, Storm MP, Porcedda P, Minieri V, Saviozzi S, Accomasso L, Cibrario Rocchietti E, Moorwood K, Circosta P, Cignetti A, Welham MJ, Giachino C (2012) High basal gammaH2AX levels sustain self-renewal of mouse embryonic and induced pluripotent stem cells. *Stem Cells* 30: 1414–1423
- Venkatesh S, Smolle M, Li H, Gogol MM, Saint M, Kumar S, Natarajan K, Workman JL (2012) Set2 methylation of histone H3 lysine 36 suppresses histone exchange on transcribed genes. *Nature* 489: 452–455
- Voss TC, Hager GL (2014) Dynamic regulation of transcriptional states by chromatin and transcription factors. *Nat Rev Genet* 15: 69–81
- Wang T, Chen K, Zeng X, Yang J, Wu Y, Shi X, Qin B, Zeng L, Esteban MA, Pan G, Pei D (2011) The histone demethylases Jhd1a/1b enhance somatic cell reprogramming in a vitamin-C-dependent manner. *Cell Stem Cell* 9: 575–587
- Wang B, Pfeiffer MJ, Schwarzer C, Arauzo-Bravo MJ, Boiani M (2014a) DNA replication is an integral part of the mouse oocyte's reprogramming machinery. *PLoS One* 9: e97199
- Wang L, Du Y, Ward JM, Shimbo T, Lackford B, Zheng X, Miao YL, Zhou B, Han L, Fargo DC, Jothi R, Williams CJ, Wade PA, Hu G (2014b) INO80 facilitates pluripotency gene activation in embryonic stem cell self-renewal, reprogramming, and blastocyst development. *Cell Stem Cell* 14: 575–591
- Wang L, Zhang J, Duan J, Gao X, Zhu W, Lu X, Yang L, Zhang J, Li G, Ci W, Li W, Zhou Q, Aluru N, Tang F, He C, Huang X, Liu J (2014c) Programming and inheritance of parental DNA methylomes in mammals. *Cell* 157: 979–991
- Wen D, Banaszyński LA, Liu Y, Geng F, Noh KM, Xiang J, Elemento O, Rosenwaks Z, Allis CD, Rafii S (2014) Histone variant H3.3 is an essential maternal factor for oocyte reprogramming. *Proc Natl Acad Sci USA* 111: 7325–7330
- Wu T, Liu Y, Wen D, Tseng Z, Tahmasian M, Zhong M, Rafii S, Stadtfeld M, Hochedlinger K, Xiao A (2014) Histone variant H2AX deposition pattern serves as a functional epigenetic mark for distinguishing the developmental potentials of iPSCs. *Cell Stem Cell* 15: 281–294
- Xie HF, Ye M, Feng R, Graf T (2004) Stepwise reprogramming of B cells into macrophages. *Cell* 117: 663–676
- Xie W, Schultz MD, Lister R, Hou Z, Rajagopal N, Ray P, Whitaker JW, Tian S, Hawkins RD, Leung D, Yang H, Wang T, Lee AY, Swanson SA, Zhang J, Zhu Y, Kim A, Nery JR, Ulrich MA, Kuan S et al (2013) Epigenomic analysis of multilineage differentiation of human embryonic stem cells. *Cell* 153: 1134–1148
- Yeap LS, Hayashi K, Surani MA (2009) ERG-associated protein with SET domain (ESET)-Oct4 interaction regulates pluripotency and represses the trophoblast lineage. *Epigenetics Chromatin* 2: 12
- Yuan P, Han J, Guo G, Orlov YL, Huss M, Loh YH, Yaw LP, Robson P, Lim B, Ng HH (2009) Eset partners with Oct4 to restrict extraembryonic trophoblast lineage potential in embryonic stem cells. *Genes Dev* 23: 2507–2520
- Zaret KS, Carroll JS (2011) Pioneer transcription factors: establishing competence for gene expression. *Genes Dev* 25: 2227–2241
- Zhang Z, Jones A, Sun CW, Li C, Chang CW, Joo HY, Dai Q, Mysliwiec MR, Wu LC, Guo Y, Yang W, Liu K, Pawlik KM, Erdjument-Bromage H, Tempst P, Lee Y, Min J, Townes TM, Wang H (2011) PRC2 complexes with JARID2, MTF2, and esPRC2p48 in ES cells to modulate ES cell pluripotency and somatic cell reprogramming. *Stem Cells* 29: 229–240
- Zhao W, Li Q, Ayers S, Gu Y, Shi Z, Zhu Q, Chen Y, Wang HY, Wang RF (2013) Jmjd3 inhibits reprogramming by upregulating expression of INK4a/Arf and targeting PHF20 for ubiquitination. *Cell* 152: 1037–1050
- Zhu J, Adli M, Zou JY, Verstappen G, Coyne M, Zhang X, Durham T, Miri M, Deshpande V, De Jager PL, Bennett DA, Houmar J, Muoio DM, Onder TT, Camahort R, Cowan CA, Meissner A, Epstein CB, Shoresh N, Bernstein BE (2013) Genome-wide chromatin state transitions associated with developmental and environmental cues. *Cell* 152: 642–654



**License:** This is an open access article under the terms of the Creative Commons Attribution 4.0 License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.