

Draft Genome Sequence of *Serratia* sp. Strain DD3, Isolated from the Guts of *Daphnia magna*

Anja Poehlein,^a Heike M. Freese,^{b*} Rolf Daniel,^a Diliana D. Simeonova^b

Genomic and Applied Microbiology and Göttingen Genomics Laboratory, Georg-August University, Göttingen, Germany^a; Laboratory of Microbial Ecology, University of Konstanz, Constance, Germany^b

* Present address: Heike M. Freese, Leibniz-Institut DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany

We report the draft genome sequence of *Serratia* sp. strain DD3, a gammaproteobacterium from the family *Enterobacteriaceae*. It was isolated from homogenized guts of *Daphnia magna*. The genome size is 5,274 Mb.

Received 11 August 2014 Accepted 15 August 2014 Published 11 September 2014

Citation Poehlein A, Freese HM, Daniel R, Simeonova DD. 2014. Draft genome sequence of *Serratia* sp. strain DD3, isolated from the guts of *Daphnia magna*. *Genome Announc.* 2(5):e00903-14. doi:10.1128/genomeA.00903-14.

Copyright © 2014 Poehlein et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Diliana D. Simeonova, diliana.simeonova@uni-konstanz.de.

Serratia sp. strain DD3 was isolated from homogenized gut samples of cultured *Daphnia magna*, grown in filtered oligotrophic water of Lake Constance and fed with the green alga *Scenedesmus obliquus* (1). Like most *Serratia* spp., strain DD3 is a psychrotolerant facultative lithotroph, exhibiting an anaerobic lifestyle. *Serratia* spp. are found in a variety of environments such as freshwater, insects, and vertebrates, including humans, and on plant surfaces where they act as opportunistic pathogens (2). A driver for *Serratia* pathogenicity may be phosphate starvation (3–5). The phosphorus requirements are usually covered by phosphate and phosphate esters. Under phosphate starvation some bacterial species assimilate alternatively reduced inorganic and organic P compounds (organophosphonates) (6, 7). The latter group are utilized from bacteria as P, C, and N sources (8–14).

For sequencing, genomic DNA of *Serratia* sp. DD3 was isolated with the PurGene core kit B (Qiagen, Hilden, Germany). Extracted DNA was used to prepare shotgun libraries for Illumina sequencing as recommended by the manufacturer (Illumina, San Diego, CA, USA). Sequencing resulted in 4,000,318 paired-end Illumina reads (112 bp) and a coverage of 79.94. The initial hybrid *de novo* assembly was performed with Ray (15) software and resulted in 123 contigs. The genome of *Serratia* sp. strain DD3 consists of a single circular chromosome. The overall G+C content was 49.16 mol%. YACOP and GLIMMER (16) software tools were used for automatic gene prediction, while RNAmmer and tRNAscan were used for identification of rRNA and tRNA genes, respectively (17, 18). The functional annotation of the protein-coding genes was carried out with the Integrated Microbial Genomes/Expert Review (IMG/ER) system (19). A total of 4,666 putative genes were identified, of which 4,595 were protein encoding. Overall, 3,958 (84.83%) of the open reading frames were assigned to functions. One complete rRNA cluster out of 23 rRNA genes and 62 tRNAs, including those for selenocysteine incorporation, were identified.

Most probably, the phosphate level regulation in strain DD3 proceeds via *phoU*, located in the genome of the strain. Phosphate is delivered into the cells through a low-affinity phosphate: sulfate

intake permease, a high-affinity Pst system (*pstSABC*), and a phosphate ester uptake systems (PTS), all found in the genome of the strain (20). Under phosphate starvation, strain DD3 assimilates phosphite or organophosphonates, but not hypophosphite or other reduced P compounds. It lacks the gene coding for phosphonate dehydrogenase (*ptxD*); therefore, phosphite presumably is oxidized from membrane-spanning alkaline phosphatase coupled to an [NiFe] hydrogenase, a process analogous to what occurs in *Escherichia coli* (13). The ability of DD3 to grow with phosphite as the single phosphorus source was proven with physiological experiments (D. D. Simeonova, unpublished data). Organophosphonate uptake and assimilation pathways identified were the 2-aminoethylphosphonate (2AEP) assimilation phosphonoacetaldehyde hydrolase-dependent (PhnWX) pathway (11), the phosphonoacetate hydrolase (PhnA) pathway (21, 22), and a complete C-P lyase operon (23), which allows the utilization of the broadest spectrum of organophosphonates, and phosphite.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [AYKS00000000](https://www.ncbi.nlm.nih.gov/nuccore/AYKS00000000). The version described in this paper is the second version, [AYKS20000000](https://www.ncbi.nlm.nih.gov/nuccore/AYKS20000000).

ACKNOWLEDGMENTS

We thank Sylke Wiechmann, Frauke-Dorothee Meyer, and Kathleen Gollnow for technical support.

This work was supported by the University of Konstanz. D.D.S. was supported partially by a grant of the Deutsche Forschungsgemeinschaft, Bonn–Bad Godesberg, Germany (SI 1300/4-1).

REFERENCES

- Lampert W. 1991. The dynamics of *Daphnia* in a shallow lake. *Verh. Int. Ver. Limnol.* 24:795–798.
- Grimont F, Grimont PAD. 2006. The genus *Serratia*, p 219–244. In Dworkin M, Falkow S, Rosenberg E, Schleifer KH, Stackebrandt E (ed), *Proteobacteria: gamma subclass*. Springer, New York, NY.
- Witney FR, Failla ML, Weinberg ED. 1977. Phosphate inhibition of secondary metabolism in *Serratia marcescens*. *Appl. Environ. Microbiol.* 33:1042–1046.
- Slater H, Crow M, Everson L, Salmond GP. 2003. Phosphate availability

- regulates biosynthesis of two antibiotics, prodigiosin and carbapenem, in *Serratia* via both quorum-sensing-dependent and -independent pathways. *Mol. Microbiol.* 47:303–320. <http://dx.doi.org/10.1046/j.1365-2958.2003.03295.x>.
5. Kuo PA, Kuo CH, Lai YK, Graumann PL, Tu J. 2013. Phosphate limitation induces the intergeneric inhibition of *Pseudomonas aeruginosa* by *Serratia marcescens* isolated from paper machines. *FEMS Microbiol. Ecol.* 84:577–587. <http://dx.doi.org/10.1111/1574-6941.12086>.
 6. Kokonova SV, Nesmeyanova MA. 2002. Phosphonates and their degradation by microorganisms. *Biochemistry (Mosc)* 67:184–195. <http://dx.doi.org/10.1023/A:1014409929875>.
 7. White AK, Metcalf WW. 2007. Microbial metabolism of reduced phosphorus compounds. *Annu. Rev. Microbiol.* 61:379–400. <http://dx.doi.org/10.1146/annurev.micro.61.080706.093357>.
 8. Metcalf WW, Wanner BL. 1991. Involvement of the *Escherichia coli* *phn* (*psiD*) gene cluster in assimilation of phosphorus in the form of phosphonates, phosphite, Pi esters, and Pi. *J. Bacteriol.* 173:587–600.
 9. Simeonova DD, Wilson MM, Metcalf WW, Schink B. 2010. Identification and heterologous expression of genes involved in anaerobic dissimilatory phosphite oxidation by *Desulfotignum phosphitoxidans*. *J. Bacteriol.* 192:5237–5244. <http://dx.doi.org/10.1128/JB.00541-10>.
 10. Villarreal-Chiu JF, Quinn JP, McGrath JW. 2012. The genes and enzymes of phosphonate metabolism by bacteria, and their distribution in the marine environment. *Front. Microbiol.* 3:19. <http://dx.doi.org/10.3389/fmicb.2012.00019>.
 11. Borisova SA, Christman HD, Metcalf ME, Zulkepli NA, Zhang JK, van der Donk WA, Metcalf WW. 2011. Genetic and biochemical characterization of a pathway for the degradation of 2-aminoethylphosphonate in *Sinorhizobium meliloti* 1021. *J. Biol. Chem.* 286:22283–22290. <http://dx.doi.org/10.1074/jbc.M111.237735>.
 12. Dumora C, Lacoste AM, Cassaigne A. 1989. Phosphonoacetaldehyde hydrolase from *Pseudomonas aeruginosa*: purification properties and comparison with *Bacillus cereus* enzyme. *Biochim. Biophys. Acta* 997: 193–198.
 13. Yang K, Metcalf WW. 2004. A new activity for an old enzyme: *Escherichia coli* bacterial alkaline phosphatase is a phosphite-dependent hydrogenase. *Proc. Natl. Acad. Sci. USA* 101:7919–7924. <http://dx.doi.org/10.1073/pnas.0400664101>.
 14. Poehlein A, Daniel R, Schink B, Simeonova DD. 2013. Life based on phosphite: a genome-guided analysis of *Desulfotignum phosphitoxidans*. *BMC Genomics* 14:753. <http://dx.doi.org/10.1186/1471-2164-14-753>.
 15. Boisvert S, Laviolette F, Corbeil J. 2010. Ray: simultaneous assembly of reads from a mix of high-throughput sequencing technologies. *J. Comput. Biol.* 17:1519–1533. <http://dx.doi.org/10.1089/cmb.2009.0238>.
 16. Tech M, Merkl R. 2003. YACOP: enhanced gene prediction obtained by a combination of existing methods. *In Silico Biol.* 3:441–451.
 17. Lagesen K, Hallin P, Rødland EA, Stærfeldt HH, Rognes T, Ussery DW. 2007. RNAMmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.
 18. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25: 0955–0964. <http://dx.doi.org/10.1093/nar/25.5.0955>.
 19. Markowitz VM, Mavromatis K, Ivanova NN, Chen IM, Chu K, Kyrpides NC. 2009. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 25:2271–2278. <http://dx.doi.org/10.1093/bioinformatics/btp393>.
 20. van Veen HW. 1997. Phosphate transport in prokaryotes: molecules, mediators and mechanisms. *Antonie van Leeuwenhoek* 72:299–315.
 21. Agarwal V, Borisova SA, Metcalf WW, van der Donk WA, Nair SK. 2011. Structural and mechanistic insights into C–P bond hydrolysis by phosphonoacetate hydrolase. *Chem. Biol.* 18:1230–1240. <http://dx.doi.org/10.1016/j.chembiol.2011.07.019>.
 22. Kulakova AN, Kulakov LA, Quinn PJ. 1997. Cloning of the phosphonoacetate hydrolase gene from *Pseudomonas fluorescens* 23F encoding a new type of carbon-phosphorus bond cleaving enzyme and its expression in *Escherichia coli* and *Pseudomonas putida*. *Gene* 195:49–53.
 23. White AK, Metcalf WW. 2004. Two C-P lyase operons in *Pseudomonas stutzeri* and their roles in the oxidation of phosphonates, phosphite, and hypophosphite. *J. Bacteriol.* 186:4730–4739. <http://dx.doi.org/10.1128/JB.186.14.4730-4739.2004>.