

CASE REPORT

Pott's puffy tumour in an immunosuppressed adult: case report and systematic review of literature

Ekta Paw^{1,*}, Chelsea TW Ong² and Venkat Vangaveti¹¹James Cook University, College of Medicine and Dentistry, Townsville, Queensland 4811, Australia and ²Mater Hospital Pimlico, Department of Surgery, Townsville, Queensland 4812, Australia*Correspondence address. College of Medicine and Dentistry, 1 James Cook Dr., Douglas, QLD 4811, Australia.
Tel: +61 408 719 233; E-mail: ektapaw@gmail.com; Twitter: @ektapaw

Abstract

This case is of a 34-year-old immunocompromised male with Pott's puffy tumour, which was treated with antibiotics and endoscopic sinus surgery. Pott's puffy tumour is typically a complication of frontal sinusitis in children and immunocompromise can be a precipitating factor. A search was conducted of Embase, Medline, CINAHL, Cochrane, Google Scholar, Informit and Scopus of 'Pott's Puffy' and variations of this phrase. Initial 804 records were reviewed by title, abstract and full text. Of the unique papers identified, 155 only included paediatric patients and 128 included adult patients. After title, abstract and full-text review of 13 papers were identified describing Pott's puffy tumour in immunocompromised patients. Immunosuppression was noted as a risk factor in only a small number of cases, with diabetes mellitus being the most common. A few cases did note immunosuppression due to newer immunomodulatory agents being used for treatment of autoimmune conditions.

INTRODUCTION

This case is of a 34-year-old immunosuppressed male who presented with Pott's puffy tumour as a complication of frontal sinusitis. Surgical incision and drainage was performed with endoscopic debridement of sinuses, in addition to treatment with antibiotics and nasal decongestants. This case highlights immunosuppression, particularly with tumour necrosis factor (TNF) inhibitors, as a potential risk factor for development of Pott's puffy tumour as a complication of frontal sinusitis.

CASE REPORT

A 34-year-old gentleman presented to the emergency department of a regional hospital with 2 days of left-sided periorbital swelling superior to the eye socket with associated erythema and tenderness. He was assessed to have normal visual acuity and

extraocular movements. He had a history of severe ankylosing spondylitis for which he was taking a TNF inhibitor, adalimumab. His blood tests showed a slightly raised C-reactive protein of 82 mg/l but otherwise no abnormality. Magnetic resonance imaging (MRI) was performed on the day of admission and noted significant left frontal sinusitis complicated by a defect of the frontal cortical bone as well as oedema and enhancement consistent with a Pott's puffy tumour (Fig. 1). There was also an abscess of the periorbital soft tissue. Immediate treatment was given in the form of phenylephrine nasal decongestants and intravenous antibiotics (amoxicillin with clavulanic acid), and the patient was transferred to a tertiary centre with ear, nose and throat surgery capacity.

Within 24 hours of arrival at the tertiary centre, an incision and drainage of the soft tissue abscess and periosteal abscess was performed. At the same time the sinuses were debrided

Received: November 8, 2020. Accepted: November 20, 2020

Published by Oxford University Press and JSCR Publishing Ltd. All rights reserved. © The Author(s) 2020.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited.For commercial re-use, please contact journals.permissions@oup.com




Figure 1: Sagittal and transverse of MRI Pott's puffy tumour




Figure 2: Intraoperative photos. Left: view of dura on endoscopic examination of frontal sinus via bony erosion. Right: complete bony occlusion of frontal recess

via endoscopic approach and a wide maxillary antrostomy and sphenoidectomy was performed. It was discovered that the frontal recess was completely occluded by a bony wall (Fig. 2). A cannula was placed into the frontal sinus for regular irrigation during the post-operative period with the other sinuses irrigated via nasal douche. Intravenous antibiotics and regular irrigation were continued for a further 5 days, and the TNF inhibitor was withheld on the advice of a rheumatologist. After this time the frontal sinus cannula was removed and the patient was discharged home. He was reviewed as an outpatient 6 days later and had a palpable soft tissue collection in the same area, which was promptly drained surgically and appeared to be a mucocele intraoperatively. None of the cultures identified a specific causative organism, although a Gram stain from the original admission showed gram negative bacilli. The patient was planned to have definitive surgery to open the bony occlusion of the frontal sinus and allow proper drainage. The patient was noted not to have any recurrence of symptoms on follow-up 1 year post-operatively.

DISCUSSION

Pott's puffy tumour is a rare clinical entity and is most commonly a complication of frontal sinusitis in children [1]. This case is

unique in that it presents in an adult with no traumatic history and immunosuppression as the only risk factor. TNF inhibitors do not appear to have a significant association with sinusitis but due to being immunosuppressants may slightly increase the risk of upper respiratory tract infections [2]. A literature review was conducted to investigate the role of immunosuppression in Pott's puffy tumour patients.

A search was conducted on 10 March 2020 of CINAHL, Cochrane Embase, Informit, Google Scholar (first 50 results only), Medline and Scopus of 'Pott's Puffy' and variations of this phrase. Initial 804 records were reviewed by title, abstract and full text (as detailed in Fig. 3) and narrowed down to 13 papers of adult Pott's puffy tumour where immunosuppression was noted in the case.

Of the unique papers identified, 155 only included paediatric patients, whereas 128 included adult patients. This would indicate that Pott's puffy tumour in adult patients is not quite as rare as previously thought. The paediatric cases were reviewed by abstract only for immunosuppression and only four were found. Two girls also on the same immunosuppressant medication as this patient (TNF inhibitor adalimumab) developed Pott's puffy tumour [3]. Another case of a 14 year old on an immunosuppressive study drug for juvenile idiopathic arthritis was published in the USA [4]. Finally one child with Type 2 diabetes mellitus in a case series was diagnosed with Pott's puffy tumour [1].

Table 1. Summary of all cases Pott's puffy tumour cases found with immunosuppression of patients

Author	Age	Sex	Presenting complaint	Immunocompromised	Intracranial spread	Pathogen	Surgical management	Complications
Adams <i>et al.</i> [5]	55	M	Pain, swelling, watery discharge	Diabetes mellitus	Left meningeal abscess	<i>Pseudomonas aeruginosa</i>	Bilateral frontal balloon catheter sinusotomy, bilateral complete ethmoidectomy, bilateral middle meatal antrostomy, right sphenoidectomy	Chronic ethmoid sinusitis and left orbital abscess requiring surgery
Akiyama <i>et al.</i> [6]	21-83	M + F		Diabetes, chronic renal failure, aplastic anaemia, breast cancer with bone metastasis	—	—	External (58.1% or endoscopic (32.9%) frontal sinus surgery, frontal bone debridement, craniotomy, simple percutaneous drainage (9.7%)	2 cases of postoperative recurrence
Chow and Szeto [7]	60	F	Headache, swelling, fever, peritonitis	End stage renal failure	—	<i>Mycobacterium tuberculosis</i>	—	—
Domville-Lewis <i>et al.</i> [8]	21	F	Headache, swelling, chemosis	Pregnancy	Right epidural empyema posterior to frontal sinus	<i>Streptococcus milleri</i>	Percutaneous drainage, unilateral uchelectomy, middle meatal antrostomy, anterior ethmoidectomy, frontal recess dissection	Caesarean section
Effat <i>et al.</i> [9]	62	F	Headache, swelling, fever, purulent discharge	Insulin-dependent diabetes, chronic renal failure on twice weekly haemodialysis	Erosion of lamina papyracea of left anterior and middle ethmoid cells and subperiosteal abscess	Mucorales	Cruciate incision and drainage, daily debridement	Fistula
Evliyaoglu <i>et al.</i> [10]	27	M	Pain headache, swelling, fever	Type 1 diabetes mellitus	Subperiosteal abscess in vertex region associated with dural thickening and irregularities of epidural space	—	Bicoronal skin incision under local, drainage of subperiosteal abscess, curette	—
Gil-Carcedo <i>et al.</i> [11]	48	M	Left hemiparesis, neck stiffness, right mydriasis, confusion, seizure	Chronic alcoholism	Epidural abscess, subdural empyema	Unclassified aerobic streptococcus	Right frontoparietal craniotomy, intraoral antrostomy and bilateral transantral ethmoidectomy, reconstruction with acrylic grafts	—
Goldfarb <i>et al.</i> [12]	58	M	Pain, swelling	Heavy smoking, diabetes mellitus	—	—	Incision and drainage	Fistula between frontal sinus and forehead skin
Husta and Reichner [13]	43	M	Altered mental status, proptosis	HIV, cocaine abuse	Subdural empyema, intracranial abscesses resulting in midline shift	<i>Streptococcus mitis</i>	Drainage of the right frontal subperiosteal abscess, left temporal craniotomy and evacuation of subdural empyema	Hypoxaemic respiratory failure requiring intubation, multiple bilateral cavitary lung abscess in bases
Lamoreau and Panciullo [14]	55	M	Left eye abscess, fever	Alcoholic cirrhosis, hepatitis C, nicotine, marijuana/other drug use	Left frontal bony erosion	Viridans like streptococci	Incision and drainage	Increased petit mal symptoms of known epilepsy
Miller [15]	60	M	Headache, swelling	Diabetes mellitus, prednisolone (wearing dose for allergy)	Erosion of anterior frontal sinus with posterior extension to meninges	<i>Streptococcus anginosus</i>	Spontaneously ruptured in hospital	—
Pasin <i>et al.</i> [16]	81	M	Painless swelling, weight loss, profound asthma	Diabetes mellitus	Erosion of left frontal bone and orbital roof	<i>Streptococcus constellatus</i>	Endoscopic sinus surgery	—
Sekine <i>et al.</i> [17]	56	M	Headache, recurrent forehead swelling, eyelid erythema	Rheumatoid arthritis, treatment (prednisolone, igitratimod, bucllamine)	—	<i>Stenotrophomonas maltophilia</i> , <i>Achromobacter xylosoxidans</i> and <i>Corynebacterium</i> spp.	Bicoronal incision and endoscopic approach for drainage. Anterolateral thigh flap reconstruction. Revision of forehead cavity with titanium plate later.	—




Figure 3: Flowchart of literature review

Of the 13 adult patients' papers comprising of both case reports and case series, 12 patients were immunosuppressed and 1 was pregnant (Table 1). The most commonly noted was diabetes in seven patients, and there was one patient with rheumatoid arthritis being treated with oral prednisolone, iguratimod and bucillamine.

Most Pott's puffy tumour cases appear to be in previously fit and well patients, so immunosuppression may play less of a role than neglect of symptoms and delayed clinical review. In this patient the location being several hours drive from the nearest hospital may be more relevant than their immunosuppression. Nonetheless, a higher index of suspicion for severe complications of sinusitis should still be considered for immunosuppressive patients.

ETHICS APPROVAL

LNR/QTHS/67480.

CONFLICT OF INTEREST STATEMENT

None declared.

REFERENCES

- Bambakidis NC, Cohen AR. Intracranial complications of frontal sinusitis in children: Pott's puffy tumor revisited. *Pediatr Neurosurg* 2001;35:82–9.
- MIMS Online. MIMS Online: Etanercept Full Product Information. Sydney: MIMS Australia; 2020. Available from: <https://www.mimsonline.com.au>.
- Christensen AE, Toftedal P. Sinogenic subdural empyema in two girls treated with adalimumab for chronic recurrent multifocal osteomyelitis (CRMO). *Pediatr Rheumatol* 2017; 15:28–9.
- Chandwani V. Pott's puffy tumour & epidural empyemas as complications of sinusitis in a teenager with jia. *J Investig Med* 2018;66:480.
- Adams J, Foster D, Stephenson L, Bourguillon P. Pott puffy tumor. *Consultant* 2017;57:251–3.
- Akiyama K, Karaki M, Mori N. Evaluation of adult Pott's puffy tumor: our five cases and 27 literature cases. *Laryngoscope* 2012;122:2382–8.
- Chow KM, Szeto CC. Headache caused by Pott's puffy tumor. *Headache* 2003;43:916.
- Domville-Lewis C, Friedland PL, Santa Maria PL. Pott's puffy tumour and intracranial complications of frontal sinusitis in pregnancy. *J Laryngol Otol* 2013;127:S35–S8.
- Effat KG, Karam M, El-Kabani A. Pott's puffy tumour caused by mucormycosis. *J Laryngol Otol* 2005;119:643–5.
- Evliyaoğlu Ç, Bademci G, Yucel E, Keskil S. Pott's puffy tumor of the vertex years after trauma in a diabetic patient: case report. *Neurocirugia* 2005;16:54–7.
- Gil-Carcedo LM, Izquierdo JM, Gonzalez M. Intracranial complications of frontal sinusitis. A report of two cases. *J Laryngol Otol* 1984;98:941–5.

12. Goldfarb A, Hocwald E, Gross M, Eliashar R. Frontal sinus cutaneous fistula: a complication of Pott's puffy tumor. *Otolaryngol Head Neck Surg* 2004;**130**:490–1.
13. Husta B, Reichner C. Pott's puffy tumor: a case with intracranial and pulmonary manifestation. *Chest* 2012;**142**: 167A.
14. Lamoreau KP, Fanciullo LM. Pott's puffy tumour mimicking preseptal cellulitis. *Clin Exp Optom* 2008;**91**:400–2.
15. Miller B. A 60-year-old man with forehead swelling. *Cleve Clin J Med* 2016;**83**:95–6.
16. Pasin F, Bonardi S, Modoni A. When the diagnosis is written on the forehead. *Eur J Internal Med* 2018;**47**:e1–2.
17. Sekine R, Omura K, Ishida K, Tanaka Y. Recurrent Pott's puffy tumor treated with anterior skull base resection with reconstruction of the anterolateral thigh flap. *J Craniofac Surg* 2019;**30**:e94–e6.