

Platelet function testing to predict hyporesponsiveness to clopidogrel in patients with chest pain seen in the emergency department

Rakesh K Sharma¹
Stephen W Erickson¹
Rohit Sharma²
Donald J Voelker¹
Hanumanth K Reddy¹
Harvinder Dod²
James D Marsh¹

¹Division of Cardiovascular Medicine, University of Arkansas for Medical Sciences, Little Rock, ²Medical Center of South Arkansas, El Dorado, AR, USA

Background: A dual antiplatelet regimen has been shown to reduce the risk of major adverse cardiovascular events after percutaneous coronary intervention. However, there is little information available on inhibition of platelet aggregation in patients with a prior coronary stent presenting with chest pain. This study evaluated the prevalence of hyporesponsiveness to clopidogrel and factors associated with this in patients presenting to our emergency department with chest pain who had previously undergone coronary stent placement and were prescribed dual antiplatelet therapy.

Methods: Responsiveness to clopidogrel was evaluated in a cohort of 533 consecutive stented patients presenting to the emergency department with chest pain. P2Y₁₂ reaction units (PRU) and percent P2Y₁₂ inhibition with clopidogrel were measured in all patients. Of 533 patients, 221 (41.6%) had PRU \geq 230. A multivariate logistic regression model was used to determine the relationship between hyporesponsiveness to clopidogrel (defined as PRU \geq 230) and several potential risk factors, ie, gender, age, race, type 1 or type 2 diabetes, hypertension, smoking, chronic renal failure, and obesity.

Results: There was a greater risk of hyporesponsiveness in African Americans than in non-African American patients (adjusted odds ratio [OR] = 2.165), in patients with type 2 diabetes than in those without (adjusted OR = 2.109), and in women than in men (adjusted OR = 1.813), as well as a greater risk of hyporesponsiveness with increasing age (adjusted OR = 1.167 per decade).

Conclusion: There was a high prevalence of hyporesponsiveness to clopidogrel in patients presenting with chest pain and a prior coronary stent. Non-insulin-dependent diabetes mellitus and African American race were the strongest predictors of hyporesponsiveness to clopidogrel, followed by gender and age.

Keywords: clopidogrel, platelet function testing, chest pain, emergency department

Introduction

Dual antiplatelet treatment has been shown to reduce the risk of major adverse cardiovascular events after percutaneous coronary intervention. Premature cessation of dual antiplatelet therapy, including aspirin and thienopyridine drugs, such as ticlopidine, clopidogrel, or prasugrel, and non-thienopyridines, such as ticagrelor, is associated with an increased risk for major cardiovascular events and stent thrombosis.¹ Dual antiplatelet therapy is recommended for 12 months after placement of a drug-eluting stent.²

Thienopyridine drugs, including ticlopidine, clopidogrel, and prasugrel, and the non-thienopyridine agent, ticagrelor, interact with P2Y₁₂, an adenosine diphosphate receptor. It is known that the pharmacodynamic response to clopidogrel

Correspondence: Rakesh K Sharma
Heart and Vascular Institute of South
Arkansas, El Dorado, AR 71730, USA
Tel +1 870 875 5540
Fax +1 870875 5568
Email rsharma@uams.edu

is less than desirable in 15%–40% of patients.^{3–6} The clinical significance of this variable response to antiplatelet agents has been debated in the medical community, and the US Food and Drug Administration has responded by issuing three sequential “boxed warnings” for clopidogrel.⁷ It is believed that a suboptimal response to an antiplatelet regimen may be associated with adverse cardiovascular and cerebrovascular outcomes.^{8,9}

During coronary intervention, there is a need for adequate P2Y₁₂ inhibition using P2Y₁₂ receptor blockers and inhibition of thromboxane production with aspirin in addition to antithrombotic regimens.¹⁰ Many trials have shown that poor inhibition of platelet aggregation may lead to a higher risk of stent thrombosis in such patients.^{8,11–13}

We report here the prevalence of hyporesponsiveness to clopidogrel assessed by platelet function assay in patients with chest pain who had previously received a coronary stent and presented in the community hospital setting. We now report the distinctive finding that a large proportion of such patients have inadequate inhibition of platelet aggregation.

Materials and methods

A platelet function assay was performed in all patients presenting to the emergency department of a community hospital with chest pain, a history of coronary artery stenting, and prescribed clopidogrel and aspirin. A cohort of 533 consecutive patients was studied prospectively, consistent with the clinical protocol of the hospital emergency department chest pain center. Characteristics of study patients are shown in Table 1.

The platelet function assay included measurement of platelet reactivity, reported as P2Y₁₂ reaction units of clopidogrel (PRU), baseline PRU (preclopidogrel), percent inhibition of P2Y₁₂, and aspirin reaction units in all patients who gave a history of taking aspirin and clopidogrel after coronary intervention. Hyporesponsiveness to clopidogrel is defined as a PRU > 230 or percent inhibition of P2Y₁₂ < 30%. The data were also analyzed using a hyporesponsiveness threshold of PRU > 208. Emergency department attendance records were reviewed for pertinent clinical factors. The protocol was approved by the institutional review board.

Platelet function assay

Assessment of platelet function was done using the commercially available VerifyNow[®] test (Accumetrics, San Diego, CA, USA) and has been described in detail elsewhere.¹⁴ Many trials, including GRAVITAS (Gauging Responsiveness with a VerifyNow Assay Impact on Thrombosis and Safety), have used a PRU > 230¹⁵ as the threshold for hyporesponsiveness.

Table 1 Patient characteristics stratified by clopidogrel hyporesponsiveness, with PRU = 230 as the cutoff

	PRU < 230	PRU ≥ 230	P value
Gender			
Male	187 (60.1%)	96 (43.2%)	<0.001
Female	124 (39.9%)	126 (56.8%)	
Age			
Mean (SD)	63.6 (12.3)	65.4 (13.2)	0.110
Race			
Caucasian	244 (78.5%)	142 (64.0%)	<0.001
African American	66 (21.2%)	75 (33.8%)	
Asian	1 (0.3%)	2 (0.9%)	
Other	0 (0.0%)	3 (1.4%)	
Type 1 diabetes			
Yes	6 (1.9%)	5 (2.3%)	1.000
No	304 (97.7%)	217 (97.7%)	
Unknown	1 (0.3%)	0 (0.0%)	
Type 2 diabetes			
Yes	76 (24.4%)	90 (40.5%)	<0.001
No	234 (75.2%)	131 (59.0%)	
Unknown	1 (0.3%)	1 (0.5%)	
Hypertension			
Yes	266 (85.5%)	194 (87.4%)	0.795
No	42 (13.5%)	28 (12.6%)	
Unknown	3 (1.0%)	0 (0.0%)	
Smoking			
Yes	136 (43.7%)	83 (37.4%)	0.153
No	173 (55.6%)	138 (62.2%)	
Unknown	2 (0.6%)	1 (0.5%)	
Chronic renal failure			
Yes	57 (18.3%)	48 (21.6%)	0.378
No	253 (81.4%)	174 (78.4%)	
Unknown	1 (0.3%)	0 (0.0%)	
Obesity			
Yes	133 (42.8%)	98 (44.1%)	0.790
No	177 (56.9%)	123 (55.4%)	
Unknown	1 (0.3%)	1 (0.5%)	

Notes: Fisher's Exact test (or Student's *t*-test for age) was used to test for dependence between hyporesponsiveness and each patient characteristic. There was a statistically significant bivariate association with gender, race, and type 2 diabetes, with the higher risk of hyporesponsiveness associated with female gender, African American race, and having type 2 diabetes. There is a marginally significant (*P* = 0.110) association with increasing age.

Abbreviations: PRU, P2Y₁₂ reaction units; SD, standard deviation.

The logic of this cutoff has been discussed previously.¹⁵ Furthermore, post hoc analysis of GRAVITAS using a PRU ≥ 208 was identified as a predictor of death, myocardial infarction, and stroke at one year, with fewer cardiovascular events in the GRAVITAS trial.¹⁶ Thus, our primary analysis was based on a PRU > 230 and the secondary analysis was done using PRU > 208.

The platelet function assay uses arachidonic acid, adenosine diphosphate (ADP), and thrombin receptor-activating peptide to assess platelet responsiveness to the P2Y₁₂ inhibitor, clopidogrel. The assay uses whole blood in a turbidometric assay and fibrinogen-coated beads to measure differences in platelet aggregation in response to various

agonists.¹⁷ Whole blood from a patient is exposed to 20 μmol ADP and 22 nmol prostaglandin E1. Prostaglandin E1 helps to increase the specificity of the P2Y12 assay by suppressing ADP-mediated P2Y1 platelet activation and aggregation. The patient's platelets agglutinate around the fibrinogen-coated beads with an increase in light transmittance. The results are reported as PRU.

The platelet function assay reported three values in this cohort of patients:

- PRU which represents the extent of inhibition of the P2Y12 ADP receptor by thienopyridines or non-thienopyridines, and the test takes advantage of different receptors on platelets stimulated by different agonists. P2Y12 subreceptors of ADP and thrombin receptors are strong platelet activators and function independently of each other.
- baseline PRU is calculated by stimulating thrombin receptors on platelets to estimate the total possible platelet aggregation in the preclopidogrel state.
- percent inhibition of P2Y12 ADP receptors is determined by the difference between baseline PRU and PRU determined after exposure to an ADP agonist; however, percent inhibition of P2Y12 may not be accurate, because

baseline PRU is not actually preclopidogrel PRU, but rather estimated by the thrombin agonist response while the patient is taking clopidogrel, and thrombin-induced platelet aggregation may not be an adequate reflection of baseline ADP-induced platelet aggregation.¹⁸

Results

A cohort of 533 consecutive patients presenting to the emergency department with chest pain and a history of being prescribed clopidogrel were studied. Two patients were excluded from analysis because they had no baseline PRU reading. Of 531 patients with baseline PRU readings, 221 (41.6%) had high platelet reactivity on treatment ($\text{PRU} \geq 230$) and 287 (54.0%) had percent P2Y12 inhibition $< 30\%$; hyporesponsiveness based on these two indicators was concordant in 439 patients (82.7%). A total of 268 patients (50.5%) had $\text{PRU} \geq 208$. Determination of hyporesponsiveness to clopidogrel based on $\text{PRU} \geq 208$ and percent P2Y12 inhibition $< 30\%$ thresholds was concordant in 456 patients (85.9%), as shown in Figure 1.

A multivariate logistic regression model was used to determine the relationship between hyporesponsiveness




Figure 1 Scatter plot of percent P2Y12 inhibition versus PRU.

Notes: Points to the right of the vertical red line are patients for whom PRU was ≥ 230 , while points below the horizontal red line are for patients in whom percent P2Y12 inhibition was < 30 . Points in the lower right quadrant ($n = 208$, 39.2%) are judged clopidogrel-hyporesponsive by both criteria, while points in the upper left quadrant ($n = 231$, 43.5%) are judged clopidogrel-responsive by both criteria. Points in the upper right and lower left quadrants ($n = 92$, 17.3%) have discordant judgments based on the two criteria. For comparison, a dotted gray line is drawn at the alternative PRU = 208 cutoff.

Abbreviation: PRU, P2Y12 reaction units.

to clopidogrel (PRU \geq 230) and several potential risk factors for variation in platelet function and ischemic heart disease. The results of this modeling are shown in Table 2. There are three predictors that are statistically significant at $P < 0.05$, and one (ie, age) at $P < 0.06$. Specifically, there was a greater risk of hyporesponsiveness in patients with type 2 diabetes than those without (adjusted odds ratio [OR] = 2.109), in African American compared with non-African American patients (adjusted OR = 2.165), and in women compared with men (adjusted OR = 1.813), and there was also a greater risk of hyporesponsiveness with increasing age (adjusted OR = 1.167 per decade). When adjusted for other risk factors, smoking was not a statistically significant predictor of hyporesponsiveness to clopidogrel.

Discussion

This study reports distinctive findings on the prevalence of inadequate platelet inhibition in patients with a coronary stent who are prescribed clopidogrel and return to a community hospital emergency department with chest pain. Depending on the definition of inadequate platelet inhibition used, 42%–54% of such patients had poor inhibition of platelet function. This is a remarkable failure of efficacy of a drug treatment strategy. Whether or not these patients prove to have an acute coronary syndrome at the time of the emergency department visit, or chest pain of other etiology, this determination of the efficacy of the current treatment strategy produces results that are concerning. The quantitative analysis of inadequate platelet inhibition varies somewhat, depending on whether percent inhibition or PRU is the metric, although concordance is reasonable (Figure 1).

Noncompliance with medication may have contributed to inadequate platelet inhibition. The incidence of

non adherence with dual antiplatelet therapy in the “real-world” international observation registry was 2% at 30-day follow-up.¹⁹ Nonetheless, the current study does provide a distinctive analysis of the incidence of adequate and inadequate platelet inhibition in a group of patients with coronary stents and chest pain, and such patients are commonly seen in emergency departments.

The individual biological response to clopidogrel is variable. Multiple studies support the hypothesis that this variable pharmacodynamic effect of clopidogrel may be responsible for atherothrombotic events following stent deployment.^{9,20–22} Diabetes and African American race were strong predictive factors. Hyporesponsiveness to clopidogrel can be secondary to several factors, including age, body mass index, diabetes, dyslipidemia, chronic renal disease, genetic polymorphism, and pharmacokinetic and pharmacodynamic interactions arising from concomitant use of other drugs.^{23,24} Failure to take the drug as prescribed certainly can produce hyporesponsiveness. Studies have found that platelets from diabetic patients are generally more reactive^{25,26} and less responsive to antiplatelet therapy.²⁷ Our findings are consistent with those observations. Several factors may explain this global hyper-reactive platelet in diabetes, such as insulin resistance, poor glycemic control, increased inflammatory status, increased response to ADP, increased reactivity on contact with collagen,²⁸ increased fibrinogen levels,²⁹ and increased production of epinephrine and thrombin receptor agonist peptide.^{28,30}

Mangicapra et al used the VerifyNow assay in diabetic patients undergoing percutaneous coronary intervention and found them to have higher platelet reactivity and worse periprocedural outcomes.³¹ Diabetic patients also have a poor response to aspirin (aspirin reaction units $>$ 550).³² In our study, patients with type 2 diabetes presenting with

Table 2 Multivariable logistic regression model of hyporesponsiveness to clopidogrel (PRU \geq 230)

Variable	Estimate	SE	Z value	P value	Adjusted OR	
					Estimate	95% CI
Intercept	−1.827	0.618	−2.957	0.0031*		
Gender (female)	0.595	0.191	3.113	0.0019*	1.813	1.248 2.641
Age (decades)	0.155	0.082	1.899	0.0575	1.167	0.996 1.372
Race (African American)	0.772	0.215	3.589	0.0003*	2.165	1.423 3.312
Type 1 diabetes	−0.109	0.692	−0.158	0.8747	0.897	0.212 3.432
Type 2 diabetes	0.746	0.206	3.618	2.97E-04*	2.109	1.410 3.166
Hypertension	−0.299	0.287	−1.041	0.2980	0.742	0.424 1.309
Smoking	−0.038	0.202	−0.190	0.8496	0.962	0.648 1.431
Chronic renal failure	0.110	0.233	0.475	0.6349	1.117	0.706 1.760
Obesity	0.015	0.194	0.076	0.9396	1.015	0.693 1.485

Notes: Parameter estimates, standard errors, P values ($H_0: \beta = 0$ versus nonzero alternative), and adjusted OR for each of nine predictors and 95% CI. * $P > 0.05$.

Abbreviations: CI, confidence interval; OR, odds ratio; SE, standard error; PRU, P2Y12 reaction units.

chest pain had a higher prevalence of hyporesponsiveness (adjusted OR = 2.1409), as shown in Table 2.

This analysis also shows a greater risk of hyporesponsiveness in African American patients than in non-African Americans (adjusted OR = 2.165). Collins et al³³ have shown the African American race to be an independent risk factor for developing stent thrombosis after placement of a drug-eluting stent. Despite higher compliance with clopidogrel in African American patients, they were nearly three times more likely to develop stent thrombosis than non-African Americans (1.71% versus 0.59%). However, no platelet function testing was done in these patients to probe any relationship between stent thrombosis and hyporesponsiveness despite higher compliance with clopidogrel. Higher stent thrombosis may be due to genetic polymorphism making African American patients less responsive to clopidogrel.

As suggested by this and other studies, mechanisms of hyporesponsiveness to clopidogrel, such as genetic polymorphism, may need to be assessed. Previous studies have demonstrated that carriers of the cytochrome P450 (CYP) 2C19 hypofunctioning allele may produce a factor leading to decreased clopidogrel metabolites, causing diminished platelet inhibition and an increased rate of cardiovascular events.^{34,35} Such polymorphism is seen more frequently in African American and East Asians than in Caucasians.³⁶ The CYP 2C19*2 allele is present in 13% of Caucasians, 18% of African Americans, and 29% in East Asians.³⁷ Rapid point-of-care genotyping at the bedside is evolving in clinical practice, which will help personalization of antiplatelet therapy.³⁸ While the importance of the platelet function assay is evolving, the platelet function test may be useful in high-risk patients, such as those with diabetes, left main stem stenosis, and diffuse atherosclerotic disease with complex coronary lesions undergoing percutaneous coronary intervention.³⁹

Limitations

Limitations of this study include its lack of clinical outcome data or biomarkers for ischemia. Because inclusion in the study was dependent on the patient stating that they were taking clopidogrel and aspirin, the level of actual, as opposed to stated, patient compliance with the antiplatelet regimen is not known. Therefore, this was not a study of drug efficacy but rather a study of platelet inhibition in the clinical syndrome of chest pain evaluated in the emergency department for patients who have been prescribed clopidogrel and aspirin.

Tailoring antiplatelet therapy according to platelet function testing

Use of the platelet function assay to guide antiplatelet therapy has received considerable attention. This study and others have identified clinical risk factors for inadequate platelet inhibition by clopidogrel. However, clinical profiling clearly has inadequate predictive value in itself, and a platelet function assay may add value. Results from the large ADAPT-DES (Assessment of Dual AntiPlatelet Therapy With Drug Eluting Stents) registry suggest that hyporesponsiveness to clopidogrel after implantation of a drug-eluting stent is an independent predictor of stent thrombosis and myocardial infarction.⁴⁰ A platelet function assay may help screen for hyporesponders and nonresponders, and could also identify patients in whom medication compliance may be the main issue.

There are several alternatives to clopidogrel for patients with a suboptimal response to antiplatelet therapy. Newer antiplatelet agents, such as prasugrel and ticagrelor, have demonstrated superiority in patients with an acute coronary syndrome.^{41,42} These new drugs are not affected by genetic variants of CYP 2C19, and are more effective, with a more predictable pharmacodynamic response, than the standard dose of clopidogrel. However, it is likely that clopidogrel will remain a valuable and widely used inhibitor of platelet function. This study demonstrates that the therapeutic goal, ie, platelet inhibition, assessed by an in vitro assay, is not achieved by a large proportion of patients prescribed clopidogrel to prevent coronary stent thrombosis. Ongoing trials will expand our understanding of dual antiplatelet regimens and the role of the platelet function assay. The use of in vitro testing may be useful to guide treatment and to identify hyporesponders or nonresponders.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Iakovou I, Schmidt T, Bonizzi E, et al. Incidence, predictors, and outcome of thrombosis after successful implantation of drug-eluting stents. *JAMA*. 2005;293(17):2126–2130.
2. Kushner FG, Hand M, Smith SC Jr, et al. 2009 focused updates: ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction (updating the 2004 guideline and 2007 focused update) and ACC/AHA/SCAI guidelines on percutaneous coronary intervention (updating the 2005 guideline and 2007 focused update) a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2009;54(23):2205–2241.
3. Serebruany VL, Steinhubl SR, Berger PB, Malinin AI, Bhatt DL, Topol EJ. Variability in platelet responsiveness to clopidogrel among 544 individuals. *J Am Coll Cardiol*. 2005;45(2):246–251.

4. Bonello L, Tantry US, Marcucci R, et al. Consensus and future directions on the definition of high on-treatment platelet reactivity to adenosine diphosphate. *J Am Coll Cardiol*. 2010;56(12):919–933.
5. Price MJ, Coleman JL, Steinhubl SR, Wong GB, Cannon CP, Teirstein PS. Onset and offset of platelet inhibition after high-dose clopidogrel loading and standard daily therapy measured by a point-of-care assay in healthy volunteers. *Am J Cardiol*. 2006;98(5):681–684.
6. Angiolillo DJ, Fernandez-Ortiz A, Bernardo E, et al. Variability in individual responsiveness to clopidogrel: clinical implications, management, and future perspectives. *J Am Coll Cardiol*. 2007;49(14):1505–1516.
7. Holmes DR Jr, Dehmer GJ, Kaul S, Leifer D, O’Gara PT, Stein CM. ACCF/AHA clopidogrel clinical alert: approaches to the FDA “boxed warning”: a report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents and the American Heart Association endorsed by the Society for Cardiovascular Angiography and Interventions and the Society of Thoracic Surgeons. *J Am Coll Cardiol*. 2010;56(4):321–341.
8. Gurbel PA, Bliden KP, Hiatt BL, O’Connor CM. Clopidogrel for coronary stenting: response variability, drug resistance, and the effect of pretreatment platelet reactivity. *Circulation*. 2003;107(23):2908–2913.
9. Matetzky S, Shenkman B, Guetta V, et al. Clopidogrel resistance is associated with increased risk of recurrent atherothrombotic events in patients with acute myocardial infarction. *Circulation*. 2004;109(25):3171–3175.
10. Anderson JL, Adams CD, Antman EM, et al. ACC/AHA 2007 guidelines for the management of patients with unstable angina/non-ST-Elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines for the Management of Patients With Unstable Angina/Non-ST-Elevation Myocardial Infarction) developed in collaboration with the American College of Emergency Physicians, the Society for Cardiovascular Angiography and Interventions, and the Society of Thoracic Surgeons endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation and the Society for Academic Emergency Medicine. *J Am Coll Cardiol*. 2007;50(7):e1–e157.
11. Cuisset T, Frere C, Quilici J, et al. High post-treatment platelet reactivity identified low-responders to dual antiplatelet therapy at increased risk of recurrent cardiovascular events after stenting for acute coronary syndrome. *J Thromb Haemost*. 2006;4(3):542–549.
12. Bonello L, Paganelli F, Arpin-Bornet M, et al. Vasodilator-stimulated phosphoprotein phosphorylation analysis prior to percutaneous coronary intervention for exclusion of postprocedural major adverse cardiovascular events. *J Thromb Haemost*. 2007;5(8):1630–1636.
13. Buonamici P, Marcucci R, Migliorini A, et al. Impact of platelet reactivity after clopidogrel administration on drug-eluting stent thrombosis. *J Am Coll Cardiol*. 2007;49(24):2312–2317.
14. Price MJ. Bedside evaluation of thienopyridine antiplatelet therapy. *Circulation*. 2009;119(19):2625–2632.
15. Price MJ, Berger PB, Angiolillo DJ, et al. Evaluation of individualized clopidogrel therapy after drug-eluting stent implantation in patients with high residual platelet reactivity: design and rationale of the GRAVITAS trial. *Am Heart J*. 2009;157(5):818–824.
16. Price MJ, Angiolillo DJ, Teirstein PS, et al. Platelet reactivity and cardiovascular outcomes after percutaneous coronary intervention: a time-dependent analysis of the Gauging Responsiveness with a VerifyNow P2Y12 assay: Impact on Thrombosis and Safety (GRAVITAS) trial. *Circulation*. 2011;124(10):1132–1137.
17. Malinin A, Pokov A, Swaim L, Kotob M, Serebruany V. Validation of a VerifyNow-P2Y12 cartridge for monitoring platelet inhibition with clopidogrel. *Methods Find Exp Clin Pharmacol*. 2006;28(5):315–322.
18. Lordkipanidze M, Pharand C, Nguyen TA, Schampaert E, Diodati JG. Assessment of VerifyNow P2Y12 assay accuracy in evaluating clopidogrel-induced platelet inhibition. *Ther Drug Monit*. 2008;30(3):372–378.
19. Mehran R. The PARIS Registry; Patterns of Non-Adherence to Anti-Platelet Regimens In Stented Patients: An Observational Single Arm Study. Proceedings of the 23rd Annual Transcatheter Cardiovascular Therapeutics Conference; 2011 Nov 7–11; San Francisco, CA, USA.
20. Gurbel PA, Bliden KP, Guyer K, et al. Platelet reactivity in patients and recurrent events post-stenting: results of the PREPARE post-stenting study. *J Am Coll Cardiol*. 2005;46(10):1820–1826.
21. Marcucci R, Gori AM, Panicia R, et al. Cardiovascular death and nonfatal myocardial infarction in acute coronary syndrome patients receiving coronary stenting are predicted by residual platelet reactivity to ADP detected by a point-of-care assay: a 12-month follow-up. *Circulation*. 2009;119(2):237–242.
22. Price MJ, Endemann S, Gollapudi RR, et al. Prognostic significance of post-clopidogrel platelet reactivity assessed by a point-of-care assay on thrombotic events after drug-eluting stent implantation. *Eur Heart J*. 2008;29(8):992–1000.
23. Sharma RK, Reddy HK, Sharma R, et al. The interaction between clopidogrel and proton pump inhibitors (PPI): is there any clinical relevance. *Clin Pharmacol*. 2011;2:155–162.
24. O’Donoghue ML, Braunwald E, Antman EM, et al. Pharmacodynamic effect and clinical efficacy of clopidogrel and prasugrel with or without a proton-pump inhibitor: an analysis of two randomised trials. *Lancet*. 2009;374(9694):989–997.
25. Leet H, Paton RC, Passa P, Caen JP. Fibrinogen binding and ADP-induced aggregation in platelets from diabetic subjects. *Thromb Res*. 1981;24(1–2):143–150.
26. Mandal S, Sarode R, Dash S, Dash RJ. Hyperaggregation of platelets detected by whole blood platelet aggregometry in newly diagnosed noninsulin-dependent diabetes mellitus. *Am J Clin Pathol*. 1993;100(2):103–107.
27. Angiolillo DJ, Bernardo E, Ramirez C, et al. Insulin therapy is associated with platelet dysfunction in patients with type 2 diabetes mellitus on dual oral antiplatelet treatment. *J Am Coll Cardiol*. 2006;48(2):298–304.
28. Ferreira IA, Mocking AI, Feijge MA, et al. Platelet inhibition by insulin is absent in type 2 diabetes mellitus. *Arterioscler Thromb Vasc Biol*. 2006;26(2):417–422.
29. Ang L, Palakodeti V, Khalid A, et al. Elevated plasma fibrinogen and diabetes mellitus are associated with lower inhibition of platelet reactivity with clopidogrel. *J Am Coll Cardiol*. 2008;52(13):1052–1059.
30. Angiolillo DJ, Bernardo E, Sabate M, et al. Impact of platelet reactivity on cardiovascular outcomes in patients with type 2 diabetes mellitus and coronary artery disease. *J Am Coll Cardiol*. 2007;50(16):1541–1547.
31. Mangiacapra F, Patti G, Peace A, et al. Comparison of platelet reactivity and periprocedural outcomes in patients with versus without diabetes mellitus and treated with clopidogrel and percutaneous coronary intervention. *Am J Cardiol*. 2010;106(5):619–623.
32. Mortensen SB, Larsen SB, Grove EL, Kristensen SD, Hvas AM. Reduced platelet response to aspirin in patients with coronary artery disease and type 2 diabetes mellitus. *Thromb Res*. 2010;126(4):e318–e322.
33. Collins SD, Torguson R, Gaglia MA Jr, et al. Does black ethnicity influence the development of stent thrombosis in the drug-eluting stent era? *Circulation*. 2010;122(11):1085–1090.
34. Mega JL, Close SL, Wiviott SD, et al. Cytochrome p-450 polymorphisms and response to clopidogrel. *N Engl J Med*. 2009;360(4):354–362.
35. Brandt JT, Close SL, Iturria SJ, et al. Common polymorphisms of CYP2C19 and CYP2C9 affect the pharmacokinetic and pharmacodynamic response to clopidogrel but not prasugrel. *J Thromb Haemost*. 2007;5(12):2429–2436.
36. Desta Z, Zhao X, Shin JG, Flockhart DA. Clinical significance of the cytochrome P450 2C19 genetic polymorphism. *Clin Pharmacokinet*. 2002;41(12):913–958.
37. Luo HR, Poland RE, Lin KM, Wan YJ. Genetic polymorphism of cytochrome P450 2C19 in Mexican Americans: a cross-ethnic comparative study. *Clin Pharmacol Ther*. 2006;80(1):33–40.

38. Roberts J, et al. Reassessment of Anti-platelet therapy Using Individualized Strategy Based on Genetic Evaluation (RAPID GENE). Proceedings of the 23rd Annual Transcatheter Cardiovascular Therapeutics Conference; 2011 Nov 7–11; San Francisco, CA, USA.
39. Sharma RK, Voelker DJ, Sharma R, Reddy HK, Dod H, Marsh JD. Evolving role of platelet function testing in coronary artery interventions. *Vasc Health Risk Manag.* 2012;8:65–75.
40. Stone GW, et al. Assessment of dual anti-platelet therapy with drug eluting stents (ADAPT-DES). Proceedings of the 23rd Annual Transcatheter Cardiovascular Therapeutics Conference; 2011 Nov 7–11; San Francisco, CA, USA.
41. Wiviott SD, Braunwald E, McCabe CH, et al. Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med.* 2007; 357(20):2001–2015.
42. Wallentin L, Becker RC, Budaj A, et al. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med.* 2009; 361(11):1045–1057.

Vascular Health and Risk Management

Publish your work in this journal

Vascular Health and Risk Management is an international, peer-reviewed journal of therapeutics and risk management, focusing on concise rapid reporting of clinical studies on the processes involved in the maintenance of vascular health; the monitoring, prevention and treatment of vascular disease and its sequelae; and the involvement of

metabolic disorders, particularly diabetes. This journal is indexed on PubMed Central and MedLine. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/vascular-health-and-risk-management-journal>

Dovepress