



Striking a balance when operating for acute type A aortic dissection

Viswajit Kandula¹^, Adham Elmously², Thomas F. X. O'Donnell¹, Virendra I. Patel¹, Hiroo Takayama¹

¹Division of Cardiac, Thoracic, and Vascular Surgery, NewYork-Presbyterian Hospital, Columbia University Irving Medical Center, New York, NY, USA; ²Department of Cardiovascular Surgery, Houston Methodist DeBakey Heart & Vascular Center, Houston Methodist Hospital, Houston, TX, USA

Correspondence to: Viswajit Kandula, MD. Division of Cardiac, Thoracic, and Vascular Surgery, NewYork-Presbyterian Hospital, Columbia University Irving Medical Center. Room 7-435, Milstein Hospital Building, 7th Floor, 177 Fort Washington Ave, New York, NY, USA. Email: vk2509@cumc.columbia.edu.

Comment on: Liu H, Zhang YY, Ding XH, *et al.* Proximal vs Extensive Repair in Acute Type A Aortic Dissection Surgery. *Ann Thorac Surg* 2023;116:270-8.

Keywords: Type A dissection; proximal repair; extensive repair

Submitted Dec 31, 2023. Accepted for publication Apr 24, 2024. Published online May 27, 2024.

doi: 10.21037/jtd-23-1945

View this article at: <https://dx.doi.org/10.21037/jtd-23-1945>

Acute type A aortic dissection (ATAAD) continues to be associated with high morbidity and mortality (1,2). These patients, if they don't die before arriving at a hospital, often suffer from concomitant life-threatening pathology such as malperfusion syndromes (including stroke, acute myocardial infarction, renal/visceral ischemia and limb ischemia), aortic regurgitation, or cardiac tamponade. Though surgical intervention, the gold standard treatment, has reduced mortality compared to non-operative management (3,4), the optimal type and extent of repair continue to be debated and remain dependent on the surgeon and surgical center a patient is brought to. Depending on the extent of the dissection, the location of the entry tear, and the patency of the false lumen, patients who undergo repair of a type A dissection are at risk for aneurysmal degeneration over time, which could lead to the development of a thoracoabdominal aneurysm—a factor that should be considered when determining the extent of the repair at the time of the initial operation. These sequelae may require subsequent intervention such as an endovascular stent placement, or in more complicated situations, an open repair of the aneurysm. More recently, total arch replacements with or without concomitant elephant trunks (conventional and frozen) have been implemented to promote false lumen

thrombosis and to reduce the risk of distal aneurysm formation.

The article titled “Proximal vs Extensive Repair in Acute Type A Aortic Dissection Surgery” by Liu *et al.* analyzed 5,510 patients who presented with ATAAD from 13 hospitals over a 5-year period to create a risk scoring model to help direct the type of repair that should be performed (5). Before delving into this article, it is important to note the excellent results of this series in context. There is a large disparity in the literature regarding surgical outcomes of repair for ATAAD as well as controversies regarding the optimal extent of repair—one that optimizes operative mortality and minimizes re-operative intervention—to achieve the ideal result. Whether one supports a more conservative surgical approach or an aggressive technique such as total arch reconstruction, few series are able to report mortality rates such as these, consistently below 10% (6-8).

The authors' propensity score matched patients and implemented the eXtreme Gradient Boosting machine learning library to create a risk prediction model to predict operative mortality in patients. Their alphabet risk model includes parameters of age, body mass index, platelet-to-leukocyte ratio, albumin, hemoglobin, serum creatinine, and preoperative malperfusion to predict operative mortality.

[^] ORCID: 0000-0002-2245-5845.

Their key finding is that beyond a certain risk probability threshold (4.5%), extensive repair is associated with higher mortality than proximal repair [odds ratio (OR), 2.164; 95% confidence interval (CI): 1.679–2.788], indicating that although a certain subset of patients may benefit from more extensive repair, a tailored strategy for repair of ATAAD leads to more favorable outcomes.

Similar to this study, our group previously studied the association between the extent of aortic replacement and the outcomes of the procedures (9). We found that distal extension of an aortic procedure is independently associated with a higher complication rate, whereas proximal extension is not. Our multivariate logistic regression further supported this finding by demonstrating that partial or total arch replacement was an independent risk factor for post-operative complications. Like the study published by Liu *et al.*, our work also suggests that the immediate post-operative risk associated with aortic replacement should be balanced against the suspected future risk of an aortic event.

Like the Society of Thoracic Surgeons (STS) risk calculator, which serves as a tool to aid in the prediction of a patient's overall risk of mortality and morbidity when undergoing coronary and/or valve surgery, the model proposed by Liu *et al.* may provide the initial groundwork for the development of a risk calculator for patients who require surgery for ATAAD. However, it is important to acknowledge that, unlike coronary and valve surgery, for which indications and approaches are standardized and overall mortality is much lower (10–12), outcomes of surgical repair of ATAAD depend much more on the team performing the procedure due to the highly variable presentation and the technical challenges surgical teams face in the operating room. The subgroup analysis (Tab. S2) comparing outcomes of low (<100 cases annually) *vs.* high volume centers (>100 cases annually) highlights this difference. In other countries, having a volume of <100 cases per year would certainly not be considered low volume. Nonetheless, this comparison demonstrated a significant reduction in mortality at high volume centers compared to low volume centers (8.1% *vs.* 11.5%, $P=0.02$). Literature suggests the outcome could be dependent even on surgeons (13), and thus it may be worth considering surgeon case volume in the final model.

While the authors included patients who underwent the full spectrum of aortic surgery repairs for ATAAD, one approach that is highly debated among aortic experts but was not examined in the present study is an endovascular approach first for patients presenting with

malperfusion—also known as fenestrated endovascular aortic repair. Additionally, not analyzed is the Zone II arch reconstruction, where the distal anastomosis is sewn proximal to the left subclavian artery at a level of the aortic arch, which is more easily accessible than a total arch approach, decreases clamp time and bypass time, and minimizes dissection around the recurrent laryngeal nerve (14). This middle ground for repair of ATAAD, which has become the preference at our center for appropriately selected patients, not only simplifying the index operation while still replacing most of the arch, but it also creates a landing zone for further endovascular repair.

Finally, follow-up studies are warranted on long-term mortality or the need for re-intervention for a follow-up staged endovascular repair for subsequent descending aneurysm formation or a type B dissection and associated morbidity from that follow-up procedure.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Journal of Thoracic Disease*. The article has undergone external peer review.

Peer Review File: Available at <https://jtd.amegroups.com/article/view/10.21037/jtd-23-1945/prf>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://jtd.amegroups.com/article/view/10.21037/jtd-23-1945/coif>). H.T. serves as an unpaid editorial board member of *Journal of Thoracic Disease* from October 2022 to January 2025. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with

the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Hirst AE Jr, Johns VJ Jr, Kime SW Jr. Dissecting aneurysm of the aorta: a review of 505 cases. *Medicine (Baltimore)* 1958;37:217-79.
2. Howard DP, Banerjee A, Fairhead JF, et al. Population-based study of incidence and outcome of acute aortic dissection and premorbid risk factor control: 10-year results from the Oxford Vascular Study. *Circulation* 2013;127:2031-7.
3. Hagan PG, Nienaber CA, Isselbacher EM, et al. The International Registry of Acute Aortic Dissection (IRAD): new insights into an old disease. *JAMA* 2000;283:897-903.
4. Trimarchi S, Nienaber CA, Rampoldi V, et al. Contemporary results of surgery in acute type A aortic dissection: The International Registry of Acute Aortic Dissection experience. *J Thorac Cardiovasc Surg* 2005;129:112-22.
5. Liu H, Zhang YY, Ding XH, et al. Proximal vs Extensive Repair in Acute Type A Aortic Dissection Surgery. *Ann Thorac Surg* 2023;116:270-8.
6. Aalberts JJ, Boonstra PW, van den Berg MP, et al. In-hospital mortality and three-year survival after repaired acute type A aortic dissection. *Neth Heart J* 2009;17:226-31.
7. Obel LM, Lindholt JS, Lasota AN, et al. Clinical Characteristics, Incidences, and Mortality Rates for Type A and B Aortic Dissections: A Nationwide Danish Population-Based Cohort Study From 1996 to 2016. *Circulation* 2022;146:1903-17.
8. Santini F, Montalbano G, Messina A, et al. Survival and quality of life after repair of acute type A aortic dissection in patients aged 75 years and older justify intervention. *Eur J Cardiothorac Surg* 2006;29:386-91.
9. Yamabe T, Zhao Y, Kurlansky PA, et al. Extent of aortic replacement and operative outcome in open proximal thoracic aortic aneurysm repair. *JTCVS Open* 2022;12:1-12.
10. Adelborg K, Horváth-Puhó E, Schmidt M, et al. Thirty-Year Mortality After Coronary Artery Bypass Graft Surgery: A Danish Nationwide Population-Based Cohort Study. *Circ Cardiovasc Qual Outcomes* 2017;10:e002708.
11. Movahed MR, Etemad S, Hashemzadeh M, et al. Persistent reduction in the age adjusted mortality rate from aortic valve surgery in the United State with elimination of gender gap in recent years. *Am J Cardiovasc Dis* 2020;10:522-7.
12. Moreira JL, Barletta PHAAS, Baucia JA. Morbidity and Mortality in Patients Undergoing Mitral Valve Replacement at a Cardiovascular Surgery Referral Service: a Retrospective Analysis. *Braz J Cardiovasc Surg* 2021;36:183-91.
13. Umana-Pizano JB, Nissen AP, Sandhu HK, et al. Acute Type A Dissection Repair by High-Volume Vs Low-Volume Surgeons at a High-Volume Aortic Center. *Ann Thorac Surg* 2019;108:1330-6.
14. Bavaria J, Vallabhajosyula P, Moeller P, et al. Hybrid approaches in the treatment of aortic arch aneurysms: postoperative and midterm outcomes. *J Thorac Cardiovasc Surg* 2013;145:S85-90.

Cite this article as: Kandula V, Elmously A, O'Donnell TFX, Patel VI, Takayama H. Striking a balance when operating for acute type A aortic dissection. *J Thorac Dis* 2024;16(5):3522-3524. doi: 10.21037/jtd-23-1945