


Optimal Graft Choice in Athletic Patients with Anterior Cruciate Ligament Injuries: Review and Clinical Insights

Katarina Sim ¹, Richard Rahardja^{1,2}, Mark Zhu¹, Simon W Young^{1,2}

¹Department of Orthopaedics, North Shore Hospital, Auckland, New Zealand; ²Department of Surgery, University of Auckland, Auckland, New Zealand

Correspondence: Katarina Sim, Department of Orthopaedics, North Shore Hospital, 124 Shakespeare Road, Takapuna, Auckland, 0620, New Zealand, Email katarina.sim@waitematahnb.govt.nz

Abstract: Anterior cruciate ligament (ACL) rupture is a common sporting-related knee injury with a potentially detrimental impact on the athlete's career, yet there is no formal consensus on the optimal graft choice for reconstructing the ruptured ACL in this specific population. Options for reconstruction include autograft, allograft, and artificial grafts. However, each has associated failure risk and donor site morbidity. Our operational definition of the athlete is a skeletally mature individual participating in high level activity with the expectation to return to pre-injury level of activity. The athlete has unique injury characteristics, post-operative expectations, and graft demands that differ to the general population. Long-term outcomes are of particular importance given ongoing mechanical demands on the reconstructed knee. Therefore, the purpose of this review is to consolidate current literature on the various ACL reconstruction graft options, with a focus on the optimal graft for returning the athlete to activity with the lowest rate of re-injury.

Keywords: anterior cruciate ligament, reconstruction, athlete, graft choice, graft failure, return to activity

Introduction

The anterior cruciate ligament (ACL) is the primary restraint to anterior tibial translation and contributes to tibial rotational stability. Rupture of the ACL commonly occurs during sporting activity involving either a non-contact or pivoting mechanism. A higher incidence is observed in the young athlete population, with surgical reconstruction often performed to restore structural stability to the knee and facilitate return to sport. Reconstructing the ACL-deficient knee may also reduce the risk of long-term complications due to instability, such as injury to the menisci and cartilage.^{1,2} Additionally, reconstruction has been shown to be economically advantageous with an increase in quality-of-life compared to rehabilitation and non-operative management.^{3,4}

The ideal graft for reconstruction of the athlete's ACL will recreate the complex anatomical and biomechanical properties of the native ligament. An adequate graft length of over 7 cm with a mid-substance diameter between 10–11 mm contributes to the mechanical properties that allow the graft to withstand the tensile load, stiffness, and strain of a native ACL with minimal structural graft-related complications.^{5–8} As this surgical technique has evolved over decades, a number of different grafts have been used by surgeons around the world. Each graft has unique features, advantages, and disadvantages, and ultimately graft choice is individualized depending on multiple factors. This includes surgeon experience and preference, tissue availability, patient activity level, occupation, comorbidities, prior surgery, extent of injury, and patient preference. Each contributes to the likelihood of post-operative success, which itself is a subjective measure unique to each patient. Aspects of surgical technique, such as graft fixation or femoral tunnel drilling method (transtibial or anteromedial portal) may also influence outcome; however, these have less influence on revision rates than graft selection and patient factors such as age.^{9,10} Better understanding of graft choices and knowledge of updated evidence will facilitate decision-making around graft choice and ultimately improve outcomes, decrease morbidity, and lower revision rates.

In the athletic population, the primary goal of ACL reconstruction (ACLR) is to return the athlete back to sport at the lowest rate of reinjury. Taking into consideration this goal, we aim to review the current literature on graft choices and outcomes and provide our recommendation on the ideal graft choice in primary ACLR in the athlete population.

Discussion

ACLR Graft Options

ACLR graft choices can be broadly categorised into autograft, allograft, and artificial grafts. The ideal graft should have similar properties to the native ligament, limit donor site morbidity, and allow for secure fixation and rapid incorporation.⁸

Autograft

Three autograft options are commonly used. The bone patella tendon bone (BPTB) autograft is classically harvested from the middle third of the patellar tendon. It produces a consistent graft diameter of around 10 mm and contains bone plugs from the patella and tibia at either end of the graft that facilitates bone-to-bone healing within the bony graft tunnel.^{11,12} This reinforces fixation stability and is comparable to fracture healing, which is faster and stronger than soft tissue healing.⁷ Typically, fixation is achieved using an interference screw which engages the bone plugs of the graft in the bony tunnel with compression, further facilitating graft healing.^{13,14} Particular consideration should be taken in skeletally immature individuals as graft harvest and fixation methods could compromise physes and increase the risk of growth arrest.^{8,12}

The hamstring tendon (HT) autograft is harvested from the semitendinosus and/or gracilis tendon to yield variable sizes and constructs.¹⁵ In contrast to the BPTB autograft, the HT autograft is a soft tissue graft without bone plugs, therefore relying on soft tissue healing in bony tunnels, which may occur more slowly and is weaker compared to bone-to-bone healing.¹¹ Furthermore, the size of a HT graft ultimately depends on the size of the semitendinosus tendon, which varies between patients. There are a variety of different HT graft constructs which can involve creating as many as eight strands between the semitendinosus and gracilis tendons, and can produce a graft diameter varying from 6 mm to more than 10 mm. Literature supports greater biomechanical strength with larger grafts, but a downside of potential impingement during knee range of motion.⁸ In contrast to the standardized interference fixation for BPTB, a variety of different fixation methods are used with the HT graft with no consensus on the optimal method between surgeons. The most common methods are interference screw, suspensory, or cross-pin fixation. Similarly, surgeons must choose whether to use an interference screw with a sheath or without, or a fixed or adjustable loop when using a suspensory fixation device. A concerning complication, thought to occur secondary to graft mobility within the tunnel, is development of a fibrous tissue layer around the graft which leads to tunnel enlargement. This is more common in the tibial tunnel, and can compromise graft integration and potentially complicate revision procedures due to the created bone defect.^{11,16} Advantages include no donor-site compromise of the extensor mechanism which can impede early rehabilitation, polyvalence (variable composition and diameter) to meet specific reconstruction requirements, and reduced likelihood of causing growth arrest in the skeletally immature population.^{15,17,18}

The quadriceps tendon (QT) autograft has recently increased in usage and popularity, with studies reporting comparable outcomes to HT autograft.^{8,19} Different harvesting techniques have been described to obtain partial- to full-thickness grafts with or without a bone block.²⁰ The range of graft subtypes allow for different sizes and thicknesses and, therefore, varying biomechanical graft properties. For example, full-thickness QT harvested with bone block without retinaculum has been demonstrated to have equivalent load to failure compared to the BPTB autograft.²⁰ In addition, a variety of fixation methods have also been described. The versatility of the QT autograft in ACLR stems from its practicality in a variety of circumstances with differing tunnel sizes, tunnel positions, and navigating around previous grafts.²⁰ A consequent challenge is the applicability of outcomes of QT autografts as a cluster and the ability to determine specific outcome measures related to any QT autograft subtype. There is a lack of robust evidence describing outcomes from a “gold standard” QT autograft ACLR procedure, particularly in our target population.

Tunnel widening can occur with any graft type, but is more common with soft tissue grafts such as the hamstring autograft.²¹ While not thought to affect short-term clinical outcomes, tunnel widening can make revision procedures more challenging and bone grafting may be required.²¹

Allograft

Allograft options for ACLR are numerous, with patella tendon, hamstring, tibialis posterior, tibialis anterior, and achilles tendon all described extensively in the literature.^{3,22–25} Allograft has its main advantages of lack of donor site morbidity, better control over graft size, shorter operative time, and improved cosmesis.^{3,26,27} Disadvantages include graft availability, cost, and, more significantly, infection risk and graft rejection.^{3,26–28} Disease transmission remains a serious complication that has been largely eliminated with development of better donor screening and testing procedures and introduction of sterilization techniques.^{7,26} However, a major drawback of sterilization techniques is the effect it has on the biomechanical properties of the allograft. All allografts have slower rates of incorporation compared to autograft, alongside a much higher failure rate of up to 25% in the active population.^{8,29} Additionally, allograft is reported to have a lower return-to-sport rate compared to autograft (43% vs 75%).³⁰ Current evidence supports its use in specific circumstances such as multi-ligament knee reconstruction, inadequate autograft tissue, or in older, less active populations.

Artificial Graft

Artificial grafts were first reported as a reconstruction graft option in the 1980s, with the main attractions of lack of donor site morbidity, shorter surgical time, and reduced risk of disease transmission. And potentially an earlier return to sport.^{27,31,32} However, early studies reported satisfactory short-term outcomes but mid-to-long term complications of immunological response, foreign-body synovitis, tunnel osteolysis, femoral and tibial fractures near tunnels, and delayed failure.^{28,31,33,34} This resulted in a decline in use of artificial grafts, although there is a recent resurgence of interest with newer generation grafts that have yielded successful outcomes when used in specific circumstances, such as in an older population.^{31,33,35,36}

Artificial graft is a broad term encompassing both synthetic and augmented grafts. They serve as scaffolds, stents, or prostheses in ACLR. Legnani et al³¹ discuss the evolution of synthetic grafts and augments over the years, with the Ligament Advanced Reinforcement System™ (LARS™) being the most recent development. LARS, composed of polyethylene terephthalate, is designed to have better tissue ingrowth, particularly in intra-articular portions of the graft.^{31,37} This addresses concerns from studies that have suggested failure of intra-articular artificial graft integration contributing to failure of early generation synthetic grafts.³⁸ However, causes of failure are multifactorial, ranging from impaired fibrovascular ingrowth secondary to surrounding foreign body reactions to mechanical factors such as fiber properties and tunnel position.³⁷

Outcomes

There are multiple subjective and objective outcome measures to assess success of primary ACLR. Objective outcome measures frequently reported in the literature are graft failure, return to activity, and contralateral ACL injury. These factors are interlinked and have a combined 23% risk of secondary ACL injury in athletes younger than 25 years old.^{39,40}

Graft Failure

Graft failure is the most common outcome analyzed in ACL research. Though it has varying definitions depending on study type, it is commonly defined as graft rupture or residual instability following ACLR. In the athletic population, this is one of the most important outcome measures, as one ACL rupture may threaten an athlete's career, but a second re-rupture may end a career.

The rate of graft failure reported in the literature ranges between 1.8–33%.^{41–43} The most commonly reported risk factors for graft failure include graft type, age at surgery, and activity level.^{27,44–46} Under 20 years old at time of surgery is the most commonly reported risk factor for graft failure.^{9,41,44,46–53} For the athlete, the above three factors compound their risk of graft failure.

Autograft

An early meta-analysis by Freedman et al, consisting of studies between 1966 and 2000, reported a lower rate of graft failure with BPTB autograft compared to HT autograft.^{9,54} Further registry studies and systematic reviews have

repeatedly highlighted lower rates of graft failure and revision surgery with BPTB autograft compared to HT and QT autografts in ACLR across subgroups of patient ages and follow-up timeframes.^{10,55–57} A New Zealand (NZ) ACL Registry study reviewing 7,155 primary ACLR identified a higher revision rate with HT autograft compared to BPTB autograft (2.7% vs 1.3%, $p<0.001$).¹⁰ Supporting this is a longitudinal study over 6 years illustrating a 2.1-times higher odds of graft failure with HT autograft compared to BPTB autograft ($p=0.004$).⁵⁸

There are few adequately powered randomized controlled trials (RCTs) comparing outcomes following BPTB and HT autograft in primary ACLR. Early RCTs yielded comparative results between the two groups.^{59–65} However, a prospective randomized study by Beynon et al⁶⁶ illustrated superior objective results with BPTB autograft reconstruction compared to double-strand HT autograft at 3 years follow-up. Though not statistically significant, these findings are consistent with those by Maletis et al⁶⁷ which also identified a greater number of patients returning to preinjury activity levels following BPTB ACLR. Since, a double-blind RCT by Mohtadi et al,⁴⁶ consisting of 330 patients, compared 2 year outcomes between BPTB and HT autograft. The study reported a significantly lower proportion of traumatic reinjuries in the BPTB group (3 of 110) compared to the HT autograft group (7 of 110).⁴⁶

Comparative data on QT grafts is limited and conflicting; however, studies report a failure rate of 2.2–4.8%, depending on the size of the study and duration of follow-up.^{68,69} Runer et al⁴⁵ performed a prospective study of 875 patients undergoing ACLR and reported a 3-times higher risk of revision surgery when using a HT autograft compared to a QT autograft (OR=2.7, $p=0.007$). Furthermore, in high activity patients with a pre-injury Tegner score of ≥ 7 , the revision rate was 11.1% in the HT autograft group compared to 5% in the QT autograft group ($p=0.01$).⁴⁵ Interestingly, the HT autograft cohort had significantly higher ipsilateral graft ruptures than contralateral ACL injuries (4.9% vs 2.3%), with this difference being more marked in individuals with higher activity levels (11.1% vs 4.2%).⁴⁵ In contrast, a Danish registry study illustrated higher graft failure (revision rate) for ACLR with QT compared to HT and BPTB (4.7 vs 2.3 vs 1.5%).⁷⁰ However, this study is limited by a lack of specificity on QT graft characteristics (ie, size, bone block use, fixation technique) and, therefore, applicability of its findings is limited. A prospective cohort study consisting of 48 patients (27 BPTB and 21 QT autograft) reported the highest rate of QT graft failure at 4.8% compared to 3.7%.⁶⁹ The main criticism is the small study sample size. Comparatively, the largest cohort study of 198 QT autograft compared to 30 BPTB autografts yielded graft failure rates of 2.0% and 3.3%, respectively.⁷¹

Current evidence suggests BPTB autograft for ACLR in the athletic population has a failure rate significantly lower than all other graft options.^{9,10,46,54–58,66,67,69,70}

Allograft

Allografts have a consistently higher failure rate compared to autografts at all stages of follow-up, with up to triple the failure rate of autograft quoted by Bottoni et al.³ Allograft failure is related to structural characteristics of the graft, sterilization techniques, and donor characteristics. Structurally, single-strand allografts have higher rates of graft failure compared to multistrand allografts.²² Infection rates are higher with unsterilized and/or aseptically processed grafts. Viral and bacterial transmission risk remains low but has a high detrimental impact and is associated with non-irradiated allografts. Gamma radiation can be used to address infection risk; however, it has a higher graft failure rate compared to autograft and fresh frozen allograft.^{25,26} Furthermore, donor characteristics influence graft failure rates with increased risk of failure of grafts from female donors over 50 years old.²²

Allograft sterilization techniques alter the mechanical properties of allografts and are broadly categorized into radiation (gamma, electron beam, x-ray) or ethylene oxide. The extent of alteration to graft mechanical properties is dependent on irradiation dose exposure.²⁵ Doses greater than 2.5 kilogray (kGy) have been shown to decrease allograft tension and, therefore, result in greater post-operative laxity and predispose to failure.²⁵ Farago et al²⁶ reviewed articles over 29 years assessing the impact of various sterilization techniques on objective tendon mechanics (ie, failure load/ultimate strength and Young's modulus of elasticity). Findings from the review support freezing followed by gamma radiation or electron beam at 14.8–28.5 kGy as the sterilization technique with the greatest biomechanical preservation.²⁶ However, allograft failure is not attributable to sterilization techniques alone, as rates of allograft failure remain higher when comparing fresh-frozen allografts with autografts.²⁶

In the young active population, allograft reconstruction is a risk factor for failure compared to autograft.^{3,27,53,72–75} A systematic review by Hayback et al⁷³ suggested the odds of ACL graft rupture decreases with every yearly increase in patient age for a study population with a minimum mean age of 18 years. Allograft use has been shown to be an independent risk factor for graft failure and the need for any subsequent surgery, with the risk of repeat ACL injury being 5.2-times greater for an allograft compared to a BPTB autograft within the first 2 years of reconstruction surgery.⁵³

Given the above risk-benefits, allograft is not recommended as a first option graft choice in isolated primary ACLR in the athletic population. They may be beneficial in certain circumstances such as for older patients, in ACL revision surgery, or cases of multi-ligamentous reconstruction.³¹

Artificial Graft

Multiple cohort studies have yielded positive short- to mid-term outcomes of the LARS™ artificial graft subtype in ACLR.^{32,76} However, there is limited evidence supporting favorable long-term outcomes of artificial grafts, even if isolated to LARS™. The large majority of studies are composed of small study populations, as expected with the infrequent use of synthetic grafts in primary ACLR. A retrospective study of 18 patients over 10 years by Tiefenboeck et al⁷⁷ reported a 27.8% artificial graft re-rupture rate. Similarly, Tulloch et al³⁸ published a 33.3% mechanical failure rate in artificial graft ACLR in a cohort of 55 patients over a median timeframe of 7.8 years.

An earlier study by Pan et al³³ compared mid-term outcomes between BPTB autograft and LARS™ in a cohort of 62 patients undergoing primary ACLR. Findings from the study did not yield statistically significant differences in functional outcomes.³³ However, demographic characteristics of the study population are not entirely applicable to that of our athlete population, with a mean age of 34 in the BPTB autograft group and 36 in the LARS™ group, and pre-operative Tegner scores consistent with recreational sport participation only.³³ A more recent systematic review of 748 studies by Fan et al²⁸ comparing BPTB autograft with synthetic and augmented graft reconstruction confirmed more favorable objective outcomes with BPTB autograft (OR=0.49; 95% CI=0.28–0.86).

Contralateral ACL Injury

Contralateral ACL injury is a major complication that is as impactful as an ipsilateral graft rupture for the young athlete following primary ACL reconstruction. It is associated with reduced function, quality-of-life, and likelihood of return to pre-injury activity.⁷⁸ Interestingly, some studies have reported that the risk of contralateral ACL injury may be higher than the risk of ipsilateral re-rupture following primary ACLR.^{42,79,80}

Risk factors for contralateral ACL injury include younger age, sex, graft selection, and level of sport played.^{52,81} Registry-based studies have consistently reported age at time of index ACLR as a statistically significant risk factor for contralateral injuries at all follow-up timeframes.^{41,42,47,49,53,82–85} Furthermore, returning to a high level of activity post-reconstruction is a risk factor for contralateral ACL injury.⁸³ In a study of 2,488 primary ACLR from the MOON cohort, Kaeding et al⁵³ reported that the odds of a contralateral ACL tear increased by 0.12 for every increased point on the Marx activity score.

Leys et al⁸⁶ performed a randomized controlled trial comparing BPTB and HT autograft in 90 patients over 15 years and reported a 2.6-times higher odds of contralateral ACL rupture with BPTB. Similar findings have been reported by ACL registry studies.^{10,42} In 17,436 ACL reconstructions recorded in the Kaiser Permanente Registry, a 1.3-times higher risk of contralateral ACL reconstruction was reported in patients with a BPTB autograft compared to a HT autograft.⁴² In a NZ ACL Registry study of 7,155 patients, those in the BPTB group had 1.9-times higher risk of contralateral ACL reconstruction compared with the HT group (adjusted HR=1.91; 95% CI=1.15–3.16, $p=0.012$).¹⁰

In contrast, a 755 patient case series with 15-year follow-up by Bourke et al⁷⁹ showed no significant difference in contralateral ACL rupture between the BPTB and HT autograft groups (hazard ratio [HR]=1.5; 95% CI=1.0–2.2). Additionally, a randomized controlled trial of 330 patients with a minimum 2 year follow-up by Mohtadi et al⁴⁶ showed no difference in the rate of contralateral injury between graft types.

It is unclear why patients with a BPTB autograft have a higher risk of contralateral injury compared to patients with a HT autograft. One explanation is that patients with a BPTB autograft are more able to return to sport and do so earlier

and at a higher level, which consequently makes the contralateral knee susceptible to injury.⁸⁷ Other possible reasons may relate to biochemical changes or altered neuromuscular function following ACLR.

Return to Activity

In young active patients, the risk of reconstruction failure increases with higher activity levels.¹ A variety of different methods are used to define return to activity. Return to activity may involve a patient returning to casual, amateur, or competitive level of sport, or analyzing scores calculated from an activity questionnaire such as the Marx Activity Questionnaire or the Tegner Activity Scale. In a meta-analysis of 48 studies performed by Ardern et al⁸⁸ comprising 5,770 participants, an 82% return to participation, 63% return to pre-injury level, and 44% return to competitive sport at final follow-up (mean time 41.5 months) were reported.

The majority of studies compare return to activity rates with BPTB and HT autografts since they are the most common autografts used in ACLR for the athlete. Isolated meta-analysis of four RCTs comparing return to preinjury level of sport following BPTB and HT autograft for ACLR showed no difference (OR=1.02, 95% CI=0.7–1.5, $p=0.92$).⁸⁹ However, the majority of publications favor BPTB autograft when return to activity is the focused outcome.^{87,90,91} In total, 7,556 patients from 69 studies were systematically reviewed by Ardern et al⁸⁹ to illustrate ACLR with BPTB was favored over HT autograft in return to pre-injury level of sport rates (OR=1.2). Supporting this is an updated systematic review of 20 articles, inclusive of 2,348 athletes by DeFazio et al,⁹¹ which identified an 81% overall return to sport rate with BPTB autograft compared to 70.6% with HT autograft.

A common limitation when comparing return to activity between BPTB and HT autografts is the effect of surgeon preference and the possible selection bias. Some surgeons may prefer to use a BPTB autograft in “higher risk” patients such as individuals participating in competitive sport. Subsequently, this may introduce bias into these studies as expectations of “return to activity” varies and is more challenging in professional athletes when compared to an amateur or casual athlete returning to sport.

Some clinicians believe that a higher rate of contralateral ACL injury following primary ACLR with a BPTB autograft is a marker of success of the BPTB autograft, as it demonstrates that the patient had returned to sport prior to their reinjury, which is the goal of the original procedure. A study from the NZ ACL Registry supports this theory by only analyzing high activity patients with a pre-injury Marx score of 13, so that the patients in both graft type groups were of equal activity levels.⁸⁷ Subsequently, this study found that BPTB patients had higher odds of returning to activity compared to HT patients. Additionally, the study identified a higher proportion of individuals with BPTB autograft were performing at their preinjury activity levels (Marx score ≥ 13) compared to HT autograft at 2 years (23.3% vs 13.3%, respectively; $p<0.001$).⁸⁷ Cohort studies have reported similar findings of higher return to sport rate in BPTB compared to HT autografts.^{40,90}

Donor Site Morbidity

Donor site morbidity is a significant concern, particularly when considering BPTB autograft in ACLR. Kartus et al⁹² report 40–60% of patients report various donor-site problems following BPTB autograft use in ACLR. The most common symptom is anterior knee and/or kneeling pain.^{8,92} Other serious, but rarer complications such as patella fracture and patella tendon rupture may also occur.^{7,8} Another commonly used measures of post-operative symptomology is the Knee Injury and Osteoarthritis Outcome Score (KOOS). A RCT by Lind et al⁹³ of 99 adults illustrated that, at 2 years follow-up, 50% of HT autograft recipients compared to 27% of QT autograft recipients had donor site symptoms as assessed using KOOS.

Anterior and Kneeling Knee Pain

Average incidence of anterior knee pain with autograft ACLR is up to 21.5%.¹⁸ There is a higher incidence of anterior knee pain and kneeling pain following ACLR using BPTB compared to all other autografts.^{7,12,18,54,92,94,95} This is consistently proven in the literature, including a recent meta-analysis of 685 patients across seven studies which identified 35.7% of the BPTB ACLR population to have anterior knee pain compared to 5.7% of QT autograft recipients.¹⁸ Furthermore, 70.8% of 100 patients in a retrospective non-randomized study evaluating donor site morbidity

following BPTB autograft ACLR had kneeling pain.⁹⁶ Symptomology can occur anywhere along the extensor mechanism and is likely to be related to graft choice and its harvesting method.^{7,8}

A prospective study by Kartus et al⁹⁷ aimed to identify donor site-related problems and compared post-operative anterior knee sensitivity following two different BPTB graft harvesting techniques. The study suggested a correlation between anterior knee sensitivity and intraoperative injury of the infrapatellar branches of the saphenous nerve.⁹⁷ This nerve travels along the inferior medial aspect of the knee joint.⁷ Thus, given its location, it is especially vulnerable to injury when harvesting the central third of the patellar tendon or when hamstring tendons are harvested through a medial incision.⁹⁸ This is further supported by literature highlighting anterior knee sensitivity and difficulty kneeling after incisions in the prepatellar area for various surgical procedures ranging from arthroscopic access to midline incisions for tibial nailing.⁹⁷ Therefore, it is important to consider the suitability and risk-benefit of a BPTB autograft in individuals whose lifestyle would be impacted by this.

Anterior knee pain may be associated with functional flexion and/or extension deficits in the postoperative period and, likewise, restoration of full extension compared with uninjured side after ACLR is essential to avoid post-operative discomfort in the anterior knee.⁹² In some studies, the two are shown to have a statistically significant risk association.^{94,98} Pain has been shown to occur 5-times more than in the absence of a deficit.^{94,98,99} This may have consequential functional limitations and, therefore, result in impairment of the athlete with higher functional demands of the extensor mechanism of the knee.

Patella-Related Complications

Though rare, patella tendon ruptures and fractures of the patella or proximal tibia in the donor knee are associated with medial-third BPTB autograft use in ACLR.¹⁰⁰ If undiagnosed or inadequately managed, these disruptions to the extensor mechanism can have devastating functional impact.

Patella Fracture

The reported incidence of patella fracture in the donor knee following BPTB autograft harvest in ACLR varies between 0–2%, with no reports of association with other graft types.¹⁰¹

A literature review by Tay et al¹⁰¹ of five case reports and eight case series, reports a cumulative 31 patella fractures following ACLR with BPTB autograft. The majority of these were sustained by indirect injuries and mostly stellate in configuration.^{101,102} The average time to injury was 11 weeks, though it can be as short as 24 days.^{101,102} The incidence of patella fractures suggests potentially altered distribution of forces acting on the iatrogenically weakened patella after graft harvest, therefore, placing it at increased risk of injury even at usual strain and contact stresses of normal activities such as stair climbing.¹⁰¹ Management of these complications was governed by displacement.^{101,102} Though complications extend rehabilitation, at full recovery, there was no significant difference in functional outcome between ACLR procedures complicated by patellar fracture.^{101–103}

Patella Tendon Rupture

Patella tendon rupture is rare, with a reported incidence of between 0.2–1.3% after BPTB harvest.^{54,96,100,102,104–108} This may not be a complication directly associated with BTPB autograft harvest alone, as earlier studies have reported this complication with other reconstructive grafts, including allograft and earlier-generation synthetic grafts.^{107,109} However, occurrences with alternate grafts has not been described in recent literature.

Patella tendon rupture usually occurs secondary to trauma such as a fall, accompanied by either a forced hyperflexion injury or a forceful quadriceps contraction.^{100,107,108} Current literature reports varying timing of occurrence following ACLR and patellar tendon rupture from days to years, with the majority occurring within the first year from surgery.^{100,103,107,108,110} The concern regarding patellar tendon rupture in a BPTB harvested knee is the technical challenge of repair and early loss of flexion.^{106,111}

A series of 13 cases of patella tendon rupture in the donor knee, from a database of 5,364 ACLR using BPTB autograft, identified varying tendon rupture sites differing from the commonly observed proximal-only tear pattern in unharvested patellar tendons.¹¹¹ The majority of post-reconstruction patellar tendon ruptures occurred from the patellar

origin medially extending to the tibial attachment laterally in a Z-shaped pattern.¹¹¹ In all cases, the donor site tendon defect was closed with running No. 0 Vicryl and the bone plug harvest defects in the patella and tibia were bone grafted with bone shavings from drilling of the femoral and tibial tunnels.¹¹¹ All cases recovered with positive long-term objective and subjective outcomes following patellar tendon repair and intensive rehabilitation.¹¹¹

Range of Motion and Strength

A meta-analysis by Zhao et al⁵⁷ evaluated donor site morbidity following HT and BPTB ACLR. They identified a significant difference, favoring HT, with regard to loss in knee extension range (16.2% vs 10.2% for BPTB vs HT, respectively, OR=1.67).⁵⁷ However, a 15-year follow-up of a RCT by Webster et al¹¹² noted that early increased extension deficit following BPTB ACLR was no longer present at 15 years. There are yet to be additional long-term studies to challenge the findings by Webster et al.

Muscle strength of the reconstructed knee remains similar following ACLR with either BPTB and HT autograft, as illustrated in a RCT by Beard et al.⁶² However, findings from a cross-sectional study highlight muscular weakness in the lower limb which has had a quadriceps tendon harvest compared to a contralateral undisturbed limb by 15% in men and 30% in women.²⁰ This is supported by Lind et al.⁹³

Donor site morbidity is a major factor contributing to hesitancy around use of BPTB autograft in ACLR. It has the potential to have significant post-operative consequences on specific populations, particularly individuals whose occupation or sporting activities involve kneeling. Fortunately, patella-related complications are rare and long-term functional outcomes are statistically indifferent following management of these complications. Balancing objective outcomes with donor site morbidity, there is robust evidence supporting use of BPTB autograft in the vast majority of the athletic population.

Lateral Extra-Articular Procedures

Adjuvant lateral extra-articular procedures (LEAPs), including lateral extra-articular tenodesis (LET) and anterolateral ligament reconstruction (ALL), are increasingly performed with ACLR to provide additional rotational stability and, therefore, reduce the risk of graft failure in high risk patients, such as the athlete.^{113–116}

A meta-analysis of 1,010 patients from six randomized controlled trials by Mao et al¹¹⁴ illustrated a lower risk of graft failure in those who had ACLR and LET compared to those who had isolated ACLR. Of significance is the STABILITY study by Getgood et al.¹¹⁷ This is the largest randomized multicenter study to date, looking at ACLR clinical failure between two ACLR groups – single-bundle hamstring tendon autograft without LET and with LET – in individuals between 14–25 years old. Recent data of 2-year outcomes illustrate statistically significant differences, with lower rates of clinical graft failure in the group receiving concurrent LET procedure (120 of 298 in the ACLR group compared to 72 of 291 in the ACLR+LET group, RRR=0.38; 95% CI=0.21–0.52; $p<0.0001$).¹¹⁷ Return to level of sport activity (Marx score) remains similar in both study groups.¹¹⁷

Similarly, the SANTI study group compared outcomes between isolated autograft ACLR to combined autograft ACLR with ALL procedure.¹¹⁸ Findings from this prospective cohort study of 270 patients showed statistically better ACL graft survivorship in the group who underwent ACLR with ALL (96.5% vs 82.6%; $p=0.0027$). Additionally, at 108 months, those who had concurrent ALL with primary ACLR had at least 2.5-times reduced graft failure rates.¹¹⁸

There is increasing evidence supporting concurrent LEAPs in primary ACLR to reduce risk of graft failure, particularly in high risk population groups such as athletes. Current limitations in evidence include comparisons between specific autograft subtypes and concurrent LEAPs. This is a developing area which will drive significant change in management of primary ACLR in the coming years.

Conclusion

In the athlete undergoing ACLR, the primary aim of treatment is to enable the highest rate of return to sport at the lowest rate of reinjury. When considering this aim, literature clearly supports the use of autograft over allograft, and more specifically the use of a BPTB autograft in primary ACLR. QT grafts are a newer development compared to other autografts and evidence supporting their use in reconstruction of the athlete's ACL is presently limited. Although the HT autograft is more commonly used globally, the BPTB autograft has been consistently reported to have the lowest rate of

failure by both RCTs and national registry studies. Additionally, there is evidence that patients with a BPTB autograft are more likely to return to sport and at a higher level. However, this comes at the expense of greater donor site morbidity with BPTB autografts, including a higher incidence of anterior knee pain, kneeling difficulty, and rare, but severe complications such as patella tendon rupture and fracture.

Individualizing graft choice remains the most important when consulting the ACL-deficient athlete, and the athlete must balance their desire and aspiration to return to sport, risk of reinjury, and donor-site morbidity.

Disclosure

Dr Simon W. Young reports non-financial support from New Zealand ACL Registry, grants, personal fees from Stryker, grants from Smith and Nephew, during the conduct of the study. The authors report no other conflicts of interest in this work.

References

1. Thorolfsson B, Svantesson E, Snaebjornsson T, et al. Adolescents have twice the revision rate of young adults after ACL reconstruction with hamstring tendon autograft: a study from the Swedish national knee ligament registry. *Orthop J Sports Med.* 2021;9(10):232596712110388. doi:10.1177/23259671211038893
2. Chalmers PN, Mall NA, Moric M, et al. Does ACL reconstruction alter natural history?: A systematic literature review of long-term outcomes. *J Bone Joint Surg.* 2014;96(4):292–300. doi:10.2106/JBJS.L.01713
3. Bottoni CR, Smith EL, Shaha J, et al. Autograft versus allograft anterior cruciate ligament reconstruction. *Am J Sports Med.* 2015;43(10):2501–2509. doi:10.1177/0363546515596406
4. Davies L, Cook J, Leal J, et al. Comparison of the clinical and cost effectiveness of two management strategies (rehabilitation versus surgical reconstruction) for non-acute anterior cruciate ligament (ACL) injury: study protocol for the ACL SNNAP randomised controlled trial. *Trials.* 2020;21(1). doi:10.1186/s13063-020-04298-y
5. Cheung SC, Allen CR, Gallo RA, Ma CB, Feeley BT. Patients' attitudes and factors in their selection of grafts for anterior cruciate ligament reconstruction. *Knee.* 2012;19(1):49–54. doi:10.1016/j.knee.2010.11.009
6. Kraeutler MJ, Wolsky RM, Vidal AF, Bravman JT. Anatomy and biomechanics of the native and reconstructed anterior cruciate ligament: surgical implications. *J Bone Joint Surg Am.* 2017;99(5):438–445. doi:10.2106/JBJS.16.00754
7. West RV, Harner CD. Graft selection in anterior cruciate ligament reconstruction. *JAAOS-Journal of the American Academy of Orthopaedic Surgeons.* 2005. May 1;13(3):197–207. doi:10.5435/00124635-200505000-00006
8. Buerba RA, Boden SA, Lesniak B. Graft selection in contemporary anterior cruciate ligament reconstruction. *JAAOS.* 2021;5(10). doi:10.5435/JAAOSGlobal-D-21-00230
9. Rahardja R, Zhu M, Love H, Clatworthy MG, Monk AP, Young SW. Factors associated with revision following anterior cruciate ligament reconstruction: a systematic review of registry data. *Knee.* 2020;27(2):287–299. doi:10.1016/j.knee.2019.12.003
10. Rahardja R, Zhu M, Love H, Clatworthy MG, Monk AP, Young SW. Effect of graft choice on revision and contralateral anterior cruciate ligament reconstruction: results from the New Zealand ACL registry. *Am J Sports Med.* 2020;48(1):63–69. doi:10.1177/0363546519885148
11. Thaanat M, Fayard JM, Sonnery-Cottet B. Hamstring tendons or bone-patellar tendon-bone graft for anterior cruciate ligament reconstruction? *Orthopaed Traumatol.* 2019;105(1):S89–S94. doi:10.1016/j.otsr.2018.05.014
12. Cerulli G, Placella G, Sebastiani E, Tei MM, Speziali A, Manfreda F. ACL reconstruction: choosing the graft. *Joints.* 2013;1(1):18–24.
13. Provencher MT, Ryu JH, Gaston T, Dewing CB. Technique: bone-patellar tendon-bone autograft ACL reconstruction in the young, active patient. *J Knee Surg.* 2011;24(2):83–92. doi:10.1055/s-0031-1280875
14. Schoderbek RJ, Treme GP, Miller MD. Bone-patella tendon-bone autograft anterior cruciate ligament reconstruction. *Clin Sports Med.* 2007;26(4):525–547. doi:10.1016/j.csm.2007.06.006
15. Chen L, Cooley V, Rosenberg T. ACL reconstruction with hamstring tendon. *Orthop Clin North Am.* 2003;34(1):9–18. doi:10.1016/S0030-5898(02)00016-0
16. Amano H, Tanaka Y, Kita K, et al. Significant anterior enlargement of femoral tunnel aperture after hamstring ACL reconstruction, compared to bone-patellar tendon-bone graft. *Knee Surg Sports Traumatol Arthrosc.* 2019;27(2):461–470. doi:10.1007/s00167-018-5098-9
17. Mouarbes D, Menetrey J, Marot V, Courtot L, Berard E, Cavaignac E. Anterior cruciate ligament reconstruction: a systematic review and meta-analysis of outcomes for quadriceps tendon autograft versus bone-patellar tendon-bone and hamstring-tendon autografts. *Am J Sports Med.* 2019;47(14):3531–3540. doi:10.1177/0363546518825340
18. Ajrawat P, Dwyer T, Whelan D, et al. A comparison of quadriceps tendon autograft with bone-patellar tendon-bone autograft and hamstring tendon autograft for primary anterior cruciate ligament reconstruction: a systematic review and quantitative synthesis. *Clin J Sport Med.* 2021;31(4):392–399. doi:10.1097/JSM.0000000000000765
19. Slone HS, Romine SE, Premkumar A, Xerogeanes JW. Quadriceps tendon autograft for anterior cruciate ligament reconstruction: a comprehensive review of current literature and systematic review of clinical results. *Arthrosc-J Arthrosc Relat Surg.* 2015;31(3):541–554. doi:10.1016/j.arthro.2014.11.010
20. Diermeier T, Tisherman R, Hughes J, et al. Quadriceps tendon anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2020;28(8):2644–2656. doi:10.1007/s00167-020-05902-z
21. Clatworthy MG, Annear P, Bulow JU, Bartlett RJ. Tunnel widening in anterior cruciate ligament reconstruction: a prospective evaluation of hamstring and patella tendon grafts. *Knee Surg Sports Traumatol Arthrosc.* 1999;7(3):138–145. doi:10.1007/s001670050138

22. Shumborski S, Salmon LJ, Monk C, Heath E, Roe JP, Pinczewski LA. Allograft donor characteristics significantly influence graft rupture after anterior cruciate ligament reconstruction in a young active population. *Am J Sports Med.* 2020;48(10):2401–2407. doi:10.1177/0363546520938777
23. Su M, Jia X, Zhang Z, et al. Medium-term (Least 5 Years) comparative outcomes in anterior cruciate ligament reconstruction using 4SHG, allograft, and LARS ligament. *Clin J Sport Med.* 2021;31(2):e101–e110. doi:10.1097/JSM.0000000000000730
24. Goetz G, de Villiers C, Sadoghi P, Geiger-Gritsch S. Allograft for Anterior Cruciate Ligament Reconstruction (ACLR): a systematic review and meta-analysis of long-term comparative effectiveness and safety. results of a health technology assessment. *Arthros, Sports Med Rehabil.* 2020;2(6):e873–e891. doi:10.1016/j.asmr.2020.07.003
25. Guo L, Yang L, Duan XJ, et al. Anterior cruciate ligament reconstruction with bone-patellar tendon-bone graft: comparison of autograft, fresh-frozen allograft, and γ -irradiated allograft. *Arthrosc- J Arthrosc Relat Surg.* 2012;28(2):211–217. doi:10.1016/j.arthro.2011.08.314
26. Farago D, Kozma B, Kiss RM. Different sterilization and disinfection methods used for human tendons – a systematic review using mechanical properties to evaluate tendon allografts. *BMC Musculoskelet Disord.* 2021;22(1). doi:10.1186/s12891-021-04296-4
27. Duchman KR, Lynch TS, Spindler KP. Graft selection in anterior cruciate ligament surgery: who gets what and why? *Clin Sports Med.* 2017;36(1):25–33. doi:10.1016/j.csm.2016.08.013
28. Fan DY, Ma J, Zhang L. Patellar tendon versus artificial grafts in anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *J Orthop Surg Res.* 2021;16(1). doi:10.1186/s13018-021-02624-x
29. Thomas Vangsness C, Garcia IA, Randal Mills C, Kainer MA, Roberts MR, Moore TM. Allograft transplantation in the knee: tissue regulation, procurement, processing, and sterilization. *Am J Sports Med.* 2003;31(3):474–481. doi:10.1177/03635465030310032701
30. Kaeding CC, Pedroza AD, Reinke EK, et al. Change in anterior cruciate ligament graft choice and outcomes over time. *Arthrosc- J Arthrosc Relat Surg.* 2017;33(11):2007–2014. doi:10.1016/j.arthro.2017.06.019
31. Legnani C, Ventura A, Terzaghi C, Borgo E, Albisetti W. Anterior cruciate ligament reconstruction with synthetic grafts. A review of literature. *Int Orthop.* 2010;34(4):465–471. doi:10.1007/s00264-010-0963-2
32. Satora W, Królikowska A, Czamara A, Reichert P. Synthetic grafts in the treatment of ruptured anterior cruciate ligament of the knee joint. *Poly Med.* 2017;47(1):55–59. doi:10.17219/pim/76819
33. Pan X, Wen H, Wang L, Ge T. Bone-patellar tendon-bone autograft versus LARS artificial ligament for anterior cruciate ligament reconstruction. *Eur J Orthopaed Surg Traumatol.* 2013;23(7):819–823. doi:10.1007/s00590-012-1073-1
34. Ventura A, Terzaghi C, Legnani C, Borgo E, Albisetti W. Synthetic grafts for anterior cruciate ligament rupture: 19-year outcome study. *Knee.* 2010;17(2):108–113. doi:10.1016/j.knee.2009.07.013
35. Gao K, Chen S, Wang L, et al. Anterior cruciate ligament reconstruction with LARS artificial ligament: a multicenter study with 3- to 5-year follow-up. *Arthrosc- J Arthrosc Relat Surg.* 2010;26(4):515–523. doi:10.1016/j.arthro.2010.02.001
36. Parchi PD, Ciapini G, Paglialonga C, et al. Anterior cruciate ligament reconstruction with LARS artificial ligament—clinical results after a long-term follow-up. *Joints.* 2018;6(2):75–79. doi:10.1055/s-0038-1653950
37. Di Benedetto P, Giardini P, Beltrame A, Mancuso F, Gisonni R, Causero A. Histological analysis of ACL reconstruction failures due to synthetic-ACL (LARS) ruptures. *Acta Biomedica.* 2020;91:136–145. doi:10.23750/abm.v91i4-S.9702
38. Tulloch SJ, Devitt BM, Porter T, et al. Primary ACL reconstruction using the LARS device is associated with a high failure rate at minimum of 6-year follow-up. *Knee Surg Sports Traumatol Arthrosc.* 2019;27(11):3626–3632. doi:10.1007/s00167-019-05478-3
39. Wiggins AJ, Grandhi RK, Schneider DK, Stanfield D, Webster KE, Myer GD. Risk of secondary injury in younger athletes after anterior cruciate ligament reconstruction. *Am J Sports Med.* 2016;44(7):1861–1876. doi:10.1177/0363546515621554
40. King E, Richter C, Jackson M, et al. Factors influencing return to play and second anterior cruciate ligament injury rates in level I athletes after primary anterior cruciate ligament reconstruction: 2-year follow-up on 1432 reconstructions at a single center. *Am J Sports Med.* 2020;48(4):812–824. doi:10.1177/0363546519900170
41. Sutherland K, Clatworthy M, Chang K, Rahardja R, Young SW. Risk factors for revision anterior cruciate ligament reconstruction and frequency with which patients change surgeons. *Orthop J Sports Med.* 2019;7(11):232596711988048. doi:10.1177/2325967119880487
42. Maletis GB, Inacio MCS, Funahashi TT. Risk factors associated with revision and contralateral anterior cruciate ligament reconstructions in the Kaiser Permanente ACLR registry. *Am J Sports Med.* 2015;43(3):641–647. doi:10.1177/0363546514561745
43. Paterno MV, Rauh MJ, Schmitt LC, Ford KR, Hewett TE. Incidence of second ACL injuries 2 years after primary ACL reconstruction and return to sport. *Am J Sports Med.* 2014;42(7):1567–1573. doi:10.1177/0363546514530088
44. Andernord D, Desai N, Björnsson H, Ylander M, Karlsson J, Samuelsson K. Patient predictors of early revision surgery after anterior cruciate ligament reconstruction: a cohort study of 16,930 patients with 2-year follow-up. *Am J Sports Med.* 2015;43(1):121–127. doi:10.1177/0363546514552788
45. Runer A, Csapo R, Hepperger C, Herbort M, Hoser C, Fink C. Anterior cruciate ligament reconstructions with quadriceps tendon autograft result in lower graft rupture rates but similar patient-reported outcomes as compared with hamstring tendon autograft: a comparison of 875 patients. *Am J Sports Med.* 2020;48(9):2195–2204. doi:10.1177/0363546520931829
46. Mohtadi N, Chan D, Barber R, Paolucci EO. Ruptures, Reinjuries, and Revisions at a Minimum 2-Year Follow-up: A Randomized Clinical Trial Comparing 3 Graft Types for ACL Reconstruction. *Clin J Sport Med.* 2016. Mar;26(2):96–107. doi:10.1097/JSM.0000000000000209. PMID: 26164058.
47. Snaebjörnsson T, Hamrin Senorski E, Sundemo D, et al. Adolescents and female patients are at increased risk for contralateral anterior cruciate ligament reconstruction: a cohort study from the Swedish National Knee Ligament Register based on 17,682 patients. *Knee Surg Sports Traumatol Arthrosc.* 2017;25(12):3938–3944. doi:10.1007/s00167-017-4517-7
48. Bram JT, Magee LC, Mehta NN, Patel NM, Ganley TJ. Anterior cruciate ligament injury incidence in adolescent athletes: a systematic review and meta-analysis. *Am J Sports Med.* 2021;49(7):1962–1972. doi:10.1177/0363546520959619
49. Andernord D, Desai N, Björnsson H, Gillén S, Karlsson J, Samuelsson K. Predictors of contralateral anterior cruciate ligament reconstruction: a cohort study of 9061 patients with 5-year follow-up. *Am J Sports Med.* 2015;43(2):295–302. doi:10.1177/0363546514557245
50. Fältström A, Hägglund M, Magnusson H, Forssblad M, Kvist J. Predictors for additional anterior cruciate ligament reconstruction: data from the Swedish national ACL register. *Knee Surg Sports Traumatol Arthrosc.* 2016;24(3):885–894. doi:10.1007/s00167-014-3406-6

51. Rahardja R, Zhu M, Love H, Clatworthy MG, Monk AP, Young SW. Rates of revision and surgeon-reported graft rupture following ACL reconstruction: early results from the New Zealand ACL registry. *Knee Surg Sports Traumatol Arthrosc.* 2020;28(7):2194–2202. doi:10.1007/s00167-019-05773-z
52. Svantesson E, Hamrin Senorski E, Baldari A, et al. Factors associated with additional anterior cruciate ligament reconstruction and register comparison: a systematic review on the Scandinavian knee ligament registers. *Br J Sports Med.* 2019;53(7):418–425. doi:10.1136/bjsports-2017-098192
53. Kaeding CC, Pedroza AD, Reinke EK, et al. Risk factors and predictors of subsequent ACL injury in either knee after ACL reconstruction: prospective analysis of 2488 primary ACL reconstructions from the MOON cohort. *Am J Sports Med.* 2015;43(7):1583–1590. doi:10.1177/0363546515578836
54. Freedman KB, D'amato MJ, Nedeff DD, Kaz A, Bach BR. Arthroscopic anterior cruciate ligament reconstruction: a metaanalysis comparing patellar tendon and hamstring tendon autografts. *Am J Sports Med.* 1966;31:2–11.
55. Persson A, Fjeldsgaard K, Gjertsen JE, et al. Increased risk of revision with hamstring tendon grafts compared with patellar tendon grafts after anterior cruciate ligament reconstruction: a study of 12,643 patients from the Norwegian cruciate ligament registry, 2004–2012. *Am J Sports Med.* 2014;42(2):285–291. doi:10.1177/0363546513511419
56. Gifstad T, Foss OA, Engebretsen L, et al. Lower risk of revision with patellar tendon autografts compared with hamstring autografts: a registry study based on 45,998 primary ACL Reconstructions in Scandinavia. *Am J Sports Med.* 2014;42(10):2319–2328. doi:10.1177/0363546514548164
57. Zhao L, Lu M, Deng M, Xing J, He L, Wang C. Outcome of bone-patellar tendon-bone vs hamstring tendon autograft for anterior cruciate ligament reconstruction: a meta-analysis of randomized controlled trials with a 5-year minimum follow-up. *Medicine.* 2020;99(48):e23476. doi:10.1097/MD.00000000000023476
58. Spindler KP, Huston LJ, Zajchek A, et al. Anterior cruciate ligament reconstruction in high school and college-aged athletes: does autograft choice influence anterior cruciate ligament revision rates? *Am J Sports Med.* 2020;48(2):298–309. doi:10.1177/0363546519892991
59. Volpi P, Cervellin M, Denti M, et al. ACL reconstruction in sports active people: transtibial DB technique with ST/G vs. transtibial SB technique with BPTB: preliminary results. *Injury.* 2010;41(11):1168–1171. doi:10.1016/j.injury.2010.09.029
60. Zaffagnini S, Bruni D, Marcheggiani Muccioli GM, et al. Single-bundle patellar tendon versus non-anatomical double-bundle hamstrings ACL reconstruction: a prospective randomized study at 8-year minimum follow-up. *Knee Surg Sports Traumatol Arthrosc.* 2011;19(3):390–397. doi:10.1007/s00167-010-1225-y
61. Björnsson H, Samuelsson K, Sundemo D, et al. A randomized controlled trial with mean 16-Year follow-up comparing hamstring and patellar tendon autografts in anterior cruciate ligament reconstruction. *Am J Sports Med.* 2016;44(9):2304–2313. doi:10.1177/0363546516646378
62. Beard DJ, Anderson JL, Davies S, Price AJ, Dodd CA. Hamstrings vs. patella tendon for anterior cruciate ligament reconstruction: a randomised controlled trial. *Knee.* 2001;8(1):45–50. doi:10.1016/s0968-0160(01)00062-x
63. Aune AK, Holm I, Risberg MA, Jensen HK, Steen H. Four-strand hamstring tendon autograft compared with patellar tendon-bone autograft for anterior cruciate ligament reconstruction. A randomized study with two-year follow-up. *Am J Sports Med.* 2001;29(6):722–728. doi:10.1177/03635465010290060901
64. Holm I, Oiestad BE, Risberg MA, Aune AK. No difference in knee function or prevalence of osteoarthritis after reconstruction of the anterior cruciate ligament with 4-strand hamstring autograft versus patellar tendon-bone autograft: a randomized study with 10-year follow-up. *Am J Sports Med.* 2010;38(3):448–454. doi:10.1177/0363546509350301
65. Kautzner J, Kos P, Hanus M, Trc T, Havlas V. A comparison of ACL reconstruction using patellar tendon versus hamstring autograft in female patients: a prospective randomised study. *Int Orthop.* 2015;39(1):125–130. doi:10.1007/s00264-014-2495-7
66. Beynon BD, Johnson RJ, Fleming BC, et al. Anterior cruciate ligament replacement: comparison of bone-patellar tendon-bone grafts with two-strand hamstring grafts. A prospective, randomized study. *J Bone Joint Surg Am.* 2002;84(9):1503–1513. doi:10.2106/00004623-200209000-00001
67. Maletis GB, Cameron SL, Tengan JJ, Burchette RJ. A prospective randomized study of anterior cruciate ligament reconstruction: a comparison of patellar tendon and quadruple-strand semitendinosus/gracilis tendons fixed with bioabsorbable interference screws. *Am J Sports Med.* 2007;35(3):384–394. doi:10.1177/0363546506294361
68. Cavaignac E, Coulin B, Tscholl P. Is quadriceps tendon autograft a better choice than hamstring autograft for anterior cruciate ligament reconstruction? A comparative study with a mean follow-up of 3.6 years. *Am J Sports Med.* 2017;45(6):1326–1332. doi:10.1177/0363546516688665
69. Kim SJ, Kumar P, Oh KS. Anterior cruciate ligament reconstruction: autogenous quadriceps tendon-bone compared with bone-patellar tendon-bone grafts at 2-year follow-up. *Arthroscopy.* 2009;25(2):137–144. doi:10.1016/j.arthro.2008.09.014
70. Lind M, Strauss MJ, Nielsen T, Engebretsen L. Quadriceps tendon autograft for anterior cruciate ligament reconstruction is associated with high revision rates: results from the Danish Knee Ligament Registry. *Knee Surg Sports Traumatol Arthrosc.* 2020;28(7):2163–2169. doi:10.1007/s00167-019-05751-5
71. Geib TM, Shelton WR, Phelps RA, Clark L. Anterior cruciate ligament reconstruction using quadriceps tendon autograft: intermediate-term outcome. *Arthroscopy.* 2009;25(12):1408–1414. doi:10.1016/j.arthro.2009.06.004
72. Hettrich CM, Dunn WR, Reinke EK, Spindler KP. The rate of subsequent surgery and predictors after anterior cruciate ligament reconstruction: two- and 6-year follow-up results from a multicenter cohort. *Am J Sports Med.* 2013;41(7):1534–1540. doi:10.1177/0363546513490277
73. Haybäck G, Raas C, Rosenberger R. Failure rates of common grafts used in ACL reconstructions: a systematic review of studies published in the last decade. *Arch Orthop Trauma Surg.* 2021. doi:10.1007/s00402-021-04147-w
74. Yang X, Wang F, He X, et al. Network meta-analysis of knee outcomes following anterior cruciate ligament reconstruction with various types of tendon grafts. *Int Orthop.* 2020;44(2):365–380. doi:10.1007/s00264-019-04417-8
75. Foster TE, Wolfe BL, Ryan S, Silvestri L, Krall Kaye E. Does the graft source really matter in the outcome of patients undergoing anterior cruciate ligament reconstruction?: an evaluation of autograft versus allograft reconstruction results: a systematic review. *Am J Sports Med.* 2010;38(1):189–199. doi:10.1177/0363546509356530
76. Bugelli G, Dell'Osso G, Ascione F, Gori E, Bottai V, Giannotti S. LARS™ in ACL reconstruction: evaluation of 60 cases with 5-year minimum follow-up. *Musculoskelet Surg.* 2018;102(1):57–62. doi:10.1007/s12306-017-0499-3

77. Tiefenboeck TM, Thurmaier E, Tiefenboeck MM, et al. Clinical and functional outcome after anterior cruciate ligament reconstruction using the LARS™ system at a minimum follow-up of 10 years. *Knee*. 2015;22(6):565–568. doi:10.1016/j.knee.2015.06.003
78. Fältström A, Hägglund M, Kvist J. Patient-reported knee function, quality of life, and activity level after bilateral anterior cruciate ligament injuries. *Am J Sports Med*. 2013;41(12):2805–2813. doi:10.1177/0363546513502309
79. Bourke HE, Salmon LJ, Waller A, Patterson V, Pinczewski LA. Survival of the anterior cruciate ligament graft and the contralateral ACL at a minimum of 15 years. *Am J Sports Med*. 2012;40(9):1985–1992. doi:10.1177/0363546512454414
80. Wright RW, Magnussen RA, Dunn WR, Spindler KP. Ipsilateral graft and contralateral ACL rupture at five years or more following ACL reconstruction: a systematic review. *J Bone Joint Surg*. 2011;93(12):1159–1165. doi:10.2106/JBJS.J.00898
81. Webster KE, Feller JA. Exploring the high reinjury rate in younger patients undergoing anterior cruciate ligament reconstruction. *Am J Sports Med*. 2016;44(11):2827–2832. doi:10.1177/0363546516651845
82. Cristiani R, Forssblad M, Edman G, Eriksson K, Stålmán A. Age, time from injury to surgery and hop performance after primary ACLR affect the risk of contralateral ACLR. *Knee Surg Sports Traumatol Arthrosc*. 2021;29:4154–4162. doi:10.1007/s00167-021-06759-6
83. Cronström A, Tengman E, Häger CK. Risk factors for contra-lateral secondary anterior cruciate ligament injury: a systematic review with meta-analysis. *Sports Med*. 2021;51(7):1419–1438. doi:10.1007/s40279-020-01424-3
84. Gallo MC, Bolia IK, Jalali O, et al. Risk factors for early subsequent (Revision or Contralateral) ACL reconstruction: a retrospective database study. *Orthop J Sports Med*. 2020;8(2):232596711990117. doi:10.1177/2325967119901173
85. Shelbourne KD, Gray T, Haro M. Incidence of subsequent injury to either knee within 5 years after anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sports Med*. 2009;37(2):246–251. doi:10.1177/0363546508325665
86. Leys T, Salmon L, Waller A, Linklater J, Pinczewski L. Clinical results and risk factors for reinjury 15 years after anterior cruciate ligament reconstruction: a prospective study of hamstring and patellar tendon grafts. *Am J Sports Med*. 2012;40(3):595–605. doi:10.1177/0363546511430375
87. Rahardja R, Love H, Clatworthy MG, Monk AP, Young SW. Higher rate of return to preinjury activity levels after anterior cruciate ligament reconstruction with a bone–patellar tendon–bone versus hamstring tendon autograft in high-activity patients: results from the New Zealand ACL registry. *Am J Sports Med*. 2021;49(13):3488–3494. doi:10.1177/03635465211044142
88. Ardern CL, Webster KE, Taylor NF, Feller JA. Return to sport following anterior cruciate ligament reconstruction surgery: a systematic review and meta-analysis of the state of play. *Br J Sports Med*. 2011;45(7):596–606. doi:10.1136/bjsm.2010.076364
89. Ardern CL, Taylor NF, Feller JA, Webster KE. Fifty-five per cent return to competitive sport following anterior cruciate ligament reconstruction surgery: an updated systematic review and meta-analysis including aspects of physical functioning and contextual factors. *Br J Sports Med*. 2014;48(21):1543–1552. doi:10.1136/bjsports-2013-093398
90. Gupta R, Kapoor A, Soni A, Khatri S, Masih GD. Anterior cruciate ligament reconstruction with bone-patellar tendon-bone graft is associated with higher and earlier return to sports as compared to hamstring tendon graft. *Knee Surg Sports Traumatol Arthrosc*. 2020;28(11):3659–3665. doi:10.1007/s00167-020-06074-6
91. DeFazio MW, Curry EJ, Gustin MJ, et al. Return to sport after ACL reconstruction with a BTB versus hamstring tendon autograft: a systematic review and meta-analysis. *Orthop J Sports Med*. 2020;8(12):232596712096491. doi:10.1177/2325967120964919
92. Kartus J, Movin T, Karlsson J. Donor-site morbidity and anterior knee problems after anterior cruciate ligament reconstruction using autografts. *Arthroscopy*. 2001;17(9):971–980. doi:10.1053/jars.2001.28979
93. Lind M, Nielsen TG, Soerensen OG, Mygind-Klavsen B, Faunø P. Quadriceps tendon grafts does not cause patients to have inferior subjective outcome after anterior cruciate ligament (ACL) reconstruction than do hamstring grafts: a 2-year prospective randomised controlled trial. *Br J Sports Med*. 2020;54(3):183–187. doi:10.1136/bjsports-2019-101000
94. Marques F, Barbosa PHB, Alves PR, et al. Anterior knee pain after anterior cruciate ligament reconstruction. *Orthop J Sports Med*. 2020;8(10):232596712096108. doi:10.1177/2325967120961082
95. Xie X, Xiao Z, Li Q, et al. Increased incidence of osteoarthritis of knee joint after ACL reconstruction with bone–patellar tendon–bone autografts than hamstring autografts: a meta-analysis of 1443 patients at a minimum of 5 years. *Eur J Orthopaed Surg Traumatol*. 2015;25(1):149–159. doi:10.1007/s00590-014-1459-3
96. Mastrokalos DS, Springer J, Siebold R, Paessler HH. Donor site morbidity and return to the preinjury activity level after anterior cruciate ligament reconstruction using ipsilateral and contralateral patellar tendon autograft: a retrospective, nonrandomized study. *Am J Sports Med*. 2005;33(1):85–93. doi:10.1177/0363546504265926
97. Kartus J, Ejerhed L, Sernert N. Comparison of traditional and subcutaneous patellar tendon harvest a prospective study of donor site-related problems after anterior cruciate ligament reconstruction using different graft harvesting techniques. *Am J Sports Med*. 2000;28:328–335.
98. Kartus J, Stener S, Lindahl S, et al. Factors affecting donor-site morbidity after anterior cruciate ligament reconstruction using bone-patellar tendon-bone autografts. *Knee Surg, Sports Traumatol, Arthrosc*. 1997;5:222–228.
99. Kartus J, Magnusson L, Stener S, Brandsson S, Eriksson BI, Karlsson J. Complications following arthroscopic anterior cruciate ligament reconstruction. A 2–5-year follow-up of 604 patients with special emphasis on anterior knee pain. *Knee Surg Sports Traumatol Arthrosc*. 1999;7(1):2–8. doi:10.1007/s001670050112
100. Benner RW, Shelbourne KD, Freeman H. Infections and patellar tendon ruptures after anterior cruciate ligament reconstruction: a comparison of ipsilateral and contralateral patellar tendon autografts. *Am J Sports Med*. 2011;39(3):519–525. doi:10.1177/0363546510388163
101. Tay G, Warrior B, Riquis G. Indirect patella fractures following ACL reconstruction: a review. *Acta Orthopaedica*. 2006;77(3):494–500. doi:10.1080/17453670610046451
102. Stein DA, Hunt SA, Rosen JE, Sherman OH. The incidence and outcome of patella fractures after anterior cruciate ligament reconstruction. *Arthroscopy*. 2002;18(6):578–583. doi:10.1053/jars.2002.30658
103. Viola R, Vianello R. Three cases of patella fracture in 1320 anterior cruciate ligament reconstructions with bone-patellar tendon-bone autograft. *Arthroscopy*. 1999;15(1):93–97. doi:10.1053/ar.1999.v15.015009
104. Chouteau J, Laptouï D, Lerat JL, Moyer B. Fracture de la patella après prélèvement os-tendon patellaire-os pour ligamentoplastie de reconstruction du ligament croisé antérieur: incidence du type de prélèvement [Patellar fracture after anterior cruciate ligament reconstruction using a bone patellar tendon bone transplant: a comparative study of two harvesting techniques]. *Revue de chirurgie orthopedique et reparatrice de l'appareil moteur*. 2008;94(6):561–565. French. doi:10.1016/j.rco.2007.12.002

105. Lee GH, McCulloch P, Cole BJ, Bush-Joseph CA, Bach BR. The incidence of acute patellar tendon harvest complications for anterior cruciate ligament reconstruction. *Arthrosc- J Arthrosc Relat Surg.* 2008;24(2):162–166. doi:10.1016/j.arthro.2007.08.020
106. Miroslav MZ, Robert S, Natasa M, Vladimir H. Reconstruction of patellar tendon rupture after anterior cruciate ligament reconstruction: a case report. *Knee.* 2008;15(5):419–422. doi:10.1016/j.knee.2008.05.009
107. Marumoto JM, Mitsunaga MM, Richardson AB, Medoff RJ, Mayfield GW. Late patellar tendon ruptures after removal of the central third for anterior cruciate ligament reconstruction. A report of two cases. *Am J Sports Med.* 1996;24(5):698–701. doi:10.1177/036354659602400524
108. Mickelsen PL, Morgan SJ, Johnson WA, Ferrari JD. Patellar tendon rupture 3 years after anterior cruciate ligament reconstruction with a central one third bone-patellar tendon-bone graft. *Arthroscopy.* 2001;17(6):648–652. doi:10.1053/jars.2001.23577
109. Bonamo JJ, Krinick RM, Sporn AA. Rupture of the patellar ligament after use of its central third for anterior cruciate reconstruction. A report of two cases. *J Bone Joint Surg Am.* 1984;66(8):1294–1297. doi:10.2106/00004623-198466080-00020
110. Salvi AE, Metelli GP, Musella G. Spontaneous patellar fracture in a soccer player occurred 3 years and 3 months after anterior cruciate ligament reconstruction with central third bone-patellar tendon-bone graft. *Knee Surg Sports Traumatol Arthrosc.* 2007;15(10):1280–1281. doi:10.1007/s00167-007-0370-4
111. Benner RW, Shelbourne KD, Urch SE, Lazarus D. Tear patterns, surgical repair, and clinical outcomes of patellar tendon ruptures after anterior cruciate ligament reconstruction with a bone-patellar tendon-bone autograft. *Am J Sports Med.* 2012;40(8):1834–1841. doi:10.1177/0363546512449815
112. Webster KE, Feller JA, Hartnett N, Leigh WB, Richmond AK. Comparison of patellar tendon and hamstring tendon anterior cruciate ligament reconstruction: a 15-year follow-up of a randomized controlled trial. *Am J Sports Med.* 2016;44(1):83–90. doi:10.1177/0363546515611886
113. Castoldi M, Magnussen RA, Gunst S, et al. A randomized controlled trial of bone-patellar tendon-bone anterior cruciate ligament reconstruction with and without lateral extra-articular tenodesis: 19-year clinical and radiological follow-up. *Am J Sports Med.* 2020;48(7):1665–1672. doi:10.1177/0363546520914936
114. Mao Y, Zhang K, Li J, Fu W. Supplementary lateral extra-articular tenodesis for residual anterolateral rotatory instability in patients undergoing single-bundle anterior cruciate ligament reconstruction: a meta-analysis of randomized controlled trials. *Orthop J Sports Med.* 2021;9(5):232596712110022. doi:10.1177/23259671211002282
115. Beckers L, Vivacqua T, Firth AD, Getgood AMJ. Clinical outcomes of contemporary lateral augmentation techniques in primary ACL reconstruction: a systematic review and meta-analysis. *J Exp Orthopaed.* 2021;8(1). doi:10.1186/s40634-021-00368-5
116. Na BR, Kwak WK, Seo HY, Seon JK. Clinical outcomes of anterolateral ligament reconstruction or lateral extra-articular tenodesis combined with primary ACL reconstruction: a systematic review with meta-analysis. *Orthop J Sports Med.* 2021;9(9):232596712110230. doi:10.1177/23259671211023099
117. Getgood AMJ, Bryant DM, Litchfield R, et al. Lateral extra-articular tenodesis reduces failure of hamstring tendon autograft anterior cruciate ligament reconstruction: 2-year outcomes from the STABILITY study randomized clinical trial. *Am J Sports Med.* 2020;48(2):285–297. doi:10.1177/0363546519896333
118. Sonnery-Cottet B, Haidar I, Rayes J, et al. Long-term graft rupture rates after combined acl and anterolateral ligament reconstruction versus isolated ACL reconstruction: a matched-pair analysis from the SANTI study group. *Am J Sports Med.* 2021;49(11):2889–2897. doi:10.1177/03635465211028990

Open Access Journal of Sports Medicine

Dovepress

Publish your work in this journal

Open Access Journal of Sports Medicine is an international, peer-reviewed, open access journal publishing original research, reports, reviews and commentaries on all areas of sports medicine. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/open-access-journal-of-sports-medicine-journal>