

# Synthesis of structurally diverse major groove DNA interstrand crosslinks using three different aldehyde precursors

Shivam Mukherjee<sup>1</sup>, Angelo Guainazzi<sup>2</sup> and Orlando D. Schärer<sup>1,2,\*</sup>

<sup>1</sup>Department of Chemistry, Stony Brook University, Stony Brook, NY 11794-3400, USA and <sup>2</sup>Department of Pharmacological Sciences, Stony Brook University, Stony Brook, NY 11794-8651, USA

Received March 3, 2014; Revised April 03, 2014; Accepted April 4, 2014

## ABSTRACT

**DNA interstrand crosslinks (ICLs) are extremely cytotoxic lesions that block essential cellular processes, such as replication and transcription. Crosslinking agents are widely used in cancer chemotherapy and form an array of structurally diverse ICLs. Despite the clinical success of these agents, resistance of tumors to crosslinking agents, for example, through repair of these lesions by the cellular machinery remains a problem. We have previously reported the synthesis of site-specific ICLs mimicking those formed by nitrogen mustards to facilitate the studies of cellular responses to ICL formation. Here we extend these efforts and report the synthesis of structurally diverse major groove ICLs that induce severe, little or no distortion in the DNA. Our approach employs the incorporation of aldehyde precursors of different lengths into complementary strands and ICL formation using a double reductive amination with a variety of amines. Our studies provide insight into the structure and reactivity parameters of ICL formation by double reductive amination and yield a set of diverse ICLs that will be invaluable for exploring structure–activity relationships in ICL repair.**

## INTRODUCTION


DNA interstrand crosslinks (ICLs) are extremely cytotoxic lesions that covalently connect two complementary strands of a DNA duplex. As DNA strand separation is essential to cellular processes, such as transcription and replication, ICLs are highly cytotoxic (1–3). Bifunctional electrophiles, such as cisplatin, nitrogen mustards (NMs) chloro ethyl nitroso ureas and mitomycin C, form ICLs and are a mainstay as frontline chemotherapeutic drugs (4). Despite the clinical success of these drugs, resistance mechanisms in-

cluding the removal of ICLs from DNA by cellular proteins contributes to the resistance of tumor cells to treatment with crosslinking agents (5). ICLs are also formed by endogenous bifunctional agents, such as malondialdehyde and formaldehyde (6–8), and such agents have likely been the evolutionary drivers for the cellular responses to ICL formation. The importance of these repair pathways is underscored by the existence of the hereditary cancer prone disorder Fanconi anemia (FA) (9,10). Cells from FA patients display exquisite sensitivity to ICL-forming agents and exposure to crosslinking agents serves as the definitive clinical diagnosis for FA.


The repair of ICLs is a complex process and the pathways involved have been subject of intense recent studies (11). These efforts have been critically dependent on the ability to generate site-specific ICL (reviewed in (12)). Breakthroughs in the synthesis of site-specific ICLs have been achieved by a number of laboratories by incorporating ICL precursors into DNA as phosphoramidites by solid phase DNA synthesis. Such approaches made use of the incorporation of nucleosides crosslinked outside of the DNA (7,13–15) and the incorporation of ICL precursors on one or two strands of DNA and the use of specific reaction to induce ICL formation (16–26). These efforts have provided access to structurally diverse ICLs formed in the major and minor grooves, on the Watson–Crick base pairing face or stacked between the two bases. ICLs induce a variety of alterations in DNA duplex structures, due to differences in the attachment sites on the DNA bases and the chemical composition of the crosslink. Recent studies have revealed that the biological responses triggered by ICL can vary depending on how they influence duplex structure (27–32).

Our own studies have been mainly concerned with major groove ICLs (17,22), the site where adducts are formed by the clinically important cisplatin and nitrogen mustards or the environmental pollutant diepoxybutane. The length of the crosslink agent bridging two N7 atoms of dG residues on opposite strands of ICLs formed in the major groove

\*To whom correspondence should be addressed. Tel: +1 631 632 7545; Fax: +1 631 632 7546; Email: orlando.scharer@stonybrook.edu  
Present address:  
Angelo Guainazzi, Helsinn Therapeutics (U.S.) Inc., Bridgewater, NJ 08807, USA.



**Figure 1.** Native and synthetic nitrogen mustard ICLs. **A.** Nitrogen mustard and the DNA ICL formed by reaction with duplex DNA. The structure of the side chain R, varies with the type of nitrogen mustard for example 'CH<sub>3</sub>' is for mechlorethamine, the first clinically used antitumor drug. **B.** Our approach to forming ICLs having possible linker lengths from 4.8 to 13.2 Å uses a combination of ICL precursors, a 7-alkylaldehyde-7-deazaguanine with C1, C2 or C3 alkyl chains in complementary strands of a DNA duplex and a double reductive amination reaction with ammonia, hydrazine or DMEDA. Note that guanine residues at the ICL are replaced with 7-deazaguanine to facilitate ICL synthesis and increase stability.



**Figure 2.** Distance between N7 position on opposing strand of two G residues in B-form DNA and the theoretical length of the alkylamine chain linker in ICLs. The lengths of the alkyl chains are indicated over the dotted lines represent the maximum possible length in a fully extended all *anti*-conformation.

lead to different degrees of distortion in DNA, dictated by length of the ICL linking the two bases (33–36).

We have synthesized ICLs using precursor molecules containing reactive aldehyde groups masked as diols that were incorporated into complementary strands of a duplex

and by using a double reductive amination reaction (Figure 1). Using aldehyde precursors connected to the base through one or two carbon alkyl chains and using a variety of amines, we found that the yield of ICL formation dramatically depended on the length of the ICL formed, and the reactivity of the amine used (17,22). Here we extend these studies through the synthesis of a new ICL precursor and a systematic assessment of reaction conditions for ICL formation, providing access to a variety of NM ICL mimics inducing moderate, mild or no distortion in the DNA duplex. We expect that these structurally diverse ICLs will be invaluable for structure–function relationships of ICL repair, extending previous studies that have started to reveal important differences in how ICLs of different structures are processed in replication-dependent and replication-independent repair (27,29,37).

## MATERIALS AND METHODS

### Synthesis of the C3-aldehyde phosphoramidite **8**


The synthesis of phosphoramidite **8**, including protocols for the synthesis and structural characterization (<sup>1</sup>H-NMR, <sup>13</sup>C-NMR, HR-MS) of each intermediate is described in detail in the Supplementary Data section.

### Oligonucleotide synthesis and purification

Oligonucleotide synthesis was carried out on an Expedite 8909 Nucleic Acid Synthesis System (Applied Biosystems) using 1 μmol 1000 Å CPG –dG and –dC column cartridges (Biosearch Technologies). An extended coupling time of 15 min was used with all the modified phosphoramidite building blocks. The following sequences were synthesized: 5'-d(GTCACTGGTAG\*ACAGCATTG) and 5'-d(CAATGCTG\*TCTACCAGTGAC), where G\* is the modified phosphoramidite having either C1, C2 or C3 linker. The trityl-on oligonucleotides were deprotected by treatment with concentrated NH<sub>4</sub>OH solution at 50°C for 12 h, then purified and detritylated using 1 μmol TOP-cartridges (Agilent Technologies) following the manufacturer's protocol. The *tert*-butyldimethylsilyl (TBDMS) group in the oligonucleotides containing the C3 aldehyde precursor was removed by treating with TEA·3HF overnight at 40°C followed by precipitation using 1-butanol. Subsequently, oligonucleotides were purified by high pressure liquid chromatography (HPLC) on a C18 column (Phenomenex Clarity 5μ Oligo-RP 50 × 10 mm) using the following elution gradient: linear 2.5–15% B over 21 min, linear 15–90% B over 25 min, isocratic 90% B till 26 min, linear 90–2.5% B till 28 min, isocratic 2.5% B till 30 min (eluent A: 0.1 M TEAA (pH 7); eluent B: CH<sub>3</sub>CN).

### Formation of ICLs by reductive amination

A solution containing two single stranded oligonucleotides (25 nmols in 125 μl 100 mM NaCl) was heated to 95°C and allowed to cool over a period of 5 h for annealing to take place. For oxidation, 10 μl 50 mM NaIO<sub>4</sub> and 15 μl 1 M sodium phosphate buffer (pH 5.4) was added and mixture was kept at 4°C overnight. Excess NaIO<sub>4</sub> was removed by




**Figure 3.** Scheme showing the synthesis of the C3 alkyl chain ICL precursor phosphoramidite **8**. Reagents, conditions and yields: (a)  $\text{PPh}_3$ ,  $\text{CBr}_4$ ,  $\text{CH}_2\text{Cl}_2$ , 3 h, 84%; (b) Dowex-50W, MeOH, 12 h,  $50^\circ\text{C}$ , 95%; (c) TBDMSCl, Imidazole, DMF, 12 h, 83%; (d)  $\text{EtMgBr}$ , THF, 3 h, 90%; (e) **4**,  $\text{Pd}(\text{PPh}_3)_4$ ,  $\text{CuI}$ ,  $\text{Et}_3\text{N}$ , DMF, 24 h,  $60^\circ\text{C}$ , 59%; (f)  $\text{H}_2$ , Pd/C,  $\text{EtOAc}/\text{MeOH}$ , 42h, 84%; (g) NaOMe, THF, 6 h, 90%; (h) DMTrCl, Py, 1.5 h, 72%; (i)  $(i\text{Pr})_2\text{NP}(\text{Cl})\text{OEtCN}$ ,  $\text{CH}_2\text{Cl}_2$ , 1.5 h, 71%; (j) solid phase DNA synthesis; (k)  $\text{NH}_3$ , 16 h,  $50^\circ\text{C}$ ; (l) TEA.3HF,  $40^\circ\text{C}$ , o.n.; (m)  $\text{NaIO}_4$ , pH = 5.4.

centrifugation through Microcon columns with a 3K cut-off (Millipore). The crosslink was formed by adding  $10\ \mu\text{l}$  5 mM aqueous solution of the amine (ammonium acetate, hydrazine or  $N,N'$ -dimethylethylene diamine (DMEDA)) and  $10\ \mu\text{l}$  of a 0.5 M solution of sodium cyanoborohydride and allowing to react in the dark at room temperature overnight. ICL formation was analyzed by gel electrophoresis on a 20% denaturing polyacrylamide gel (7 M urea). To purify and isolate the ICL-containing oligonucleotide the gel was visualized by UV shadowing and the crosslinked oligonucleotide was excised from the gel and the DNA was extracted by electroelution using the Elutrap device (Schleicher & Schuell). After purification the isolated yields of ICLs were in the range from 5 to 12 nmols.

## RESULTS AND DISCUSSION

### Strategy for the synthesis of structurally diverse ICLs

We have previously reported that NM-like ICLs can be synthesized by incorporation of two aldehyde precursors on complementary strands of DNA and crosslink formation using a double reductive amination reaction (17,22). In the course of these studies, we noted that the efficiency of ICL formation is dependent upon the length of the ICL as well as the reactivity of the amine used in the reaction (Figure 1B). This is illustrated by our attempts to synthesize an ICL isosteric to those formed by NMs that contain a 5-atom chain in the crosslink and induce a bend of about  $20^\circ$  in the DNA ( $n = 1$ , Figure 1). We observed that formation of this crosslink was unsuccessful with two aldehyde precursors with a C2 alkyl chain and ammonia as the amine, while it could be formed with a C2 and C1 alkyl and hydrazine as an amine.



**Figure 4.** ICL formation by reductive amination using C1, C2 and C3 precursors. Analysis of ICL formation of the reaction of duplexes containing C1, C2 and C3 precursors in the presence of amines and  $\text{NaBH}_3\text{CN}$ . Reactions were analyzed by denaturing polyacrylamide gel electrophoresis and stained with methylene blue. The sequences used were 5'-d(GTCACTGGTAG\*ACAGCATTG) and 5'-d(CAATGCTG\*TCTACCAGTGAC), where G\* represents the modified guanine having either C1, C2 or C3 diol. As a control for each set of ICL formation, reactions in which the amine or  $\text{NaBH}_3\text{CN}$  was omitted were performed. The precursor and amine used are indicated over each gel.  $\text{NH}_3$ , ammonia; HY, hydrazine; DM, DMEDA.

By contrast, an ICL that does not induce a distortion can be formed with a number of different amines (17,22). These examples show that the length of the ICL to be formed and the reactivity of the amine (hydrazine is more nucleophilic than ammonia) determine the success of ICL formation in our approach. We wished to more fully exploit our approach to generate structurally diverse major groove ICLs by double reductive amination and to study the parameters of reactivity that govern crosslink formation.

To be able to access ICLs spanning a broad range of distances across the major groove, we set out to synthesize a new ICL precursor with a three-carbon alkylaldehyde (C3) precursor. Together with the previously synthesized C1 and C2 aldehyde precursors and using ammonia, hydrazine and  $\text{N,N}'$ -dimethylethylenediamine (DMEDA) this should allow us to generate a set of structurally diverse ICLs. We de-

fine here the length of an ICL based on the fully extended alkylamine chain spanning the distance between the two crosslinked deazaguanine residues (illustrated in Figure 2). The distance between the atoms at the 7 position of dG where the ICL is attached is approximately 8.9 Å in B-form DNA, so any ICL that spans a length of 8.9 Å or less will induce some form of distortion in the DNA.

#### Synthesis of a three-carbon alkyl aldehyde chain precursor


The synthesis of the C3 precursor **8** started with a Sonogashira cross-coupling reaction between 7-Iodo-7-deazaguanosine (**5**) (17) with the protected alkynediol **4** (Figure 3). Note that **4** was synthesized starting from the protected (*S*)-glyceraldehyde **1**, which was converted to the dibromo-olefin **2** (38) and reprotected as TBDMS-ether **3**, compatible for solid phase DNA synthesis. Treatment of **3** with  $\text{EtMgBr}$  gave acetylene **4** (39). The alkyne group in **6** was reduced by passing hydrogen over palladium catalyst. Subsequently, protecting groups on the 5'- and 3'- positions of the sugar were exchanged using standard reagents and conditions to generate the phosphoramidite **8**. Using solid phase DNA synthesis, **8** was incorporated in two complementary 20-mer strands containing a 5'-G\*NC sequence, where G\* denotes the aldehyde-bearing residue. The oligonucleotides were deprotected—first with concentrated  $\text{NH}_3$  to remove the standard base and phosphate protecting groups and then with TEA·3HF to remove the TBDMS groups, generating the free diol. The oligonucleotides were purified by HPLC and the presence of the diol in the modified residues was confirmed by MALDI-TOF mass spectrometry measurements (see Supplementary Data). In addition, 20-mer oligonucleotides with the C1 and C2 aldehyde precursors on complementary strands were also synthesized and purified according to our earlier published protocol (17,22).

#### ICL formation using three different length alkyl aldehyde precursors

We studied ICL formation using the three different length precursors (C1, C2 and C3) and three amines (ammonia, hydrazine and DMEDA) in a systematic manner (Figure 4). ICLs were in 5'-G\*NC sequences, the preferred site of ICL formation by nitrogen mustards (40,41). As previously discussed, the efficiency of ICL formation is dependent on both the reactivity of the amine as well as the distortion induced in the DNA. Duplex bearing diol-containing ICL precursors were annealed, the diol oxidized to the aldehyde using sodium periodate and ICL formation induced by incubation with one of the three amines in the presence of sodium cyanoborohydride. The identity of the ICLs formed was confirmed by MALDI-TOF measurements (see Supplementary Data). With a duplex having C1 precursors on each strand, ICL formation, indicated by the appearance of a band with slower mobility on a denaturing polyacrylamide gel, was only observed with DMEDA, yielding a 8.4 Å linkage (Figure 4, lanes 3). By contrast, we did not observe the formation of the 4.8 Å or 6.0 Å ICLs with two C1 precursors and ammonia and/or hydrazine, respectively

Precursors	C1			C2			C3			C3			C3					
Amines	NH <sub>3</sub>	HY	DM	NH <sub>3</sub>	HY	DM	NH <sub>3</sub>	HY	DM	NH <sub>3</sub>	HY	DM	NH <sub>3</sub>	HY	DM			
Bridge Length (Å)	4.8	6.0	8.4	6.1	7.2	9.6	7.2	8.4	10.8	7.2	8.4	10.8	8.4	9.6	12.0	9.6	10.8	13.2
% ICL	-	-	+	-	+	++	-	+	++	-	++	+++	++	++	+++	++	+++	+++

**Figure 5.** Overview of the reaction with all the ICL reactions tested. Reactions with the C1, C2 and C3 precursors and the three different amines (NH<sub>3</sub>, HY and DMEDA), the theoretical bridge lengths and relative qualitative yields are indicated. '+' low yield, '++' moderate yield and '+++' for high yield. As staining with methylene blue does not allow for the quantitative determination of the ratio of ss- versus ds-DNA, qualitative yields are indicated.



**Figure 6.** Models of structures of nitrogen mustard ICLs. **A.** C1/C2/HY ICL isosteric to a native nitrogen mustard ICL with a bridge length of 7.2 Å inducing a 20° bend in the DNA as predicted by molecular modeling studies (22); **B.** C2/C3/NH<sub>3</sub> ICL with a bridge length of 8.4 Å, inducing a distortion in the DNA. **C.** C3/C3/NH<sub>3</sub> ICL with a bridge length of 9.6 Å without DNA distortion. The structure in (A) was calculated using molecular dynamic simulations (22), (B) and (C) were manually generated using VMD 1.9 (42).

(Figure 4, lanes 1 and 2). ICL formation was dependent on the presence of the reducing agent and the amine and omission of either one prevented ICL formation (Figure 4, lanes 4 and 5). In a duplex containing a C1 and a C2 aldehyde precursor on complementary strands, ICL formation was observed with hydrazine and DMEDA (Figure 4, lanes 7 and 8), but not with ammonia (Figure 4, lane 6), as previously reported (22). The 7.2 Å ICL containing hydrazine is the shortest ICL that we were able to form in our approach and it is isosteric to the native nitrogen mustard ICL. Having two C2 precursors gave ICLs with hydrazine and DMEDA (Figure 4, lanes 17 and 18), but not with ammonia (Figure 4, lane 16). These results confirmed that an ICL isostructural to that formed by NM with a linker length of 7.2 Å and inducing a bend in the DNA of about 20° can be formed by double reductive amination with hydrazine, but not ammonia, in line with the higher reactivity of hydrazine.

The reactivity patterns of ICL formation were further investigated with the newly synthesized C3 aldehyde ICL precursor paired to a complementary strand with C1 or C2 precursor. As was expected from the results with the C2/C2 precursors (Figure 4, lanes 16–18), ICL formation with the C1/C3 pair did not yield an ICL upon reaction with ammonia, showing that a 7.2 Å ICL cannot be formed with ammonia, irrespective of the position of carbon and nitrogen atoms in the ICL. As expected, ICLs were formed with C1/C3 using hydrazine and DMEDA (Figure 4, lanes 12 and 13) as an amine.

The question then arose whether the combination of C3 and C2 precursors would be able to yield an ICL with NH<sub>3</sub>, with an 8.4 Å linkage. This was indeed the case, and ammonia, hydrazine and DMEDA all formed ICLs with the C2/C3 oligonucleotides (Figure 4, lanes 21–23). This result shows that the minor distortion in the DNA of the 8.4 Å

ICL allowed the reductive amination to occur with ammonia. A band of lesser intensity comigrating with the ICL band was also formed in the absence of an amine with the C2/C3 duplex (Figure 4, lane 24). Although the identity of this band remains to be determined, we speculate that it is the result of a reductive amination reaction of one of the aldehyde groups with an exocyclic amine in the complementary strand of DNA. As expected based on these observations, reactions of the duplex containing two C3 aldehyde precursors, yielded ICL with all of the three amines with lengths of 9.6 Å and more (lanes 26–28, Figure 4).

#### Summary and conclusion: Distance-dependent ICL formation

We used oligonucleotides with 7-deazaguanine residues having alkyl aldehyde chains of different lengths (C1, C2, C3) at the 7 position and studied ICL formation with ammonia, hydrazine and DMEDA using a reductive amination reaction. The efficiency of ICL formation was found to be correlated with the length of the ICL and the reactivity of the amine (summarized in Figure 5). We were able to form ICLs with bridge lengths ranging from 7.2 Å, for which our molecular modeling studies predict a bend of about 20° in the DNA duplex (22), to those of 10.8 Å and more, which our preliminary nuclear magnetic resonance (NMR) experiments show are free of distortion (AG, T. Zaliznyak, C. de los Santos, ODS, unpublished data). ICL formation was found to be most efficient with nondistorting ICLs (10.8–13.2 Å), followed by those with minor (8.4–9.6 Å) and moderate distortion (7.2–8.4 Å) (Figure 5). The higher nucleophilicity and reactivity of hydrazine allowed for the formation of more distorted ICLs. The major groove ICLs reported here inducing no, minor and moderate distortion in DNA duplexes (Figure 6) will be invaluable for advancing

studies elucidating structure–function relationships in ICL repair.

## SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

## ACKNOWLEDGMENTS

We are grateful to Dr. Arthur Campbell for help with Figure 6 and Robert Rieger for help with the HR-MS and MALDI-TOF spectral analyses supported by grant NIH/NCRR 1 S10 RR023680-1.

## FUNDING

National Cancer Institute (NCI) grant [R01 CA165911 to O.D.S.]. Source of open access funding: NCI [CA165911]. *Conflict of interest statement.* None declared.

## REFERENCES

- Lawley,P.D. and Phillips,D.H. (1996) DNA adducts from chemotherapeutic agents. *Mutat. Res.*, **355**, 13–40.
- Noll,D.M., Mason,T.M. and Miller,P.S. (2006) Formation and repair of interstrand cross-links in DNA. *Chem. Rev.*, **106**, 277–301.
- Rajski,S.R. and Williams,R.M. (1998) DNA cross-linking agents as antitumor drugs. *Chem. Rev.*, **98**, 2723–2795.
- Schärer,O.D. (2005) DNA interstrand crosslinks: natural and drug-induced DNA adducts that induce unique cellular responses. *Chembiochem*, **6**, 27–32.
- McHugh,P.J., Spanswick,V.J. and Hartley,J.A. (2001) Repair of DNA interstrand crosslinks: molecular mechanisms and clinical relevance. *Lancet Oncol.*, **2**, 483–490.
- Chaw,Y.F., Crane,L.E., Lange,P. and Shapiro,R. (1980) Isolation and identification of cross-links from formaldehyde-treated nucleic acids. *Biochemistry*, **19**, 5525–5531.
- Harwood,E.A., Sigurdsson,S.T., Edfeldt,N.B.F., Reid,B.R. and Hopkins,P.B. (1999) Chemical synthesis and preliminary structural characterization of a nitrous acid interstrand cross-linked duplex DNA. *J. Am. Chem. Soc.*, **121**, 5081–5082.
- Stone,M.P., Cho,Y.J., Huang,H., Kim,H.Y., Kozekov,I.D., Kozekova,A., Wang,H., Minko,I.G., Lloyd,R.S., Harris,T.M. *et al.* (2008) Interstrand DNA cross-links induced by alpha,beta-unsaturated aldehydes derived from lipid peroxidation and environmental sources. *Acc. Chem. Res.*, **41**, 793–804.
- Langevin,F., Crossan,G.P., Rosado,I.V., Arends,M.J. and Patel,K.J. (2011) Fancd2 counteracts the toxic effects of naturally produced aldehydes in mice. *Nature*, **475**, 53–58.
- Moldovan,G.L. and D'Andrea,A.D. (2009) How the fanconi anemia pathway guards the genome. *Annu. Rev. Genet.*, **43**, 223–249.
- Clauson,C., Schärer,O.D. and Niedernhofer,L. (2013) Advances in understanding the complex mechanisms of DNA interstrand cross-link repair. *Cold Spring Harb. Perspect. Biol.*, **5**, a012732.
- Guainazzi,A. and Schärer,O.D. (2010) Using synthetic DNA interstrand crosslinks to elucidate repair pathways and identify new therapeutic targets for cancer chemotherapy. *Cell. Mol. Life Sci.*, **67**, 3683–3697.
- Noll,D.M., Noronha,A.M. and Miller,P.S. (2001) Synthesis and characterization of DNA duplexes containing an N(4)C-ethyl-N(4)C interstrand cross-link. *J. Am. Chem. Soc.*, **123**, 3405–3411.
- Wilds,C.J., Noronha,A.M., Robidoux,S. and Miller,P.S. (2004) Mispair-aligned N3T-alkyl-N3T interstrand cross-linked DNA: synthesis and characterization of duplexes with interstrand cross-links of variable lengths. *J. Am. Chem. Soc.*, **126**, 9257–9265.
- Wilds,C.J., Xu,F. and Noronha,A.M. (2008) Synthesis and characterization of DNA containing an N-1–2'-deoxyinosine-ethyl-N-3-thymidine interstrand cross-link: a structural mimic of the cross-link formed by 1,3-bis-(2-chloroethyl)-1-nitrosourea. *Chem. Res. Toxicol.*, **21**, 686–695.
- Alzeer,J. and Schärer,O.D. (2006) A modified thymine for the synthesis of site-specific thymine-guanine DNA interstrand crosslinks. *Nucleic Acids Res.*, **34**, 4458–4466.
- Angelov,T., Guainazzi,A. and Schärer,O.D. (2009) Generation of DNA interstrand cross-links by post-synthetic reductive amination. *Org. Lett.*, **11**, 661–664.
- Dooley,P.A., Tsarouhtsis,D., Korbel,G.A., Nechev,L.V., Shearer,J., Zegar,I.S., Harris,C.M., Stone,M.P. and Harris,T.M. (2001) Structural studies of an oligodeoxynucleotide containing a trimethylaldehyde cross-link in a 5'-(CpG) motif: model of a malondialdehyde cross-link. *J. Am. Chem. Soc.*, **123**, 1730–1739.
- Dutta,S., Chowdhury,G. and Gates,K.S. (2007) Interstrand cross-links generated by abasic sites in duplex DNA. *J. Am. Chem. Soc.*, **129**, 1852–1853.
- Erlanson,D.A., Chen,L. and Verdine,G.L. (1993) DNA methylation through a locally unpaired intermediate. *J. Am. Chem. Soc.*, **115**, 12583–12584.
- Ferentz,A.E., Keating,T.E. and Verdine,G.L. (1993) Synthesis and characterization of disulfide cross-linked oligonucleotides. *J. Am. Chem. Soc.*, **115**, 9006–9014.
- Guainazzi,A., Campbell,A.J., Angelov,T., Simmerling,C. and Schärer,O.D. (2010) Synthesis and molecular modeling of a nitrogen mustard DNA interstrand crosslink. *Chem. Eur. J.*, **16**, 12100–12103.
- Guan,L. and Greenberg,M.M. (2009) DNA interstrand cross-link formation by the 1,4-dioxobutane abasic lesion. *J. Am. Chem. Soc.*, **131**, 15225–15231.
- Hentschel,S., Alzeer,J., Angelov,T., Schärer,O.D. and Luedtke,N.W. (2012) Synthesis of DNA interstrand cross-links using a photocaged nucleobase. *Angew Chem. Int. Edit.*, **51**, 3466–3469.
- Kobertz,W.R. and Essigmann,J.M. (1997) Solid-phase synthesis of oligonucleotides containing a site-specific psoralen derivative. *J. Am. Chem. Soc.*, **119**, 5960–5961.
- Kozekov,I.D., Nechev,L.V., Moseley,M.S., Harris,C.M., Rizzo,C.J., Stone,M.P. and Harris,T.M. (2003) DNA interchain cross-links formed by acrolein and crotonaldehyde. *J. Am. Chem. Soc.*, **125**, 50–61.
- Ho,T.V., Guainazzi,A., Derkunt,S.B., Enoiu,M. and Schärer,O.D. (2011) Structure-dependent bypass of DNA interstrand crosslinks by translesion synthesis polymerases. *Nucleic Acids Res.*, **39**, 7455–7464.
- Minko,I.G., Harbut,M.B., Kozekov,I.D., Kozekova,A., Jakobs,P.M., Olson,S.B., Moses,R.E., Harris,T.M., Rizzo,C.J. and Lloyd,R.S. (2008) Role for DNA polymerase kappa in the processing of N2-N2-guanine interstrand cross-links. *J. Biol. Chem.*, **283**, 17075–17082.
- Räschle,M., Knipscheer,P., Enoiu,M., Angelov,T., Sun,J.C., Griffith,J.D., Ellenberger,T.E., Schärer,O.D. and Walter,J.C. (2009) Mechanism of replication-coupled DNA interstrand crosslink repair. *Cell*, **136**, 969–980.
- Smeaton,M.B., Hlavin,E.M., McGregor Mason,T., Noronha,A.M., Wilds,C.J. and Miller,P.S. (2008) Distortion-dependent unhooking of interstrand cross-links in mammalian cell extracts. *Biochemistry*, **47**, 9920–9930.
- Smeaton,M.B., Hlavin,E.M., Noronha,A.M., Murphy,S.P., Wilds,C.J. and Miller,P.S. (2009) Effect of cross-link structure on DNA interstrand cross-link repair synthesis. *Chem. Res. Toxicol.*, **22**, 1285–1297.
- Yamanaka,K., Minko,I.G., Takata,K., Kolbanovskiy,A., Kozekov,I.D., Wood,R.D., Rizzo,C.J. and Lloyd,R.S. (2010) Novel enzymatic function of DNA polymerase nu in translesion DNA synthesis past major groove DNA-perturbed and DNA-DNA cross-links. *Chem. Res. Toxicol.*, **23**, 689–695.
- Dong,Q., Barskt,D., Colvin,M.E., Melius,C.F., Ludeman,S.M., Moravek,J.F., Colvin,O.M., Bigner,D.D., Modrich,P. and Friedman,H.S. (1995) A structural basis for a phosphoramidate mustard-induced DNA interstrand cross-link at 5'-d(GAC). *Proc. Natl. Acad. Sci. U.S.A.*, **92**, 12170–12174.
- Millard,J.T., McGowan,E.E. and Bradley,S.Q. (2012) Diepoxybutane interstrand cross-links induce DNA bending. *Biochimie*, **94**, 574–577.
- Rink,S.M. and Hopkins,P.B. (1995) A mechlorethamine-induced DNA interstrand cross-link bends duplex DNA. *Biochemistry*, **34**, 1439–1445.
- Sawyer,G.A., Frederick,E.D. and Millard,J.T. (2004) Flanking sequences modulate diepoxide and mustard cross-linking efficiencies at the 5'-GNC site. *Chem. Res. Toxicol.*, **17**, 1057–1063.

37. Enoiu, M., Jiricny, J. and Schärer, O.D. (2012) Repair of cisplatin-induced DNA interstrand crosslinks by a replication-independent pathway involving transcription-coupled repair and translesion synthesis. *Nucleic Acids Res.*, **40**, 8953–8964.
38. Gooding, O.W., Beard, C.C., Jackson, D.Y., Wren, D.L. and Cooper, G.F. (1991) Enantioselective formation of functionalized 1,3-disubstituted allenes—synthesis of alpha-allenic omega-carbomethoxy alcohols of high optical purity. *J. Org. Chem.*, **56**, 1083–1088.
39. Jiang, B. and Ma, P. (1995) An improved synthesis of (+)-3,4-O-isopropylidene butyne. *Synth. Commun.*, **25**, 3641–3645.
40. Millard, J.T., Raucher, S. and Hopkins, P.B. (1990) Mechlorethamine cross-links deoxyguanosine residues at 5'-GNC sequences in duplex DNA fragments. *J. Am. Chem. Soc.*, **112**, 2459–2460.
41. Rink, S.M., Solomon, M.S., Taylor, M.J., Rajur, S.B., McLaughlin, L.W. and Hopkins, P.B. (1993) Covalent structure of a nitrogen mustard-induced DNA interstrand cross-link—an N7-to-N7 linkage of deoxyguanosine residues at the duplex sequence 5'-d(GNC). *J. Am. Chem. Soc.*, **115**, 2551–2557.
42. Humphrey, W., Dalke, A. and Schulten, K. (1996) VMD: visual molecular dynamics. *J. Mol. Graph.*, **14**, 33–38, 27–38.