

A new species of *Aegyptobia* and redescription of *Tenuipalpus szarvasensis* Bozai, 1970 (Acari, Tenuipalpidae)

Jenő Kotschán¹, Géza Ripka²

1 Plant Protection Institute, Centre for Agricultural Research, Hungarian Academy of Sciences, H-1525 Budapest, P.O. Box 102, Hungary **2** National Food Chain Safety Office, Directorate of Plant Protection, Soil Conservation and Agri-environment, Department of Pest Management Development and Coordination, H-1118 Budapest, Budaörsi út 141–145, Hungary

Corresponding author: Jenő Kotschán (kotschan.jeno@agrar.mta.hu)

Academic editor: V. Pesic | Received 21 June 2018 | Accepted 5 August 2018 | Published 19 September 2018

<http://zoobank.org/BE84A4DE-2A3D-4306-8166-097810EB84BF>

Citation: Kotschán J, Ripka G (2018) A new species of *Aegyptobia* and redescription of *Tenuipalpus szarvasensis* Bozai, 1970 (Acari, Tenuipalpidae). ZooKeys 785: 99–115. <https://doi.org/10.3897/zookeys.785.27684>

Abstract

A new tenuipalpid mite species, *Aegyptobia bozaii* **sp. n.**, is described from Central-Hungary on leaves of the endemic Hungarian statice *Limonium gmelinii* subsp. *hungaricum* (Plumbaginaceae) based on females, nymphs and larva. The previously described endemic flat mite, *Tenuipalpus szarvasensis* Bozai, 1970 is re-described. This species had been treated as a junior synonym of *Tenuipalpus cheladzeae* Gomelauri, 1960, but our new investigation shows that the two species are not the same.

Keywords

tenuipalpids, mites, flat mite, taxonomy, Hungary

Introduction

Tenuipalpid mites are a diverse group of plant-feeding mites found in most regions of the world. Several species are pests, especially within *Brevipalpus*, but pest species are also found in *Dolichotetranychus*, *Raoiella* and *Tenuipalpus*. The family has received considerable attention in some parts of the world, but the majority of the Central

European countries have been scarcely investigated. Hungary is no exception, with only 19 recorded species (Kontschán and Ripka 2017). However, numerous rare and endemic flat mites might remain undiscovered in natural ecosystems in Hungary; and disturbed ecosystems may harbor several unrecorded or possibly new species.

The aim of our paper is to describe a new tenuipalpid species from Hungary and to redescribe the endemic Hungarian flat mite *Tenuipalpus szarvasensis* Bozai, 1970, which we also remove from its synonymy with *Tenuipalpus cheladzeae* Gomelauri, 1960 by Mitrofanov and Strunkova (1979).

Material and methods

Specimens of the new species (*Aegyptobia bozaii* sp. n.) were collected in a pasture close the border of the village Farnos (Central-Hungary) from the leaves of an endemic Hungarian plant (*Limonium gmelinii* subsp. *hungaricum*). The specimens were placed into lactic acid for a week and then slide-mounted in Keifer's F-medium (in 2014) and Hoyer medium (in 2017). The holotype and some paratypes of the new species are stored in the Hungarian Natural History Museum and other paratypes in the Arachnida collection of the Natural History Museum of Geneva (Switzerland).

The type specimens of *Tenuipalpus szarvasensis* Bozai, 1970 were loaned from the Hungarian Natural History Museum.

All specimens were investigated using a Leica 1000 scientific microscope; the illustrations were made with the aid of a drawing tube on this microscope. Pictures were made with a VHX-5000 with 20–200× objective (Keyence Co., Osaka, Japan) digital microscope. All measurements and scales are given in micrometers.

Result

Aegyptobia bozaii sp. n.

<http://zoobank.org/CF433BD9-1455-4EE2-84F3-DFF3ABCC858C>

Figures 1–26

Material examined. Holotype: female, Hungary, Pest county, Farnos, 47°22'30"N, 19°52'08"E, 10 m a.s.l, from the leaves of the Hungarian statice, *Limonium gmelinii* subsp. *hungaricum*, 2 August 2014, Ripka, G. coll. Paratypes: one female, three deutonymphs, three protonymphs and one larva, locality and date same as for holotype. Other paratypes: four females, Hungary, Farnos, 47°22'30"N, 19°52'08"E, 10 m a.s.l, from the leaves of *Limonium gmelinii* subsp. *hungaricum*, 2 August 2017, Kontschán, J. and Ripka, G. coll.

Diagnosis (based on female). Tarsal claws uncinata. Anterior margin of prodorsal shield with paired projections, prodorsum weakly sculptured, with few irregular lines. Opisthosoma with polygonal reticulation. Propodosomal and opisthosomal setae




Figure 1. Dorsal view of *Aegyptobia bozaii* sp. n., holotype, female.

smooth and simple; seta *f2* present. Length of dorsal setae 6–14. Dorsal opisthosomal pores close to *e1*. Rostrum extending to middle of tibia I. Genital flap smooth. Intercoxal area between *3a* and *4a* smooth.

Description (females; $n = 6$). Idiosoma reddish-brown (Figure 27), oval in shape, body measured from *v2* to *h1* 240–245; from tip of rostrum 260–266; width 143–147 near setae *sc2*; distance between setae *sc2* 120–125; length of legs I–IV (without coxa), leg I 105–110, leg II 80–87, leg III 72–78, leg IV 80–88.

Dorsum (Figure 1): Anterior margin of prodorsal shield with paired projections, depth of notch 7–8. Propodosoma finely lineate. Opisthosoma with polygonal reticulations; polygons longitudinally elongate medially, transversally elongate anterolaterally. Propodosomal and opisthosomal setae simple and smooth. Opisthosomal pores present close to *e1*. Prodorsal setae *v2* shorter than half distance between their bases.

Length of dorsal setae: *v2* 11–12, *sc1* 12–14, *sc2* 11–13, *c1* 8–9, *c2* 8–9, *c3* 9–10, *d1* 9–10, *d2* 8–9, *d3* 7–8, *e1* 8–9, *e2* 7–8, *e3* 8–9, *f2* 7–8, *f3* 6–8, *h1* 6–7, *h2* 7–8.

Venter (Figure 2): Surface of ventral idiosoma smooth, except lateral to *ag*, genital and anal plates where longitudinal striations visible. Genital and anal plates smooth. Length of ventral setae, *1a* 53–57, *3a* 11–12, *4a* 8–9, *1b* 7–8, *2b* 8–9, *3b* 8–9, *4b* 9–10, *1c* 11–12, *2c* 10–13, *ag* 8–9, *g1* 9–10, *g2* 7–8. Pseudanal setae, all 6–8. All ventral setae simple and smooth.

Gnathosoma (Figure 3): Rostrum extending to middle of tibia I; palp setal counts: tarsus with one solenidion and two eupathidia, tibia with two setae, genu without seta and femur with one simple dorsal seta. All setae smooth. Subcapitulum with setae *m* (4–5).

Legs (Figures 4–8): Setal formula for leg I–IV (coxae to tarsi): 3-1-4-3-3-9, 2-1-4-3-3-9, 2-2-2-1-3-5, 2-1-1-0-3-5. A supplementary lateral (*l*) seta present on femora I. Solenidia on tarsi I and II 8–10 long, broad, leaf-like. Tarsal claws uncinatate and empodium pad-like.

Deutonymph (*n* = 3; Figures 9–15). Idiosoma oval in shape, body measured from *v2* to *h1* 190–200; width 130–140 near setae *sc2*.

Dorsum (Figure 9) covered with a few striae, all setae short, simple and needle-like. Length of all setae 5–7.

Venter covered with very few striae with one pair of setae *1a*, *1b*, *2b*, *3a*, *3b*, *3c*, *4a* and *4b*, one pair of aggenital, one pair of genital and three pairs of anal setae, all simple and smooth *1a* 15–16, other setae on venter 5–7 (Figure 10). Palp setal counts: tarsus with one solenidion and two eupathidia, tibia with two setae, genu without seta and femur with one simple dorsal seta (Figure 11). Legs as Figures 12–15 and Table 1.

Protonymph (*n* = 3; Figures 16–21). Idiosoma oval in shape, body measured from *v2* to *h1* 149–155; width 94–100 near setae *sc2*.

Dorsum (Figure 16). Surface without striae, all setae short, simple and needle-like. All setae 4–6 in length.

Venter covered with very few striae with one pair of setae *1a*, *1b*, *2b*, *3a* and *3b*, one pair of aggenital and three pairs of anal setae, all simple and smooth. *1a* 13–15, other setae on venter 5–6 (Figure 17). Legs as Figures 18–21 and Table 1.


Larva (*n* = 1; Figures 22–26). Idiosoma oval in shape, body measured from *v2* to *h1* 109; width 82 near setae *sc2*.

Dorsum (Figure 22). Covered with a few striae, all setae short, simple and needle-like. Length of setae: *v2* 30–31; *sc1* 30–37; *sc2* 28–36; *c1* 29–38; *c2* 26–39; *c3* 31–37; *d1* 5; *d3* 37–43; *e1* 3; *e3* 40–54; *f2* 3–4; *f3* 64–66; *h1* 2–3; *h2* 2–5.

Venter covered with few striae with one pair of setae *1a* and three pairs of anal setae, all simple and smooth (Figure 23). Legs as Figures 24–26 and Table 1.

Etymology. We dedicate the new species to Dr. József Bozai, former Hungarian tenuipalpid specialist.

Notes on the host. The host plant, Hungarian statice (*Limonium gmelinii* subsp. *hungaricum*) (Plumbaginaceae), is an endemic subspecies occurring on salt meadows in Central-Hungary. The mites appeared to prefer leaves of the host plant that were lying close to the surface of the soil. The alkali steppe where the host plant was found



Figures 2–8. *Aegyptobia bozaii* sp. n., holotype, female **2** Ventral view of idiosoma **3** Ventral view of gnathosoma **4** Ventral view of leg I **5** Ventrolateral view of tarsus I **6** Ventral view of leg II **7** Ventral view of leg III **8** Ventral view of leg IV.





Figure 9. Dorsal view of *Aegyptobia bozaii* sp. n., paratype, deutonymph.



is hot and dry in summer, typical habitat for tenuipalpid species, which prefer warm and dry conditions. Up until now, only one species has been reported from *Limonium* plants: *Capedulia maritima* Gerson & Smith Meyer, 1980 was found on the roots of *Limonium meyeri* in Israel (Ueckermann et al. 2018).

Remarks. The new species has uncinuate claws and therefore belongs to the *Aegyptobia tragardhi* species group (Khanjani et al. 2008). It is very similar to *A. iranensis* Khanjani et al., 2008 and *A. wainsteini* Bagdasarian, 1962 based on the claw-like empodium, the slender prodorsal setae, the deeply emarginated notch and the medially smooth prodorsum. The most important differences among three species are summarized in Table 2.


Only one species, *Aegyptobia wainsteini* Bagdasarian, 1962, was previously reported from Hungary from a *Biota orientalis* tree (Cupressaceae) close to the town Kecskemét (Bozai 1969). Other new occurrences have not been given since this first report. The two *Aegyptobia* species reported from Hungary differ in the shape of the dorsal setae, which are short and smooth in *A. bozaii* sp. n. and longer and finely pilose on *Aegyptobia wainsteini*.




10



11




12



13



14



15

Figures 10–15. *Aegyptobia bozaii* sp. n., paratype, deutonymph **10** Ventral view of idiosoma **11** Ventral view of palp **12** Ventral view of leg I **13** Ventral view of leg II **14** Ventral view of leg III **15** Ventral view of leg IV.

Table I. Development of leg setae (after Seeman and Beard 2011).

	Cx I			Cx II		Cx III		Cx IV		Tr I	Tr II	Tr III		Tr IV	Fe I*			Fe II			Fe III			
	1a	1b	1c	2b	2c	3a	3b	4a	4b	v'	v''	l'	v'	v''	d	v'	bv''	d	v'	bv''	d	ev'		
Larva	+	+				+									+	+	+	+	+	+	+	+	+	
PN	+	+	+	+		+	+					+			+	+	+	+	+	+	+	+	+	
DN	+	+	+	+	+	+	+	+	+	+	+	+	+		+	+	+	+	+	+	+	+	+	
Adult	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	Fe IV	Ge I			Ge II			Ge III	Ti I**				Ti II				Ti III			Ti IV				
	ev'	l'	d	l''	l'	d	l''	l'	d	l'	v'	v''	d	l'	v'	v''	d	v'	v''	d	v'	v''		
Larva		+			+				+	+	+	+								+	+	+	+	+
PN	+	+			+			+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
DN	+	+	+	+	+	+	+	+	+	?	+	+	+	?	+	+	+	+	+	+	+	+	+	+
Adult	+	+	+	+	+	+	+	+	+	?	+	+	?	+	+	+	+	+	+	+	+	+	+	+
	Ta I-II					Ta III					Ta IV													
	u'	u''	p'	p''	tc'	tc''	ft'	ft''	ω	u'	u''	p'	p''	tc'	tc''	ft'	u'	u''	tc'	tc''	ft'			
Larva	+	+	+	+			+	+	+	+	+					+								
PN	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+				+	
DN	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Adult	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	

* The new species has a supplementary lateral (*l'*) seta on femora I.

** The setae *l'* on tibiae I and II in adult female and deutonymphs are not visible.





Figure 16. Dorsal view of *Aegyptobia bozaii* sp. n. paratype, protonymph.

Table 2. Distinguishing characteristics among *Aegyptobia bozaii*, *A. iranensis* and *A. wainsteini*.

Character	<i>Aegyptobia bozaii</i>	<i>Aegyptobia iranensis</i>	<i>Aegyptobia wainsteini</i>
Surface between <i>c1</i> and <i>d1</i>	with large reticulations	with large reticulations	smooth
Distance between setae <i>v2</i>	three times longer than length of <i>v2</i>	two times longer than length of <i>v2</i>	same as length of <i>v2</i>
Setae <i>3a</i>	1/2 the distance <i>3a-3a</i>	two times the distance <i>3a-3a</i>	1/2 the distance <i>3a-3a</i>



Figures 17–21. *Aegyptobia bozaii* sp. n., paratype, protonymph **17** Ventral view of idiosoma **18** Ventral view of leg I **19** Ventral view of leg II **20** Ventral view of leg III **21** Ventral view of leg IV.




Figure 22. Dorsal view of *Aegyptobia bozai* sp. n., paratype, larva.

Tenuipalpus szarvasensis Bozai, 1970


Tenuipalpus szarvasensis Bozai, 1970: 367.

Tenuipalpus cheladzeae Gomelauri, 1960 as senior synonym of *T. szarvasensis* by Mitrofanov and Strunkova 1979: 51.

Tenuipalpus cheladzeae: Kontschán and Ripka 2017.

Material examined. Holotype: female, HNHM Astig-242, Szarvas, 8 October 1968, from *Picea excelsa* Lk. No. 1250, Bozai, J. coll.

Diagnosis (based on female). Anterior margin of prodorsal shield with forked projection; prodorsum smooth medially, with some striae laterally; anterolateral projections carrying setae *sc2* weakly formed. Opisthosoma smooth anteriorly, with posteromedial reticulation and posterolateral longitudinal striation. Propodosomal setae as follows: *v2* short and smooth, *sc1* broad and obovate, *sc2* long and phylliform. Opisthosomal setae: *c1*, *c3* and *d1* broad, long, oblanceolate, *d3* short and oblanceolate, *e1* short and smooth, *h1*, *f1*, *f2* and *e3* long, oblanceolate, *h2* very long and smooth. Rostrum extending to middle of tibia I. Genital flap smooth. Intercoxal area



Figures 23–26. *Aegyptobia bozaii* sp. n., paratype, larva **23** Ventral view of idiosoma **24** Ventral view of leg I **25** Ventral view of leg II **26** Ventral view of leg III.

between *3a* and *4a* smooth, *1a* and *4a* very long, *1b*, *2b*, *2c*, *3a*, *3b*, *4b*, *ag*, *g1*, *g2* short. Legs with large, broad and phylliform, smooth and pilose setae.

Description (female holotype). Colorization of idiosoma not observable in the holotype. Idiosoma (Figure 27) pentagonal in shape, body measured from *v2* to *h1* 305; from tip of rostrum 350; width between setae *sc2* 190.



Figure 27. Photos of the female of *Aegyptobia bozaii* sp. n., paratype.




Figure 28. Habitat and details of the host plant. **a** Habitat **b** Host plant **c–d** Leaves of host plant.

Dorsum (Figure 29): Anterior margin of prodorsal shield with paired projections, depth of notch 16. Prodorsum smooth medially, with some striae laterally; anterolateral projections carrying setae *sc2* weakly formed. Opisthosoma smooth anteriorly, with posteromedial reticulation and posterolateral longitudinal striation. Propodosomal setae as follows: *v2* short and smooth, *sc1* broad and obovate, *sc2* long and phylliform. Opisthosomal setae: *c1*, *c3* and *d1* broad, long, oblanceolate, *d3* short and oblanceolate, *e1* short and smooth, *h1*, *f1*, *f2* and *e3* long, oblanceolate, *h2* very long and smooth. Rostrum extending to middle of tibia I. Opisthosomal pores between *d1* and *e1*. Length of dorsal setae: *v2* 9–10, *sc1* 30–32, *sc2* 65, *c1* 45–46, *c3* 32–34, *d1* 38–40, *d3* 12–14, *e1* 12, *e2* 45–47, *f2* and *f3* 45–46, *h1* 36–37, *h2* 105–110.

Venter (Figure 30): Very few striations observable in the holotype, only a few longitudinal striations visible posterior to *g1–g2*. Genital and anal plates smooth. Length of ventral setae, *1a* 120–122, *3a* 26–27, *4a* 130–133, *1b* 16, *2b* 16–17, *3b* 18–19, *4b*




Figure 29. Dorsal view of *Tenuipalpus szarvasensis* Bozai, 1970, holotype, female.

13, 1c and 2c 24–25, ag 12, g1–g2 14–16. Pseudanal setae, all 12–13. All ventral setae simple and smooth.


Gnathosoma: Rostrum extending to middle of tibia I; palp setal counts as in Figure 31.

Legs: Setal formula for leg I–IV (coxae to tarsi): 3-1-4-2-5-9, 2-1-4-2-4-9, 2-2-2-1-3-5, 2-1-1-0-3-5. Shape of the setae on legs illustrated on Figures 32–35.

Notes. Bozai's (1970) noted in his detailed description of *Tenuipalpus szarvasensis*, that the species is very similar to *T. cheladzeae*, but he mentioned some easy to observe differences (like shape and length of setae *c1*, *d1*) between these two species. Despite these known differences, Mitrofanov and Strunkova (1979) synonymized the name under *Tenuipalpus cheladzeae*. Mitrofanov and Strunkova (1979) did not study the type specimens of *T. szarvasensis*, therefore their opinion was questionable.

This year, we studied the types of Bozai's *T. szarvasensis* in order to confirm Bozai's hypothesis that *T. szarvasensis* differs from Gomelauri's *T. cheladzeae*. The differences are presented in Table 3 and are illustrated in Figures 36, 37.

Tenuipalpus is the largest genus of flat mites, but very few are known from Pinaceae. Apart from the above-mentioned two species, the only other species is *T. hondurensis* Evans, in Evans et al. (1993) (Mesa et al. 2009), which is considerably different from both the above species. *Tenuipalpus cupressoides* Smith, Meyer




Figures 30–35. *Tenuipalpus szarvasensis* Bozai, 1970, holotype, female **30** Ventral view of idiosoma **31** Dorsal view of palp **32** Ventral view of leg I **33** Ventral view of leg II **34** Ventral view of leg III **35** Ventral view of leg IV.

Table 3. Distinguishing characters between *Tenuipalpus cheladzeae* and *T. szarvasensis*.

Character	<i>T. cheladzeae</i>	<i>T. szarvasensis</i>
Shape of <i>v2</i>	apically rounded	apically pointed
Shape of <i>sc1</i>	short (12–14) and bulbiform	long (30–32) and phylliform
Shape of <i>c1</i> and <i>d1</i>	short (<i>c1</i> 10–11, <i>d1</i> 9–11) and bulbiform	long (<i>c1</i> 45–46, <i>d1</i> 38–40) and phylliform

& Gerson, 1980 is very similar to *T. szarvasensis*, but the shape of *v2* and *e2* and the length of *sc1*, *c1* and *d1* are different. In addition, the host plant, *Cupressus sempervirens* of *T. cupressoides* belongs to the family Cupressaceae and not to Pinaceae (Ueckermann et al. 2018).



Figures 36–37. Illustrated differences between *T. szarvasensis* (36) and *T. cheladzeae* (37) Drawing of *T. cheladzeae* modified after Mitrofanov and Strunkova (1979).

Acknowledgements

This study was supported by the NKFIH (OTKA) 108663 and by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. We are very grateful to Dr. Owen D. Seeman for the correction of the previous version of the manuscript and his linguistic corrections.

References

- Bagdasarian AT (1962) Contribution to the fauna of false spider mites from Armenia. *Izvestiia Akademii Nauk SSR, Moscow* 15: 49–58.
- Bozai J (1969) Új lapostestű atkák Magyarországon (Tenuipalpidae, Acari). (Some tenuipalps (Acari) new for the fauna of Hungary.) *Folia Entomologica Hungarica* 22: 117–130.
- Bozai J (1970) *Tenuipalpus szarvasensis* sp. n., a new mite species from Hungary (Acari: Tenuipalpidae). *Acta Zoologica Academiae Scientiarum Hungaricae* 16(3–4): 367–369.
- Evans G, Cromroy HL, Ochoa R (1993) The Tenuipalpidae of Honduras (Tenuipalpidae: Acari). *Florida Entomologist* 76(1): 126–155. <https://doi.org/10.2307/3496021>
- Gerson U, Smith Meyer MKP (1980) A new species of *Capedulia* (Acari: Tenuipalpidae) from a marine habitat in Israel. *Israel Journal of Entomology* 14: 9–12.

- Gomelaury LA (1960) New species in the genus *Tenuipalpus* from the Georgia, Gruzia, SSR (Acarina: Tenuipalpidae). Reports of the Academy of Science of the Georgian SSR. Soobschcheniya Akademii Nauk Gruzinskoi 24(1): 77–79.
- Khanjani M, Gotoh T, Varandi HB (2008) A new species of the genus *Aegyptobia* Sayed (Acari: Prostigmata, Tenuipalpidae) from Iran. Systematic & Applied Acarology 13(1): 65–69. <https://doi.org/10.11158/saa.13.1.7>
- Kontschán J, Ripka G (2017) Checklist of the Hungarian spider mites and flat mites (Acari: Tetranychidae and Tenuipalpidae). Systematic & Applied Acarology 22(8): 1199–1225. <https://doi.org/10.11158/saa.22.8.6>
- Mesa NC, Ochoa R, Welbourn WC, Evans G, Moraes GJ de (2009) A catalog of the Tenuipalpidae (Acari) of the World with a key to genera. Zootaxa 2098: 1–185.
- Mitrofanov VI, Strunkova ZI (1979) Opredelitelj kleshchej-ploszkotelok. Izdateljsztvo „Donis” Dushanbe, 148 pp.
- Seeman OD, Beard JJ (2011) A new species of *Aegyptobia* (Acari: Tenuipalpidae) from Myrta-ceae in Australia. Systematic & Applied Acarology 16: 73–89. <https://doi.org/10.11158/saa.16.1.10>
- Ueckermann EA, Palevsky E, Gerson U, Recht E, Theron PD (2018) The Tenuipalpidae (Acari: Trombidiformes) of Israel. Acarologia 58(2): 483–525.