


RESEARCH ARTICLE

Indicators of abdominal size relative to height associated with sex, age, socioeconomic position and ancestry among US adults

Henry S. Kahn*, Kai McKeever Bullard

Division of Diabetes Translation, Centers for Disease Control and Prevention, Atlanta, Georgia, United States of America

* hkahn@cdc.gov



Abstract

Background/Objectives

The supine sagittal abdominal diameter (SAD) and standing waist circumference (WC) describe abdominal size. The SAD/height ratio (SADHtR) or WC/height ratio (WHtR) may better identify cardiometabolic disorders than BMI (weight/height²), but population-based distributions of SADHtR and WHtR are not widely available. Abdominal adiposity may differ by sociodemographic characteristics.

Subjects/Methods

Anthropometry, including SAD by sliding-beam caliper, was performed on 9894 non-pregnant adults ≥ 20 years in the US National Health and Nutrition Examination Surveys of 2011–2014. Applying survey design factors and sampling weights, we estimated nationally representative SADHtR and WHtR distributions by sex, age, educational attainment, and four ancestral groups.

Results

The median (10th percentile, 90th percentile) for men's SADHtR was 0.130 (0.103, 0.165) and WHtR 0.569 (0.467, 0.690). For women, median SADHtR was 0.132 (0.102, 0.175) and WHtR 0.586 (0.473, 0.738). Medians for SADHtR and WHtR increased steadily through age 79. The median BMI, however, reached maximum values at ages 40–49 (men) or 60–69 (women) and then declined. Low educational attainment, adjusted for age and ancestry, was associated with elevated SADHtR more strongly than elevated BMI. While non-Hispanic Asians had substantially lower BMI compared to all other ancestral groups (adjusted for sex, age and education), their relative reductions in SADHtR and WHtR, were less marked.

Conclusions

These cross-sectional data are consistent with monotonically increasing abdominal adipose tissue through the years of adulthood but decreasing mass in non-abdominal regions

OPEN ACCESS

Citation: Kahn HS, Bullard KMck (2017) Indicators of abdominal size relative to height associated with sex, age, socioeconomic position and ancestry among US adults. PLoS ONE 12(3): e0172245. doi:10.1371/journal.pone.0172245

Editor: Clemens Fürnsinn, Medical University of Vienna, AUSTRIA

Received: October 28, 2016

Accepted: February 1, 2017

Published: March 1, 2017

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: Public-use data are available from the National Center for Health Statistics at http://www.cdc.gov/nchs/nhanes/search/nhanes11_12.aspx and http://www.cdc.gov/nchs/nhanes/search/nhanes13_14.aspx.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

beyond middle age. They suggest also that visceral adipose tissue, estimated by SADHtR, expands differentially in association with low socioeconomic position. Insofar as Asians have lower BMIs than other populations, employing abdominal indicators may attenuate the adiposity differences reported between ancestral groups. Documenting the distribution and sociodemographic features of SADHtR and WHtR supports the clinical and epidemiologic adoption of these adiposity indicators.

Introduction

Clinical assessments of adiposity commonly depend on the body mass index (BMI, weight/height²) as a convenient indicator of fatness relative to height [1]. While BMI is more useful than body weight alone, its dependence on non-specified “mass” may obscure differing contributions made by fat and lean tissues [2] or fail to recognize the health consequences attributable to variations in body fat distribution [3, 4]. Despite these limitations for clinical screening and physiologic research, BMI has served those who monitor population-based, categorical “overweight” and “obesity” in adults across time and different cultures [5, 6]. For over three decades, population reference data on BMI in the United States have been published periodically from the National Health and Nutrition Examination Survey (NHANES). Most recently, NHANES has reported the detailed BMI distribution for adult women and men during survey years 2011–2014 [7].

Reports based on the BMI describe an increasing prevalence of obesity worldwide [8]. Serial surveys obtained across at least two decades have documented disproportionate increases in the standing waist circumference (WC) for a given level of BMI [9–11]. Since WC is correlated with abdominal adipose tissue, total body fat, and with various markers of cardiometabolic risk [12], any increases in WC that are disproportionate (i.e., beyond increases in total body mass) may carry adverse health consequences for the population. In order to simplify the concept of abdominal adiposity, WC was transformed into the WC/height ratio (WHtR) more than two decades ago [13, 14] so as to acknowledge the variation in adult stature [15].

The supine sagittal abdominal diameter (SAD, “abdominal height”), also correlated with abdominal adipose tissue, has been proposed as an alternative body dimension to the WC. The SAD, which can be measured with a sliding-beam caliper, may estimate the expansion of the visceral (intra-abdominal) adipose compartment more specifically than the WC [16–20]. SAD has been transformed similarly into the SAD/height ratio (SADHtR) [21, 22].

Early studies reported that inter- and intra-observer variations in measurement of WC or SAD were comparable and small in comparison to the variation between patients [23, 24]. A more recent report suggested that SAD (measured supine with flexed hips) had higher reliability than supine WC especially among those with BMI ≥ 26 kg/m² [25]. However, each of these 3 studies was limited by its small study sample.

The clinical use of SADHtR [26, 27] or WHtR [26–30] can improve recognition of cardiometabolic risk when compared to the use of BMI. Despite growing interest in low-cost assessments of abdominal adiposity, current population distributions of SADHtR and WHtR are not widely available. This paper summarizes community-based estimates (reference data) of how these two adiposity indicators are currently distributed among adults (ages 20+ years) in the United States by sex and decades of age. We also point out variations in abdominal adiposity associated with low educational attainment, low family income, and four major ancestral groups. Our report concludes with commentary that acknowledges the few, community-based

distributions of adult SADHtR and WHtR that were published in earlier years and from other countries.

Participants and methods

Anthropometric data were obtained from the NHANES, an ongoing, nationally representative, cross-sectional survey of the resident civilian, non-institutionalized, US population [31]. Participants chosen for NHANES undergo home interviews followed by standardized anthropometric and laboratory assessments in mobile examination centers. The NHANES protocol was approved by the Research Ethics Review Board of the National Center for Health Statistics; participants provided informed consent.

Our report covers non-pregnant adults (age ≥ 20 years) examined in 2011–2014. Weight, height and WC (standing position, by tape measure just above the uppermost lateral border of the ilium) were obtained by established methods [32]. WHtR was calculated as WC [cm]/height [cm]. BMI was calculated as weight [kg]/height [m]². During 2011–2014 NHANES did not measure circumferences of the hip or midhigh.

The SAD measurement used a portable, sliding-beam caliper (Holtain, Ltd, Wales, UK) [32]. Supine participants rested on a lightly padded exam table with their hips in flexed position as the examiner marked the level of their iliac crests. The lower arm of the caliper was then positioned under the small of the back, and the upper arm was raised above the belly in alignment with their iliac-crest level. The examiner asked the participant to inhale gently, slowly let the air out, and then relax. The examiner then lowered the caliper's upper arm, letting it lightly touch the abdomen but without compressing it. The SAD value was read directly from a centimeter scale on the caliper shaft, recorded in duplicate to the nearest 0.1 cm. For 94.2% of adults we defined SAD as the mean of two initial measurements; for 5.8% (with initial SAD discrepancy >0.5 cm) we used the mean of up to 4 measurements [33]. SADHtR was calculated as SAD [cm]/height [cm].

Of 10,785 non-pregnant adults attending the examination centers, 10,636 provided BMI values, 9,907 provided SADHtR, and 10,116 provided WHtR. We restricted our primary descriptive sample to include only the 9,894 adults who had values for both SADHtR and WHtR (91.7% of examination attendees). These examinees represented approximately 208.5 million US adults. Multivariable regression models were restricted to 9,882 of these who had values also for BMI, as well as SADHtR and WHtR (91.6% of attendees).

Sociodemographic covariates

As derived from self-reported race and Hispanic origin (ethnicity) we identified five, mutually exclusive, ancestral groups as non-Hispanic white, non-Hispanic black, Hispanic, non-Hispanic Asian, and "others" [34]. For comparison with persons who attended some college or higher education, we categorized low educational attainment as either less than high school completion or high school graduation (may include passing the General Educational Development tests). For comparison with persons having a family income of at least 200% of poverty guidelines, we categorized low income as either $<100\%$ of poverty guidelines or 100% to $<200\%$ of poverty guidelines; an additional category of "unknown" income was included in these models [35].

Statistical methods

NHANES selected participants through a complex, multistage-probability design requiring a sampling weight for each participant. We used SAS (release 9.3; SAS Institute Inc.) and SUDAAN (PROC DESCRIPT and PROC REGRESS, release 11.1; RTI International) to

account for the complex design and sampling weights so that characteristics of the represented population could be correctly described.

To describe the associations of socioeconomic variables with adiposity we created sex-specific, multivariable, linear regression models having outcomes of SADHtR, WHtR or BMI. All multivariate models were age-adjusted in 7 categories (decades) of which the highest was 80+ years. As generated by these linear regression models, a beta coefficient (sometimes called *regression coefficient*) estimates the amount by which the outcome differs for persons in a designated socioeconomic category (e.g., “less than high-school completion” or “high school graduation”) compared to persons in the reference category (e.g., “persons who attended some college or higher education”). In these separate models we employed the adjusted Wald F statistic to estimate the confidence assigned to the contrasts (summary effect) within the 3-level variables for education or for family income regarding their influence on the observed value for the adiposity indicator. The adjusted Wald F is based on the Wald chi-squared with adjusted denominator degrees of freedom. A higher adjusted Wald F value means a higher likelihood of the variable showing statistical significance.

To describe how each of four major ancestral groups differed from the remaining US population, we built a series of similar linear regression models having outcomes of SADHtR, WHtR or BMI. As generated by these linear regression models, a beta coefficient estimates the amount by which the outcome differs for persons in a designated ancestral group (e.g., “Non-Hispanic whites”) compared to persons in the reference category (e.g., “sum of all the ancestries excluding Non-Hispanic whites”). As used here, the adjusted Wald F statistic estimates the confidence assigned to each 2-level variable of ancestral contrasts.

Results

For SADHtR, among US men (sample N = 4,949) the population mean was 0.1326 (SE 0.0007) and median was 0.130 (10th percentile = 0.103; 90th percentile = 0.165). Among women (sample N = 4,945) the SADHtR population mean was 0.1357 (SE 0.0007) and median was 0.132 (10th percentile 0.102; 90th percentile = 0.175). Greater detail for SADHtR percentiles 5 through 95 is available from Tables A and B in [S1 File](#).

For WHtR, among US men the population mean was 0.5770 (SE 0.0023) and median was 0.569 (10th percentile = 0.467; 90th percentile = 0.690). Among women the WHtR population mean was 0.5972 (SE 0.0024) and median was 0.586 (10th percentile 0.473; 90th percentile = 0.738). Greater detail for WHtR percentiles is available from Tables C and D in [S1 File](#).

Age and sex

[Fig 1](#) demonstrates that for both abdominal adiposity indicators the women had a wider distribution of values than the men across the complete adult age range. The sex difference (men < women) in median values was less prominent for SADHtR than for WHtR. When median values of SADHtR and WHtR in each age group were normalized with reference to the sex-specific median of men or women at all ages (20+ y), these two indicators demonstrated a roughly parallel increase with age from 20–29 through 70–79 y ([Fig 2](#)). The relative increment, however, across these six age decades was slightly steeper for SADHtR than WHtR. Age-related variation in BMI showed a different pattern. After rises in early adulthood (similar to relative rises in abdominal indicators), the men’s BMI was lower after age 50, then more so after about 79 y. For women’s BMI there was a marked increase preceding age 30–39 (steeper than the relative rises in abdominal indicators), a more gradual increase up to age 60–69, followed thereafter by steep declines ([Fig 2](#)).




Fig 1. Distributions of [A] SAD/height ratio (SADHtR) and [B] waist circumference/height ratio (WHtR) by age group and sex (men on left, women on right). Boxes indicate population percentiles 25, 50, 75; whiskers identify p10 and p90.

doi:10.1371/journal.pone.0172245.g001




Fig 2. Age-specific medians plotted by decade for SADHtR, WHtR, and BMI. Median values were normalized with reference to the sex-specific median [1.00] estimated for the full population of [A] men or [B] women at ages 20+ years.

doi:10.1371/journal.pone.0172245.g002

Socioeconomic position

Compared to those who attained education beyond high school, after adjustments for age and ancestry, men with less than high school completion had elevated SADHtR, but no significant difference in WHtR or BMI (Fig 3). Men who attained only high school graduation had greater SADHtR and WHtR, but not greater BMI. The summary effect on adiposity of men’s low educational attainment was greater when estimated by SADHtR (adjusted Wald F = 6.2, p = 0.006) than by the alternative indicators. For women, low education was associated with elevation of all three adiposity indicators, but the summary effect was greatest when estimated by SADHtR (adjusted Wald F = 24.9, p<0.0001) or by WHtR (adjusted Wald F = 24.7, p<0.0001).

In a set of similar statistical models we compared adiposity among persons with low family income (<100% of poverty guidelines or 100% to <200% of poverty guidelines) to those with family income ≥200% of poverty guidelines. These models included an additional category of “unknown” family income (7% of participants). Among men in the lowest income group, BMI was significantly lower than the reference income category, but for this most impoverished group there were no significant differences for SADHtR or WHtR (S1 Fig). Men at 100% to <200% of poverty guidelines had elevated SADHtR and WHtR, but no difference in their BMI. Among women, both categories of low family income were associated with elevations of all adiposity indicators, with the greatest summary effects seen for SADHtR (adjusted Wald F = 27.6, p<0.0001) and WHtR (adjusted Wald F = 26.8, p<0.0001).




Fig 3. Sex-specific beta coefficients (95% confidence range) associated with low educational attainment (less than high-school completion or high school graduation) compared with persons who attended some college or higher education. Separate models, each adjusted for age and ancestral groups, were prepared for [top] SADHtR, [middle] WHtR, and [bottom] BMI. F indicates the adjusted Wald F statistic, a summary effect describing confidence that contrasts in educational attainment (3-levels) have influenced the observed value for the adiposity indicator.

doi:10.1371/journal.pone.0172245.g003

Ancestral groups

Within our represented US adult population the identified ancestral proportions were non-Hispanic white (66.4%), non-Hispanic black (11.1%), Hispanic (14.6%), non-Hispanic Asian (5.2%), and others (2.7%). Across three age strata, we have displayed the distributions of SADHtR, WHtR, and BMI found in the four largest ancestral groups for men (Fig 4) and women (Fig 5).

Non-Hispanic white men, compared to men remaining in the total of all other groups, had no significant differences in mean SADHtR, WHtR, or BMI (Table 1). For non-Hispanic white women, however, the means of all three of the adiposity indicators were modestly lower than those found for the total of all other women; this relative reduction was greatest for SADHtR (adjusted Wald $F = 31.5$; Table 1).

For non-Hispanic black men (*vs* all others) we found a reduced mean for WHtR, but no difference for SADHtR or BMI. For non-Hispanic black women, the means of all three indicators were elevated compare to those for all other women, with the greatest increment observed for SADHtR (adjusted Wald $F = 212.2$; Table 1). As estimated by their interquartile ranges or by their ranges between percentiles 10 and 90, we noted that the variation in BMI was wider for non-Hispanic black men (Fig 4) and women (Fig 5) than the variation seen within each of the other three major ancestral groups. The ranges in WHtR and SADHtR among non-Hispanic blacks (*vs* all others) also showed wider variation, but less markedly than the variations in BMI.

Hispanic men and women (*vs* all others) had elevated mean values of all adiposity indicators, with a more pronounced increment for the abdominal indicators than for the BMI (Table 1).

Non-Hispanic Asian men and women (*vs* all others) had lower mean values of all adiposity indicators, with BMI showing the most prominent decrement. In models that included both sexes (with sex adjustment), the non-Hispanic Asians had lower mean BMI compared to the total of all other ancestries (adjusted Wald $F = 520.0$), along with less marked decrements in SADHtR (adjusted Wald $F = 233.8$) and WHtR (adjusted Wald $F = 357.4$).

Discussion

Anthropometric reference data for adults are empirical, normative values associated with a given place and time. Regarding the SADHtR, a population distribution was first reported nationally from a survey in 2000–2001 of older Finnish adults (ages 30+ years) who were almost entirely of white, northern European ancestry [36]. We are unaware of any other population-based descriptions of SADHtR. From both Finland and the US these cross-sectional data, stratified by sex, demonstrate the same pattern of continuous increases with ageing through at least 69 y. In other adult samples of diverse ancestries, abdominal imaging studies have shown likewise that visceral adipose tissue increases with age more than the subcutaneous adipose depot [37, 38]. Taken together, these observations describe a generalized, monotonic increase of abdominal adipose tissue, especially in the visceral depot, through nearly all the years of adulthood. The relative declines in BMI beyond middle age (Fig 3), however, indicate a likelihood among the elderly of decreasing mass in non-abdominal regions.

Compared to the earlier Finnish report of SADHtR, the more recent US data reveal larger SADHtR values obtained for the same age strata. Within each sex, the US SADHtR values approximated the Finnish SADHtR values found for the next oldest decade of age. These contrasts could be explained by the rising adiposity trend worldwide during recent years or by inherent cultural and environmental differences between Finland and the US.




Fig 4. Men's distributions of SAD/height ratio (SADhR), waist circumference/height ratio (WHtR), and body mass index by 4 ancestral groups and 3 age groups. Boxes indicate population percentiles 25, 50, 75; whiskers identify p10 and p90. NHW = Non-Hispanic whites; NHB = Non-Hispanic blacks; Hispan = Hispanics; Asian = Non-Hispanic Asians.

doi:10.1371/journal.pone.0172245.g004




Fig 5. Women’s distributions of SAD/height ratio (SADhtR), waist circumference/height ratio (WHtR), and body mass index by 4 ancestral groups and 3 age groups. Boxes indicate population percentiles 25, 50, 75; whiskers identify p10 and p90. NHW = Non-Hispanic whites; NHB = Non-Hispanic blacks; Hispan = Hispanics; Asian = Non-Hispanic Asians.

doi:10.1371/journal.pone.0172245.g005

Table 1. Adiposity differences among US adults (according to SADHtR, WHtR, or BMI) for four ancestral groups, each compared to all other ancestries.

		<u>Beta coefficient</u>	<u>SE beta</u>	<u>Adjusted Wald F (ancestral contrast)</u>
Non-Hispanic Whites (vs all others)				
<i>Men –</i>	SADHtR	-0.0011	0.0008	1.8
	WHtR	0.004	0.004	1.4
	BMI, kg/m ²	0.08	0.24	0.1
<i>Women –</i>	SADHtR	-0.0062	0.0011	31.5***
	WHtR	-0.015	0.004	15.0**
	BMI, kg/m ²	-0.91	0.28	10.4*
Non-Hispanic Blacks (vs all others)				
<i>Men –</i>	SADHtR	0.0006	0.0009	0.4
	WHtR	-0.020	0.004	32.1***
	BMI, kg/m ²	0.14	0.26	0.3
<i>Women –</i>	SADHtR	0.0138	0.0009	212.2***
	WHtR	0.033	0.004	90.4***
	BMI, kg/m ²	3.43	0.28	150.9***
Hispanics (vs all others)				
<i>Men –</i>	SADHtR	0.0049	0.0013	14.9**
	WHtR	0.023	0.005	19.6**
	BMI, kg/m ²	1.00	0.35	7.9*
<i>Women –</i>	SADHtR	0.0053	0.0015	12.6*
	WHtR	0.022	0.005	16.3**
	BMI, kg/m ²	0.87	0.35	6.1
Non-Hispanic Asians (vs all others)				
<i>Men –</i>	SADHtR	-0.0113	0.0009	147.2***
	WHtR	-0.044	0.003	207.8***
	BMI, kg/m ²	-3.66	0.21	317.9***
<i>Women –</i>	SADHtR	-0.0145	0.0012	153.5***
	WHtR	-0.053	0.004	218.4***
	BMI, kg/m ²	-4.96	0.27	341.6***

* p < 0.01;
 ** p < 0.001;
 *** p < 0.0001

Models are adjusted for age (7 categories) and education (3 categories).

doi:10.1371/journal.pone.0172245.t001

Regarding the WHtR, adult population statistics were published for the United States in 1999–2004 (ages 20+; N = 12,936). In that earlier time period the WHtR for adult US men was 0.565 and women was 0.575, but these summary values did not show stratifications by age [39]. Subsequent adult population statistics on WHtR have been reported from China (in 2005–2007, ages 35–70; N = 43,841) [40], Canada (in 2007–2009, ages 20–69; N = 10,605) [9], and Colombia (in 2010, ages 20–64; N = 83,220) [41]. Of these large reported surveys, only the Colombian publication included breakdowns by age group. A comparison of these various reports is complicated by variation in anthropometric method or sample frames that differ by age or urban-rural status [42]. However, the limited data are consistent with previous observations that the WC, as well as the WHtR, has increased more among women than men during recent decades [9–11].

We are unaware of any previous report relating SADHtR to a marker of socioeconomic position, but adult WHtR has been associated inversely with educational attainment in one prior study [43]. SAD alone (without division by height), however, was inversely associated with educational attainment among Finnish adults, more strongly for women than men [36]. Similarly, adult WC alone was inversely associated with education, especially for women, in large studies from four other countries [44–47]. Our report confirms that a similar inverse association with education is found when SADHtR or WHtR is the adiposity indicator (Fig 3). We demonstrate further that either abdominal adiposity indicator had a stronger inverse association than BMI with educational attainment.

Associations between low income and adiposity are infrequently reported, perhaps because of difficulties in ascertaining the income level of individual participants. In this regard the NHANES calculation of poverty-income ratio—despite the absence of income information in about 7% of participants—demonstrates that women’s poverty had strongly inverse associations with SADHtR and WHtR. For men, however, we found that the lowest income level was associated with no change in abdominal adiposity but with a decrement of about 1 BMI unit (S1 Fig). Why the men’s pattern differs so greatly from the women’s merits further study.

A unique contribution of our report is its demonstration of substantial differences between ancestral groups in the empirical distributions of SADHtR and WHtR (Figs 4 [men] and 5 [women]). Among these box-and-whisker plots we present also an updated description of ancestral differences in the BMI distribution. A Scientific Statement from the American Heart Association recently addressed these distinctions with regard to BMI [48]. Given that some ancestral groups have BMI distributions substantially larger or smaller than the majority non-Hispanic white population, the authors questioned the value of applying universal BMI threshold values, such as 25 or 30 kg/m², to the assessment of diverse adult populations. Our models adjusted for age and education confirm that the ancestral differences in BMI are robust, especially as they apply to the lower BMI values that distinguish the non-Hispanic Asian population from all others (Table 1). The Asian differences are also substantial with regard to WHtR and SADHtR, although the distinctions appear to be attenuated (reduced Wald F statistic for ancestral contrast) when compared to the Asian differences in BMI. Some authorities have recommended that for clinical screening purposes a WHtR boundary should be set simply at 0.5, and that this threshold would be suitable for all ethnic groups [49]. We found among US adults, however, that the application of this universal screening threshold for WHtR would designate more than half of non-Hispanic Asians, and larger proportions of the remaining population, as being at increased health risk (Figs 4 and 5). Thus, the proposed WHtR threshold of 0.5 may be impractical for application in the contemporary United States [28]. To our knowledge, no universal threshold values for SADHtR have been proposed. Any anthropometric thresholds recommended for specific ancestral groups might still misclassify risk status because of confounding by sex [50], socioeconomic position, and the ambiguities inherent in terms such as “Asian”, “Hispanic”, “black”, and “white” [48].

A limitation of our report is related to the variety of available protocols for measurement of WC. The NHANES protocol, adopted also by the US National Institutes of Health and some Canadian organizations, specifies that the tape measure should be applied just above the uppermost lateral border of the ilium [32]. An alternative WC protocol in wide use, recommended by the World Health Organization (WHO), specifies that the tape measure should be applied midway between the inferior margin of the last rib and the crest of the ilium [6]. The reproducibility of the adult WC measured by either of these protocols is high, but the resulting WC values may differ slightly, most notably for women [51]. These alternative definitions of the WC result in small differences of the WHtR values that may influence the prevalence estimates of “high WHtR” or the WHtR percentile positions. Among our cited distributions of

adult WHtR obtained from outside the US, the recent Chinese article [40] depended on WC measured by the NHANES protocol while the Canadian [9] and Colombian [41] articles used the WHO protocol.

The anthropometric protocol for SAD measurement in Finland's survey was defined by the iliac crests [36], the same as our US protocol in NHANES. Evaluations of four different SAD protocols concluded that this site for measuring SAD was the best correlated with markers of cardiometabolic risk [52, 53].

We understand there are many reasons why BMI will continue to be calculated and reported as a primary index of weight status and health risk. In settings where a high quality scale is available, the simplicity of standardized weight measurement has its appeal. Whether patients and research participants will reliably disrobe and put down their various heavy objects is another question. It is likely, however, that some clinicians, researchers, and epidemiologists will come to prefer the SADHtR and WHtR as inexpensive markers more specifically focused on abdominal adiposity. Agreement on a standardized, common protocol for WC measurement will encourage practitioners to familiarize themselves with meaningful use of a tape measure and calculation of the WHtR. The NHANES success with standardized caliper measurements of SAD should build confidence that the SAD can also be obtained easily and interpreted by calculation of the SADHtR. The documentation of how WHtR and SADHtR have recently been distributed among US adults may help to promote the adoption of these abdominal adiposity indicators.

Supporting information

S1 File. Adult distributions (SADHtR, WHtR) by sex.

Table A. SAD/height ratio (SADHtR) for men aged 20+

Table B. SAD/height ratio (SADHtR) for women aged 20+

Table C. Waist circumference/height ratio (WHtR) for men aged 20+

Table D. Waist circumference/height ratio (WHtR) for women aged 20+.

(DOCX)

S1 Fig. Sex-specific beta coefficients (95% confidence range) associated with low family income (<100% of poverty guidelines or 100% to <200% of poverty guidelines) compared to persons with family income \geq 200% of poverty guidelines. Separate models, each adjusted for age and ancestral groups, were prepared for [top] SADHtR, [middle] WHtR, and [bottom] BMI. F indicates the adjusted Wald F statistic, a summary effect describing confidence that contrasts in family income (3-levels) have influenced the observed value for the adiposity indicator. These models include a category of persons (7%) with unknown family income. (TIF)

Acknowledgments

The authors acknowledge the participants in the 2011–2014 NHANES, and the efforts of the NHANES field staff. The findings and conclusions in this article are those of the authors and do not necessarily reflect the official position of the Centers for Disease Control and Prevention.

Author Contributions

Conceptualization: HSK KMB.

Data curation: HSK KMB.

Formal analysis: HSK KMB.

Methodology: HSK.

Project administration: HSK.

Resources: HSK.

Software: KMB.

Supervision: HSK.

Validation: HSK KMB.

Visualization: HSK.

Writing – original draft: HSK.

Writing – review & editing: HSK KMB.

References

1. Garrow JS, Webster J. Quetelet's index (W/H²) as a measure of fatness. *Int J Obes*. 1985; 9, 147–153.
2. Cornier MA, Despres JP, Davis N, Grossniklaus DA, Klein S, Lamarche B, et al. Assessing adiposity: A scientific statement from the American Heart Association. *Circulation*. 2011; 124, 1996–2019. doi: [10.1161/CIR.0b013e318233bc6a](https://doi.org/10.1161/CIR.0b013e318233bc6a) PMID: [21947291](https://pubmed.ncbi.nlm.nih.gov/21947291/)
3. Manolopoulos KN, Karpe F, Frayn KN. Gluteofemoral body fat as a determinant of metabolic health. *Int J Obes (Lond)*. 2010; 34, 949–959.
4. Tchernof A, Despres JP. Pathophysiology of human visceral obesity: an update. *Physiol Rev*. 2013; 93, 359–404. doi: [10.1152/physrev.00033.2011](https://doi.org/10.1152/physrev.00033.2011) PMID: [23303913](https://pubmed.ncbi.nlm.nih.gov/23303913/)
5. Kuczmarski RJ, Flegal KM. Criteria for definition of overweight in transition: Background and recommendations for the United States. *Am J Clin Nutr*. 2000; 72, 1074–1081. PMID: [11063431](https://pubmed.ncbi.nlm.nih.gov/11063431/)
6. World Health Organization Expert Committee on Physical Status. Physical Status: The Use and Interpretation of Anthropometry—Report of a WHO Expert Committee. Geneva: WHO technical report series; 1995 [updated 30 March 2012; Accessed 13 January 2017]; <http://helid.digicollection.org/en/d/Jh0211e/#Jh0211e>.
7. Fryar CD, Gu Q, Ogden CL, Flegal KM. Anthropometric reference data for children and adults: United States, 2011–2014. *Vital and Health Statistics, Series 3*. 2016 [Accessed 13 January 2017]; Number 39 (August 2016): https://www.cdc.gov/nchs/data/series/sr_03/sr03_039.pdf.
8. NCD Risk Factor Collaboration. Trends in adult body-mass index in 200 countries from 1975 to 2014: A pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet*. 2016 [Accessed 13 January 2017]; 387 (10026): Available from: <http://www.sciencedirect.com/science/article/pii/S014067361630054X>.
9. Shields M, Tremblay MS, Connor Gorber S, Janssen I. Measures of abdominal obesity within body mass index categories, 1981 and 2007–2009. *Health Rep*. 2012; 23, 33–38.
10. Albrecht SS, Gordon-Larsen P, Stern D, Popkin BM. Is waist circumference per body mass index rising differentially across the United States, England, China and Mexico? *Eur J Clin Nutr*. 2015; 69, 1306–1312. doi: [10.1038/ejcn.2015.71](https://doi.org/10.1038/ejcn.2015.71) PMID: [25944180](https://pubmed.ncbi.nlm.nih.gov/25944180/)
11. Freedman DS, Ford ES. Are the recent secular increases in the waist circumference of adults independent of changes in BMI? *Am J Clin Nutr*. 2015; 101, 425–431. doi: [10.3945/ajcn.114.094672](https://doi.org/10.3945/ajcn.114.094672) PMID: [25733625](https://pubmed.ncbi.nlm.nih.gov/25733625/)
12. Tanamas SK, Lean ME, Combet E, Vlassopoulos A, Zimmet PZ, Peeters A. Changing guards: time to move beyond body mass index for population monitoring of excess adiposity. *QJM*. 2016 [Accessed 13 January 2017]; 109 (7): Available from: <http://qjmed.oxfordjournals.org/content/109/7/443>.
13. Hsieh SD, Yoshinaga H. Abdominal fat distribution and coronary heart disease risk factors in men—waist/height ratio as a simple and useful predictor. *Int J Obes Relat Metab Disord*. 1995; 19, 585–589. PMID: [7489031](https://pubmed.ncbi.nlm.nih.gov/7489031/)
14. Ashwell M, Cole TJ, Dixon AK. Ratio of waist circumference to height is strong predictor of intra-abdominal fat. *BMJ*. 1996; 313, 559–560.

15. Schneider HJ, Klotsche J, Silber S, Stalla GK, Wittchen HU. Measuring abdominal obesity: effects of height on distribution of cardiometabolic risk factors risk using waist circumference and waist-to-height ratio. *Diabetes Care*. [Letter]. 2011; 34, e7.
16. van der Kooy K, Leenen R, Seidell J, Deurenberg P, Visser M. Abdominal diameters as indicators of visceral fat: comparison between magnetic resonance imaging and anthropometry. *Br J Nutr*. 1993; 70, 47–58. PMID: [8399118](#)
17. Jensen MD, Kanaley JA, Reed JE, Sheedy PF. Measurement of abdominal and visceral fat with computed tomography and dual-energy x-ray absorptiometry. *Am J Clin Nutr*. 1995; 61, 274–278. PMID: [7840063](#)
18. Sampaio LR, Simoes EJ, Assis AM, Ramos LR. Validity and reliability of the sagittal abdominal diameter as a predictor of visceral abdominal fat. *Arq Bras Endocrinol Metabol*. 2007 [Accessed 13 January 2017]; 51 (6): Available from: <http://www.scielo.br/pdf/abem/v51n6/a13v51n6.pdf>.
19. Kullberg J, von Below C, Lonn L, Lind L, Ahlstrom H, Johansson L. Practical approach for estimation of subcutaneous and visceral adipose tissue. *Clin Physiol Funct Imaging*. 2007; 27, 148–153. doi: [10.1111/j.1475-097X.2007.00728.x](#) PMID: [17445065](#)
20. Yim JY, Kim D, Lim SH, Park MJ, Choi SH, Lee CH, et al. Sagittal abdominal diameter is a strong anthropometric measure of visceral adipose tissue in the Asian general population. *Diabetes Care*. 2010; 33, 2665–2670. doi: [10.2337/dc10-0606](#) PMID: [20843976](#)
21. Richelsen B, Pedersen SB. Associations between different anthropometric measurements of fatness and metabolic risk parameters in non-obese, healthy, middle-aged men. *Int J Obes*. 1995; 19, 169–174.
22. Kumlin L, Dimberg L, Marin P. Ratio of abdominal sagittal diameter to height is strong indicator of coronary risk. *BMJ*. 1997; 314, 830.
23. Williamson DF, Kahn HS, Worthman CM, Burnette JC, Russell CM. Precision of recumbent anthropometry. *Am J Hum Biol*. 1993; 5, 159–167.
24. Rasmussen MH, Andersen T, Breum L, Hilsted J, Gotzsche PC. Observer variation in measurements of waist-hip ratio and the abdominal sagittal diameter. *Int J Obes Relat Metab Disord*. 1993; 17, 323–327. PMID: [8392497](#)
25. Nordhamn K, Sodergren E, Olsson E, Karlstrom B, Vessby B, Berglund L. Reliability of anthropometric measurements in overweight and lean subjects: consequences for correlations between anthropometric and other variables. *Int J Obes Relat Metab Disord*. 2000; 24, 652–657.
26. Carlsson AC, Riserus U, Engstrom G, Arnlov J, Melander O, Leander K, et al. Novel and established anthropometric measures and the prediction of incident cardiovascular disease: a cohort study. *Int J Obes (Lond)*. 2013; 37, 1579–1585.
27. Kahn HS, McKeever Bullard K. Beyond body mass index: Advantages of abdominal measurements for recognizing cardiometabolic disorders. *Am J Med*. 2016; 129, 74–81.e72. doi: [10.1016/j.amjmed.2015.08.010](#) PMID: [26302146](#)
28. Savva SC, Lamnisos D, Kafatos AG. Predicting cardiometabolic risk: waist-to-height ratio or BMI. A meta-analysis. *Diabetes Metab Syndr Obes*. 2013 [Accessed 13 January 2017]; 6: Available from: <http://dx.doi.org/10.2147/DMSO.S34220>.
29. Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: Systematic review and meta-analysis. *Obes Rev*. 2012; 13, 275–286. doi: [10.1111/j.1467-789X.2011.00952.x](#) PMID: [22106927](#)
30. Ashwell M, Mayhew L, Richardson J, Rickayzen B. Waist-to-height ratio is more predictive of years of life lost than body mass index. *PLoS One*. 2014 [Accessed 13 January 2017]; 9 (9): Available from: <http://dx.doi.org/10.1371/journal.pone.0103483>.
31. CDC/National Center for Health Statistics. About the National Health and Nutrition Examination Survey. Hyattsville, MD: NCHS; 2014 [updated 6 November 2015; Accessed 13 January 2017]; http://www.cdc.gov/nchs/nhanes/about_nhanes.htm.
32. National Center for Health Statistics. Anthropometry Procedures Manual—National Health and Nutrition Examination Survey (NHANES). Hyattsville, MD: NCHS; 2013 [Accessed 13 January 2017]; http://www.cdc.gov/nchs/data/nhanes/nhanes_13_14/2013_Anthropometry.pdf.
33. National Center for Health Statistics. NHANES 2013–2014 Data Documentation, Codebook, and Frequencies—Body Measures (BMX_H) Hyattsville, MD: NCHS; 2015 [Accessed 13 January 2017]; http://wwwn.cdc.gov/Nchs/Nhanes/2013-2014/BMX_H.htm.
34. Ogden CL, Carroll MD, Fryar CD, Flegal KM. Prevalence of obesity among adults and youth: United States, 2011–2014. *NCHS data brief*. 2015 [Accessed 13 January 2017]; (219): Available from: <http://www.cdc.gov/nchs/data/databriefs/db219.htm>. PMID: [26633046](#)

35. National Center for Health Statistics. NHANES 2013–2014 Data Documentation, Codebook, and Frequencies—Demographic Variables and Sample Weights (DEMO_H). Hyattsville, MD: NCHS; 2015 [Accessed 13 January 2017]; https://wwwn.cdc.gov/Nchs/Nhanes/2013-2014/DEMO_H.htm.
36. Kahn HS, Rissanen H, Bullard KM, Knekt P. The population distribution of the sagittal abdominal diameter (SAD) and SAD/height ratio among Finnish adults. *Clin Obes.* 2014; 4, 333–341. doi: [10.1111/cob.12078](https://doi.org/10.1111/cob.12078) PMID: [25826163](https://pubmed.ncbi.nlm.nih.gov/25826163/)
37. Sugihara M, Oka R, Sakurai M, Nakamura K, Moriuchi T, Miyamoto S, et al. Age-related changes in abdominal fat distribution in Japanese adults in the general population. *Intern Med.* 2011 [Accessed 13 January 2017]; 50 (7): Available from: https://www.jstage.jst.go.jp/article/internalmedicine/50/7/50_7_679/_pdf.
38. Demerath EW, Rogers NL, Reed D, Lee M, Choh AC, Siervogel RM, et al. Significant associations of age, menopausal status and lifestyle factors with visceral adiposity in African-American and European-American women. *Ann Hum Biol.* 2011; 38, 247–256. doi: [10.3109/03014460.2010.524893](https://doi.org/10.3109/03014460.2010.524893) PMID: [21175300](https://pubmed.ncbi.nlm.nih.gov/21175300/)
39. Flegal KM, Shepherd JA, Looker AC, Graubard BI, Borrud LG, Ogden CL, et al. Comparisons of percentage body fat, body mass index, waist circumference, and waist-stature ratio in adults. *Am J Clin Nutr.* 2009; 89, 500–508. doi: [10.3945/ajcn.2008.26847](https://doi.org/10.3945/ajcn.2008.26847) PMID: [19116329](https://pubmed.ncbi.nlm.nih.gov/19116329/)
40. Peng Y, Li W, Wang Y, Bo J, Chen H. The cut-off point and boundary values of waist-to-height ratio as an indicator for cardiovascular risk factors in chinese adults from the PURE Study. *PLoS One.* 2015 [Accessed 13 January 2017]; 10 (12): Available from: <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0144539>.
41. Ramirez-Velez R, Correa-Bautista JE, Martinez-Torres J, Meneses-Echavez JF, Gonzalez-Ruiz K, Gonzalez-Jimenez E, et al. LMS tables for waist circumference and waist-height ratio in Colombian adults: analysis of nationwide data 2010. *Eur J Clin Nutr.* 2016 [Accessed 13 January 2017]: Available from: <http://www.nature.com/ejcn/journal/vaop/ncurrent/full/ejcn201646a.html>.
42. Xi B, Liang Y, He T, Reilly KH, Hu Y, Wang Q, et al. Secular trends in the prevalence of general and abdominal obesity among Chinese adults, 1993–2009. *Obes Rev.* 2012; 13, 287–296. doi: [10.1111/j.1467-789X.2011.00944.x](https://doi.org/10.1111/j.1467-789X.2011.00944.x) PMID: [22034908](https://pubmed.ncbi.nlm.nih.gov/22034908/)
43. Kowalkowska J, Poinhos R, Franchini B, Afonso C, Correia F, Pinhao S, et al. General and abdominal adiposity in a representative sample of Portuguese adults: dependency of measures and socio-demographic factors' influence. *Br J Nutr.* 2016; 115, 185–192. doi: [10.1017/S0007114515004055](https://doi.org/10.1017/S0007114515004055) PMID: [26489926](https://pubmed.ncbi.nlm.nih.gov/26489926/)
44. Hermann S, Rohrmann S, Linseisen J, May AM, Kunst A, Besson H, et al. The association of education with body mass index and waist circumference in the EPIC-PANACEA study. *BMC Public Health.* 2011 [Accessed 13 January 2017]; 11: Available from: <http://pubmedcentralcanada.ca/picrender.cgi?accid=PMC3070651&blobtype=pdf>.
45. Sardinha LB, Santos DA, Silva AM, Coelho-e-Silva MJ, Raimundo AM, Moreira H, et al. Prevalence of overweight, obesity, and abdominal obesity in a representative sample of Portuguese adults. *PLoS One.* 2012 [Accessed 13 January 2017]; 7 (10): Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3485315/pdf/pone.0047883.pdf>.
46. Yoon JM, Kim JH, Suh BS, Park SM. Educational disparities in distribution of cardiovascular risk factors and quality of care in Korean adults: Korean National Health and Nutrition Survey IV. *Korean J Fam Med.* 2013 [Accessed 13 January 2017]; 34 (1): Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3560337/pdf/kjfm-34-27.pdf>.
47. Alves RF, Faerstein E. Educational inequality in the occurrence of abdominal obesity: Pro-Saude Study. *Rev Saude Publica.* 2015 [Accessed 13 January 2017]; 49: Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4587820/pdf/0034-8910-rsp-S0034-89102015049005786.pdf>.
48. Rao G, Powell-Wiley TM, Ancheta I, Hairston K, Kirley K, Lear SA, et al. Identification of obesity and cardiovascular risk in ethnically and racially diverse populations: A scientific statement from the American Heart Association. *Circulation.* 2015; 132, 457–472. doi: [10.1161/CIR.0000000000000223](https://doi.org/10.1161/CIR.0000000000000223) PMID: [26149446](https://pubmed.ncbi.nlm.nih.gov/26149446/)
49. Ashwell M, Gibson S. A proposal for a primary screening tool: 'Keep your waist circumference to less than half your height'. *BMC Med.* 2014 [Accessed 13 January 2017]; 12: Available from: <http://www.biomedcentral.com/1741-7015/12/207>.
50. Klimentidis YC, Arora A, Zhou J, Kittles R, Allison DB. The genetic contribution of West-African ancestry to protection against central obesity in African-American men but not women: Results from the ARIC and MESA studies. *Front Genet.* 2016 [Accessed 13 January 2017]; 7: Available from: <http://journal.frontiersin.org/article/10.3389/fgene.2016.00089/full>.
51. Mason C, Katzmarzyk PT. Variability in waist circumference measurements according to anatomic measurement site. *Obesity (Silver Spring).* 2009; 17, 1789–1795.

52. de Almeida Paula HA, de Cassia Lanes Ribeiro R, de Lima Rosado LEFP, Abranches MV, do Carmo Castro Franceschini S. Relationship between waist circumference and supine abdominal height measured at different anatomical sites and cardiometabolic risk factors in older women. *J Hum Nutr Diet.* 2012; 25, 563–568. doi: [10.1111/j.1365-277X.2012.01267.x](https://doi.org/10.1111/j.1365-277X.2012.01267.x) PMID: [23173640](https://pubmed.ncbi.nlm.nih.gov/23173640/)
53. Anunciacao PC, Ribeiro RCL, Pereira MQ, Comunian M. Different measurements of waist circumference and sagittal abdominal diameter and their relationship with cardiometabolic risk factors in elderly men. *J Hum Nutr Diet.* 2014 [Accessed 13 January 2017]: Available from: [http://onlinelibrary.wiley.com/enhanced/doi/10.1111/jhn.12201/](http://onlinelibrary.wiley.com/doi/10.1111/jhn.12201/).