

Poster presentation

Open Access

## Open label multicenter study of once weekly Etanercept 0.8 mg/kg in active polyarticular Juvenile idiopathic arthritis (JIA)

G Horneff\*<sup>1</sup>, K Minden<sup>2</sup>, I Foeldvari<sup>3</sup>, J Kuemmerle-Deschner<sup>4</sup>, A Thon<sup>5</sup>, H Girschick<sup>6</sup> and HI Huppertz<sup>7</sup>

Address: <sup>1</sup>Asklepios Clinic, Sankt Augustin, Germany, <sup>2</sup>Charite, Berlin, Germany, <sup>3</sup>Private office, Hamburg, Germany, <sup>4</sup>University Hospital, Tuebingen, Germany, <sup>5</sup>University Hospital, Hannover, Germany, <sup>6</sup>University Hospital, Wuerzburg, Germany and <sup>7</sup>Prof. Hess Kinderklinik, Bremen, Germany

\* Corresponding author

from 15<sup>th</sup> Paediatric Rheumatology European Society (PreS) Congress  
London, UK. 14–17 September 2008

Published: 15 September 2008

*Pediatric Rheumatology* 2008, **6**(Suppl 1):P39 doi:10.1186/1546-0096-6-S1-P39

This abstract is available from: <http://www.ped-rheum.com/content/6/S1/P39>

© 2008 Horneff et al; licensee BioMed Central Ltd.

### Background

In Europe Etanercept is licensed for the treatment of resistant polyarticular JIA at a dosage of 0.4 mg/kg bw. twice weekly in children older than 4 years.

### Objectives

To evaluate the safety and efficacy of Etanercept once weekly 0.8 mg/kg up to 50 mg in a formal trial.

### Methods

At each study site an independent ethics committee approved the protocol, and each patient's parent gave written informed consent (EudraCT No. 2007-000255-34). 20 patients 4 to 17 years old were included and received 0.8 mg/kg bw. of etanercept subcutaneously once weekly for 12 weeks "Active" polyarticular disease was defined by the presence of five or more active joints. PedACR30/50/70 criteria were calculated.

Safety assessments were based on adverse events (AE) reported.

### Results

15 of 20 JIA patients, 16 girls and 4 boys, mean age 12.9 years, disease duration 4.1 years, already have completed the 12 week study period. The mean dosage was 0.80 +/- 0.04 mg/kg Etanercept. Concomitant treatments were kept stable 3 months before and throughout the study and consisted of NSAID (n = 20), prednisone (n = 4), meth-

otrexate (n = 12), leflunomide (n = 2), sulfasalazine (n = 1). A PedACR 30/50/70 response was reached by 73%/26%/10% of patients after 4 weeks, 86%/73%/40% after 8 weeks and 92%/92%/79% after 12 weeks of treatment. There were 33 AEs but no SAE: 9 minor infections 12 injection site reactions and 12 other AEs. There was no drop out.

### Conclusion

These data indicate that once weekly application of Etanercept at double dosage of 0.8 mg/kg bodyweight up to 50 mg per injection is safe and efficacious in polyarticular JIA patients.