



Crystal structure of (1,3-di-*tert*-butyl- η^5 -cyclopentadienyl)trimethylhafnium(IV)

Adrián Pérez-Redondo,* Víctor Varela-Izquierdo and Carlos Yélamos

Departamento de Química Orgánica y Química Inorgánica, Universidad de Alcalá, Campus Universitario, ES 28871 Alcalá de Henares (Madrid), Spain. *Correspondence e-mail: adrian.perez@uah.es

Received 17 March 2015; accepted 23 March 2015

Edited by M. Weil, Vienna University of Technology, Austria


The molecule of the title organometallic hafnium(IV) compound, $[\text{Hf}(\text{CH}_3)_3(\text{C}_{13}\text{H}_{21})]$ or $[\text{HfMe}_3(\eta^5\text{-C}_5\text{H}_3\text{-1,3-}^t\text{Bu}_2)]$, adopts the classical three-legged piano-stool geometry for monocyclopentadienylhafnium(IV) derivatives with the three methyl groups bonded to the Hf(IV) atom at the legs. The C atoms of the two *tert*-butyl group bonded to the cyclopentadienyl (Cp) ring are 0.132 (5) and 0.154 (6) Å above the Cp least-squares plane. There are no significant intermolecular interactions present between the molecules in the crystal structure.

Keywords: crystal structure; hafnium; cyclopentadienyl ligand; organometallic compound.

CCDC reference: 1055619

1. Related literature

The synthesis of the compound was described by Cuenca *et al.* (1996). For the structures of related Hf^{IV} derivatives and a comparison of Hf—C bond lengths, see: Itagaki *et al.* (2009); Schäfer *et al.* (2013); Shah *et al.* (1996); Swenson *et al.* (2000).



2. Experimental

2.1. Crystal data

$[\text{Hf}(\text{CH}_3)_3(\text{C}_{13}\text{H}_{21})]$

$M_r = 400.89$

Monoclinic, $P2_1/n$
 $a = 13.238$ (3) Å
 $b = 9.613$ (2) Å
 $c = 14.486$ (3) Å
 $\beta = 109.63$ (2)°
 $V = 1736.3$ (7) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 5.99$ mm^{−1}
 $T = 200$ K
 $0.42 \times 0.14 \times 0.11$ mm

2.2. Data collection

Nonius KappaCCD diffractometer
 Absorption correction: multi-scan
 (Blessing, 1995)
 $T_{\min} = 0.297$, $T_{\max} = 0.531$

29484 measured reflections
 3136 independent reflections
 2471 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.100$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.073$
 $S = 1.12$
 3136 reflections

164 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 1.42$ e Å^{−3}
 $\Delta\rho_{\min} = -1.35$ e Å^{−3}

Table 1

Selected geometric parameters (Å, °).

C1—Hf1	2.198 (6)	C12—Hf1	2.500 (4)
C2—Hf1	2.211 (6)	C13—Hf1	2.524 (4)
C3—Hf1	2.213 (6)	C14—Hf1	2.484 (5)
C11—Hf1	2.519 (5)	C15—Hf1	2.468 (5)
C1—Hf1—C2	103.7 (2)	C2—Hf1—C3	102.4 (3)
C1—Hf1—C3	99.7 (2)		

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *DIRAX* (Duisenberg *et al.*, 2000); data reduction: *EVALCCD* (Duisenberg *et al.*, 2003); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014/7* (Sheldrick, 2015); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *publCIF* (Westrip, 2010).

Acknowledgements

We are grateful to the Factoría de Cristalización (CONSO-LIDER-INGENIO 2010 CSD2006-00015) for support of this research.

Supporting information for this paper is available from the IUCr electronic archives (Reference: WM5139).

References

- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
- Cuenca, T., Montejano, C. & Royo, P. (1996). *J. Organomet. Chem.* **514**, 93–96.
- Duisenberg, A. J. M., Hooft, R. W. W., Schreurs, A. M. M. & Kroon, J. (2000). *J. Appl. Cryst.* **33**, 893–898.
- Duisenberg, A. J. M., Kroon-Batenburg, L. M. J. & Schreurs, A. M. M. (2003). *J. Appl. Cryst.* **36**, 220–229.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Itagaki, K., Hasumi, S., Fujiki, M. & Nomura, K. (2009). *J. Mol. Catal. A Chem.* **303**, 102–109.
- Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Schäfer, S., Bauer, H., Becker, J., Sun, Y. & Sitzmann, H. (2013). *Eur. J. Inorg. Chem.* pp. 5694–5700.
- Shah, S. A. A., Dorn, H., Voigt, A., Roesky, H. W., Parisini, E., Schmidt, H.-G. & Noltemeyer, M. (1996). *Organometallics*, **15**, 3176–3181.

Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.

Swenson, D. C., Guo, Z., Crowther, D. J., Baenziger, N. C. & Jordan, R. F.
(2000). *Acta Cryst.* **C56**, e313–e314.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2015). E71, m100–m101 [doi:10.1107/S205698901500585X]

Crystal structure of (1,3-di-*tert*-butyl- η^5 -cyclopentadienyl)trimethylhafnium(IV)

Adrián Pérez-Redondo, Víctor Varela-Izquierdo and Carlos Yélamos

S1. Synthesis and crystallization

The title compound was synthesized according to a literature procedure (Cuenca *et al.*, 1996). Crystals were obtained from the resultant oil by removing the volatile components of a *n*-hexane solution.

S2. Refinement

H atoms attached to sp^2 C-atoms were placed geometrically, with C—H = 0.95 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. Methyl H-atoms were refined using a rotating-group model, with C—H = 0.98 Å, and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure of $[\text{Hf}(\eta^5\text{-1,3-Bu}_2\text{C}_5\text{H}_3)\text{Me}_3]$ with displacement ellipsoids drawn at the 50% probability level. Hydrogen atoms are omitted for clarity.

(1,3-Di-*tert*-butyl- η^5 -cyclopentadienyl)trimethylhafnium(IV)*Crystal data*[Hf(CH₃)₃(C₁₃H₂₁)] $M_r = 400.89$ Monoclinic, $P2_1/n$ $a = 13.238$ (3) Å $b = 9.613$ (2) Å $c = 14.486$ (3) Å $\beta = 109.63$ (2)° $V = 1736.3$ (7) Å³ $Z = 4$ $F(000) = 792$ $D_x = 1.534$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 117 reflections

 $\theta = 3\text{--}21^\circ$ $\mu = 5.99$ mm⁻¹ $T = 200$ K

Prism, colourless

 $0.42 \times 0.14 \times 0.11$ mm*Data collection*

Nonius KappaCCD

diffractometer

Radiation source: Enraf Nonius FR590

Horizontally mounted graphite crystal
monochromatorDetector resolution: 9 pixels mm⁻¹

CCD scans

Absorption correction: multi-scan
(Blessing, 1995) $T_{\min} = 0.297$, $T_{\max} = 0.531$

29484 measured reflections

3136 independent reflections

2471 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.100$ $\theta_{\max} = 25.2^\circ$, $\theta_{\min} = 3.3^\circ$ $h = -15 \rightarrow 15$ $k = -11 \rightarrow 10$ $l = -17 \rightarrow 16$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.032$ $wR(F^2) = 0.073$ $S = 1.12$

3136 reflections

164 parameters

0 restraints

0 constraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: structure-
invariant direct methodsHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.027P)^2 + 1.470P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 1.42$ e Å⁻³ $\Delta\rho_{\min} = -1.35$ e Å⁻³Extinction correction: *SHELXL2014/7*
(Sheldrick, 2015)

Extinction coefficient: 0.0113 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0972 (5)	−0.0651 (7)	0.2994 (4)	0.0639 (18)
H1A	0.0395	−0.1199	0.3098	0.096*
H1B	0.1367	−0.1231	0.2675	0.096*
H1C	0.0664	0.015	0.2576	0.096*
C2	0.3457 (5)	0.0906 (7)	0.4060 (5)	0.0643 (18)
H2A	0.4003	0.0178	0.4168	0.096*

H2B	0.376	0.1704	0.4483	0.096*
H2C	0.3221	0.1199	0.3372	0.096*
C3	0.1226 (5)	0.2003 (6)	0.4579 (5)	0.069 (2)
H3A	0.0907	0.2447	0.3937	0.103*
H3B	0.1737	0.2644	0.5024	0.103*
H3C	0.066	0.177	0.4848	0.103*
C11	0.2768 (4)	−0.0503 (5)	0.6213 (4)	0.0337 (11)
C12	0.3349 (4)	−0.1313 (5)	0.5761 (3)	0.0338 (11)
H12	0.4093	−0.1215	0.5863	0.041*
C13	0.2663 (4)	−0.2296 (5)	0.5131 (3)	0.0326 (11)
C14	0.1638 (4)	−0.2058 (5)	0.5167 (4)	0.0347 (11)
H14	0.1007	−0.2555	0.4809	0.042*
C15	0.1698 (4)	−0.0956 (5)	0.5824 (4)	0.0361 (12)
H15	0.1111	−0.058	0.5979	0.043*
C16	0.3228 (4)	0.0569 (6)	0.7029 (4)	0.0464 (14)
C17	0.2339 (6)	0.1215 (8)	0.7320 (5)	0.085 (2)
H17A	0.1948	0.0483	0.753	0.127*
H17B	0.1845	0.1712	0.6758	0.127*
H17C	0.2646	0.1869	0.786	0.127*
C18	0.3969 (6)	−0.0205 (6)	0.7922 (5)	0.076 (2)
H18A	0.4239	0.0443	0.8472	0.113*
H18B	0.4574	−0.0601	0.7764	0.113*
H18C	0.357	−0.0955	0.8102	0.113*
C19	0.3869 (6)	0.1672 (6)	0.6724 (5)	0.077 (2)
H19A	0.3404	0.2161	0.6144	0.116*
H19B	0.4462	0.1232	0.6571	0.116*
H19C	0.4158	0.2338	0.7261	0.116*
C20	0.3013 (4)	−0.3463 (5)	0.4585 (4)	0.0425 (13)
C21	0.3666 (6)	−0.4500 (6)	0.5375 (5)	0.071 (2)
H21A	0.4269	−0.4014	0.585	0.106*
H21B	0.3938	−0.5248	0.5064	0.106*
H21C	0.3204	−0.4896	0.5714	0.106*
C22	0.2040 (5)	−0.4224 (6)	0.3889 (5)	0.0629 (18)
H22A	0.1602	−0.4593	0.426	0.094*
H22B	0.2283	−0.4992	0.3571	0.094*
H22C	0.1611	−0.3574	0.3389	0.094*
C23	0.3706 (5)	−0.2918 (7)	0.4017 (5)	0.0697 (19)
H23A	0.3286	−0.2277	0.3507	0.105*
H23B	0.3953	−0.3699	0.3714	0.105*
H23C	0.4327	−0.2427	0.4465	0.105*
Hf1	0.20704 (2)	0.00819 (2)	0.44138 (2)	0.03781 (13)

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.065 (4)	0.068 (4)	0.049 (4)	−0.013 (3)	0.007 (3)	0.015 (3)
C2	0.064 (4)	0.077 (4)	0.054 (4)	−0.018 (3)	0.022 (3)	0.006 (3)
C3	0.057 (4)	0.051 (4)	0.088 (5)	0.007 (3)	0.011 (4)	0.018 (3)

C11	0.037 (3)	0.034 (2)	0.028 (3)	−0.002 (2)	0.008 (2)	−0.002 (2)
C12	0.026 (2)	0.042 (3)	0.032 (3)	0.003 (2)	0.008 (2)	0.003 (2)
C13	0.036 (3)	0.034 (3)	0.028 (3)	0.003 (2)	0.009 (2)	0.000 (2)
C14	0.031 (3)	0.034 (3)	0.040 (3)	−0.005 (2)	0.013 (2)	0.002 (2)
C15	0.033 (3)	0.038 (3)	0.043 (3)	0.000 (2)	0.019 (2)	0.002 (2)
C16	0.054 (3)	0.045 (3)	0.037 (3)	−0.006 (3)	0.012 (3)	−0.010 (2)
C17	0.090 (5)	0.093 (5)	0.075 (5)	0.001 (4)	0.031 (4)	−0.048 (4)
C18	0.089 (5)	0.067 (4)	0.042 (4)	−0.001 (3)	−0.016 (4)	−0.012 (3)
C19	0.104 (6)	0.070 (4)	0.055 (4)	−0.047 (4)	0.024 (4)	−0.018 (3)
C20	0.041 (3)	0.047 (3)	0.039 (3)	0.004 (2)	0.013 (2)	−0.007 (2)
C21	0.090 (5)	0.060 (4)	0.057 (4)	0.033 (4)	0.017 (4)	−0.008 (3)
C22	0.071 (4)	0.050 (4)	0.062 (4)	−0.007 (3)	0.016 (3)	−0.026 (3)
C23	0.072 (5)	0.071 (4)	0.079 (5)	−0.001 (3)	0.043 (4)	−0.027 (4)
Hf1	0.03159 (16)	0.04226 (18)	0.03505 (17)	−0.00414 (9)	0.00523 (10)	0.00804 (10)

Geometric parameters (Å, °)

C1—Hf1	2.198 (6)	C15—H15	0.95
C1—H1A	0.98	C16—C17	1.511 (9)
C1—H1B	0.98	C16—C19	1.513 (8)
C1—H1C	0.98	C16—C18	1.529 (8)
C2—Hf1	2.211 (6)	C17—H17A	0.98
C2—H2A	0.98	C17—H17B	0.98
C2—H2B	0.98	C17—H17C	0.98
C2—H2C	0.98	C18—H18A	0.98
C3—Hf1	2.213 (6)	C18—H18B	0.98
C3—H3A	0.98	C18—H18C	0.98
C3—H3B	0.98	C19—H19A	0.98
C3—H3C	0.98	C19—H19B	0.98
C11—C12	1.402 (7)	C19—H19C	0.98
C11—C15	1.406 (6)	C20—C23	1.517 (8)
C11—C16	1.531 (7)	C20—C22	1.530 (7)
C11—Hf1	2.519 (5)	C20—C21	1.545 (7)
C12—C13	1.412 (6)	C21—H21A	0.98
C12—Hf1	2.500 (4)	C21—H21B	0.98
C12—H12	0.95	C21—H21C	0.98
C13—C14	1.395 (6)	C22—H22A	0.98
C13—C20	1.531 (7)	C22—H22B	0.98
C13—Hf1	2.524 (4)	C22—H22C	0.98
C14—C15	1.409 (7)	C23—H23A	0.98
C14—Hf1	2.484 (5)	C23—H23B	0.98
C14—H14	0.95	C23—H23C	0.98
C15—Hf1	2.468 (5)		
Hf1—C1—H1A	109.5	H18A—C18—H18B	109.5
Hf1—C1—H1B	109.5	C16—C18—H18C	109.5
H1A—C1—H1B	109.5	H18A—C18—H18C	109.5
Hf1—C1—H1C	109.5	H18B—C18—H18C	109.5

H1A—C1—H1C	109.5	C16—C19—H19A	109.5
H1B—C1—H1C	109.5	C16—C19—H19B	109.5
Hf1—C2—H2A	109.5	H19A—C19—H19B	109.5
Hf1—C2—H2B	109.5	C16—C19—H19C	109.5
H2A—C2—H2B	109.5	H19A—C19—H19C	109.5
Hf1—C2—H2C	109.5	H19B—C19—H19C	109.5
H2A—C2—H2C	109.5	C23—C20—C22	109.6 (5)
H2B—C2—H2C	109.5	C23—C20—C13	111.8 (4)
Hf1—C3—H3A	109.5	C22—C20—C13	110.9 (4)
Hf1—C3—H3B	109.5	C23—C20—C21	109.6 (5)
H3A—C3—H3B	109.5	C22—C20—C21	108.5 (5)
Hf1—C3—H3C	109.5	C13—C20—C21	106.3 (4)
H3A—C3—H3C	109.5	C20—C21—H21A	109.5
H3B—C3—H3C	109.5	C20—C21—H21B	109.5
C12—C11—C15	106.1 (4)	H21A—C21—H21B	109.5
C12—C11—C16	126.5 (4)	C20—C21—H21C	109.5
C15—C11—C16	127.2 (5)	H21A—C21—H21C	109.5
C12—C11—Hf1	73.0 (3)	H21B—C21—H21C	109.5
C15—C11—Hf1	71.6 (3)	C20—C22—H22A	109.5
C16—C11—Hf1	124.1 (4)	C20—C22—H22B	109.5
C11—C12—C13	109.8 (4)	H22A—C22—H22B	109.5
C11—C12—Hf1	74.5 (3)	C20—C22—H22C	109.5
C13—C12—Hf1	74.6 (3)	H22A—C22—H22C	109.5
C11—C12—H12	125.1	H22B—C22—H22C	109.5
C13—C12—H12	125.1	C20—C23—H23A	109.5
Hf1—C12—H12	117.6	C20—C23—H23B	109.5
C14—C13—C12	106.8 (4)	H23A—C23—H23B	109.5
C14—C13—C20	127.3 (4)	C20—C23—H23C	109.5
C12—C13—C20	125.7 (4)	H23A—C23—H23C	109.5
C14—C13—Hf1	72.3 (3)	H23B—C23—H23C	109.5
C12—C13—Hf1	72.7 (3)	C1—Hf1—C2	103.7 (2)
C20—C13—Hf1	123.8 (3)	C1—Hf1—C3	99.7 (2)
C13—C14—C15	108.3 (4)	C2—Hf1—C3	102.4 (3)
C13—C14—Hf1	75.4 (3)	C1—Hf1—C15	113.1 (2)
C15—C14—Hf1	72.8 (3)	C2—Hf1—C15	138.5 (2)
C13—C14—H14	125.9	C3—Hf1—C15	90.0 (2)
C15—C14—H14	125.9	C1—Hf1—C14	88.1 (2)
Hf1—C14—H14	117.8	C2—Hf1—C14	136.9 (2)
C11—C15—C14	108.9 (4)	C3—Hf1—C14	116.4 (2)
C11—C15—Hf1	75.6 (3)	C15—Hf1—C14	33.1 (2)
C14—C15—Hf1	74.1 (3)	C1—Hf1—C12	128.3 (2)
C11—C15—H15	125.5	C2—Hf1—C12	88.6 (2)
C14—C15—H15	125.5	C3—Hf1—C12	126.7 (2)
Hf1—C15—H15	116.7	C15—Hf1—C12	53.7 (2)
C17—C16—C19	111.0 (6)	C14—Hf1—C12	53.8 (2)
C17—C16—C18	107.6 (6)	C1—Hf1—C11	142.6 (2)
C19—C16—C18	108.9 (5)	C2—Hf1—C11	106.0 (2)
C17—C16—C11	110.3 (5)	C3—Hf1—C11	95.6 (2)

C19—C16—C11	111.3 (5)	C15—Hf1—C11	32.7 (2)
C18—C16—C11	107.5 (5)	C14—Hf1—C11	54.5 (2)
C16—C17—H17A	109.5	C12—Hf1—C11	32.5 (2)
C16—C17—H17B	109.5	C1—Hf1—C13	96.4 (2)
H17A—C17—H17B	109.5	C2—Hf1—C13	104.6 (2)
C16—C17—H17C	109.5	C3—Hf1—C13	144.1 (2)
H17A—C17—H17C	109.5	C15—Hf1—C13	54.1 (2)
H17B—C17—H17C	109.5	C14—Hf1—C13	32.3 (2)
C16—C18—H18A	109.5	C12—Hf1—C13	32.6 (2)
C16—C18—H18B	109.5	C11—Hf1—C13	54.3 (2)
C15—C11—C12—C13	−2.5 (6)	Hf1—C14—C15—C11	−68.3 (3)
C16—C11—C12—C13	172.8 (5)	C13—C14—C15—Hf1	67.7 (3)
Hf1—C11—C12—C13	−66.9 (3)	C12—C11—C16—C17	−179.8 (6)
C15—C11—C12—Hf1	64.5 (3)	C15—C11—C16—C17	−5.4 (8)
C16—C11—C12—Hf1	−120.2 (5)	Hf1—C11—C16—C17	86.6 (6)
C11—C12—C13—C14	2.1 (5)	C12—C11—C16—C19	56.5 (8)
Hf1—C12—C13—C14	−64.7 (3)	C15—C11—C16—C19	−129.2 (6)
C11—C12—C13—C20	−173.5 (5)	Hf1—C11—C16—C19	−37.1 (6)
Hf1—C12—C13—C20	119.6 (5)	C12—C11—C16—C18	−62.7 (7)
C11—C12—C13—Hf1	66.8 (3)	C15—C11—C16—C18	111.6 (6)
C12—C13—C14—C15	−1.0 (5)	Hf1—C11—C16—C18	−156.3 (5)
C20—C13—C14—C15	174.6 (5)	C14—C13—C20—C23	134.0 (5)
Hf1—C13—C14—C15	−66.0 (3)	C12—C13—C20—C23	−51.3 (7)
C12—C13—C14—Hf1	65.0 (3)	Hf1—C13—C20—C23	41.1 (6)
C20—C13—C14—Hf1	−119.4 (5)	C14—C13—C20—C22	11.3 (7)
C12—C11—C15—C14	1.8 (6)	C12—C13—C20—C22	−174.0 (5)
C16—C11—C15—C14	−173.4 (5)	Hf1—C13—C20—C22	−81.6 (5)
Hf1—C11—C15—C14	67.3 (3)	C14—C13—C20—C21	−106.5 (6)
C12—C11—C15—Hf1	−65.4 (3)	C12—C13—C20—C21	68.2 (7)
C16—C11—C15—Hf1	119.3 (5)	Hf1—C13—C20—C21	160.6 (4)
C13—C14—C15—C11	−0.6 (6)		
