

Methodology article

Open Access

Cytokine generation, promoter activation, and oxidant-independent NF- κ B activation in a transfectable human neutrophilic cellular model

Thornin Ear and Patrick P McDonald*

Address: Pulmonary Division, Faculty of Medicine, Université de Sherbrooke and Centre de recherche du CHUS, Sherbrooke, Qc, Canada

Email: Thornin Ear - thornin.ear@USherbrooke.ca; Patrick P McDonald* - patrick.mcdonald@USherbrooke.ca

* Corresponding author

Published: 11 April 2008

Received: 6 November 2007

BMC Immunology 2008, 9:14 doi:10.1186/1471-2172-9-14

Accepted: 11 April 2008

This article is available from: <http://www.biomedcentral.com/1471-2172/9/14>

© 2008 Ear and McDonald; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Human neutrophils are key players of innate immunity, and influence inflammatory and immune reactions through the production of numerous cytokines and chemokines. Despite major advances in our understanding of this important functional response of neutrophils, the short lifespan of these cells and their resistance to transfection have always been an obstacle to the detailed dissection of signaling pathways and effector responses that is often possible in other cell types.

Results: Here, we report that granulocytic differentiation of human PLB-985 cells with DMSO yields cells that are neutrophil-like with respect to surface markers, acquisition of responsiveness to physiological neutrophil stimuli (fMLP, LPS), cytokine expression and production profile, and transcription factor activation profile (NF- κ B, C/EBP, AP-1, STAT). We also show that granulocytic PLB-985 cells can be reliably transfected by nucleofection in a rapid and efficient manner. Indeed, we overexpressed several proteins and luciferase constructs into these cells. In particular, overexpression of a dominant negative I κ B- α confirmed the central role of NF- κ B in the production of cytokines by granulocytes. Moreover, the use of PLB-985 granulocytes in which the NADPH oxidase is inactive due to the targeted disruption of a key component (gp91phox) revealed that NF- κ B activation and κ B-dependent responses are independent of endogenous reactive oxygen intermediates in these cells. Antioxidant studies performed in primary human neutrophils support this conclusion.

Conclusion: Our results unveil a new facet of the NF- κ B system of human granulocytes, and pave the way for deciphering signal transduction pathways and promoter activation in these cells.

Introduction

Human polymorphonuclear neutrophils are terminally differentiated cells that represent about 60% of all circulating leukocytes. Beside their notorious role as professional phagocytes, neutrophils can also express a wide array of cytokines and chemokines in response to physio-

logical stimuli [1]. In this regard, a mounting body of evidence shows that neutrophil-derived cytokines and chemokines play an important role in several inflammatory and immune reactions *in vivo* [2-8]. Although much progress has been made in understanding the various facets of neutrophil biology, studies of the signal transduc-

tion mechanisms leading to functional activation of neutrophils have remained hampered by the fact that these cells are very refractory to transfection. This owes much to the fact that their lifespan merely averages 16–24 h in the bloodstream [9]. Thus, the characterization of a cell line that can be differentiated into a neutrophilic phenotype, and that can be transfected efficiently and reliably, would prove a most useful research tool to extend our understanding of cytokine generation and upstream processes in human granulocytes.

Various approaches have already been employed to differentiate the human pro-myeloid cell line PLB-985 into neutrophil-like cells [10-17]. However, the functional properties of granulocytic PLB-985 cells have only been defined for some cellular responses, such as degranulation and the associated respiratory burst [17], or arachidonic acid metabolism [14]. Thus, much remains to be determined concerning the suitability of differentiated PLB-985 to serve as a cellular model for primary neutrophils, especially in the case of more recently described neutrophil functional responses, such as the production of inflammatory cytokines and chemokines, and the underlying activation of discrete transcription factor families. This being said, PLB-985 cells represent a potentially attractive model since they are transfectable. Indeed, a few groups have stably transfected PLB-985 cells prior to granulocytic differentiation for specific purposes [14,18-20]. However, stable transfections are not always feasible, as the overexpressed proteins can interfere with granulocytic differentiation, a process which typically spans 4 to 6 days. Transient transfection of PLB-985 cells, on the other hand, has only been achieved in isolated instances, and with mitigated success [16,21]. This owes much to drawbacks inherent to the transfection techniques used, i.e. electroporation and cationic liposome-mediated transfection. Indeed, electroporation of PLB-985 cells reportedly results in high mortality rates while only achieving moderate transfection efficiency [16,21], whereas the liposome-based approach had a low transfection efficiency and required that the cells be pretreated with TPA for 4 h prior to transfection [21]. This in itself is problematic, because TPA is a differentiating agent that commits the cells to the monocytic lineage [10]. In contrast, a recent paper, which was published while we were completing the current study, reported the efficient transfection of PLB-985 cells with the nucleofection technique [22], an approach which entails lower mortality rates.

In the present study, we ascertained that granulocytic PLB-985 cells respond like primary neutrophils in terms of inflammatory cytokine production and transcription factor activation. We then optimized the transient transfection of granulocytic PLB-985 cells using the nucleofection technique, which yields high transfection efficiencies, rel-

atively low mortality rates, and rapid overexpression of various proteins. We also show that this approach lends itself well to promoter activation studies in neutrophil-like PLB-985 cells. We finally applied this approach to confirm the implication of NF- κ B in inflammatory cytokine production by human granulocytes, and to elucidate the long-standing issue of whether endogenous reactive oxygen intermediates (ROI)¹ influence transcription factor activation and downstream processes in cells that are heavy ROI producers, such as neutrophilic granulocytes.

Results

Granulocytic differentiation of PLB-985 cells

We initially differentiated PLB-985 cells using either DMSO or DMF, as both agents were reported to promote granulocytic differentiation [10-13,15,17]. Culture of PLB-985 cells with either agent resulted in a gradual inhibition of proliferation (that was most pronounced with DMSO). By day 5 of differentiation, over 80% of the cells had acquired a typically neutrophilic morphology i.e. lobular nuclei, and all of the cells had an increased granularity (Figure 1A), as reported previously [16,17]. Differentiating PLB-985 cells also displayed an augmented expression of cell surface markers such as the integrin, CD11b – a phenomenon that was especially evident in DMSO-differentiated cells (Figure 1B and 1C). Similarly, neutrophil-like PLB-985 cells acquired the ability to generate IL-8 in response to LPS and to the chemoattractant, fMLP (Figure 1D), in keeping with the observation that granulocytic differentiation correlates with the onset of CD14 and fMLP receptor surface expression [17,23]. Again, the acquisition of fMLP and LPS responsiveness was most marked in DMSO-differentiated cells (Figure 1D). Together, the above observations indicate that DMSO treatment leads to a more complete differentiation of PLB-985 cells into a neutrophil-like phenotype. As a result, we utilized DMSO as a differentiating agent in all subsequent experiments.

Cytokine expression profile of granulocytic PLB-985 cells

We next determined the extent to which the cytokine expression pattern of DMSO-differentiated PLB-985 cells conforms to that of primary neutrophils. To this end, granulocytic PLB-985 cells were stimulated with either LPS or TNF for increasing lengths of time, and cytokine release was measured. As shown in Figure 2A, neutrophil-like PLB-985 cells generate IL-8, Mip-1 α , Mip-1 β , and TNF α with kinetics that are very reminiscent of those observed in human neutrophils [24-26]. Another similarity between granulocytic PLB-985 cells and human neutrophils is the delayed IL-8 production kinetics observed in response to LPS (compared to those elicited by TNF), which reflects a requirement for endogenous TNF α in LPS-treated neutrophils [27]. Accordingly, stimulation of




Figure 1
Acquisition of granulocytic features following differentiation of PLB-985 cells. (A) Cells were differentiated for 5 days in the presence of DMF or DMSO, or left undifferentiated, prior to FACS determination of their size (forward scatter) and granularity (side-scatter). (B) Cells were differentiated for 5 days in the presence of DMF or DMSO, or left undifferentiated, prior to staining with an anti-CD11b Ab (black trace), with an isotype-matched control Ab (grey trace) or without a first Ab (dotted trace) and subsequent incubation with a FITC-conjugated 2nd Ab. CD11b surface expression was then determined by FACS analysis; a minimum of 10,000 cells were processed for each sample. Results are representative of at least five independent experiments. (C) Results obtained as described above were compiled and expressed as mean \pm s.e.m. of at least 5 independent experiments. For comparison, 100% of human neutrophils express CD11b on their surface. (D) Cells were differentiated for 5 days in the presence of DMF or DMSO, or left undifferentiated, prior to stimulation for 6 h at 37°C with fMLP or LPS. Culture supernatants were then collected and analyzed by ELISA. Mean \pm s.e.m. of at least 6 independent experiments.




Figure 2
Cytokine expression profile of neutrophil-like PLB-985 cells. (A) DMSO-differentiated PLB-985 cells (day 5) were cultured in the absence ("ctrl") or presence of either 100 ng/ml LPS or 100 U/ml TNF α for the indicated times. When IP-10 was measured, the cells were instead cultured for 9 h with the above stimuli, in the presence or absence of 100 U/ml IFN γ . Culture supernatants were then harvested and their cytokine content determined in ELISA. Results are representative of at least 4 independent experiments. (B) DMSO-differentiated PLB-985 cells (day 5) were pre-treated with or without MG-132 (15 μ M, 30 min, 37°C), and stimulated in the absence ("ctrl") or presence of either 30 nM fMLP, 100 ng/ml LPS or 100 U/ml TNF α for 30 min at 37°C. Total RNA was extracted and cytokine mRNA expression was analyzed in RPA. Results shown in this figure are representative of at least 4 independent experiments.

granulocytic PLB-985 cells with LPS in the presence of a neutralizing anti-TNF antibody reduced IL-8 secretion by $48 \pm 5\%$ (mean \pm s.d., $n = 2$) at the 12-h time point. More strikingly, the generation of IP-10 by granulocytic PLB-985 cells (Figure 2A) shares the same singular induction characteristics which we originally reported for primary neutrophils [28], i.e. the requirement for a co-stimulation using IFN γ with either LPS or TNF α . When the expression of the corresponding cytokine genes was investigated, LPS and TNF were found to be good inducers in differentiated PLB-985 cells (Figure 2B and data not shown), again in keeping with the induction characteristics of the same cytokine and chemokine genes observed in human neutrophils. Thus, it appears that DMSO-differentiated PLB-985 cells have a resolutely neutrophilic cytokine expression profile.

Transcription factor binding profile of granulocytic PLB-985 cells

The induction of inflammatory cytokines is to a large extent under the control of discrete transcription factors (mainly from the NF- κ B, C/EBP, AP-1, and STAT families), and we indeed observed that the expression of most cytokines investigated in granulocytic PLB-985 cells can be inhibited by MG-132 (Figure 2B and data not shown), a proteasome inhibitor which prevents the proteasome-dependent degradation of I κ B- α and subsequent NF- κ B activation [29,30]. This led us to characterize the binding of the aforementioned transcription factors in this cellular model. As shown in Figure 3A, neither LPS nor TNF α could increase the binding of AP-1 or C/EBP complexes over constitutive levels, be it in granulocytic or undifferentiated PLB-985 cells. By comparison, TNF α potently activated NF- κ B independently of the PLB-985 differentiation stage, and LPS also proved to be a strong inducer in granulocytic PLB-985 cells (Figure 3A, left panel). The inability of LPS to activate NF- κ B in undifferentiated PLB-985 cells (Figure 3A, left panel) is consistent with the lack of detectable CD14 in these cells [23]. The major inducible NF- κ B DNA-binding complex of granulocytic PLB-985 cells was found to be specific, since it was competed out by a 25-fold excess of unlabeled probe in the binding mix, but unaffected by a 25-fold excess of unlabeled oligonucleotide in which the NF- κ B sequence was mutated (Figure 3C, last 2 lanes). To identify the constituents of this specific NF- κ B complex, supershift assays were performed. As shown in Figure 3C, the inducible NF- κ B band was efficiently supershifted by antibodies against p50 and RelA, suggesting that it represents the prototypical p50/RelA homodimer, as in primary human neutrophils [31].

We also investigated inducible STAT binding in response to stimuli which potently induce this response in neutrophils. For this purpose, we used two different oligonucleotide probes. The first is the gamma response region

(GRR) of the CD64 gene, which can detect binding of STAT1-, STAT3- and STAT5-containing complexes [32]. The other probe is the human serum-inducible element (hSIE) found within the *c-fos* promoter, and which binds STAT1- and STAT3-containing complexes with very high affinity [33]. As shown in Figure 3B, IFN γ proved to be a good inducer of STAT binding in undifferentiated PLB-985 cells, and even more so in granulocytic PLB-985 cells. By comparison, G-CSF and GM-CSF weakly induced a hSIE-binding activity (Figure 3B, right panel). In the particular case of GM-CSF, it is noteworthy that this inducible activity was only observed in neutrophil-like PLB-985 cells, consistent with the observation that the GM-CSF receptor is barely expressed in undifferentiated PLB-985 cells and increases considerably during granulocytic differentiation [17]. The identity of the major inducible STAT-containing complex (i.e. that observed in IFN γ -stimulated cells) was finally investigated in supershift assays. Antibodies against STAT1 completely supershifted the hSIE complex, whereas antibodies directed against other STAT isoforms were without effect (Figure 3D and data not shown), suggesting that the inducible complex represents STAT1 homodimers, as we initially reported in primary human neutrophils [34,35].

Together, the transcription factor binding characteristics of granulocytic PLB-985 cells closely match those observed in human neutrophils [31,34,36]. Moreover, supershift analyses showed that the inducible NF- κ B and STAT complexes of granulocytic PLB-985 cells have the same subunit composition as those of primary neutrophils, i.e. p50/RelA heterodimers in the case of the NF- κ B complex, STAT1 homodimers in the case of IFN γ -induced complexes (Figure 3C, 3D).

Transient overexpression of proteins in granulocytic PLB-985 cells

Having ascertained that granulocytic PLB-985 cells represent a suitable cellular model for the study of inflammatory cytokine generation in primary neutrophils, we turned our attention to the introduction of constructs into these cells. Although isolated studies have shown that it is possible to transiently transfect PLB-985 cells using electroporation, transfection efficiency is reportedly modest, and the mortality rates are very high [16,21]. We obtained similar results when we introduced pcDNA3.1 vector containing a full-length E-GFP insert granulocytic PLB-985 by electroporation under a variety of settings. We subsequently used the nucleofection technique, which proved far superior to electroporation in a number of ways. As shown in Figure 4A, granulocytic PLB-985 cells nucleofected with the same E-GFP vector consistently displayed a large increase in fluorescence, relative to cells nucleofected with vector alone. In 8 independent experiments, transfection efficiency was $73 \pm 4\%$ at 6 h postnucleofec-




Figure 3

Transcription factor binding profile of undifferentiated and neutrophil-like PLB-985 cells. (A) Undifferentiated or DMSO-differentiated PLB-985 cells (day 5) were incubated for 15 min at 37°C in the absence ("-") or presence of 100 ng/ml LPS or 100 U/ml TNF α , and nuclear extracts were prepared and analyzed in EMSA (1 μ g/lane) using NF- κ B, C/EBP, or AP-1 oligonucleotide probes. (B) Undifferentiated or DMSO-differentiated PLB-985 cells (day 5) were incubated for 15 min at 37°C in the absence ("-") or presence of 100 U/ml IFN γ , 1 nM GM-CSF, or 1000 U/ml G-CSF, and nuclear extracts were prepared and analyzed in EMSA (1 μ g/lane) using GRR or hSIE oligonucleotide probes. (C) Nuclear extracts from TNF-stimulated granulocytic PLB-985 cells (day 5) were incubated for 30 min in binding buffer in the presence of antibodies to p50, p52, c-Rel, RelA, or an isotype-matched antibody ("IgG"), or in the presence of a 10- or 25-fold molar excess of unlabeled NF- κ B oligonucleotide probe ("cold"), or in the presence of a 25-fold molar excess of unlabeled NF- κ B probe featuring a mutated NF- κ B site ("mut"), prior to the addition of labeled NF- κ B probe and subsequent EMSA analysis. (D) Nuclear extracts from IFN γ -stimulated granulocytic PLB-985 cells (day 5) were incubated for 30 min in binding buffer in the absence ("-") or presence of antibodies to STAT1, STAT3, or an isotype-matched antibody ("IgG"), prior to the addition of labeled hSIE/m67 probe and subsequent EMSA analysis. Experiments shown in this figure are representative of at least three.




Figure 4

Overexpression of E-GFP and β -galactosidase in neutrophil-like PLB-985 cells. (A) DMSO-differentiated PLB-985 (day 5) were nucleofected with a vector encoding E-GFP or empty vector (pcDNA3.1). Cells were harvested 6 h after nucleofection and processed for FACS analysis. A minimum of 10,000 cells was analyzed per sample. Results are representative of 8 independent experiments. (B) DMSO-differentiated PLB-985 (day 5) were nucleofected (squares) or electroporated (triangles) with an E-GFP vector and cultured for the indicated times prior to propidium iodide staining and FACS analysis. Results are expressed as the proportion of GFP-expressing cells in the population that excludes PI staining. Mean \pm s.e.m. of 5 independent experiments. (C) DMSO-differentiated PLB-985 (day 5) were nucleofected (squares) or electroporated (triangles) with an E-GFP vector and cultured for the indicated times prior to determination of cell survival. Mean \pm s.e.m. of 5 independent experiments. (D) DMSO-differentiated PLB-985 (day 5) were nucleofected with a β -galactosidase vector and cultured for 6 h. Cells were then lysed and processed for their enzymatic activity, using 1-O-[2-nitrophenyl]- β -D-galactopyranoside as a substrate. Results are of two independent experiments.

tion, which proved to be the optimal time (Figure 4B). By comparison, electroporation of the same vector into granulocytic PLB-985 cells required much longer to reach an optimal expression efficiency that was lower than the one obtained by nucleofection (Figure 4B). Like electroporation, nucleofection is known to cause irreversible damages to a fraction of the cell population, even under optimized conditions. As shown in Figure 4C, the survival rate of granulocytic PLB-985 cells following nucleofection was $56 \pm 4\%$ at 6 h ($n = 8$) and gradually decreased to about 20% 24 h. By comparison, electroporation of the same vector resulted in much higher initial mortality rates (Figure 4C). Thus, nucleofection proved to be a rapid, reliable, efficient, and reasonably gentle method of overexpressing a protein into granulocytic PLB-985 cells.

Having successfully introduced E-GFP into granulocytic PLB-985 cells, we next determined whether our protocol could be used for the transient overexpression of proteins that feature an enzymatic activity. For this purpose, the cells were nucleofected with a plasmid encoding β -galactosidase. As shown in Figure 4D, a β -galactosidase activity was readily detectable in transfected granulocytic PLB-985 cells. We next attempted to transfect a more physiologically relevant protein. For this purpose, we used a plasmid encoding wild type PKC α . As shown in Figure 5A, granulocytic PLB-985 cells had largely increased PKC α cellular levels 6 h after nucleofection. When we examined the effect of overexpressing PKC α on the cytokine expression profile of PLB-985 cells, however, no significant change was noted relative to mock-transfected cells (data not shown). We also overexpressed a dominant negative I κ B- α mutant (S32,36A) in neutrophil-like PLB-985 cells. Although this did not result in any apparent change in the overall expression of I κ B- α , cells transfected with the dominant negative mutant were rendered resistant to inducible I κ B- α degradation (Figure 5B). Another functional consequence of introducing the S32,36A mutant was a profound inhibition of inflammatory cytokine production by granulocytic PLB-985 cells (Figure 5C), which confirms that these cytokines are under the control of NF- κ B, as observed in primary neutrophils [37].

Transient transfection of granulocytic PLB-985 cells with promoter-reporter constructs

We next investigated whether it would be possible to carry out promoter studies in granulocytic PLB-985 cells. For this purpose, the cells were nucleofected with luciferase constructs under the control of NF- κ B or AP-1 enhancer elements. As shown in Figure 6A, cell stimulation with either LPS or TNF α resulted in a marked induction of NF κ B-driven promoter activity, whereas a no significant induction was observed with the AP-1-driven construct. These results prompted us to investigate the ability of differentiated PLB-985 cells to activate an actual chemokine

promoter. To this end, the cells were nucleofected with a construct containing the IL-8 promoter coupled to a luciferase reporter gene. As shown in Figure 6B, both LPS and TNF stimulation led to a robust activation of the IL-8 promoter. These results agree well with data from primary neutrophils, in which IL-8 gene transcription and NF- κ B (but not AP-1) are potently activated in response to LPS or TNF α [31,36,38,39]. We finally sought to confirm the impact of NF- κ B on IL-8 promoter activation. To this end, granulocytic PLB-985 cells were transfected with luciferase constructs featuring either a wild-type proximal IL-8 promoter, or a variant mutated within the κ B site. As shown in Figure 6C, the mutant IL-8 promoter was mostly resistant to stimulation with TNF α ; similar results were obtained in LPS-stimulated cells (not shown). This is consistent with the data presented in Figure 5C, and is highly reminiscent of the essential role played by NF- κ B in IL-8 gene induction in neutrophils [36].

Role of endogenous reactive oxygen intermediates (ROI) in transcription factor activation and downstream processes in human granulocytes

Endogenously generated ROI have been implicated in the regulation of NF- κ B activation in several cellular models (reviewed in [40,41]). In neutrophils, however, the role of endogenous ROI in transcription factor activation and cytokine generation remains unclear, despite the fact that neutrophils probably produce more ROI than any other cell type. To address this issue, we used X-CGD PLB-985 cells, in which the NADPH oxidase is inactive due to the targeted disruption of a key component, gp91phox [42]. Indeed, Figure 7A shows that whereas granulocytic PLB-985 cells respond to TNF α , LPS, or to more classical NADPH oxidase activators (such as PMA), granulocytic X-CGD cells completely lack this ability, in agreement with previous studies [19,42,43]. We next investigated the ability of LPS and TNF α to activate NF- κ B in DMSO-differentiated X-CGD PLB-985 cells. We additionally examined the ability of IFN γ to induce STAT1 binding to DNA in these cells, since this process can also be affected by endogenous reactive oxygen species in some instances [44,45]. As shown in Figure 7B, NF- κ B was robustly activated by LPS and TNF α , to an extent comparable to that observed in wild-type PLB-985 cells (Figure 3A). Similarly, STAT1 binding to hSIE in response to IFN γ was potently induced, both in X-CGD granulocytic PLB-985 cells (Figure 7B) and in their wild-type counterparts (Figure 3A). Thus, activation of these two key transcription factors does not appear to be influenced by the production of endogenous ROI in human granulocytes. We next investigated whether the transcriptional activity of promoter constructs might nevertheless be affected in DMSO-differentiated X-CGD PLB-985 cells. To this end, the cells were transfected with either pNF κ B-Luc or pIL8-Luc, and stimulated with LPS or TNF α . As shown in Figure 7C, each




Figure 5
Transient overexpression of proteins in neutrophil-like PLB-985 cells. (A) DMSO-differentiated PLB-985 (day 5) were nucleofected with a plasmid encoding human PKC α or with vector alone (lane 1). The cells were cultured for 6 or 12 h after nucleofection, and processed for immunoblot analysis of their PKC α content. Membranes were then reblotted with a p38 MAPK antibody (as a loading control). (B) DMSO-differentiated PLB-985 (day 5) were nucleofected with a plasmid encoding a dominant negative form of human I κ B- α (S32,36A), or with empty vector ("mock"). The cells were cultured for 6 h after nucleofection, and stimulated for 45 min with 100 ng/ml LPS or 100 U/ml TNF α , prior to immunoblot determination of their I κ B- α or actin content. (C) DMSO-differentiated PLB-985 (day 5) were nucleofected with a plasmid encoding human I κ B- α (S32,36A), or with empty vector ("mock"). The cells were cultured for 6 h after nucleofection, and stimulated for another 6 h in the absence ("resting") or presence of 100 ng/ml LPS or 100 U/ml TNF α . Culture supernatants were then harvested and analyzed in ELISA for the indicated cytokines. Results shown in this figure are representative of three independent experiments.

construct was activated to a similar extent in X-CGD granulocytes as in the parental cell line, in agreement with our NF- κ B binding data in wild-type vs X-CGD granulocytic PLB-985 cells. Finally, we examined the release of three

κ B-dependent chemokines (IL-8, Mip-1 α , Mip-1 β) in DMSO-differentiated X-CGD PLB-985 cells stimulated with LPS or TNF α . Expectedly, all three chemokines were released with similar amplitude and kinetics as in wild-




Figure 6

Transient transfection of neutrophil-like PLB-985 cells with promoter-reporter constructs. (A) Granulocytic PLB-985 cells (day 5) were nucleofected with luciferase constructs driven by 5 repeated NF-κB elements ("pNFκB-Luc"), or by 7 repeats of the consensus AP-1 element ("pAPI-Luc"). The cells were then cultured for 6 h at 37°C in the presence or absence of 100 ng/ml LPS or 100 U/ml TNFα, prior to cell lysis and luciferase activity measurement. Mean ± s.e.m. of at least 3 independent experiments. Transfections made using empty vectors yielded values that did not differ significantly from background, whether in resting or stimulated cells (not shown). (B) Granulocytic PLB-985 cells (day 5) were nucleofected with a luciferase construct driven by the IL-8 promoter ("pIL-8-Luc"), and cultured for 6 h at 37°C in the presence or absence of 100 ng/ml LPS or 100 U/ml TNFα, prior to cell lysis and luciferase activity measurement. Mean ± s.e.m. from 8 independent experiments. (C) Granulocytic PLB-985 cells (day 5) were nucleofected with luciferase constructs encoding either either a wild-type ("wt") IL-8 promoter or a variant mutated in the NF-κB site ("mut κB"). The cells were then cultured for 6 h at 37°C in the presence or absence of 100 U/ml TNFα, prior to cell lysis and luciferase activity measurement. Mean ± s.e.m. from 2 independent experiments.




Figure 7

Responsiveness of neutrophil-like X-CGD PLB-985 cells and their parental counterparts. (A) Granulocytic (day 5) X-CGD PLB-985 cells (grey bars) or parental controls (open bars) were incubated for 30 min in the presence or absence of 100 ng/ml LPS, 100 U/ml TNF α , or 25 nM PMA (as a positive control), prior to the determination of superoxide generation by the cytochrome c reduction assay. Mean \pm s.e.m. from at least 3 independent experiments. (B) Granulocytic (day 5) X-CGD PLB-985 cells (right panel) or parental controls (left panel) were incubated for 15 min at 37°C in the absence ("-") or presence of 100 ng/ml LPS, 100 U/ml TNF α , or 100 U/ml IFN γ , and nuclear extracts were prepared and analyzed in EMSA using a NF- κ B or hSIE oligonucleotide probe. This experiment is representative of three. (C) Granulocytic (day 5) X-CGD PLB-985 cells (grey bars) or parental controls (open bars) were nucleofected with luciferase constructs driven by 5 repeated NF- κ B elements ("pNF κ B-Luc"), or by the IL-8 promoter ("pIL-8-Luc"), and cultured for 6 h at 37°C in the presence or absence of 100 ng/ml LPS or 100 U/ml TNF α , prior to cell lysis and luciferase activity measurement. Mean \pm s.e.m. from 3 independent experiments.

type PLB-985 cells (data not shown). Thus, NF-κB activation and downstream responses do not appear to be under the control of endogenous ROI in human granulocytes.

To ascertain whether these results properly reflect the behavior of primary neutrophils, we cultured the latter cells in the presence of the antioxidant, N-acetyl cysteine,

prior to stimulation with LPS or TNFα, and assessment of the IKK cascade activation. As shown in Figure 8, N-acetyl cysteine pre-incubation failed to affect the inducible phosphorylation of IKKα/β and RelA, or the inducible degradation of IκB-α, in human neutrophils. Accordingly, the inducible DNA binding of NF-κB was also unaffected under the same conditions (Figure 8B). Identical results were obtained when neutrophils were pretreated with a different antioxidant, PDTC, as reported before [46,37]. By comparison, N-acetyl cysteine pretreatment abolished superoxide generation from fMLP- or PMA-stimulated neutrophils (data not shown). Together, these results strongly suggest that NF-κB activation and κB-dependent responses (such as inflammatory cytokine generation) are independent of endogenous ROI in human neutrophils.

Discussion

In this study, we characterized a transfectable cellular model whose behavior closely corresponds to that of primary human neutrophils, both phenotypically and functionally. Indeed, PLB-985 cells that had been differentiated along the granulocytic lineage displayed neutrophil-like surface markers, and became responsive to physiological neutrophil stimuli such as fMLP, LPS, and GM-CSF. More importantly, transcription factor activation profile and cytokine expression kinetics of granulocytic PLB-985 cells were found to faithfully match those observed in primary neutrophils. Perhaps most strikingly, the generation of IP-10 by granulocytic PLB-985 cells shares the same singular induction characteristics that we originally reported for primary neutrophils [28], i.e. the requirement for a co-stimulation using IFNγ with either LPS or TNFα. Whereas neutrophils are notoriously refractory to transfection, this limitation was overcome in granulocytic PLB-985 cells, thereby making it possible to study promoter activation, as well as the impact of overexpressed proteins on cytokine production and related events. This allowed us to confirm the central role of NF-κB in the production of cytokines by granulocytes, and to show that κB-dependent responses are independent of endogenous reactive oxygen intermediates in these cells.

From a technical perspective, our data shows that the nucleofection approach (as opposed to electroporation or lipofection) is particularly well suited for the transient overexpression of various proteins into PLB-985 granulocytes. Indeed, the high transfection efficiency and relatively moderate mortality rates allowed us to successfully introduce E-GFP, β-galactosidase, wt PKCα, and dn IκB-α into these cells. As this study was nearing completion, Boulven *et al.* similarly overexpressed two mutant PI 3-kinase subunits in differentiated PLB-985 cells using a nucleofection approach similar to the one described herein [22]. Although their nucleofector settings and nucleofection buffers were different than those used




Figure 8
Effect of N-acetyl cysteine on the IKK/NF-κB pathway in human neutrophils. (A) Neutrophils were pre-treated with culture medium containing 20 mM N-acetyl cysteine ("NAC") or diluent control for 30 min at 37°C, and further incubated in the absence ("ctrl") or presence of 100 ng/ml LPS or 100 U/ml TNFα. Samples were then processed for SDS-PAGE and immunoblot analysis. Note: actin and IκB are shown on the same blot, as the membrane was first immunoblotted for IκB-α, and then re-blotted for actin. (B) Neutrophils were pretreated with N-acetyl cysteine or diluent as described above, and further incubated in the absence or presence of 100 ng/ml LPS or 100 U/ml TNFα. Nuclear extracts were then prepared and analyzed in EMSA using a NF-κB oligonucleotide probe. The experiments depicted are representative of at least two.

herein, they yielded comparable transfection efficiencies (T. Ear, unpublished data), thereby further validating the nucleofection approach. From a more functional perspective, our overexpression of dn I κ B- α prevented its degradation in response to physiological stimuli, thereby severely impairing the generation of several inflammatory cytokines that are known to be under the control of NF- κ B. The fact that dn I κ B- α overexpression yielded a greater than 50% inhibition of IL-8, Mip-1 α and Mip-1 β production is especially relevant, considering that the E-GFP overexpression experiments indicated that about 70% of the cells actually express the transfected material. Thus, dn I κ B- α must have been very effective in inhibiting downstream cellular responses in those cells which overexpressed the mutant protein. At any rate, these results confirm the pivotal role of NF- κ B in the inducible generation of inflammatory cytokines in human granulocytes, in keeping with our previous observations made in primary neutrophils using various pharmacological inhibitors of the NF- κ B pathway [37,47]. Collectively, our overexpression experiments foreshadow exciting new advances in our understanding of neutrophil biology, insofar as they make possible the future dissection of the various signal transduction pathways controlling cytokine production and transcription factor activation in human granulocytes.

The outcome of the experiments in which we introduced luciferase constructs into granulocytic PLB-985 cells also confirmed and extended several observations made in primary neutrophils. In particular, the inducibility of NF- κ B-driven luciferase constructs (but not of AP1-driven constructs) agrees well with the observation that NF- κ B can be activated in neutrophils stimulated with LPS or TNF α [31,34], whereas AP-1 cannot [36,39]. In this regard, it is also noteworthy that the activation of the NF- κ B-driven luciferase construct represents the first direct demonstration of the ability of a transcription factor to deactivate a downstream gene in human granulocytes. A more compelling example is our demonstration that LPS and TNF α promote the transactivation of the IL-8 promoter in granulocytic PLB-985 cells, as the IL-8 gene is known to be under the control of NF- κ B in human neutrophils [37]. Accordingly, we showed that a proximal IL-8 promoter mutated within its κ B site became mostly unresponsive to stimulation. Again, it is worth noting that the activation of the IL-8 promoter in PLB-985 granulocytes confirms our previous observations made in primary neutrophils, which had shown that the IL-8 gene can be transcriptionally activated by the same stimuli, as determined in nuclear run-on analyses [38] and primary transcript PCR [47]. From a more general standpoint, the above considerations establish that the nucleofection of PLB-985 granulocytes described herein paves the way for detailed

promoter studies in human granulocytes – an enterprise which had heretofore remained elusive.

In a variation of the aforementioned promoter studies, we applied the approach described herein to elucidate the long-standing issue of whether endogenously generated ROI contribute to transcription factor activation and cytokine production in granulocytes, as already reported for several other cellular models [40,41]. For this purpose, we used DMSO-differentiated X-CGD PLB-985 cells, in which the ROI-generating NADPH oxidase complex is inactive. Following LPS or TNF stimulation, these X-CGD PLB-985 cells were found to behave in strikingly similar fashion to their wild-type counterparts, be it in terms of promoter activation (using pNF κ B-Luc or pIL8-Luc), transcription factor activation (NF- κ B, STAT) and inflammatory cytokine generation (IL-8, Mip-1 α , Mip-1 β). These results strongly indicate that NF- κ B and STAT activation, as well as downstream responses, are independent of endogenous ROI in human granulocytes. This conclusion is further supported by experiments made in primary neutrophils, in which a powerful antioxidant (N-acetyl cysteine) similarly failed to affect the activation of the IKK/I κ B- α /NF- κ B cascade [46]. This being said, it could be argued that since LPS and TNF are not known as strong NADPH oxidase activators, then it stands to reason that endogenous ROI should play little or no role in cellular processes. In this regard, Selmezy and colleagues reported that TNF secretion in response to opsonized zymosan was nearly abolished in granulocytic (DMF-differentiated) X-CGD cells, compared to parental controls [48]. However, this was attributable to the much lower levels of surface CD16 found in these X-CGD granulocytes [48]. On the opposite, ROI are abundantly produced in phagocytosing neutrophils, and it was reported that NF- κ B activation under these conditions was unchanged in the presence of various oxidant scavengers (exogenous catalase, superoxide dismutase, or methionine), and that conversely, exogenous H₂O₂ failed to activate NF- κ B [49]. In agreement with these results, we also observed that exogenous H₂O₂ (up to 1 mM) does not induce NF- κ B or STAT activation in nonphagocytosing neutrophils (our unpublished data). It is perhaps precisely because neutrophils are such prolific producers of ROI that they are so well protected from their adverse effects. In this regard, the specific activity of catalase was described to be at least 4-fold higher in neutrophils vs all other phagocytes, and neutrophil function was reportedly unaffected by 0.5 mM exogenous H₂O₂ over several hours [50]. Similarly, it was found that among human blood cells, neutrophils uniquely express high levels of methionine-sulfoxide-reductase enzymes [51]. Whatever the case may be, it has now become quite clear that neither endogenous ROI, nor exogenously provided ROI (such as H₂O₂) significantly affect NF- κ B activation and downstream processes (such as

inflammatory cytokine production) in human neutrophils. Studies are now under way to further decipher the intricacies of transcription factor regulation in human granulocytes.

Conclusion

In this report, we characterized a transfectable cellular model whose behavior closely corresponds to that of primary human neutrophils, both phenotypically and functionally, with a particular emphasis on a prominent functional response of these cells, i.e. inflammatory cytokine production, as well as some key underlying processes such as transcription factor activation. Using this model, we confirmed the pivotal role of NF- κ B in the onset of cytokine production, and further showed that NF- κ B activation is independent of endogenous oxygen-derived intermediates. Because such studies were heretofore impossible to carry out in primary human neutrophils, the approach which we describe is likely to enable significant advances in our understanding of various aspects of neutrophil biology.

Methods

Antibodies and reagents

Antibodies raised against NF- κ B/Rel proteins, I κ B- α , C/EBP isoforms, PKC α , and actin were from Santa Cruz Biotechnology (Santa Cruz, CA, USA), and the anti-CD11b was from BD Biosciences (Mississauga, Canada). Ficoll-Paque, T4 polynucleotide kinase and poly (dI-dC) were from Amersham-Pharmacia (Baie d'Urfé, Qc, Canada); radionucleotides were from NEN (Boston, MA, USA). Endotoxin-free (< 2 pg/ml) RPMI 1640 and FCS were from Sigma (St-Louis, MO, USA) and Wisent (St-Bruno, Qc, Canada), respectively. Recombinant human cytokines were from R&D Systems (Minneapolis, MN, USA), and UltraPure LPS (from *E. coli* 0111:B4) was from InvivoGen (San Diego, CA, USA). Acetylated BSA, diisopropyl fluorophosphate (DFP), dimethyl formamide (DMF), dimethyl sulfoxide (DMSO), N-formyl-methionyl-phenylalanine (fMLP), and phenylmethanesulphonyl fluoride (PMSF) were from Sigma-Aldrich (St. Louis, MO, USA). Aprotinin, 4-(2-aminomethyl)benzenesulfonyl fluoride (AEBSF), leupeptin, Nutridoma-SP, and pepstatin A were from Roche (Laval, Qc, Canada). All other reagents were of the highest available grade, and all buffers and solutions were prepared using pyrogen-free clinical grade water.

Plasmids

A plasmid encoding β -galactosidase in the pCMV β vector was from Clontech (Mountain View, CA, USA). Plasmids containing luciferase constructs under the control of 5 repeated NF- κ B elements (pNF κ B-Luc) or 7 repeated AP-1 elements (pAP1-Luc) were from Stratagene (La Jolla, CA, USA). Plasmids containing luciferase constructs encoding

the full IL-8 promoter (-1498 to +44), a proximal IL-8 promoter (-162 to +44), or a version of the latter mutated within its κ B site [52], were a kind gift from Dr. Allan R. Brasier (University of Texas Medical Branch). A construct encoding a dominant negative form of I κ B- α (S32/36A) was obtained from Dr. Christian Jobin (University of North Carolina at Chapel Hill), and a construct encoding wild type PKC α was from Dr. Gilles Dupuis (Université de Sherbrooke); both constructs were subcloned into pcDNA3.1. Similarly, cDNA sequences corresponding to GFP (ex 488 nm, em 507 nm) or β -galactosidase were respectively excised from the commercial vectors, pEGFP-N1 and pCMVb (Clontech), and subcloned into pcDNA3.1. Following amplification in DH-5 α bacteria, all plasmids were purified using Maxiprep kits featuring EndoFree buffers for endotoxin removal (Qiagen).

Cell isolation and culture

Neutrophils were isolated from the peripheral blood of healthy donors as described previously [46]; each blood donor gave informed consent under a protocol that had been duly approved by the ethics committee of our Research center (comité d'éthique humaine du Centre de recherche du CHUS). As determined by Wright staining and nonspecific esterase cytochemistry, the final neutrophil suspensions consistently contained fewer than 0.5% mononuclear cells; neutrophil viability exceeded 98% after up to 4 h in culture, as determined by trypan blue exclusion. The myelomonoblastic PLB-985 cell line was purchased from the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (Braunschweig, Germany). X-CGD PLB-985 cells, which feature a targeted disruption of the gene encoding the essential NAPDPH subunit, gp91phox [42], were a kind gift from Dr. Mary Dinuer (Indiana University, Indianapolis, IN). All cells were cultured at 37°C under a humidified 5% CO₂ atmosphere in RPMI 1640 containing 10% FCS, 100 U/ml penicillin and 100 μ g/ml streptomycin (hereafter referred to as complete RPMI medium). To induce granulocytic differentiation, 1.25% DMSO was added to the culture medium, which was refreshed every second day. Alternatively, culture medium was sometimes supplemented with 0.5% DMF, 1% Nutridoma SP, and 0.5% FCS to induce granulocytic differentiation, as described by Pedruzzi *et al.* [17]. Cytospins from control and DMSO- or DMF-differentiated cells were submitted to Wright staining for counting and morphological characterization.

Cytofluorimetric analyses

Cells were washed twice in PBS and 5×10^5 cells were incubated on ice for 30 min with anti-CD11b or isotype-matched control antibodies (0.25 μ g/ml). After washing 3 times with PBS, the FITC-conjugated 2nd antibodies (0.5 μ g/ml) were added and left to incubate for 30 min in the darkness before washing. Stained cells were analyzed

(minimum of 10,000 cells) on a FACScan instrument (Becton Dickinson) using the CELLQuest software. For GFP expression, differentiated PLB-985 cells were washed twice in PBS before being analyzed by FACScan.

Transient transfections

On the 5th day of granulocytic differentiation, PLB-985 cells were washed twice in pre-warmed PBS, and processed for transfection. When using the nucleofection technique, the cells were resuspended (5×10^6 cells/100 μ l) in prewarmed Human Dendritic Cell Nucleofector Solution (Amaxa Biosystems, Köln, Germany) containing 5 μ g of the plasmid of interest. Cells were incubated for 5 min at room temperature, transferred into a 2 mm-gap electroporation cuvette, and transfected in a Nucleofector instrument (Amaxa Biosystems) using the preconfigured Q-01 or U-15 settings. Nucleofected cells were washed once with pre-warmed RPMI 1640 medium containing 10% FBS, and then cultured in complete RPMI medium. When using the electroporation technique, day 5 granulocytic PLB-985 cells (5×10^6 cells/condition) were washed twice with pre-warmed PBS and resuspended in 400 μ l of electroporation buffer (20 mM HEPES, 137 mM NaCl, 0.7 mM Na_2HPO_4) containing 5 μ g of plasmid. The cells were transferred into a 4 mm-gap electroporation cuvette, and electroporated (270 V, 960 μ F) using a Bio-Rad Gene Pulser instrument. Electroporated cells were washed with pre-warmed RPMI medium containing 10% of FBS, and cultured in a complete RPMI medium. These electroporation conditions represent optimized settings for maximal transfection efficiency and survival rate.

Assays of β -galactosidase activity

Nucleofected cells were cultured for 6 h, washed twice with PBS, and resuspended in a buffer containing 40 mM TrisBase (pH 7.5), 150 mM EDTA, and 150 mM NaCl containing protease inhibitors. Cells were disrupted by three freeze/thaw cycles followed by one cycle of sonication (3 s, maximal power, on ice). Samples were cleared by centrifugation (10 min, 15,000 \times g), and the supernatants were incubated at 37°C using 1-O-[2-nitrophenyl]- β -D-galactopyranoside as a substrate in a sodium phosphate buffer (pH 7.0), according to the supplier's instructions (Promega Technical bulletin #094).

Luciferase assays

Nucleofected cells were cultured for 6 h in the presence or absence of stimuli, washed twice with PBS, and disrupted in Reporter Lysis Buffer following the manufacturer's instructions (Promega Corp., Madison, WI, USA). The lysates were cleared by centrifugation (12 000 g, 10 min), and the resulting supernatants were diluted using Luciferase Assay Reagent (Promega). Luciferase activity was then measured in a Sirius luminometer (Berthold Detection Systems, Pforzheim, Germany).

Denaturing electrophoreses and immunoblots

Differentiated PLB-985 cells were resuspended in ice-cold PBS supplemented with protease inhibitors (10 μ g/ml aprotinin, leupeptin, and pepstatin; 1 mM PMSF; 0.5 mM DFP) and phosphatase inhibitors (10 mM NaF, 1 mM Na_3VO_4 , 10 mM $\text{Na}_4\text{P}_2\text{O}_7$). A small aliquot was taken prior to centrifugation (300 g, 5 min, 4°C) for subsequent protein content determination, and an equal volume boiling sample buffer (2 \times) was added. Samples were briefly vortexed and immediately placed in boiling water for a further 3 min. Samples thus prepared were sonicated to disrupt chromatin, and stored at -20°C prior to analysis. All samples were electrophoresed on denaturing gels prepared according to the method of Laemmli [53]; equal loading was ascertained by adjusting sample volumes based on their respective protein content. Following SDS-PAGE, proteins were transferred onto nitrocellulose membranes, which were stained with Ponceau Red, destained, and then processed for immunoblot analysis, as previously described [31].

Electrophoretic mobility shift assays (EMSA)

Cells were resuspended in ice-cold relaxation buffer (10 mM PIPES pH 7.30, 10 mM NaCl, 3.5 mM MgCl_2 , 0.5 mM EGTA, 0.5 mM EDTA, 1 mM DTT) supplemented the aforementioned protease and phosphatase inhibitors. Nuclear extracts were then prepared using a nitrogen bomb procedure, which we described previously [31,34]. The nuclear extracts were subsequently analyzed in EMSA for NF- κ B, GRR, hSIE, and AP-1 binding as described earlier [31,34,36]. Except for the inclusion of 1 mM MgCl_2 in the binding buffer, the binding conditions used for the C/EBP probe, 5'-tgcagaTTGCGCAATctgca-3', were identical to those used for NF- κ B binding.

Isolation of RNA and Ribonuclease protection assays

Neutrophils were incubated in the presence or absence of stimuli or inhibitors for the desired times, as indicated. Total RNA was extracted following a slightly modified Chomczynski & Sacchi procedure [54], and analyzed by ribonuclease protection assay as previously described [36], using multiprobe templates hCK3 or hCK5 from BD-Pharmingen (Mississauga, Ont, Canada).

ELISA analysis of secreted proteins

Cells were cultured in 12-well plates at 37°C under a 5% CO_2 atmosphere, in the presence or absence of stimuli and/or inhibitors, for the indicated times. Culture supernatants were carefully collected, snap-frozen in liquid nitrogen, and stored at -70°C. Cytokine concentrations were determined in in-house sandwich ELISA assays, using commercially available capture and detection antibody pairs (R&D Systems, BD-PharMingen). Detection limits using these assays varied between 3 and 10 pg/ml.

Measurement of NADPH oxidase activity

Superoxide production by granulocytic PLB-985 cells was measured in whole cells by monitoring the reduction of cytochrome c in a ThermoMax microtiter plate reader (Molecular Devices). Briefly, cells cultured in PBS (supplemented with 5% FCS and 100 μ M cytochrome c) were plated in a 96-well tissue culture-treated plate (2 \times 10⁵/well) and incubated at 37°C for the indicated times in the presence or absence of stimuli, prior to reading absorbance at 550 nm, and calculating the extent of superoxide generation.

Abbreviations

AP-1, activator protein-1; C/EBP, CCAAT enhancer binding protein; GM-CSF, granulocyte-macrophage colony-stimulating factor; LPS, lipopolysaccharide; Mip-1, Macrophage inflammatory protein-1; NF- κ B, nuclear factor- κ B; ROI, reactive oxygen intermediates; STAT, signal transducers and activators of transcription; TNF, tumor necrosis factor

Authors' contributions

TE performed most of the experiments, analyzed the data, and drafted the manuscript. PPMcD conceived and designed the research project, provided the necessary funding, and edited the final manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by grants to PPMcD from the Canadian Arthritis Society, the Canadian Institutes of Health Research (CIHR), and the Canadian Foundation for Innovation. PPMcD is a Scholar of the Fonds de la recherche en santé du Québec (FRSQ). We also wish to thank Ms Emilie Blais-Charron for excellent technical support, and Mr Alexandre Cloutier for his help with some of the gel shift experiments.

References

- Cassatella MA: **Neutrophil-derived proteins: Selling cytokines by the pound.** *Adv Immunol* 1999, **73**:369-509.
- Xing Z, Jordana M, Kirpalani H, Driscoll KE, Schall TJ, Gauldie J: **Cytokine expression by neutrophils and macrophages in vivo: Endotoxin induces tumor necrosis factor- α , macrophage inflammatory protein-2, interleukin-1 β , and interleukin-6 but not RANTES or transforming growth factor- β 1 mRNA expression in acute lung inflammation.** *Am J Respir Cell Mol Biol* 1994, **10**:148-153.
- Ivey CL, Williams FM, Collins PD, Jose PJ, Williams TJ: **Neutrophil chemoattractants generated in two phases during reperfusion of ischemic myocardium in the rabbit. Evidence for a role for C5a and interleukin-8.** *J Clin Invest* 1995, **95**:2720-2728.
- Bennouna S, Bliss SK, Curriel TJ, Denkers EY: **Cross-talk in the innate immune system: Neutrophils instruct recruitment and activation of dendritic cells during microbial infection.** *J Immunol* 2003, **171**(11):6052-6058.
- Bennouna S, Denkers EY: **Microbial antigen triggers rapid mobilization of TNF- α to the surface of mouse neutrophils transforming them into inducers of high-level dendritic cell TNF- α production.** *J Immunol* 2005, **174**(8):4845-4851.
- von Stebut E, Metz M, Milon G, Knop J, Maurer M: **Early macrophage influx to sites of cutaneous granuloma formation is dependent on MIP-1 α/β released from neutrophils recruited by mast cell-derived TNF α .** *Blood* 2003, **101**(1):210-215.
- Wipke BT, Allen PM: **Essential role of neutrophils in the initiation and progression of a murine model of rheumatoid arthritis.** *J Immunol* 2001, **167**(3):1601-1608.
- Jonsson H, Allen P, Peng SL: **Inflammatory arthritis requires Foxo3a to prevent Fas ligand-induced neutrophil apoptosis.** *Nat Med* 2005, **11**(6):666-671.
- Mary JY: **Normal human granulopoiesis revisited. I. Blood data.** *Biomed Pharmacother* 1984, **38**(1):33-43.
- Tucker KA, Lilly MB, Heck L Jr., Rado TA: **Characterization of a new human diploid myeloid leukemia cell line (PLB-985) with granulocytic and monocytic differentiating capacity.** *Blood* 1987, **70**(2):372-378.
- Shen Q, Chada S, Whitney C, Newburger PE: **Regulation of the human cellular glutathione peroxidase gene during in vitro myeloid and monocytic differentiation.** *Blood* 1994, **84**(11):3902-3908.
- Monczak Y, Trudel M, Lamph WW, Miller WH Jr.: **Induction of apoptosis without differentiation by retinoic acid in PLB-985 cells requires the activation of both RAR and RXR.** *Blood* 1997, **90**(9):3345-3355.
- Lowenthal A, Levy R: **Essential requirement of cytosolic phospholipase A2 for activation of the H(+) channel in phagocyte-like cells.** *J Biol Chem* 1999, **274**(31):21603-21608.
- Dana R, Leto TL, Malech HL, Levy R: **Essential requirement of cytosolic phospholipase A2 for activation of the phagocyte NADPH oxidase.** *J Biol Chem* 1998, **273**:441-445.
- Pessach I, Levy R: **The NADPH oxidase diaphorase activity in permeabilized human neutrophils and granulocytic like PLB-985 cells.** *Adv Exp Med Biol* 2000, **479**:107-114.
- Dana RR, Eigsti C, Holmes KL, Leto TL: **A regulatory role for ADP-ribosylation factor 6 (ARF6) in activation of the phagocyte NADPH oxidase.** *J Biol Chem* 2000, **275**(42):32566-32571.
- Pedruzzi E, Fay M, Elbim C, Gaudry M, Gougerot-Pocidalo MA: **Differentiation of PLB-985 myeloid cells into mature neutrophils, shown by degranulation of terminally differentiated compartments in response to N-formyl peptide and priming of superoxide anion production by granulocyte-macrophage colony-stimulating factor.** *Br J Haematol* 2002, **117**(3):719-726.
- Mankelov TJ, Pessach E, Levy R, Henderson LM: **The requirement of cytosolic phospholipase A2 for the PMA activation of proton efflux through the N-terminal 230-amino-acid fragment of gp91phox.** *Biochem J* 2003, **374**(Pt 2):315-319.
- Hiraoka W, Vazquez N, Nieves-Neira W, Chanock SJ, Pommier Y: **Role of oxygen radicals generated by NADPH oxidase in apoptosis induced in human leukemia cells.** *J Clin Invest* 1998, **102**(11):1961-1968.
- Gaudreault E, Thompson C, Stankova J, Rola-Pleszczynski M: **Involvement of BLT1 endocytosis and Yes kinase activation in leukotriene B4-induced neutrophil degranulation.** *J Immunol* 2005, **174**(6):3617-3625.
- Hao LJ, Yang D, Fujii Y, Yamauchi A, Suzuki N, Kikuchi H, Kaneda Y, Nakamura M: **Phorbol ester-potentiated liposomal transfection to monocytic PLB-985 cells.** *J Biochem (Tokyo)* 2000, **128**(6):989-998.
- Boulven I, Levasseur S, Marois S, Pare G, Rollet-Labelle E, Naccache PH: **Class IA phosphatidylinositide 3-kinases, rather than p110 γ , regulate formyl-methionyl-leucyl-phenylalanine-stimulated chemotaxis and superoxide production in differentiated neutrophil-like PLB-985 cells.** *J Immunol* 2006, **176**(12):7621-7627.
- Liberty IF, Raichel L, Hazan-Eitan Z, Pessach I, Hadad N, Schlaeffer F, Levy R: **Cytosolic phospholipase A2 is responsible for prostaglandin E2 and leukotriene B4 formation in phagocyte-like PLB-985 cells: Studies of differentiated cPLA2-deficient PLB-985 cells.** *J Leukoc Biol* 2004, **76**(1):176-184.
- Cassatella MA, Bazzoni F, Calzetti F, Guasparri I, Rossi F, Trinchieri G: **Interferon- γ transcriptionally modulates the expression of the genes for the high affinity IgG-Fc receptor and the 47-kDa cytosolic component of NADPH oxidase in human polymorphonuclear leukocytes.** *J Biol Chem* 1991, **266**(33):22079-22082.
- Kasama T, Strieter RM, Standiford TJ, Burdick MD, Kunkel SL: **Expression and regulation of human neutrophil-derived macrophage inflammatory protein 1 α .** *J Exp Med* 1993, **178**:63-72.

26. Kasama T, Strieter RM, Lukacs NW, Burdick MD, Kunkel SL: **Regulation of neutrophil-derived chemokine expression by IL-10.** *J Immunol* 1994, **152**:3559-3569.
27. Cassatella MA, Meda L, Bonora S, Ceska M, Constantin G: **Interleukin-10 inhibits the release of proinflammatory cytokines from human polymorphonuclear leukocytes. Evidence for an autocrine role of TNF and IL-1 β in mediating the production of IL-8 triggered by lipopolysaccharide.** *J Exp Med* 1993, **178**:2207-2211.
28. Cassatella MA, Gasperini S, Calzetti F, Bertagnin A, Luster AD, McDonald PP: **Regulated production of the interferon- γ -inducible protein-10 (IP-10) chemokine by human neutrophils.** *Eur J Immunol* 1997, **27**:111-115.
29. Palombella VJ, Rando OJ, Goldberg AL, Maniatis T: **The ubiquitin-proteasome pathway is required for processing the NF- κ B precursor protein and the activation of NF- κ B.** *Cell* 1994, **78**(5):773-785.
30. Zavrski I, Krebbel H, Wildemann B, Heider U, Kaiser M, Possinger K, Sezer O: **Proteasome inhibitors abrogate osteoclast differentiation and osteoclast function.** *Biochem Biophys Res Commun* 2005, **333**(1):200-205.
31. McDonald PP, Bald A, Cassatella MA: **Activation of the NF- κ B pathway by inflammatory stimuli in human neutrophils.** *Blood* 1997, **89**:3421-3433.
32. Wehinger J, Gouilleux F, Groner B, Finke J, Mertelmann R, Weber-Nordt RM: **IL-10 induces DNA binding activity of three STAT proteins (Stat1, Stat3, and Stat5) and their distinct combinatorial assembly in the promoters of selected genes.** *FEBS Lett* 1996, **394**(3):365-370.
33. Zhong Z, Wen Z, Darnell JE Jr.: **Stat3: A STAT family member activated by tyrosine phosphorylation in response to epidermal growth factor and interleukin-6.** *Science* 1994, **264**(5155):95-98.
34. McDonald PP, Bovolenta C, Cassatella MA: **Activation of distinct transcription factors in neutrophils by bacterial LPS, interferon- γ , and GM-CSF and the necessity to overcome the action of endogenous proteases.** *Biochemistry* 1998, **37**:13165-13173.
35. Bovolenta C, Gasperini S, McDonald PP, Cassatella MA: **High affinity receptor for IgG (Fc γ RI/CD64) gene and STAT protein binding to the IFN γ response region (GRR) are regulated differentially in human neutrophils and monocytes by IL-10.** *J Immunol* 1998, **160**:911-919.
36. Cloutier A, Ear T, Borissevitch O, Larivee P, McDonald PP: **Inflammatory cytokine expression is independent of the c-Jun N-terminal kinase/AP-1 signaling cascade in human neutrophils.** *J Immunol* 2003, **171**(7):3751-3761.
37. Cloutier A, Ear T, Blais-Charron E, Dubois CM, McDonald PP: **Differential involvement of NF- κ B and MAP kinase pathways in the generation of inflammatory cytokines by human neutrophils.** *J Leukoc Biol* 2007:567-577.
38. Cassatella MA, Gasperini S, Calzetti F, McDonald PP, Trinchieri G: **Lipopolysaccharide-induced interleukin-8 gene expression in human granulocytes: Transcriptional inhibition by interferon- γ .** *Biochem J* 1995, **310**:751-735.
39. Tamassia N, Le Moigne V, Calzetti F, Donini M, Gasperini S, Ear T, Cloutier A, Martinez FO, Fabbri M, Locati M, Mantovani A, McDonald PP, Cassatella MA: **The MyD88-independent pathway is not mobilized in human neutrophils stimulated via TLR4.** *J Immunol* 2007, **178**(11):7344-7356.
40. Janssen-Heininger YM, Poynter ME, Baeuerle PA: **Recent advances towards understanding redox mechanisms in the activation of nuclear factor kappaB.** *Free Radic Biol Med* 2000, **28**(9):1317-1327.
41. Arrigo AP: **Gene expression and the thiol redox state.** *Free Radic Biol Med* 1999, **27**(9-10):936-944.
42. Zhen L, King AA, Xiao Y, Chanock SJ, Orkin SH, Dinanuer MC: **Gene targeting of X chromosome-linked chronic granulomatous disease locus in a human myeloid leukemia cell line and rescue by expression of recombinant gp91 phox.** *Proc Natl Acad Sci U S A* 1993, **90**(21):9832-9836.
43. Yu L, Cross AR, Zhen L, Dinanuer MC: **Functional analysis of NADPH oxidase in granulocytic cells expressing a Δ 488-497 gp91 (phox) deletion mutant.** *Blood* 1999, **94**(7):2497-2504.
44. Kaur N, Lu B, Monroe RK, Ward SM, Halvorsen SW: **Inducers of oxidative stress block ciliary neurotrophic factor activation of Jak/STAT signaling in neurons.** *J Neurochem* 2005, **92**(6):1521-1530.
45. Pawate S, Shen Q, Fan F, Bhat NR: **Redox regulation of glial inflammatory response to lipopolysaccharide and interferon- γ .** *J Neurosci Res* 2004, **77**(4):540-551.
46. Ear T, Cloutier A, McDonald PP: **Constitutive nuclear expression of the I κ B kinase complex and its activation in human neutrophils.** *J Immunol* 2005, **175**(3):1834-1842.
47. Tamassia N, Calzetti F, Ear T, Cloutier A, Gasperini S, Bazzoni F, McDonald PP, Cassatella MA: **Molecular mechanisms underlying the synergistic induction of CXCL10 by LPS and IFN γ in human neutrophils.** *Eur J Immunol* 2007, **37**:2627-2634.
48. Selmecky Z, Szelenyi J, Nemet K, Vizi ES: **The inducibility of TNF α production is different in the granulocytic and monocytic differentiated forms of wild type and CGD-mutant PLB-985 cells.** *Immunol Cell Biol* 2003, **81**(6):472-479.
49. Vollebregt M, Hampton MB, Winterbourn CC: **Activation of NF- κ B in human neutrophils during phagocytosis of bacteria independently of oxidant generation.** *FEBS Lett* 1998, **432**(1-2):40-44.
50. Pietarinen-Runtti P, Lakari E, Raivio KO, Kinnula VL: **Expression of antioxidant enzymes in human inflammatory cells.** *Am J Physiol Cell Physiol* 2000, **278**(1):C118-25.
51. Achilli C, Ciana A, Rossi A, Balduini C, Minetti G: **Neutrophil granulocytes uniquely express, among human blood cells, high levels of methionine-sulfoxide-reductase enzymes.** *J Leukoc Biol* 2008, **83**(1):181-189.
52. Vlahopoulos S, Boldogh I, Casola A, Brasier AR: **Nuclear factor- κ B-dependent induction of interleukin-8 gene expression by tumor necrosis factor- α : Evidence for an antioxidant sensitive activating pathway distinct from nuclear translocation.** *Blood* 1999, **94**(6):1878-1889.
53. Laemmli UK: **Cleavage of structural proteins during the assembly of the head of bacteriophage T4.** *Nature* 1970, **227**:680-685.
54. Chomczynski P, Sacchi N: **Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction.** *Anal Biochem* 1987, **162**:156-159.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

