



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contents lists available at ScienceDirect

European Journal of Internal Medicine

journal homepage: www.elsevier.com/locate/ejim

Letter to the Editor

Standardized PaO₂/FiO₂ ratio in COVID-19: Added value or risky assumption. Author's reply

ARTICLE INFO

Keywords

Standardized PaO₂

COVID-19

Acute respiratory failure

Hypoxemia

Dear Editor,

We have read with great interest the comment of Luciano Gattinoni and coworkers who carefully reviewed our paper “Standardized PaO₂/FiO₂ ratio in COVID-19: Added value or risky assumptions” recently published in the European Journal of Internal Medicine [1, 2]. We agree with most of their comments on the potential limitations of the standardized PaO₂/FiO₂ approach. There are, however, a few points that we would like to further discuss and clarify.

First, we do not entirely agree on the need to systematically correct the alveolar partial pressure of oxygen (PAO₂) using the respiratory quotient (RQ). This is usually measured at the mouth (respiratory exchange ratio, R, V'CO₂/V'O₂) and therefore it is extremely difficult, if not impossible, to measure during helmet CPAP or high oxygen flow supplementation (HOF) and, because of the contribution/changes in lung gas stores, R does not reflect changes in RQ. Thus, the correction for R is impractical not reflecting the metabolic state (RQ), and as pointed out in a previous work by Gattinoni and his group “the impact of the PaCO₂/R ratio on alveolar PO₂ (PAO₂) is less dramatic unless extracorporeal CO₂ removal is in use” [3].

Second, as pointed out by Gattinoni and coworkers, the advantage of standardized PaO₂/FiO₂ over PaO₂/FiO₂ is that standardized PaO₂/FiO₂, (i.e. PaO₂ corrected for the PaCO₂), takes in account the contribution of the respiratory effort sustained by the patient which has been demonstrated to be a predictor of poor outcome in acute respiratory failure. We would like to add that it also reflects the need to increase PAO₂ to maintain an adequate arterial oxygenation (i.e. PaO₂) at the cost of a left shift in arterial Hb-O₂ dissociation curve due to respiratory alkalosis. This is a well-known compensation mechanism that occurs for example at high altitude where the need to preserve arterial oxygenation (i.e., increased Hb-O₂ affinity at the lung due to increased pH) prevails on the diffusion of oxygen, from peripheral arterial capillaries at tissue levels. Obviously, other factors, such as body temperature, Hb levels and cardiac output should be taken in account to have a clear picture of the O₂ transport efficiency.

Third, we do agree that standardized PaO₂/FiO₂ ratio should be used with caution when tracking patients over time, because of the influence

of dead space of shunt fractions that may change along the illness. This, in fact, was not the aim of our study in which we tested the superiority of the standardized PaO₂/FiO₂ ratio versus the PaO₂/FiO₂ ratio in predicting the outcome failure and mortality measured at the time of admission to the Pulmonology unit. It should be pointed out, however, that also the use of the “classical” PaO₂/FiO₂ ratio has limitations not only as a single observation but also when it is used for tracking patients over time. In particular, the PaO₂/FiO₂ ratio is significantly influenced by the fraction of inspired oxygen (FiO₂). Of notice when the patient is breathing FiO₂≤0.6 the possible contribution of V'/Q' mismatch and diffusion limitation from shunt cannot be ruled out. By contrast when breathing FiO₂>0.6 the contribution of shunt mechanisms becomes clearer [4].

In conclusion, although we do agree with Gattinoni and coworkers in their analysis on the need to consider all the variables that may influence gas exchange impairment in severe ARF, we believe that the standard PaO₂/FiO₂ ratio should be utilized in the prognostic evaluation of patients with COVID-19 acute respiratory failure.

Declaration of Competing Interest

The authors declare they have no conflict of interest.

Acknowledgements

We thank all the other authors of the work “Standardizing PaO₂ for PaCO₂ in P/F ratio predicts in-hospital mortality in acute respiratory failure due to Covid-19: A pilot prospective study”, Letizia D'Antoni, Paolo Carbonara, Federico Daniele, Roberto Dongilli, Roberto Flore, Angela Maria Grazia Pacilli, Lara Pisani, Corina Tomsa, Maria Laura Vega, and Vito Marco Ranieri for their contribution and their expertise throughout the study.

Author's contribution: IP draft preparation. IP, SN and PP co-write the manuscript. SN and PP supervised the final revision.

<https://doi.org/10.1016/j.ejim.2021.10.015>

Received 30 September 2021; Accepted 12 October 2021

Available online 23 October 2021

0953-6205/© 2021 European Federation of Internal Medicine. Published by Elsevier B.V. All rights reserved.

References

- [1] Gattinoni L, Busana M, Camporota L. Standardised PaO₂/FiO₂ ratio in COVID-19: Added value or risky assumptions? *Eur J Intern Med* 2021;92:31–3. <https://doi.org/10.1016/j.ejim.2021.09.004>.
- [2] Prediletto I, D'Antoni L, Carbonara P, Daniele F, Dongilli R, Flore R, Pacilli AMG, Pisani L, Tomsa C, Vega ML, Ranieri VM, Nava S, Palange P. Standardizing PaO₂ for PaCO₂ in P/F ratio predicts in-hospital mortality in acute respiratory failure due to COVID-19: A pilot prospective study. *Eur J Intern Med* 2021;92:48–54. <https://doi.org/10.1016/j.ejim.2021.06.002>.
- [3] Gattinoni L, Vassalli F, Romitti F. Benefits and risks of the P/F approach. *Intensive Care Med* 2018;44(12):2245–7. <https://doi.org/10.1007/s00134-018-5413-4>.
- [4] Aboab J, Louis B, Jonson B, Brochard L. Relation between PaO₂/F_IO₂ ratio and F_IO₂: a mathematical description. *Applied Physiology in Intensive Care Medicine*. Berlin, Heidelberg: Springer; 2006. https://doi.org/10.1007/3-540-37363-2_11.

Irene Prediletto^{a,b}, Stefano Nava^{a,b,*}, Paolo Palange^{c,1}

^a *Alma Mater Studiorum University of Bologna, Department of Clinical, Integrated and Experimental Medicine (DIMES), Bologna, Italy*

^b *IRCCS Azienda Ospedaliero Universitaria di Bologna, University Hospital Sant'Orsola-Malpighi - Respiratory and Critical Care Unit, Bologna, Italy*

^c *Department of Public Health and Infectious Disease, Sapienza University of Rome – Italy. Pulmonology, Respiratory and Critical Care Unit, Policlinico Umberto I Hospital, Rome, Italy.*

* Corresponding Author at: IRCCS Azienda Ospedaliero Universitaria di Bologna, Sant'Orsola-Malpighi University Hospital. Alma Mater Studiorum University of Bologna, Department of Clinical, Integrated and Experimental Medicine, Bologna, Italy. Address: Via Massarenti 9, 40138 Bologna, Italy.

¹ on behalf of the authors of “Standardizing PaO₂ for PaCO₂ in P/F ratio predicts in-hospital mortality in acute respiratory failure due to Covid-19: a pilot prospective study”.