



Leptin Signaling in the Carotid Body Regulates a Hypoxic Ventilatory Response Through Altering TASK Channel Expression

Fang Yuan^{1,2}, Hanqiao Wang³, Jiaqi Feng¹, Ziqian Wei¹, Hongxiao Yu¹, Xiangjian Zhang^{2,4}, Yi Zhang^{1,2*} and Sheng Wang^{1,2*}

¹ Department of Physiology, Hebei Medical University, Shijiazhuang, China, ² Hebei Key Laboratory of Vascular Homeostasis and Hebei Collaborative Innovation Center for Cardio-Cerebrovascular Disease, Shijiazhuang, China, ³ Department of Sleep, Third Hospital of Hebei Medical University, Shijiazhuang, China, ⁴ Department of Neurology, Second Hospital of Hebei Medical University, Shijiazhuang, China

OPEN ACCESS

Edited by:

Rodrigo Iturriaga,
Pontificia Universidad Católica de
Chile, Chile

Reviewed by:

Andrea Porzionato,
Università degli Studi di Padova, Italy
Silvia V. Conde,
Faculdade de Ciências Médicas,
Universidade Nova de Lisboa,
Portugal
Ana Obeso,
University of Valladolid, Spain

*Correspondence:

Yi Zhang
yizhanghebmu@163.com
Sheng Wang
wangsheng@hebmu.edu.cn

Specialty section:

This article was submitted to
Integrative Physiology,
a section of the journal
Frontiers in Physiology

Received: 17 January 2018

Accepted: 06 March 2018

Published: 27 March 2018

Citation:

Yuan F, Wang H, Feng J, Wei Z, Yu H,
Zhang X, Zhang Y and Wang S (2018)
Leptin Signaling in the Carotid Body
Regulates a Hypoxic Ventilatory
Response Through Altering TASK
Channel Expression.
Front. Physiol. 9:249.
doi: 10.3389/fphys.2018.00249

Leptin is an adipose-derived hormone that plays an important role in the regulation of breathing. It has been demonstrated that obesity-related hypoventilation or apnea is closely associated with leptin signaling pathways. Perturbations of leptin signaling probably contribute to the reduced sensitivity of respiratory chemoreceptors to hypoxia/hypercapnia. However, the underlying mechanism remains incompletely understood. The present study is to test the hypothesis that leptin signaling contributes to modulating a hypoxic ventilatory response. The respiratory function was assessed in conscious obese Zucker rats or lean littermates treated with an injection of leptin. During exposure to hypoxia, the change in minute ventilation was lower in obese Zucker rats than chow-fed lean littermates or high fat diet-fed littermates. Such a change was abolished in all groups after carotid body denervation. In addition, the expression of phosphorylated signal transducers and activators of transcription 3 (pSTAT3), as well as putative O₂-sensitive K⁺ channels including TASK-1, TASK-3 and TASK-2 in the carotid body, was significantly reduced in obese Zucker rats compared with the other two phenotype littermates. Chronic administration of leptin in chow-fed lean Zucker rats failed to alter basal ventilation but vigorously increased tidal volume, respiratory frequency, and therefore minute volume during exposure to hypoxia. Likewise, carotid body denervation abolished such an effect. In addition, systemic leptin elicited enhanced expression of pSTAT3 and TASK channels. In conclusion, these data demonstrate that leptin signaling facilitates hypoxic ventilatory responses probably through upregulation of pSTAT3 and TASK channels in the carotid body. These findings may help to better understand the pathogenic mechanism of obesity-related hypoventilation or apnea.

Keywords: leptin, hypoventilation, hypoxic ventilatory response, carotid body, TASK channels, STAT3

INTRODUCTION

Leptin, a peptide hormone secreted mainly by adipocytes, regulates multiple physiological functions including metabolism, cardiovascular activity, and breathing (Grill et al., 2002; Bassi et al., 2015). Leptin's role in controlling breathing has been implicated in recent studies, with its participation in sleep-related breathing disorders including obesity hypoventilation syndrome

(OHS) and obstructive sleep apnea (Malhotra and White, 2002). In animal models, leptin deficient mice exhibit impaired ventilatory responses to CO₂ which can be rescued by leptin replacement therapy, in favor of facilitation of breathing by leptin (O'Donnell et al., 2000). Leptin receptors (ob-Rs) are composed of six isoforms termed from ob-Ra to -Rf, with the long form of ob-Rb mediating the majority of leptin's intracellular signal transduction (Tartaglia, 1997). Although the role of leptin signaling pathways in mediating various physiological actions has been investigated intensively, the molecular mechanism underlying its action on breathing remains incompletely understood.

The peripheral respiratory chemoreflex serves as a homeostatic regulatory mechanism by which enough oxygen must be supplied to the organism when challenged by hypoxia through altering respiratory amplitude and frequency. The carotid body (CB) chemoreceptors, located near the fork of the carotid artery, are activated shortly after exposure to hypoxia and then send information to the nucleus tractus solitarius and high integrative centers, with the outcome of adaptive ventilatory responses (Gonzalez-Martin et al., 2011; Ciriello and Caverson, 2014). Accumulated evidence indicates the presence of ob-Rb in the CB cells (Porzionato et al., 2011; Messenger et al., 2013), and that leptin signaling contributes to CB-mediated ventilatory responses (Olea et al., 2015; Ribeiro et al., 2017). However, it remains controversial whether the CB mediates the acute effect of leptin on hypoxic ventilatory response (HVR) because leptin's role may involve the change in gene expression and protein synthesis, requiring hours even days for full effects (Hall et al., 2010). We thereby predicted that the stimulatory effect of leptin on HVR may require chronic activation of CBs, but such a confirmation is yet to be put forward.

In the CBs, leptin signaling pathways involve the downstream signaling proteins of ob-R signal transducers and activators of transcription 3 (STAT3), suppressor of cytokine signaling 3 (SOCS3), and extracellular-signal-regulated kinase 1/2 (ERK1/2) (Messenger et al., 2013; Moreau et al., 2015), reminiscent of modulatory effects of these molecules on carotid chemoreceptor sensitivity. Emerging evidence has shown that the chemosensitivity of glomus cells in CBs requires two-pore K⁺ channels including TWIK-related acid-sensitive K (TASK)-1 channels and acid-sensitive ion channels (Trapp et al., 2008; Tan et al., 2010). However, very little is known concerning whether the activation of the ob-R and downstream signaling molecules modulates the sensitization of CB chemoreceptors via affecting these ion channels.

We sought to address herein whether the leptin signaling pathway in the CB contributes to regulating HVR and the possible mechanism involved. We utilized whole body plethysmography (WBP) to assess HVR in obese Zucker rats (ob-R deficiency) or in lean littermate controls treated with injections of leptin. The main findings suggest that chronic application of leptin contributes to facilitation of HVRs probably through upregulation of phosphorylated STAT3 (pSTAT3) and TASK channel expression.

MATERIALS AND METHODS

Animals

The experiments were carried out in 12~20-week-old male obese Zucker rats (OZR) and lean littermates (LZR) obtained from the Charles River Laboratories (USA). Animals, synchronized for a 12:12 h light-dark cycle (lights on at 8 am, lights off at 8 pm), were housed individually and allowed to move freely in standard plastic cages in a climate-controlled room (22 ± 1°C). Food and water were provided *ad libitum* for LZRs and OZRs. In some cases, a group of LZRs were placed on a high-fat diet (HFD, 45% kcal/g fat, Research Diets D12451) for 8 weeks and used as a simple obesity control (LHZR). The LHZR and OZR groups were weight-matched to determine the effect of simple obesity-induced mechanical resistance on ventilation. Body weight was measured once a week ($n = 20$ for each phenotype). All experiments were performed in accordance to ethical guidelines of the Animal Protection Association and were approved by Animal Care and Ethical Committee of Hebei Medical University. When the animal experiments were completed, an overdose of intraperitoneal injection of sodium pentobarbital (> 200 mg/kg) was carried out for euthanasia.

Breathing Measurement

Breathing was studied by WBP in conscious, freely moving rats (EMKA Technologies, France) as described previously (Kumar et al., 2015; Fu et al., 2017). In brief, rats were placed in the WBP chamber on the day before the testing protocol (2 h acclimation period). For acute hypoxia, rats were exposed to 10% O₂ (balance N₂) for up to ~7 min by a gas mixture devices (1,500 ml/min, GSM-3, CWE, USA). Ventilatory flow signals were recorded, amplified, digitized and analyzed using IOX 2.7 (EMKA Technologies) to determine breathing parameters over sequential 20 s epochs (~50 breaths) during periods of behavioral quiescence and regular breathing. Minute volume (V_E; ml/min/g) was calculated as the product of the respiratory frequency (f_R, breaths/min) and tidal volume (V_T; ml/kg), normalized to rat body weight (g). To further confirm the CB-mediated effect of leptin, breathing parameters were also measured in rats with carotid body denervation (CBD). The carotid sinus nerves were sectioned as depicted before (Kumar et al., 2015). Shortly, anesthesia was induced with 4% halothane in 100% O₂ and maintained by reducing the inspired halothane concentration to 1.5~1.8%. The depth of anesthesia was assessed by an absence of the corneal and hindpaw withdrawal reflex. Body temperature of all mice was maintained at 37°C using a temperature-controlled heating pad. To prevent any functional regeneration of chemosensory fibers, the carotid sinus nerves were removed completely from the cranial pole of the CB until reaching the branch to the glossopharyngeal nerve. The wound was carefully sutured and disinfected with 10% of povidone iodine. Conscious chemodenervated rats were exposed to ventilatory challenge 5–7 days after recovery. No significant weight loss was observed. All the three groups of rats (LZR, LHZR and OZR) were submitted to surgery.

To examine whether hypoventilation resulted in retention of CO₂ in obese rats, arterial blood gas was measured using an

OPTI-CCA blood gas analyzer (OPTI Medical Systems, USA) at a steady state in halothane-anesthetized, paralyzed rats. General anesthesia was induced with 4% halothane in room air as depicted above. Arterial blood (200 μ l per sample) was drawn from the femoral artery in the three animal groups. Arterial blood measurements of interest included partial pressure of arterial O₂ (P_aO₂), partial pressure of CO₂ (P_aCO₂) and pH.

Plasma Leptin Levels and Hypodermic Leptin Injections

Measurements of plasma levels of leptin were performed at room air (21% O₂) in anesthetized rats. After general anesthesia as described above, whole blood samples were taken through a cardiac puncture. Blood samples were drawn into collection tubes containing the anticoagulant EDTA (Sigma-Aldrich, USA) and kept on ice. After centrifugation, the plasma was stored at -80°C for leptin analysis by ELISA kit (#ab100773, Abcam, USA), an *in vitro* enzyme-linked immunosorbent assay for the quantitative measurement as previously described (Panetta et al., 2017). The assay was read using a power wave XS2 plate reader (Biotek Instruments, USA).

To confirm whether the chronic activation of leptin signaling pathways played a part in the HVR, subcutaneous injections of leptin (60 μ g/kg) or equal volume of vehicle (saline) were carried out once daily for 7 days in CBI LZR ($n = 8$ for each group), and breathing parameters were measured after 7 day injections during exposure to room air or hypoxia. To further confirm the CB's role, subcutaneous injections of leptin or saline were performed 7 days after the carotid sinus nerves were sectioned in each group ($n = 8$ for both).

Carotid Body Protein Extracts and Western Blot Analysis

Rats were deeply anesthetized by isoflurane (2–3%) inhalation and then decapitated. The carotid bifurcation was exposed and both CBs were removed and cleaned. The CB ($n = 8$) were pooled from 4 rats in each group and homogenized in 100 μ l of RIPA buffer solution (150 mM NaCl, 1 mM EDTA, 1% Triton-X 100, 50 mM Tris-HCl at pH of 7.5) with a protease inhibitor cocktail (Roche Diagnostics, Canada). The homogenate was centrifuged at 4°C for 20 min at 13200 rpm. The resultant supernatant was retained as the protein preparation. Equal concentrations of extracted proteins normalized by colorimetric BCA Protein Assay (Pierce Corp., USA). After denaturation, the protein (~ 30 μ g) in each lane was fractionated in 10% polyacrylamide gel and then transferred onto apolyvinylidene fluoride membrane. The membranes were blocked with bovine serum albumin and incubated at 4°C overnight with primary antibodies anti-ob-R (1:2000, #ab5593, Abcam, USA), anti-TASK-1 (1:200, #APC024, Alomone labs, Israel), anti-TASK-2 (1:200, #APC037, Alomone labs, Israel), anti-TASK-3 (1:200, #APC044, Alomone labs, Israel), anti-pSTAT3 (1:2000, #9145, Cell Signaling Technology, USA), anti-STAT3 (1:1000, #9139, Cell Signaling Technology, USA), and anti- β -actin (1:3,000, #T0022, Affinity Biosciences, USA). The membranes were then incubated with corresponding secondary antibodies

for 1 h at room temperature. The reaction was visualized using the enhanced chemiluminescence (ECL) method, and the bands were analyzed by ImageJ software (NIH, USA). The protein contents were normalized to β -actin. See Supplementary Material for original gel images.

Data Acquisition and Processing

Data are expressed as mean \pm SEM. Unless indicated otherwise, two-tailed unpaired *t*-test, one-way ANOVA with Dunnett's or Tukey's *post-hoc* test and two-way ANOVA with Bonferroni *post-hoc* test were used to compare significant difference between different groups. Differences within or between groups with *P*-values of <0.05 were considered significant.

RESULTS

Reduced Basal Ventilation in OZR

Adult OZR exhibit many abnormal physiological attributes due to the deficiency of the ob-R, representing a good animal model to study the obesity-related hypoventilation as observed in humans. We thus addressed leptin's role using this phenotype and the littermate control rat. First, we measured the body weight of three groups of rats ($n = 20$ for each group). The averaged body weight was larger in LHZRs (492 ± 30 g) and OZR (512 ± 49 g) than LZR (382 ± 14 g, $P < 0.01$ vs. LHZR or OZR). In addition, weight gain was accompanied by higher levels of blood plasma leptin level in OZR (15.28 ± 0.55 μ g/L) in relative to LHZRs (7.45 ± 0.50 μ g/L, $P < 0.01$ vs. OZR) or LZR (7.34 ± 0.38 μ g/L, $P < 0.01$ vs. OZR; $P > 0.05$, vs. LHZR).

Baseline breathing parameters were measured in the three groups of animals while breathing room air (21% O₂). Compared with the LZR, V_E and V_T were considerably lower in OZR and LHZRs ($P < 0.01$ for both, vs. LZR, **Figures 1A,C**), whereas the OZR has a faster f_R than the other two groups ($P < 0.01$ for both). Of interest, although no remarkable difference in basal V_E was observed between OZR and LHZRs, V_T and f_R were comparable ($P < 0.01$ for both, **Figure 1B**), indicative of a different breathing pattern. To evaluate whether hypoventilation resulted in hypercapnia or respiratory acidosis in obese animals, the arterial blood gas was measured at a steady-state. Apparently, hypercapnia, acidosis and normal P_aO₂ were observed in LHZRs and OZR (**Table 1**). Therefore, both OZR and LHZRs exhibit basal hypoventilation, overweight and hypercapnia, with the exception of hyperleptinemia in OZR but not LHZRs.

Impairment of HVR in OZR

To address whether the ob-R deficiency yielded a diminished HVR, hypoxia was achieved while inhaling 10% O₂ to activate peripheral respiratory chemoreflex. When acutely challenged with hypoxia, all three phenotypes displayed robust increases in f_R and V_E except for V_T, with the smallest change in the HVR in OZR (**Figures 1A–C**). In addition, the hypoxia-stimulated increment of V_E was far less in OZR compared to the other two groups ($P < 0.01$, **Figure 1E**). Interestingly, LZR and LHZRs displayed a similar change in V_E during hypoxia




FIGURE 1 | The OZR exhibits impaired HVR. **(A–C)** Effect of hypoxia on V_T , f_R , and V_E in the three groups of rats. **(D)** The stimulatory effect of hypoxia on V_E in three groups of CBD rats. **(E)** Changes in V_E during exposure to 10% O_2 between CBI and CBD rats. $n = 20$ for each group, $**P < 0.01$ as drawn by two-way ANOVA with Bonferroni *post-hoc* test. V_T , tidal volume; f_R , respiratory frequency; V_E , minute ventilation; CBI, carotid body innervation; CBD, carotid body denervation.

TABLE 1 | Blood gas analysis during exposure to room air.

Group	P_aCO_2 (mmHg)	P_aO_2 (mmHg)	pH
LZR	37.9 ± 1.8	102.5 ± 5.9	7.32 ± 0.02
LHZR	$51.6 \pm 1.0^{**}$	93.8 ± 2.7	$7.24 \pm 0.01^*$
OZR	$50.9 \pm 0.6^{**}$	$87.6 \pm 2.8^*$	$7.22 \pm 0.01^*$

LZR, lean Zucker rats; LHZR, HFD-fed LZR; OZR, obese Zucker rats; $n = 12$ for each group; $*P < 0.05$, $**P < 0.01$ by one-way ANOVA with Tukey's *post-hoc* test, vs. LZRs.

($P > 0.05$, **Figure 1E**). In spite of similar degree of body weight, OZRs exhibited far more severe hypoventilation in response to hypoxia compared to LHZRs ($P < 0.01$, **Figures 1A–C**). However, exposure to hypoxia caused no significant difference in increments of V_E in all three groups of rats after sectioning carotid sinus nerves ($P > 0.05$, **Figures 1D,E**), in favor of the involvement of CBs in such an effect. Collectively, the ob-R

deficiency (OZR), instead of simple obesity (LHZR), plays a predominant role in the impaired HVR.

Downregulation of pSTAT3 and TASK Channels in OZR CBs

The pSTAT3/STAT3 signaling has been implicated in mediating major effects of the ob-R. To determine the expression level of ob-R and STAT3, the quantitative analysis was made using Western blot in the three groups ($n = 4$ for each group). Compared with LZRs and LHZRs, pSTAT3/STAT3 (**Figures 2C,D**) were remarkably downregulated in the CBs of OZRs ($P < 0.01$), reliably correlating pSTAT3/STAT3 expression with the ob-R deficiency (**Figures 2A,B**). Several lines of evidence demonstrated that the chemosensitivity is associated with TASK-1 and TASK-3 in CB glomus cells and TASK-2 in retrotrapezoid nucleus neurons (Trapp et al., 2008; Gestreau et al., 2010; Wang et al., 2013). The reduced sensitivity of CBs to hypoxia in OZRs




FIGURE 2 | Downregulation of pSTAT3/STAT3 and TASK-1,-2,-3 channels in CBs. Gel images and bar charts showing quantitative analysis of expression of ob-Rb (A,B), pSTAT3/STAT3 (C,D), and TASK-1,-2,-3 (E-H) in the CB ($n = 8$) from each group ($n = 4$). * $P < 0.05$, ** $P < 0.01$ by one-way ANOVA with Dunnett's *post-hoc* test. ob-Rb, leptin receptor type b; STAT3, signal transducers and activators of transcription 3; pSTAT3, phosphorylated STAT3.

was probably associated with these ion channels. Evidently, the expression level of TASK-1, TASK-2, TASK-3 in OZR CBs was lower in relative to the other two groups ($P < 0.05 \sim 0.01$, Figures 2E-H). However, no statistical significance in these

channel expression was found between LZRs and LHZRs ($P > 0.05$ for all, Figures 2E-H). Hence, the ob-R deficiency contributes to reduced expression of pSTAT3 and TASK channels.

Facilitation of HVR by Chronic Application of Leptin

To examine the effect of activation of ob-Rs on the HVR, subcutaneous injections of leptin (60 $\mu\text{g}/\text{kg}$) or equal volume of normal saline were carried out once daily for 7 days in LZRs ($n = 8$ for each group), and breathing parameters were measured at different time points separated by 7 days. As shown in **Table 2**, compared with the vehicle control ($8.2 \pm 0.4 \mu\text{g}/\text{L}$), the plasma levels of leptin was raised to $13.1 \pm 0.6 \mu\text{g}/\text{L}$ ($P < 0.01$) over 7-day treatment and restored to control level after 1 week. Chronic administration of leptin for 7 days produced no significant change in body weight ($P > 0.05$, data not shown) and basal breathing parameters (V_T , f_R and V_E) in relative to the vehicle control (**Figures 3A–C**). Neither $P_a\text{CO}_2$ nor blood pH changed significantly in leptin-injected rats (data not shown). During exposure to 10% O_2 , V_T , f_R , and V_E were all increased in either leptin- or saline-injected LZRs ($P < 0.05\sim 0.01$) but the change in V_E was greater in leptin-injected rats in relative to the vehicle controls (**Figure 3E**). The stimulatory effect of leptin on HVR persisted for at least 2 weeks (**Table 3**). After bilaterally sectioning carotid sinus nerves, leptin-induced increase in V_E was abolished (**Figures 3D,E**). Collectively, chronic administration of leptin potentiated the HVR.

Effect of Leptin on Expression of pSTAT3 and TASK Channels

To investigate the possible mechanism underlying exogenous application of leptin action on the HVR, we tested the expression of ob-R and the downstream pSTAT3 and TASK-1, TASK-2, TASK-3 channels in CBs after injection of leptin for 7 days. The findings indicated that leptin administration caused greater upregulation of ob-R and of pSTAT3 ($P < 0.01$, $n = 4$ for each group, **Figures 4A,B**). Furthermore, leptin also enhanced expression of TASK-1, TASK-2, TASK-3 ($P < 0.01$, $n = 4$ for each group, **Figures 4C–E**). The results suggest that the stimulatory effect of leptin on HVR are closely associated with enhanced expression of pSTAT3 and TASK channels, which may contribute to the regulation of CB's chemosensitivity.

DISCUSSION

We demonstrate herein that the ob-R deficiency, rather than simple obesity, not only reduces baseline ventilation but also inhibits the HVR, with decreased pSTAT3 expression in CBs.

Chronic administration of leptin has no marked effects on basal ventilation but considerably enhances the HVR, accompanied by increased expression of pSTAT3. Additionally, either ob-R deficiency or leptin administration is reliably associated with changes in expression of TASK-1, TASK-2, and TASK-3 channels. These findings suggest that leptin signaling in the CB contributes to potentiation of HVR probably through enhancement of pSTAT3 and TASK channels expressions.


The obese Zucker rat represents a good model of ob-R deficiency and manifests relatively early onset obesity (Bray and York, 1971). One line of early evidence shows that respiratory system compliance was significantly lower in the OZR compared with the lean phenotype, and that resting ventilatory parameters (uncorrected for body weight) were similar between obese and lean animals, with similar ventilatory response to hypoxia between two phenotypes (Farkas and Schlenker, 1994). In the present study, we compared the difference between obese and lean phenotypes using the previously described method (Kumar et al., 2015; Fu et al., 2017) to normalize breathing parameters to body weight. Interestingly, with this analysis method, the OZRs exhibited fast f_R , reduced V_T and thus lower V_E during exposure to room air. This outcome interprets occurrence of hypercapnia and respiratory acidosis observed in OZRs. Based on these attributes, the OZR resembles an animal model of leptin resistance. The LHZR, a simple obesity control carrying genotype of the LZR, displayed hypercapnia, respiratory acidosis, relatively normal serum leptin, basal hypoventilation and moderate response to hypoxia, probably representing a model of simple obesity rather than leptin resistance. In obese patients, a higher level of leptin is found to cause an increase in basal ventilation associated with excess body mass, with OHS patients exhibiting an even higher serum leptin level than eucapnic individuals matched for body mass index (Phipps et al., 2002). In animal models, HFD rats exhibit an unchanged (Olea et al., 2014) or enhanced basal V_E (Ribeiro et al., 2017). Importantly, Ribeiro et al. found 3 weeks of HFD blunted leptin responses to hypoxia in the CB, probably due to development of CB leptin resistance, suggesting at least 3 weeks required for the establishment of leptin resistance. However, in the present study, hyperleptinaemia and leptin resistance did not occur in the LHZR most likely because of the genetic manipulation which makes the difference and the hypoventilation thus appears to be a restrictive ventilatory pattern. Another explanation may be supported by previous findings suggesting that elevation of leptin levels is a consequence of hypoxia and not of fat accumulation (Tatsumi et al., 2005). Taken together, our findings support that the impaired basal ventilation and HVR in OZRs is ascribed mainly to ob-R deficiency rather than mere obesity. We did not measure chest wall mechanics and compare the difference in chest wall impedance between two obese rats, whereas it would be expected that obesity-induced increase in chest wall impedance must play a relatively small part in such effects. This also helps better understanding of the results that ΔV_E induced by hypoxia is insignificant between LZRs and LHZRs.

Leptin's actions involve fast or slow onset, thus requiring minutes, several hours, even days before major changes occur. Its acute effects on breathing has been implicated in prior studies

TABLE 2 | Blood plasma levels of leptin in response to chronic application of saline and leptin.

Day	Saline group (Leptin, $\mu\text{g}/\text{L}$)	Leptin group (Leptin, $\mu\text{g}/\text{L}$)
0 day	7.8 ± 0.5	7.6 ± 0.5
7th day	8.2 ± 0.4	$13.1 \pm 0.6^{**}$
14th day	7.6 ± 0.5	9.3 ± 0.6
21st day	8.0 ± 0.6	8.6 ± 1.0

$n = 8$ for each group; $**P < 0.01$ by two-tailed unpaired t-test, vs. saline.

**TABLE 3 |** V_E measured during exposure to room air and hypoxia.

Day	O_2 level	Saline group (V_E ml/min/g)	Leptin group (V_E ml/min/g)
0 day	Air	469 ± 41	451 ± 21
	10% O_2	810 ± 24	819 ± 26
7th day	Air	479 ± 26	470 ± 36
	10% O_2	819 ± 44	1117 ± 97**
14th day	Air	463 ± 17	481 ± 32
	10% O_2	819 ± 32	1036 ± 88**
21st day	Air	468 ± 36	438 ± 24
	10% O_2	771 ± 75	914 ± 48

V_E , minute ventilation, $n = 8$ for each group, $**P < 0.01$ by two-way ANOVA with Bonferroni *post-hoc* test, vs. Saline.

(Chang et al., 2013; Olea et al., 2015; Pye et al., 2015; Ribeiro et al., 2017). For example, Olea et al. have found that acute application of leptin in anaesthetized animals augmented basal

V_E and potentiated the ischemic hypoxia-induced V_E in a dose-dependent manner (Olea et al., 2015). More recently, Ribeiro et al. reported that leptin increases V_E both in basal and hypoxic conditions in control rats but such effects were blunted in high fat diet fed rats (Ribeiro et al., 2017). In contrast, acute application of leptin in isolated CB type I cells failed significantly to alter the resting membrane potential and acidification-induced depolarization was unaffected by leptin, thereby suggesting that acute leptin stimulation did not alter CB's chemosensitivity (Pye et al., 2015). In addition to acute effects, chronic treatment with leptin *in vivo* also has been shown to potentiate respiratory chemoreflex (Bassi et al., 2014). Acute or chronic actions may be mediated by different leptin signaling pathways. Chronic administration of leptin in the present study did not augment basal ventilation but potentiated HVR in conscious rats, an effect persisting for ≥ 7 days. This appears to play an essential role in reinforcing ventilation to supply more oxygen when challenged by hypoxia. The plasma levels of leptin after 7 days treatment are quite similar to those quantified in plasma for the




FIGURE 4 | Stimulatory effect of leptin on pSTAT3/STAT3 and TASK-1,-2,-3 channel protein expression in CBs. Gel images and bar charts showing quantitative analysis of leptin-stimulated expression of ob-Rb (A) pSTAT3/STAT3 (B) and TASK-1,-2,-3 (C-E) in the CB ($n = 8$) of each group ($n = 4$). $**P < 0.01$ by two-tailed unpaired t -test.

OZR, taken together with the enhancement of HVR with 7 day leptin administration and the impairment of HVR in OZR rats, indicates that 7 day stimulatory effects of leptin did not result in leptin resistance.

The reason why the basal ventilation was herein not enhanced by the chronic application of leptin would be attributable to animal's state (anesthetized vs. conscious) and the dose of leptin administered. The dose of leptin applied to our animals was chosen based on what was chronically applied previously (Wjidan et al., 2015), and lower than that used in prior reports examining the acute effects of leptin on cardiovascular (Rahmouni et al., 2002; Rahmouni and Morgan, 2007) and respiratory functions (O'Donnell et al., 2000; Bassi et al., 2012). Moreover, in the case of potentially therapeutic utilizations, this dose would be expected to specifically exert a respiratory but not excessive

cardiovascular effects. Higher concentrations of leptin would be expected to saturate plasma carrier molecules and some of unbound leptin would be degraded. Furthermore, since the amount of leptin to cross blood-brain barrier is relied on receptor-mediated transport mechanism (Morris and Rui, 2009), access to the brain should be limited somehow.

Leptin's intracellular signal transduction has been extensively investigated, with exception of molecular mechanisms underlying its effect on respiratory chemosensitivity. It remains poorly understood how the activation of leptin signaling affects O_2 -sensitive channels which may determine CB's chemosensitivity. Along with recent studies indicating that the chemosensitivity of CBs is closely associated with TASK-1 and TASK-3 channels (Trapp et al., 2008; Tan et al., 2010), TASK-2 channels has also been evidenced to set central respiratory CO_2 and O_2 sensitivity

(Gestreau et al., 2010). In addition, activation of ob-Rs appears to regulate ion channels including ATP-sensitive K⁺ channels and voltage-gated K⁺ channels (Gavello et al., 2016). Although STAT3, SOCS3, and ERK1/2 may mediate leptin's role in the CBs, the critical information is lacking to data concerning modulatory effects of these molecules on TASK channels. In the present study, we did not directly address how the activation of ob-Rs and downstream signaling proteins regulated TASK-1, TASK-3, and TASK-2 channels, but notably, the altered expression levels of these channels would be expected to be attributable to leptin signaling and contribute to leptin-stimulated facilitation of the HVR. Future work is required for revealing such mechanisms.

In summary, leptin signaling participates in setting CB's O₂ sensitivity probably through the modulation of TASK-1, TASK-3, and TASK-2 channels and thus contributes to the potentiation of HVR. This line of cellular evidence extends our understanding of molecular mechanism of leptin action on breathing, shedding light on the etiology of obesity-related hypoventilation or apnea.

REFERENCES

- Bassi, M., Furuya, W. I., Menani, J. V., Colombari, D. S., do Carmo, J. M., da Silva, A. A., et al. (2014). Leptin into the ventrolateral medulla facilitates chemorespiratory response in leptin-deficient (ob/ob) mice. *Acta Physiol.* 211, 240–248. doi: 10.1111/apha.12257
- Bassi, M., Giusti, H., Leite, C. M., Anselmo-Franci, J. A., do Carmo, J. M., da Silva, A. A., et al. (2012). Central leptin replacement enhances chemorespiratory responses in leptin-deficient mice independent of changes in body weight. *Pflugers Arch.* 464, 145–153. doi: 10.1007/s00424-012-1111-1
- Bassi, M., Werner, I. F., Zoccal, D. B., Menani, J. V., Colombari, E., Hall, J. E., et al. (2015). Control of respiratory and cardiovascular functions by leptin. *Life Sci.* 125, 25–31. doi: 10.1016/j.lfs.2015.01.019
- Bray, G. A., and York, D. A. (1971). Genetically transmitted obesity in rodents. *Physiol. Rev.* 51, 598–646. doi: 10.1152/physrev.1971.51.3.598
- Chang, Z., Ballou, E., Jiao, W., McKenna, K. E., Morrison, S. F., and McCrimmon, D. R. (2013). Systemic leptin produces a long-lasting increase in respiratory motor output in rats. *Front. Physiol.* 4:16. doi: 10.3389/fphys.2013.00016
- Ciriello, J., and Caverson, M. M. (2014). Carotid chemoreceptor afferent projections to leptin receptor containing neurons in nucleus of the solitary tract. *Peptides* 58, 30–35. doi: 10.1016/j.peptides.2014.05.014
- Farkas, G. A., and Schlenker, E. H. (1994). Pulmonary ventilation and mechanics in morbidly obese Zucker rats. *Am. J. Respir. Crit. Care Med.* 150, 356–362. doi: 10.1164/ajrccm.150.2.8049815
- Fu, C., Xue, J., Wang, R., Chen, J., Ma, L., Liu, Y., et al. (2017). Chemosensitive Phox2b-expressing neurons are crucial for hypercapnic ventilatory response in the nucleus tractus solitarius. *J. Physiol.* 595, 4973–4989. doi: 10.1113/JP274437
- Gavello, D., Carbone, E., and Carabelli, V. (2016). Leptin-mediated ion channel regulation: PI3K pathways, physiological role, and therapeutic potential. *Channels* 10, 282–296. doi: 10.1080/19336950.2016.1164373
- Gestreau, C., Heitzmann, D., Thomas, J., Dubreuil, V., Bandulik, S., Reichold, M., et al. (2010). Task2 potassium channels set central respiratory CO₂ and O₂ sensitivity. *Proc. Natl. Acad. Sci. U.S.A.* 107, 2325–2330. doi: 10.1073/pnas.0910059107
- Gonzalez-Martin, M. C., Vega-Agapito, M. V., Conde, S. V., Castaneda, J., Bustamante, R., Olea, E., et al. (2011). Carotid body function and ventilatory responses in intermittent hypoxia. Evidence for anomalous brainstem integration of arterial chemoreceptor input. *J. Cell Physiol.* 226, 1961–1969. doi: 10.1002/jcp.22528
- Grill, H. J., Schwartz, M. W., Kaplan, J. M., Foxhall, J. S., Breining, J., and Baskin, D. G. (2002). Evidence that the caudal brainstem is a target for the inhibitory effect of leptin on food intake. *Endocrinology* 143, 239–246. doi: 10.1210/endo.143.1.8589
- Hall, J. E., da Silva, A. A., do Carmo, J. M., Dubinon, J., Hamza, S., Munusamy, S., et al. (2010). Obesity-induced hypertension: role of sympathetic nervous system, leptin, and melanocortins. *J. Biol. Chem.* 285, 17271–17276. doi: 10.1074/jbc.R110.113175
- Kumar, N. N., Velic, A., Soliz, J., Shi, Y., Li, K., Wang, S., et al. (2015). Regulation of breathing by CO₂ requires the proton-activated receptor GPR4 in retrotrapezoid nucleus neurons. *Science* 348, 1255–1260. doi: 10.1126/science.aaa0922
- Malhotra, A., and White, D. P. (2002). Obstructive sleep apnoea. *Lancet* 360, 237–245. doi: 10.1016/S0140-6736(02)09464-3
- Messenger, S. A., Moreau, J. M., and Ciriello, J. (2013). Effect of chronic intermittent hypoxia on leptin and leptin receptor protein expression in the carotid body. *Brain Res.* 1513, 51–60. doi: 10.1016/j.brainres.2013.03.022
- Moreau, J. M., Messenger, S. A., and Ciriello, J. (2015). Effects of angiotensin II on leptin and downstream leptin signaling in the carotid body during acute intermittent hypoxia. *Neuroscience* 310, 430–441. doi: 10.1016/j.neuroscience.2015.09.066
- Morris, D. L., and Rui, L. (2009). Recent advances in understanding leptin signaling and leptin resistance. *Am. J. Physiol. Endocrinol. Metab.* 297, E1247–E1259. doi: 10.1152/ajpendo.00274.2009
- O'Donnell, C. P., Tankersley, C. G., Polotsky, V. P., Schwartz, A. R., and Smith, P. L. (2000). Leptin, obesity, and respiratory function. *Respir. Physiol.* 119, 163–170. doi: 10.1016/S0034-5687(99)00111-5
- Olea, E., Agapito, M. T., Gallego-Martin, T., Rocher, A., Gomez-Niño, A., Obeso, A., et al. (2014). Intermittent hypoxia and diet-induced obesity: effects on oxidative status, sympathetic tone, plasma glucose and insulin levels, and arterial pressure. *J. Appl. Physiol.* (1985) 117, 706–719. doi: 10.1152/jappphysiol.00454.2014
- Olea, E., Ribeiro, M. J., Gallego-Martin, T., Yubero, S., Rigual, R., Masa, J. F., et al. (2015). The carotid body does not mediate the acute ventilatory effects of leptin. *Adv. Exp. Med. Biol.* 860, 379–385. doi: 10.1007/978-3-319-18440-1_43
- Panetta, P., Berry, A., Bellisario, V., Capoccia, S., Raggi, C., Luoni, A., et al. (2017). Long-term sex-dependent vulnerability to metabolic challenges in prenatally stressed rats. *Front. Behav. Neurosci.* 11:113. doi: 10.3389/fnbeh.2017.00113
- Phipps, P. R., Starratt, E., Catterson, I., and Grunstein, R. R. (2002). Association of serum leptin with hypoventilation in human obesity. *Thorax* 57, 75–76. doi: 10.1136/thorax.57.1.75
- Porzionato, A., Rucinski, M., Macchi, V., Stecco, C., Castagliuolo, I., Malendowicz, L. K., et al. (2011). Expression of leptin and leptin receptor

AUTHOR CONTRIBUTIONS

FY, JF, and HW acquired the data; FY, JF, and ZW analyzed and interpreted data; FY, XZ, and HY drafted the manuscript; FY, SW, and YZ were responsible for study concept and design; YZ and SW obtained research funding.

ACKNOWLEDGMENTS

This study was supported by grants from the National Natural Science Foundation of China (31271223, 31671184) and from the Province Natural Science Foundation of Hebei (H2016206477). The experiments comply with the current laws of the country in which they were performed.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphys.2018.00249/full#supplementary-material>

- isoforms in the rat and human carotid body. *Brain Res.* 1385, 56–67. doi: 10.1016/j.brainres.2011.02.028
- Pye, R. L., Dunn, E. J., Ricker, E. M., Jurcsisn, J. G., Barr, B. L., and Wyatt, C. N. (2015). Acutely administered leptin increases $[Ca^{2+}]_i$ and BK Ca currents but does not alter chemosensory behavior in rat carotid body type I cells. *Adv. Exp. Med. Biol.* 860, 61–67. doi: 10.1007/978-3-319-18440-1_8
- Rahmouni, K., Haynes, W. G., Morgan, D. A., and Mark, A. L. (2002). Selective resistance to central neural administration of leptin in agouti obese mice. *Hypertension* 39, 486–490. doi: 10.1161/hy0202.102836
- Rahmouni, K., and Morgan, D. A. (2007). Hypothalamic arcuate nucleus mediates the sympathetic and arterial pressure responses to leptin. *Hypertension* 49, 647–652. doi: 10.1161/01.HYP.0000254827.59792.b2
- Ribeiro, M. J., Sacramento, J. F., Gallego-Martin, T., Olea, E., Melo, B. F., Guarino, M. P., et al. (2017). High fat diet blunts the effects of leptin on ventilation and on carotid body activity. *J. Physiol.* doi: 10.1113/JP275362. [Epub ahead of print].
- Tan, Z. Y., Lu, Y., Whiteis, C. A., Simms, A. E., Paton, J. F., Chapleau, M. W., et al. (2010). Chemoreceptor hypersensitivity, sympathetic excitation, and overexpression of ASIC and TASK channels before the onset of hypertension in SHR. *Circ. Res.* 106, 536–545. doi: 10.1161/CIRCRESAHA.109.206946
- Tartaglia, L. A. (1997). The leptin receptor. *J. Biol. Chem.* 272, 6093–6096. doi: 10.1074/jbc.272.10.6093
- Tatsumi, K., Kasahara, Y., Kurosu, K., Tanabe, N., Takiguchi, Y., and Kuriyama, T. (2005). Sleep oxygen desaturation and circulating leptin in obstructive sleep apnea-hypopnea syndrome. *Chest* 127, 716–721. doi: 10.1378/chest.127.3.716
- Trapp, S., Aller, M. I., Wisden, W., and Gourine, A. V. (2008). A role for TASK-1 (KCNK3) channels in the chemosensory control of breathing. *J. Neurosci.* 28, 8844–8850. doi: 10.1523/JNEUROSCI.1810-08.2008
- Wang, S., Benamer, N., Zanella, S., Kumar, N. N., Shi, Y., Bévéngut, M., et al. (2013). TASK-2 channels contribute to pH sensitivity of retrotrapezoid nucleus chemoreceptor neurons. *J. Neurosci.* 33, 16033–16044. doi: 10.1523/JNEUROSCI.2451-13.2013
- Wjidan, K., Ibrahim, E., Caszo, B., Gnanou, J., and Singh, H. (2015). Dysregulation of glucose homeostasis following chronic exogenous administration of leptin in healthy sprague-dawley rats. *J. Clin. Diagn. Res.* 9, OF06–OF09. doi: 10.7860/JCDR/2015/15594.7003

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Yuan, Wang, Feng, Wei, Yu, Zhang, Zhang and Wang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.