



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

CHAPTER OUTLINE

Infections of the Liver	322	Necrotizing Enterocolitis	333
Microbiology	322	Pathology and Pathogenesis	334
Pathogenesis	323	Microbiology	334
Clinical Manifestations	324	Clinical Manifestations	335
Diagnosis	324	Diagnosis	336
Prognosis	324	Treatment	337
Treatment	325	Prevention	337
Splenic Abscess	325	Prognosis	338
Infections of the Biliary Tract	325	Endocarditis	338
Infections of the Adrenal Glands	326	Pericarditis	340
Appendicitis	327	Mediastinitis	341
Microbiology	327	Esophagitis	342
Pathogenesis	327	Infections of Endocrine Organs	342
Clinical Manifestations	327	Infections of the Salivary Glands	342
Diagnosis	328	Infections of the Skin and Subcutaneous Tissue	342
Prognosis	328	Pathogenesis	343
Treatment	329	Microbiology	344
Peritonitis	329	Epidemiology	344
Microbiology	329	Clinical Manifestations	344
Pathogenesis	330	Diagnosis	347
Clinical Manifestations	331	Differential Diagnosis	347
Diagnosis	332	Treatment	348
Prognosis	332	Prevention	348
Treatment	333	Conjunctivitis and Other Eye Infections	349

INFECTIONS OF THE LIVER

Bacterial infection of the hepatic parenchyma frequently is recognized as multiple, small inflammatory foci (hepatic microabscesses) observed as an incidental finding in infants dying with sepsis. Diffuse hepatocellular damage, often in conjunction with infection of several organ systems, may be present after transplacental passage of microorganisms to the fetal circulation. Liver involvement rarely may take the form of a solitary purulent abscess. Metastatic focal infections of the liver associated with bacteremia resolve with antimicrobial therapy, are not recognized, or are found only at postmortem examination. Rarely, they are clinically apparent as solitary [1] or multiple [2] large abscesses diagnosed during life.

Although metastatic infections are rare, it is difficult to ascertain their true incidence. In a survey of more than 7500 autopsies of children performed from 1917-1967, Dehner and Kissane [3] found only 3 neonates with multiple, small, pyogenic hepatic abscesses, whereas a review of approximately 4900 autopsies [4] performed at Los Angeles Children's Hospital from 1958-1978 revealed 9 infants with pyogenic hepatic abscesses [5]. Among 175,000 neonates admitted from 1957-1977 to Milwaukee Children's Hospital, 2 died with hepatic microabscesses [6]; 3 patients

with hepatic microabscesses were seen among 83,000 pediatric patients admitted to New York Hospital from 1945-1983 [7]; and one case of hepatic microabscesses was reported at the University of Texas Medical Branch in Galveston from 1963-1984 [8]. Most reviewers who have discussed postmortem observations of infants dying with neonatal sepsis have not described the occurrence of such secondary sites of infection [9-16] or have presented them as an occasional ancillary finding [17,18].

Solitary hepatic abscesses in newborns have also been reported rarely. About 30 such cases have been described [1,3,4,19-41]. These infections frequently are associated with prematurity and umbilical vein catheterization,* whereas solitary abscesses may occur because of bacteremia. Murphy and Baker [42] described a solitary abscess after sepsis caused by *Staphylococcus aureus*.

MICROBIOLOGY

The etiologic agents in the infants described by Dehner and Kissane [3], Moss and Pysher [5], Chusid [6], and included *Escherichia coli*, *S. aureus*, *Pseudomonas aeruginosa*,

*References [5,6,21,22,25,28,32,34-41].

Klebsiella species, *Enterobacter* species, and *Listeria monocytogenes*. The causative bacteria of solitary abscesses are generally the bacteria colonizing the umbilical stump [43], including *S. aureus* (11 cases), *E. coli* alone (3 cases), *E. coli* with *S. aureus* (2 cases) or enterococcus (1 case), *Enterobacter* species (3 cases), *Klebsiella pneumoniae* alone (2 cases), *K. pneumoniae* with *Proteus* species (1 case), *P. aeruginosa* (1 case), *Staphylococcus epidermidis* with group F streptococcus (1 case), and *Streptococcus pyogenes* (1 case). In three infants, the abscesses were described as “sterile.” [19,28,31] Although one of these infants had received penicillin for 9 days before surgical drainage, it is possible that in all three cases the abscesses were caused by anaerobic bacteria that failed to grow. The presence of gas in seven abscesses [25,28,34,35,39] may indicate infection with anaerobes, a frequent cause of liver abscess in adults [44].

The most common cause of intrauterine bacterial hepatitis, congenital listeriosis, characteristically involves the liver and adrenals (see Chapter 13). Typical lesions are histologically sharply demarcated areas of necrosis (miliary granulomatosis) or microabscesses containing numerous pleomorphic gram-positive bacilli [15]. Descriptions in the early 1900s of miliary necrosis of the liver related to “gram-positive argentophilic rodlike organisms” probably also represented infections with *L. monocytogenes*, which was not isolated and identified until 1926 [26].

Intrauterine tuberculosis results from maternal bacillemia with transplacental dissemination to the fetal bloodstream (see Chapter 18). Because the liver is perfused by blood with a high oxygen content [45] and is the first organ that encounters tubercle bacilli, it is often severely involved [15,44,46]. The presence of primary liver foci is considered evidence for congenital tuberculous infection as a result of hematogenous spread through the umbilical vein [47]. Closed-needle biopsy may be less accurate in the diagnosis of hepatic granulomas, and open biopsy may be required to confirm liver and regional node involvement [48]. Although generalized fetal infection may also arise through aspiration of contaminated amniotic fluid, the lesions acquired in this manner are usually most prominent in the lungs. In addition to hepatomegaly, a clinical picture of fever with elevated serum IgM and chorioretinitis (e.g., choroid tubercles) may be similar to that caused by other congenital infectious agents [49]. In a review by Abughal and coworkers [49], positive sites of culture for tuberculosis included liver (8 of 9), gastric aspirate (18 of 23), tracheal aspirate (7 of 7), ear (5 of 6), and cerebrospinal fluid (3 of 10). Noncaseating granulomatous hepatitis, thought to be caused by a hypersensitivity reaction related to bacille Calmette-Guérin vaccination, has also been described in a neonate [50], but histologic and bacteriologic studies performed on liver biopsy specimens failed to identify the presence of acid-fast bacilli or bacille Calmette-Guérin organisms.

Bacterial infection of the fetal liver rarely has been reported in association with maternal tularemia [51], anthrax [52], typhoid fever [53], and brucellosis [54]. It is uncertain whether the isolation of bacteria from the livers of stillborn fetuses is significantly associated with their clinical course [55,56].

Treponema pallidum is the spirochete most commonly associated with transplacental hepatic infection (see

Chapter 16). Pathologic changes in the liver, which may be found in 95% of infants dying with congenital syphilis [57], include diffuse hepatitis or focal areas of inflammation, both frequently accompanied by increased connective tissue and enlargement of the liver [15,57–59]. Involvement of liver has also been documented, on the basis of isolation of organisms or their identification in histologic sections, in newborns with intrauterine infection caused by various *Leptospira* species (*Leptospira icterohaemorrhagiae* [60,61], *Leptospira pomona* [62], *Leptospira canicola* [63], *Leptospira kasman* [64]).

Transplacental infection of the fetus with *Borrelia recurrentis* causes little or no inflammation of liver parenchyma or biliary epithelium despite the presence of numerous spirochetes in the sinusoids [65–68]. Congenital infection has been suggested with *Borrelia burgdorferi* [69] (cause of Lyme disease); hepatic, central nervous system, and cardiac lesions may be observed, and widely disseminated lesions were reported to occur in other tissues. This single case is controversial, however, and the American Academy of Pediatrics does not accept congenital Lyme disease because no “causal relationship between maternal Lyme disease and abnormalities of pregnancy or congenital disease caused by *B. burgdorferi* has been documented conclusively” [68a]; no evidence exists that Lyme disease can be transmitted via human milk.

PATHOGENESIS

Infectious agents may reach the liver of the fetus or newborn by one of several pathways: transplacental or transoral intrauterine infection, extension of thrombophlebitis of the umbilical vein, through the hepatic artery during the course of a systemic bacteremia, pyelophlebitis owing to a focus of infection in the drainage of the portal vein (mesenteric or splenic veins), direct invasion from contiguous structures or because of trauma or surgical inoculation, and extension up the biliary passages in cases of suppurative cholangitis. Abscesses with no apparent focus of infection seem to be common in newborns compared with older children [30]. Three such cases, all in infants with solitary hepatic abscesses, have been described [23,24,31]. Descriptions of the surgical findings, together with the nature of the lesions, suggest that an umbilical vein infection, obscured by the large collection of purulent material in the abscess, was the probable pathogenesis in all infants.

The mode of infection usually determines the pattern of hepatic involvement. Intense and prolonged seeding of the liver parenchyma, such as that which occurs in conjunction with intrauterine infection or neonatal sepsis, almost invariably results in diffuse hepatocellular damage or multiple small inflammatory lesions [3,5,6]. Umbilical vein thrombophlebitis may cause an abscess of the falciform ligament [70] or extend into a single branch of the portal vein to produce a solitary pyogenic abscess,* or it can lead to disseminated foci of infection through dislodgment of septic emboli [6,71–74].

The frequent use of umbilical catheters has been associated with an increase in the numbers of infants

*References [6,21,22,26,29,32,33].

with solitary [5,6,20–22,32,34–40] or multiple [5,75,76] hepatic abscesses. In three large series, including almost 500 infants who died after placement of umbilical vein catheters, 29 infants were found to have purulent infections of hepatic vessels or parenchyma [37,75,77]. Use of venous catheters for infusion of hypertonic or acidic solutions may provide a necrotic focus for abscess formation [21,32,34–36,76,77], and prolonged [5,22,32,77] or repeated [63] catheterization of a necrotic umbilical stump provides an ideal pathway for introduction of pathogenic organisms. It has been postulated that some hepatic abscesses have been caused by infusion of contaminated plasma [28] or by the use of nonsterile umbilical catheters [75].

Although neonatal liver abscesses usually are caused by hematogenous dissemination of bacteria through the hepatic artery or umbilical vein, examples of infection arising from various other sources have been described. Solitary abscesses have followed a presumed portal vein bacteremia caused by amebic colitis [19,20]. Direct invasion of adjacent liver parenchyma from purulent cholecystitis [24] or postoperative perihepatic abscesses [5] also has been observed. Ascending cholangitis, the most frequent cause of hepatic purulent infections in adults [30], has not been implicated in the causes of newborn infections.

Disease caused by embryonic anatomic errors is unique to newborns. Shaw and Pierog [78] described a newborn with umbilical herniation of a pedunculated supernumerary lobe of the liver; histologic examination showed numerous small foci of early abscess formation. Although signs and symptoms of sepsis appeared at 18 days of age, possibly the result of bacterial spread from the liver to the umbilical vein, the infant improved and ultimately recovered after removal of the polypoid mass on the 19th day of life.

Descriptions of “umbilical sepsis” and “acute interstitial hepatitis” recorded by Morison [79] seem to indicate that his patients had acquired bacterial infections of umbilical vessels with widespread extension into portal tracts. Although mild periportal parenchymal necrosis was observed in a few infants, hepatocellular damage was minimal or absent in most. Similar lesions have been found in infants dying with sepsis [80] and infantile diarrhea [81].

CLINICAL MANIFESTATIONS

Multiple hepatic abscesses and diffuse hepatitis related to neonatal sepsis or transplacental fetal infection are usually recognized only at autopsy. Very few clinical manifestations referable to hepatocellular damage are evident before death. The signs and symptoms associated with these conditions are those of the underlying sepsis or of secondary metastatic complications, such as meningitis, pneumonitis, or peritonitis.*

Solitary abscesses are indolent in terms of their development and clinical presentation. Although the suppurative umbilical focus or umbilical catheterization responsible for the introduction of microorganisms can usually be traced to the 1st week of life, evidence of hepatic involvement is usually not apparent before the 2nd or 3rd

week. The abscess frequently becomes a source for the hematogenous dissemination of microorganisms so that most infants have signs and symptoms of a bacteremia. Despite intense infection of the underlying vessels, the umbilical stump usually shows no evidence of inflammation or purulent discharge. The presence of hepatomegaly, a finding commonly associated with neonatal sepsis, also offers little aid in establishing a definitive diagnosis. In one half of infants for whom physical findings are clearly described, a well-delineated, often fluctuant or tender mass could be palpated in the epigastrium or right upper quadrant. On a few occasions, this mass was noticed by the infant's mother several days before the onset of systemic symptoms. Abscesses occur in the right or left lobe of the liver with almost equal frequency and are generally 3 cm or more in diameter at the time of surgical exploration.

DIAGNOSIS

Hematologic studies are of little value in establishing a diagnosis; leukocyte counts and sedimentation rates may be normal or elevated. The serum levels of liver enzymes may also be normal [25,38] or elevated [5,23,36].

Abdominal radiographs are usually normal or show nonspecific displacement of the lower edge of the liver. In five infants, diagnosis was suspected from plain x-ray films by the presence of gas within the hepatic shadow [28,32,34,39]. Radiologic findings that commonly accompany hepatic abscess in older children, such as an altered contour of the diaphragm, right pleural effusion, and platelike atelectasis [82], are rarely present in neonates.

Ultrasonography should be the initial imaging study in newborns with clinical evidence of a hepatic abscess [83–86]. If ultrasound is negative, and the diagnosis is still strongly suspected, more sensitive techniques such as computed tomography (CT) or magnetic resonance imaging (MRI) should be performed [83–88]. Enhancement with contrast agents may increase the definition of smaller abscesses. Because congenital cysts, arteriovenous malformations, and tumors with central necrosis or hemorrhage can mimic hepatic abscess, the diagnosis should always be confirmed by aspiration of purulent material at laparotomy or by means of percutaneous drainage with ultrasound or CT guidance [84,89,90].

PROGNOSIS

The prognosis for infants with diffuse liver involvement related to fetal or neonatal sepsis depends on the underlying condition because hepatic function is rarely compromised sufficiently to determine the outcome. In most cases, pathologic changes in the liver are unsuspected before postmortem examination.

Of 24 infants with solitary hepatic abscesses whose course was described, 8 died. Two infants died before antibiotics were available [26], and the death of another was ascribed to cecal perforation [20]. Four newborns died with sepsis caused by organisms that were identical to the organisms isolated from the abscess [21,25,33,34]. Prematurity was undoubtedly a major contributing factor in two of these deaths [21,25].

*References [2,3,29,33,72,75].

TREATMENT

Newborns with a solitary hepatic abscess have traditionally been treated with open surgical drainage in conjunction with antibiotic therapy. Percutaneous catheter drainage is less invasive and often is the preferred first treatment. Several investigators have described the use of percutaneous drainage of intrahepatic abscesses and cysts, guided by CT or ultrasonography, in neonates [41,75,90] and children [7,84,89]. When combined with antibiotic therapy and monitored by ultrasound to ensure resolution, this treatment has been highly effective. It is questionable whether drainage contributed to recovery other than by aiding the selection of antibiotic coverage. Subsequently, patients have been successfully treated with empirical antibiotic therapy alone [91,92]. Conservative medical management in infants has been described in only two neonates and a 5-month-old infant [33,37,78].

The risk of bacteremia and disseminated infection is high in neonates, and the need to identify infecting organisms to guide antibiotic coverage is of greater urgency in the first weeks of life. It is appropriate to ascertain a microbiologic diagnosis with radiographically guided aspiration or drainage of hepatic abscess in a newborn. When proper equipment (e.g., CT, ultrasonography) and experienced personnel are available, this can be attempted percutaneously [89,90]. When they are unavailable, open surgical drainage should be performed. Empirical antibiotic therapy should be reserved only for infants for whom it is believed that the risk of open or closed drainage would exceed the potential benefits.

If purulent material is obtained, initial antibiotic therapy can be selected on the basis of Gram stain. In addition to *S. aureus* and the aerobic enteric organisms commonly associated with hepatic abscesses, anaerobic bacteria have been suspected as the cause of infection in numerous patients.* If foul-smelling pus is aspirated or if Gram-stained smears show organisms with the characteristic morphology of anaerobes [33], metronidazole, β -lactam and β -lactamase inhibitor combinations (e.g., piperacillin and tazobactam), clindamycin, or imipenem should be included in the initial regimen. Cultures of blood, cerebrospinal fluid, and urine should also be considered before initiation of therapy.

If empirical antibiotic therapy is required, it must be adequate for infections caused by *S. aureus*, enteric organisms, and anaerobic bacteria. The combination oxacillin, gentamicin, and clindamycin is appropriate. In nurseries where methicillin-resistant *S. aureus* (MRSA) or methicillin-resistant *S. epidermidis* infections have been a problem, substitution of vancomycin for oxacillin can provide coverage for these organisms. Gentamicin (and other aminoglycosides) and vancomycin levels must be monitored and dosages adjusted as necessary. Extended-spectrum cephalosporins (e.g., cefotaxime, cefepime, ceftazidime) and carbapenems (e.g., meropenem) may be used for enteric organisms and *Pseudomonas* species, often obviating the need for aminoglycosides. β -lactam and β -lactamase inhibitor combination drugs (e.g., piperacillin and tazobactam or ampicillin

and sulbactam) may provide coverage for many enteric organisms and anaerobic bacteria.

Definitive therapy is based on results of bacteriologic cultures that identify the bacteria and its antibiotic susceptibility. Adequate anaerobic transport and culture techniques must be available if meaningful information is to be obtained. Duration of treatment is based on clinical response, cessation of drainage, and resolution of the abscess cavity as determined by serial ultrasound examinations. Parenteral therapy should be maintained for at least 2 weeks and longer term therapy may be administered when necessary. In older children with multiple abscesses or in children for whom surgery is not feasible, therapy for 6 weeks or more has been recommended.

SPLENIC ABSCESS

Similar to hepatic abscesses, splenic abscesses have been rarely described in infants [93]. Only 1 of 55 splenic abscesses occurred in an infant younger than 6 months. *S. aureus*, *Candida* species, and streptococci were the most frequent causes. In 20 of 48 cases, hepatic abscesses coexisted with splenic abscess. In the single infant case, torsion of the splenic vessels was present, whereas in older children, other distant infections of hematologic conditions (e.g., hemoglobinopathy, hematogenous malignancy) were the associated comorbid conditions.

INFECTIONS OF THE BILIARY TRACT

The development of ultrasonography has provided a safe and rapid means for evaluating the neonatal biliary tract. Consequently, an increasing number of reports have appeared describing ultrasound changes seen in the first month of life, with hydrops [94,95], cholelithiasis [95–100], and transient distention of the gallbladder associated [94,95,99,101–104] or unassociated [99,102,103,105–108] with sepsis. Ultrasound criteria for separating normal from pathologically enlarged gallbladders and biliary tracts in neonates have also been described [109,110].

Despite advanced technology and increased surveillance, cholecystitis in the neonate is observed infrequently. The literature has documented about 25 cases, of which 9 were seen in association with an epidemic of neonatal enteritis caused by *Salmonella enteritidis* [111]. Of the remaining infants, 16 were the subjects of isolated case reports [24,103,112–123], and 3 died of other causes with inflammatory changes in the gallbladder described as an incidental finding at autopsy [81,95,124]. A tissue diagnosis of “chronic cholecystitis” was established in an infant whose biliary disease apparently began at 6 days of age [125].

The pathogenesis of this condition is uncertain; all but three cases [99,118,122] of cholecystitis in the newborn period have been acalculous. It is postulated that sepsis, dehydration, prolonged fasting (e.g., total parenteral nutrition), congenital obstruction, or a stone impacted in the cystic duct leads to biliary stasis and acute distention of the gallbladder. In most cases, resolution of the primary process permits restoration of the flow of bile and relief of distention. In some cases, prolonged obstruction leads to hydrops [94]. Cholecystitis rarely follows, perhaps because of a direct toxic effect of retained bile or because of ischemia related to elevated intraluminal

*References [25,28,32,33,35,39].

pressure. Bacterial invasion by fecal flora is probably a secondary phenomenon [114,115,126]. Organisms that have been isolated from gallbladder contents or tissue include *E. coli* [114–116,123], *Serratia marcescens* [103,117], *Pseudomonas* species [115], *Streptococcus faecalis* [123], viridans streptococci [121], *S. aureus* [123], and *Clostridium welchii* [123]. “Gram-positive cocci” were identified by Gram stain in one patient [113].

Infants with cholecystitis may become ill at any time during the first weeks of life; most cases are diagnosed in the 3rd or 4th week. The typical clinical picture is one of sepsis together with signs of peritoneal inflammation and a palpable tender right upper quadrant or epigastric mass. Diarrhea frequently accompanies these findings. Although ultrasonography and radionuclide scintigraphy are helpful in suggesting the presence of gallbladder enlargement or inflammation, diagnosis can be confirmed only by surgical exploration [94,102,103,106,108]. Treatment consists of cholecystectomy or tube cholecystotomy combined with systemic antimicrobial therapy based on Gram stain, culture, and susceptibility studies. If a T tube is placed in the gallbladder, a cholangiogram should be obtained to confirm patency of the biliary system before the tube is removed.

Changes compatible with a diagnosis of ascending cholangitis have been described in histologic sections of liver specimens from infants who died with diarrhea accompanied by hepatocellular injury with cholestasis [91]. Bacteria were also identified in the biliary tree of 2 of 178 premature infants who died after placement of an umbilical venous catheter for an exchange transfusion or for delivery of parenteral fluids [75]. The reasons for this association, if any, are unclear. An infant with spontaneous cholangitis caused by *Enterobacter agglomerans*, presenting as a fever of unknown origin at 3 weeks of age, has also been reported [127].

Severe inflammation and fibrosis of extrahepatic bile ducts and diffuse changes in the portal tracts, resembling changes found in biliary atresia, were found in a premature infant who died 3 hours after birth of listeriosis [128]. The investigator postulated that occult prenatal infections with *L. monocytogenes* might be a rare cause of ascending cholangitis manifesting as idiopathic biliary atresia at birth.

INFECTIONS OF THE ADRENAL GLANDS

Multiple adrenal microabscesses are occasionally found as metastatic lesions associated with neonatal sepsis. These abscesses are particularly characteristic of neonatal listeriosis (see Chapter 13). Solitary adrenal abscesses are rare, however; only about 25 such cases have been described [17,129–150].

The spectrum of organisms responsible for adrenal abscesses is the same as that seen in neonatal sepsis and includes *E. coli* (seven cases) [17,129,130,135–138], group B streptococci (GBS) (four cases) [138–141], *Proteus mirabilis* (three cases) [131,132,144], *S. aureus* [142,143], *Bacteroides* species [133,145], and two cases each of *Streptococcus pneumoniae* with *Bacteroides* species [134]; *Peptostreptococcus* species [146] was recovered from one

case. Drainage of foul-smelling pus at surgery suggests that anaerobic bacteria may have been present in two infants from whom *E. coli* and *S. aureus* were isolated [136,142]. Cultures were not obtained from four patients [147–150].

Fourteen abscesses were located on the right side, seven were located on the left, and three [138,139,147] were bilateral. Three fourths of the infants were male. The same laterality and sex predominance are seen with adrenal hemorrhage in the newborn [147,150–152], and it has been postulated that formation of an adrenal abscess requires a preexisting hematoma as a nidus for bacterial seeding [137,138]. This theory of pathogenesis is supported further by clinical observations [134,135,139,146,147] and by objective evidence (e.g., curvilinear calcifications [130,132] documenting the presence of hemorrhage before development of an abscess [134,138,142,145,150].

Most infants with adrenal abscess have presented in the 3rd or 4th week of life with signs of sepsis and an abdominal or flank mass. A history of difficult delivery or intrapartum asphyxia was observed in about one half of these infants, and significant maternal fever or infection during labor was observed in about one fourth [138,140,141,150]. Although a few infants are afebrile when first evaluated, a palpable mass is almost always present. Abscesses are usually 6 to 8 cm in diameter, with some containing 200 mL of pus [133] and measuring 12 cm in diameter [134] or crossing the midline [146].

Laboratory studies are helpful in the evaluation of a possible adrenal abscess. Most infants exhibit a leukocytosis; about one third are anemic with a history of prolonged neonatal jaundice, both of which are features associated with adrenal hemorrhage. Urinary excretion of catecholamines and their metabolites (particularly vanillylmandelic acid and homovanillic acid), which is usually increased with neuroblastoma, is normal. Because most infants with adrenal abscess are seen for evaluation of possible sepsis, a blood culture, lumbar puncture, urine culture, and chest radiograph should be obtained.

Ultrasonography has become a widely accepted modality for initial evaluation of all neonatal abdominal masses. With the presence of an adrenal abscess, ultrasound examination can help to define the extent and cystic nature of the lesion and often can show movable necrotic debris in the abscess cavity.* With serial examinations, abscesses can be distinguished from masses associated with liquefying hematoma, adrenal cyst, hydronephrosis of an obstructed upper pole duplication, or necrotic neuroblastoma [138,140,150,153,154]. Intravenous pyelography shows downward displacement of the kidney and compression of the upper calyces, which confirms the presence of a suprarenal mass.† A round, suprarenal, radiopaque halo or rim with central lucency, which is characteristic of adrenal abscess, may also be seen on early films [137,139,143], but is not pathognomonic [138]. Intravenous pyelography adds little diagnostic information to that provided by ultrasound studies. Experience with radionuclide scanning [140,142,143], CT [138,144], and

*References [132,137,138,141,142,144–148].

†References [130–132,134,136,138,141,142,144–146,149].

MRI [126] in this condition is limited, but these modalities are likely to be as useful as ultrasonography.

Whatever diagnostic methods are used, concern about persisting signs of sepsis and the possible presence of an adrenal neoplasm usually encourage early efforts to establish a diagnosis. In the past, recommended management has been incision and drainage or resection of the abscess [134,138,141,144,150]. Needle aspiration under ultrasound guidance, combined with placement of a catheter for drainage and irrigation, has proved to be a useful alternative method [88,130,131,143] and is likely to supplant open drainage as the preferred method. Antibiotic therapy should be based on Gram stain, culture, and susceptibility studies of abscess fluid and should be continued for 10 to 14 days, provided that drainage can be established.

The adrenals are infected in about 15% of infants with congenital syphilis [57,58]. In addition to the presence of spirochetes, the most frequent and characteristic change is an extraordinary amount of cellular connective tissue in the capsule.

APPENDICITIS

Acute appendicitis is extremely rare in infants younger than 4 weeks of age. Reviews of more than 25,000 cases of appendicitis in infants and children in Great Britain [155], Ireland [156], Norway [157], Germany [158], and the United States [159–164] revealed only 8 infants who presented during the neonatal period. Pediatric surgery centers in Germany [165], Boston [166], Cleveland [167], Chicago [168], and Detroit [169] found only four cases of neonatal appendicitis. Since the condition was first described by Albrecht in 1905 [170,171] and Diess in 1908 [172], approximately 65 cases of neonatal suppurative appendicitis have been reported in the literature with sufficient details to permit characterization of the clinical features [155,160,161,164,173–210]. Infants with appendicitis caused by other conditions, such as Hirschsprung disease [211,212], necrotizing enterocolitis (NEC) [213], or incarceration in an inguinal hernia [214,215], have not been included in this discussion. An additional 25 to 30 cases that have been reported with incomplete clinical observations, listed in series of patients with neonatal peritonitis (see “Peritonitis”) or mentioned in other review articles but unavailable for analysis, are also not included.

Inflammation of the appendix is more common in newborn boys than newborn girls. In reports in which the sex was stated, 40 cases occurred in boys, and 17 cases occurred in girls. Prematurity also seems to be a predisposing factor: 23 of the 49 infants whose birth weights were recorded weighed less than 2500 g at birth. The incidence of appendicitis in infants of multiple births (six twins and one triplet) seems to be higher than would be expected on the basis of low birth weight alone.

MICROBIOLOGY

Because obstruction of the appendiceal lumen is responsible for almost all cases of appendicitis [166], it is intuitive that gram-negative enteric organisms resident in the

bowel are usually isolated from the peritoneal fluid or periappendiceal pus of about 75% of infants. Specific etiologic agents include *E. coli*, *Klebsiella* species, *Enterobacter* species, *Pseudomonas* species, *Proteus* species, untyped streptococci, *S. aureus*, and *Bacteroides* species. These bacterial species have also been isolated from the peritoneal fluid of older children with appendicitis [163,166,216]. Attempts at isolation of anaerobic bacteria have been rarely described.

A single case of perforated amebic appendicitis with secondary bacterial peritonitis and multiple hepatic abscesses in a premature infant born in Great Britain has been reported. *Entamoeba histolytica* observed in the wall of the necrotic appendix was presumably acquired from the infant's father, who was a carrier [20]. A patient with gangrenous appendicitis associated with *Rhizopus oryzae* has also been reported [217]. It was postulated that the fungus colonized the infant's gut by transfer from an adhesive bandage used to secure an endotracheal tube.

PATHOGENESIS

Obstruction of the appendiceal lumen has been generally accepted as the primary cause of appendicitis in all age groups. The relative rarity of this condition in the first month of life is probably related to factors that serve to decrease the likelihood of obstruction, including a wide-based, funnel-shaped appendix; the predominantly liquid and soft solid diet given to infants; the absence of prolonged periods in the upright position; and the infrequency of infections that cause hyperplasia of the appendiceal lymphoid tissue [163,218,219].

The causes of luminal obstruction in the newborn period, when recognized, are often extrinsic to the appendix itself. Reports of appendicitis caused by the presence of ectopic pancreatic tissue [160], a fecalith [174], or meconium plug [167] are unusual exceptions. In 1911, it was suggested that sharp angulation of the appendix, bent on itself in the narrow retrocolic space, may be an important cause of obstruction [220]; however, this anatomy, found in 11 neonates with inflammatory changes in the appendix and noted among 200 consecutive autopsies in infants younger than 3 months at death, has not been repeated in the past 80 years.

Inflammation of the appendix with perforation has been described as the presenting illness in several infants with neonatal Hirschsprung disease [211,213]. The association of these two conditions has been attributed to functional obstruction, increased intraluminal pressure, and fecal trapping that occur proximal to aganglionic segments. Suppurative appendicitis related to incarceration and strangulation of the cecum within an inguinal or scrotal hernia has been found in numerous infants [214,215].

CLINICAL MANIFESTATIONS

The onset of neonatal appendicitis generally occurs during the first 2 weeks of life. Only 3 of 54 infants with this condition presented between the 1st and 10th day. The reasons for this phenomenon are unclear, particularly in view of the relatively even distribution of cases during the remainder of the 1st year of life [164]. Five cases of

“prenatal” appendicitis have been described [221–225]. Of the four available for analysis, only one showed definite evidence of a suppurative process in the appendix and signs of bowel obstruction clearly present at birth [221]; however, cultures and Gram stain of the pus found at surgery were free of bacteria. Poisoning by mercuric chloride was suspected in one [223] of the remaining three cases, and the other two, who were said to have prenatal rupture of the appendix, were asymptomatic until the 2nd [221] and 12th [225] days of life.

The signs of neonatal appendicitis correspond to the signs of any of the various forms of intestinal obstruction that occur during the newborn period (Table 10–1) [225]. Prominent early findings include abdominal distention; progressive and frequently bilious vomiting; and evidence of pain, as manifested by persistent crying, irritability, or “colic.” Clinical features such as diarrhea, constipation, lethargy, or refusal to feed may also be evident, but are too nonspecific to be helpful in establishing a diagnosis. The presence or absence of fever is an unreliable sign in appendicitis as in other forms of neonatal infection; temperature has been recorded as normal or subnormal in more than 50% of newborns with this condition. Abdominal tenderness and guarding are inconsistent findings and, when present, are rarely localized to the appendiceal area. Physical signs of sufficient specificity to indicate acute inflammation of the appendix are generally absent until late in the course of the illness, when gangrene and rupture may result in the formation of a localized intra-abdominal abscess or cellulitis of the anterior abdominal wall. Erythema or edema, or both, of the right lower quadrant has been observed in several patients. The presence of this finding, particularly when accompanied by a palpable mass in the right iliac fossa, indicates bowel perforation with peritonitis and should suggest a preoperative diagnosis of NEC or appendicitis (see “Necrotizing Enterocolitis”).

DIAGNOSIS

The diagnosis of appendicitis in a neonate is usually determined at surgery performed for evaluation of abdominal distention and suspected peritonitis. With the

high incidence of prematurity associated with early appendicitis, bowel perforation from NEC has been a common preoperative consideration [206]. The two conditions can coexist, and in some cases, the appendix may participate in the process of ischemic necrosis and perforation [205,213].

Laboratory studies are of little value in establishing a diagnosis of appendicitis in a newborn. White blood cell counts of less than 10,000/mm³ were found in 10 of 30 infants. Urinalyses are usually normal, although ketonuria, which reflects diminished caloric intake; hematuria; and proteinuria may be seen. Because bacteremia may accompany appendiceal perforation and peritonitis, a blood culture and evaluation for metastatic infection with lumbar puncture and chest radiography should be performed. The value of paracentesis for diagnosis of bowel perforation and peritoneal infection is discussed later (see “Necrotizing Enterocolitis”).

Radiologic examinations are occasionally helpful, but in most cases serve only to confirm a clinical impression of small bowel obstruction. The presence of an increased soft tissue density displacing loops of intestine from the right iliac fossa generally indicates appendiceal perforation with abscess formation and is perhaps the most reliable sign of acute appendicitis in the neonate. Extraluminal gas may be localized briefly to the right lower quadrant after rupture of the appendix [211]. The rapid development of an extensive pneumoperitoneum obscures the site of origin of the escaping gas in most infants within a short time [226]. Ultrasonography may aid in detection of a periappendiceal abscess [83], but is not helpful in establishing an early diagnosis of appendicitis because it lacks sensitivity and specificity.

PROGNOSIS

The overall mortality rate from appendicitis in the newborn is high, but is improving. Eight of the newborns in the last 12 reported cases have survived, whereas of 60 infants with this condition for whom the outcome was recorded, 38 (64%) died. Survival was unrelated to birth weight. Among factors responsible for mortalities, three seem to be of primary importance: delay in diagnosis, a high incidence of perforation, and the rapid onset of diffuse peritonitis after appendiceal rupture.

Perforation has been identified at surgery or autopsy in 70% of newborns with acute appendicitis. The relative frequency of this complication has been attributed to delays in establishing a diagnosis and to certain anatomic features of the appendix in young infants that predispose it to early necrosis and rupture. These features include a meager blood supply that renders the organ more vulnerable to ischemia; a cecum that is relatively smaller and less distensible than that of adults, forcing a greater intraluminal pressure on the appendix; and the presence of a thin muscularis and serosa that readily lose their structural integrity under the combined effects of ischemia and increased internal pressure [163,181,182,191].

After the appendix ruptures, infants are unable to contain infection efficiently at the site of origin. Rapid dissemination of spilled intestinal contents produces a diffuse peritonitis within hours because of the small size

TABLE 10–1 Signs of Intra-abdominal Neonatal Appendicitis in 55 Infants

Sign	Incidence (%)
Abdominal distention	90
Vomiting	60
Refusal of feedings	40
Temperature $\geq 38^{\circ}$ C	40
Temperature 37–38° C	30
Temperature $\leq 37^{\circ}$ C	30
Pain (crying, restlessness)	30
Lethargy	30
Erythema/edema of right lower quadrant	25
Mass in right lower quadrant	20
Diarrhea	20
Passage of bloody stools	20

of the infant's omentum, which fails to provide an efficient envelope for escaping material; the relatively longer and more mobile mesenteries, which favor widespread contamination; and the small size of the peritoneal cavity, which also permits access of infected material to areas distant from the site of perforation [160,166,181,182]. Peritonitis, accompanied by sepsis and by the massive outpouring of fluids, electrolytes, and proteins from inflamed serosal surfaces, is generally the terminal event in neonatal appendicitis. Deterioration of the infant's condition is often extremely rapid; failure to recognize the underlying illness and to institute appropriate therapy promptly is inevitably followed by a fatal outcome.

TREATMENT

Surgical intervention is essential for survival of young infants with appendicitis. Because vomiting, diarrhea, and anorexia frequently accompany this condition, restoration of fluid and electrolyte balance is a major factor in ensuring a favorable outcome. Loss of plasma into the bowel wall and lumen of the dilated intestine may require additional replacement with whole blood, plasma, or an albumin equivalent. Optimal preparation often necessitates a delay of several hours, but is a major determining factor in the success of any surgical procedure done during the neonatal period.

The preoperative use of antibiotics has been recommended in infants with intestinal obstruction to achieve therapeutic blood levels of drug before the time of incision and possible contamination [167,227,228]. Perforation, fecal spillage, and peritonitis occur so early in the course of neonatal appendicitis that almost all infants with this condition require treatment before surgery. After the diagnosis of gangrenous or perforated appendicitis has been established and surgery has been performed, parenteral antibiotic therapy should be continued for a minimum of 10 days. The combination of clindamycin (or metronidazole), gentamicin (or extended-spectrum cephalosporins), and ampicillin provides adequate coverage against most enteric pathogens and can be used for initial empirical therapy. Alternatively, β -lactam and β -lactamase inhibitor combinations such as piperacillin and tazobactam or carbapenem antibiotics (e.g., imipenem or meropenem) can be used alone for broad coverage of enteric bacteria, *Pseudomonas* species, and anaerobic bacteria. Until the infant is able to tolerate alimentation, careful attention to postoperative maintenance of body fluids, electrolyte balance, nutrition, and correction of blood and plasma losses is vital to survival (see "Peritonitis" and "Necrotizing Enterocolitis").

PERITONITIS

Peritonitis in the newborn is most commonly associated with perforation of the gastrointestinal tract, ruptured omphaloceles, or wound infections that follow abdominal surgery [229,230]. It has been estimated that 20% to 40% of gastrointestinal surgical problems in the neonatal period are complicated by bacterial peritonitis (see "Necrotizing Enterocolitis") [185,231]. At pediatric surgical centers in the United States [231–233], Great Britain

[230,234], Hungary [235], Germany [236,237], France [238], and Zimbabwe [239], 1 to 10 cases per year have been reported in retrospective analyses of peritonitis diagnosed during the first month of life. Among almost 3000 infants admitted to a neonatal intensive care unit (NICU) in Liverpool in 1981–1982, there were 6 cases of peritonitis, all from NEC perforation of the gastrointestinal tract [240]. Peritonitis was present in 4 (all of low birth weight) of 501 infants on whom consecutive autopsies were performed from 1960–1966 at St. Christopher's Hospital for Children in Philadelphia. These cases represented approximately 3% of all patients with inflammatory lesions associated with death in this age group [241]. Potter [15] considered the peritoneum "one of the most frequent points of localization" in infants dying with sepsis. Among 121 such infants autopsied from 1976–1988 at St. Mary's Hospital in Manchester, England, generalized peritonitis was found in 9 (7.4%) [230].

A preponderance of boys (2.5:1) [190,239,240] and a high incidence of prematurity (33%) [231,234–236] have been found in unselected series of infants with this condition. These features are probably less a characteristic of bacterial peritonitis in the newborn than of the primary surgical and septic conditions that are responsible for its occurrence (particularly NEC). There seems to be a female preponderance among newborns with primary peritonitis [235,242]. A high incidence of congenital anomalies not involving the intestinal tract has also been observed among neonates with peritonitis [231,236,243,244].

MICROBIOLOGY

The condition that permits bacteria to colonize the peritoneal surface determines the nature of the infecting organisms. Most infants in whom rupture of a viscus and fecal spillage have caused peritonitis are infected by bacteria considered to be part of the normal enteric microflora; however, prior use of antimicrobial agents and colonization patterns within a nursery are important factors in determining which organisms predominate. Although a mixed flora of two to five species can often be recovered [243], single isolates have been reported in a third of infants with peritonitis [245,246]. The predominant aerobic organisms usually include *E. coli*, *Klebsiella* species, *Enterobacter* species, *Pseudomonas* species, *Proteus* species, coagulase-negative and coagulase-positive staphylococci, ungrouped streptococci, *Enterococcus*, and *Candida* [230,231,237,239,246–248].

Techniques adequate for the isolation of anaerobic organisms have been used infrequently. In a series of 43 consecutive infants with gastrointestinal perforation and bacterial growth from peritoneal fluid, a mixed aerobic-anaerobic flora was isolated with *Bacteroides* species as the predominant anaerobes [243]; remaining specimens grew aerobic or facultative organisms alone, and no culture yielded only anaerobes. In that series and others, the same organisms were frequently isolated from the peritoneal cavity and blood [232,243,244,248].

In contrast to fecal flora isolated from infants with gastrointestinal perforation, gram-positive organisms predominated among neonates with "idiopathic primary peritonitis." This condition is caused by sepsis in most

cases, but it also has often been associated with omphalitis. Specific organisms in one representative series included *S. pneumoniae* (three cases); ungrouped β -hemolytic streptococcus (three cases); and *S. aureus*, *Pseudomonas* species, and *E. coli* (one case each) [231]. Gram-positive cocci were also the major isolates in other series of peritonitis associated with hematogenous dissemination of organisms or extension from a peripheral suppurative focus.* Many of the cases caused by *S. aureus* occurred before the advent of antibiotics or during the worldwide pandemic of staphylococcal disease in the late 1950s, whereas streptococci, particularly GBS, have been a prominent cause in recent years [242,249–253].

Rarely, peritonitis may be caused by *Candida albicans* in pure culture or mixed with gram-negative enteric organisms [230,254]. Because clinical findings in this condition are not different from the findings of bacterial peritonitis, the diagnosis is usually established by blood or peritoneal fluid culture. Severe hypothermia has been described as a possible predisposing cause of bowel perforation and peritonitis owing to *Candida* [255]. In addition to well-recognized risk factors, such as prematurity, antibiotic therapy, and parenteral nutrition with deep venous catheters, NEC may be a significant risk factor for systemic candidiasis, in which it was observed in 37% of 30 infants [256]. Only a single infant in this series had a positive culture for *Candida* species from the peritoneum, however. Peritoneal catheters or peritoneal dialysis may also be a risk for direct inoculation of *Candida* organisms into the peritoneal space, which occurred in 1 of 26 children [257] (see Chapter 33).

PATHOGENESIS

Acute bacterial peritonitis may occur whenever bacteria gain access to the peritoneal cavity, through intestinal perforation, by extension from a suppurative focus, or by the hematogenous route. Intrauterine peritonitis owing to *L. monocytogenes* has been reported [230]; however, cases of “fetal peritonitis” described in earlier reports were actually examples of meconium peritonitis caused by intrauterine intestinal perforation [258,259]. Although bacterial colonization of the gastrointestinal tract in the first days of life may lead to infection in this condition, it is an aseptic peritonitis in its initial stages. A similar condition with focal perforation of the ileum or colon occurring postnatally has been described in infants with very low birth weight. Blue-black discoloration of the abdomen, caused by meconium staining of the tissues of the underlying skin, may be the first physical finding in these infants. Clinical, radiographic, and histopathologic evidence of infection or inflammation was notably absent in most cases [228].

Conditions that predispose to neonatal peritonitis are outlined in Table 10–2. Among almost 400 newborns with peritonitis studied from 1959–1978, perforation of the intestinal tract was responsible for 72% of cases, with ruptured omphaloceles or gastroschisis responsible for 12%, hematogenous dissemination or “primary” peritonitis

TABLE 10–2 Etiology of Bacterial Peritonitis in the Neonatal Period

Gastrointestinal perforation [191,231,232,235–240,243,245,249,262,263]
Necrotizing enterocolitis
Ischemic necrosis
Spontaneous focal gastrointestinal perforation [233,246,249,261,262,264]
Volvulus
Hirschsprung disease
Meconium ileus (cystic fibrosis) [231,266]
Postoperative complications
Congenital anomalies
Internal hernia
Catheter-associated vascular thrombosis [230]
Indomethacin therapy (enteral or parenteral) [267,268]
Trauma
Feeding tubes [269]
Rectal thermometers, catheters, enema [275–279]
Intrauterine exchange transfusion [231,273]
Paracentesis of ascites fluid
Meconium peritonitis with postnatal bacterial contamination [243,259,260]
Peptic ulcer: stomach, duodenum, ectopic gastric mucosa
Acute suppurative appendicitis
Infection
<i>Shigella</i> or <i>Salmonella</i> enterocolitis [274–276]
Congenital luetic enteritis with necrosis [58]
Ruptured omphalocele or gastroschisis
Postoperative: anastomotic leaks, wound dehiscence, wound contamination
Primary peritonitis
Prenatal sepsis: listeriosis, syphilis [58], tuberculosis [46–49]
Neonatal sepsis [10,33,231,232,240,252–254,263,278]
Suppurative omphalitis [26,72,240,244,250,251,263,277]
Transmural migration (theory) [265,279]

responsible for 12%, and omphalitis and postoperative complications responsible for 2% each.* In a comprehensive review of neonatal peritonitis, Bell [229,243] described common sites and causes of gastrointestinal perforation and their relative frequencies (Figs. 10–1 and 10–2).

No cases of neonatal peritonitis in recent years have been attributed to microorganisms entering the peritoneal cavity by traversing the bowel wall through the lymphatics or within macrophages (i.e., transmural migration). Evidence for the existence of this pathway is theoretical and is based primarily on retrospective analyses of pathologic data in humans with supporting observations made on laboratory animals [262,263]. Further confirmation is necessary before the transmural pathway can be accepted as an established source of peritoneal colonization by bacteria.

*References [10,26,33,71,242,244–253].

*References [190,229,230,236,260,261].




FIGURE 10-1 Causes of perforation in 60 neonates. duo. perf., duodenal perforation; strict., stricture. (From Bell M^J. *Peritonitis in the newborn—current concepts. Pediatr Clin North Am* 32:1181, 1985.)




FIGURE 10-2 Sites of perforation in 60 neonates. (From Bell M^J. *Peritonitis in the newborn—current concepts. Pediatr Clin North Am* 32:1181, 1985.)

CLINICAL MANIFESTATIONS

Neonatal peritonitis is a disease primarily of the first 10 days of life; numerous infants have evidence of peritoneal infection within the first 24 hours [231,232,239,244]. An analysis of etiologic factors responsible for peritonitis in the newborn provides a ready explanation for this observation (see Table 10-2). Most cases of NEC [243,264] and spontaneous gastric perforation [232,243,248,260] occur within the 1st week. Ruptured omphaloceles and gastroschisis often develop early infections, and in infants with congenital obstruction, the onset of alimentation during the first 12 to 24 hours accentuates distention and ischemic necrosis of the bowel wall, which leads to early intestinal perforation. Exchange transfusions are performed most frequently within the first 1 or 2 days of life and may be followed by enterocolitis within 4 to 24 hours in infants in whom perforation ultimately occurs [265,266]. Neonatal sepsis, with potential peritoneal seeding of microorganisms, is more frequent during the first 48 hours of life than during any subsequent period [267].

The various signs and symptoms present in a young infant with peritonitis were summarized most succinctly by Thelander [248] in 1939:

“The little patient looks sick. He is cyanotic; the respirations are rapid and grunting; the abdomen is distended, and the abdominal wall, the flanks and the scrotum or vulva are usually edematous. Frequently brawny induration of the edematous area, which may resemble erysipelas, is also present. Food is taken poorly or not at all. Vomiting is frequent and persistent. The vomitus contains bile and may contain blood. The stools are either

TABLE 10-3 Signs of Bacterial Peritonitis in the Neonate*

Sign	Incidence (%)
Abdominal distention	85
Shock	80
Vomiting	70
Constipation	60
Hypothermia	60
Respiratory distress	55
Fever	15
Diarrhea	15

*Data are based on patients described in references 232, 240, and 243. Redness, edema, and induration of the anterior abdominal wall, noted in only one series [243], are also recognized as characteristic signs.

absent or scant; some mucus or blood may be passed. The temperature may be subnormal, but varying degrees of fever have been reported. The blood count is of little or no value. The hemoglobin content may be very high, which probably indicates only dehydration. The leukocytes may or may not respond with a rise.”

Although the review by Thelander [248] was limited to neonates with perforation of the intestinal tract, subsequent reports have corroborated the presence of these findings in infants with peritonitis resulting from a wide variety of causes.* Not all of the symptoms described may be encountered in any one patient; however, some are always present (Table 10-3).

The large overlap between signs of neonatal peritonitis and sepsis can make it difficult to differentiate the two on the basis of clinical findings. Signs of intestinal obstruction such as abdominal distention and vomiting, which are seen in 10% to 20% of newborns with sepsis [9,17,244], may reflect a coexistent unrecognized peritonitis. Because the early use of antibiotics often cures hematogenous peritonitis in infants with septicemia, the diagnosis may be missed in infants who survive. Peritonitis unassociated with perforation was found at postmortem examination in 4 of 20 infants with sepsis in 1933 [10], 9 of 73 premature infants dying with septicemia from 1959-1964 [9], and 9 of 121 infants with septicemia dying from 1976-1988 [229].

DIAGNOSIS

Ultrasonography [83] or abdominal radiographs taken in the erect and recumbent positions showing free intraperitoneal fluid can be helpful in the diagnosis of peritonitis, and sometimes the only evidence of perforation is apparent on these imaging studies. Absence of definition of the right inferior hepatic margin, increased density of soft tissue, and the presence of “floating” loops of bowel have been recorded as positive signs of ascites [226,268]. Diagnostic paracentesis can be useful in determining whether the fluid is caused by bacterial peritonitis [242,251,269,270], hemo-peritoneum, chylous ascites [271], or bile peritonitis [272].

*References [190,231,233-235,239,243,248].

The left lateral (“left-side down”) decubitus film is of great value in showing small amounts of intraperitoneal gas [243]. Although pneumoperitoneum can be caused by mediastinal air dissecting from the chest into the abdomen [273,274], free gas in the peritoneal cavity usually indicates intestinal perforation. An associated pneumatosis intestinalis should suggest the diagnosis of NEC, but is not specific for this condition. Several patterns of intraperitoneal gas distribution have been described [83,265-276]: the air-dome sign, falciform ligament sign, football sign, lucent-liver sign, saddlebag sign, and gas in the scrotum. Absence of a gastric air-fluid level on an erect abdominal radiograph, with a normal or decreased amount of gas in the small and large bowel, strongly favors a diagnosis of gastric perforation [276]. This finding is almost always accompanied by pneumoperitoneum.

In equivocal cases, metrizamide contrast studies of the bowel can be helpful in establishing a diagnosis of intestinal perforation [245,273]. Serial abdominal transillumination with a bright fiberoptic light is a useful bedside method for the early detection of ascites or pneumoperitoneum in the newborn [277].

Failure to show free air in the peritoneal cavity does not rule out a diagnosis of perforation, particularly if air swallowing has been reduced or prevented through orotracheal intubation, nasogastric suction, or use of neuromuscular blocking agents [269,273,278]. In some cases, the amount of gas in the bowel lumen is so small that even if perforation occurs, the gas could escape detection. Alternatively, small leaks may become walled off and the free air reabsorbed [273,279,280]. In three large series of infants with peritonitis in whom a patent site of perforation was found at surgery, pneumoperitoneum was absent in 35% to 75% [231,243,244].

Radiographic evidence of intestinal obstruction, although a common cause or consequence of peritonitis, lacks sufficient specificity to be a consistent aid to diagnosis. A diffuse granular appearance of the abdomen, with one or more irregular calcific densities lying within the bowel lumen or in the peritoneal cavity, should suggest a diagnosis of meconium peritonitis with possible bacterial superinfection [259].

PROGNOSIS

Prematurity, pulmonary infections, shock, and hemorrhage related to perforation of the intestinal tract, sepsis, and disseminated intravascular coagulopathy are often the factors responsible for the death of neonates, who may concurrently have peritonitis diagnosed at surgery or at postmortem examination. For this reason, case-fatality rates often represent the mortality rate among newborns dying with, rather than because of, infection of the peritoneal cavity [230,236,243].

Before 1970, the incidence of fatalities was exceedingly high when peritonitis was associated with gastrointestinal perforation; mortality rates of 70% were observed in large series.* Heightened awareness of conditions associated with perforation, more rapid diagnosis, and improved

*References [190,231,234,235,237-239,244,280].

surgical management have led to a doubling of survivors in recent years [236,237]. The cause of perforation seems to influence the likelihood of survival, with spontaneous gastric perforation having the lowest mortality rate (10%) and perforation of the duodenum caused by a feeding tube the highest mortality rate (50%); NEC (40%) and all other causes (25%) occupy intermediate positions [243].

As survival rates have improved, the number of nonlethal complications after perforation has increased proportionally. In one review, two thirds of surviving infants had significant postoperative complications pertaining to infection (e.g., bacteremia, wound infection, intra-abdominal abscess) or gastrointestinal tract dysfunction (e.g., esophageal reflux, obstruction, stomal stenosis) [243]. Secondary surgical procedures to correct these problems were required in more than half of the infants. Parenteral hyperalimentation for nutritional support during the recovery period was required in 60% of infants.

The mortality rate among neonates with peritonitis from causes other than perforation of the bowel, such as sepsis [190,231,235,239], omphalitis [239,281], or a ruptured omphalocele [231,236,239], although high in the past, has not been reassessed in recent years [243]. Early diagnosis and institution of appropriate surgical therapy are major factors in reducing the mortality rate [243]. It has been shown that infants operated on within 24 hours after the onset of symptoms have survival rates almost double the rates of infants operated on between 24 and 48 hours and 2½ times higher than the rate for infants whose surgery was delayed more than 48 hours [244]. Factors with an apparent adverse influence on prognosis include low birth weight [231,236,239,243,261], low birth weight for gestational age [229], congenital malformations [236], male sex [230], and initial serum pH of less than 7.30 [230].

TREATMENT

The treatment of bacterial peritonitis is directed primarily toward correction of the causative condition [230]. Careful attention to preoperative preparation of the infant is essential to survival. As soon as bowel obstruction or perforation is diagnosed, continuous nasogastric suction should be instituted for decompression and prevention of aspiration pneumonitis. Diagnostic needle paracentesis is also useful for relief of pneumoperitoneum and may facilitate exchange of gas by reducing the intra-abdominal pressure. Shock, dehydration, and electrolyte disturbances should be corrected through parenteral administration of appropriate electrolyte solutions, plasma, or plasma substitutes. If blood is discovered in fluid recovered by gastric suction or abdominal paracentesis, use of whole blood, packed red blood cells, or other fluids may be necessary to correct hypovolemia. Persistent bleeding must be evaluated for disseminated intravascular coagulation or thrombocytopenia, or both, and treated accordingly. Hypothermia, which frequently accompanies neonatal peritonitis, should be corrected before induction of anesthesia. Infants who are unable to tolerate oral or tube feedings within 2 or 3 postoperative days should be started on parenteral hyperalimentation.

If a diagnosis of peritonitis is established at the time of paracentesis or surgery, aerobic and anaerobic cultures of peritoneal contents should be taken before initiation of antibiotic therapy. Parenteral administration of a combination of gentamicin or an extended-spectrum cephalosporin and clindamycin and ampicillin should be continued for 7 to 10 days [215,228]. Other antibiotics that provide a broad spectrum against enteric organisms, *Pseudomonas* species, enterococci, and anaerobic organisms include β-lactam and β-lactamase inhibitor compounds and carbapenems. In the event of a poor clinical response, culture and susceptibility studies of the infecting organisms should be used as guides for modifying therapy.

Leakage of intestinal contents sometimes results in formation of a localized abscess, rather than contamination of the entire peritoneal cavity. Management of infants with such an abscess should include antimicrobial therapy and surgical drainage of the abscess by the most convenient route.

NECROTIZING ENTEROCOLITIS

NEC with necrosis of the bowel wall is a severe, often fatal disease that has been occurring with increasing frequency in recent years. The average annual NEC mortality rate is 13.1 per 100,000 live births; black infants, particularly boys, are three times more likely to die of NEC than white infants, and mortality rates are highest in the southern United States [280,282–291]. NEC occurs in about 5% of infants admitted to NICUs; however, the incidence varies widely among centers and from year to year at the same institution [292–305]. NEC predominantly affects infants with birth weights less than 2000 g [282,284,298,213,306–312]; in several series, the frequency in infants weighing less than 1500 g was 10% to 15%.* Only 5% to 10% of all cases of classic NEC occur in term infants [291,318–320]. It has been stated that occasional cases of NEC may be the price to be paid for the benefits of modern neonatal intensive care [321].

Although most reports of NEC emanate from the United States, Canada, and Great Britain, the condition occurs worldwide in countries maintaining NICUs. In a review of NEC in two NICUs of academic centers, NEC increased the risk for death (odds ratio 24.5), infections (odds ratio 5.7), and need for central intravenous line placement (odds ratio 14.0) [322]. Infants with surgical and medical NEC had lengths of stay of 60 days and 22 days greater than infants without NEC with additional costs of \$186,200 and \$73,700, resulting in additional hospital charges of \$216,666 per surviving infant.

NEC affects an estimated 7% to 14% of all low birth weight preterm infants [323] and is a leading cause of death and morbidity in the NICU with no change in incidence in the past 20 years [324,325]. More recent reports have estimated that about 9000 cases occur annually in the United States with a case-fatality rate of 15% to 30% [326].

*References [282–285,299,300,304,311,313–317].




FIGURE 10-3 Pathogenesis of mucosal injury leading to necrotizing enterocolitis (NEC). (Adapted from Walsb MC, Kliegman RM. Necrotizing enterocolitis. *Pediatric Basics* 40:5, 1985.)

PATHOLOGY AND PATHOGENESIS

Bowel wall necrosis of variable length and depth is the characteristic feature of NEC, with perforation in one third of affected infants generally in the terminal ileum or cecum, where microcirculation is poor.* The pathogenesis of NEC is not established, but most investigators agree that the initiating event is some form of stress to the immature gastrointestinal tract, which leads to disruption of the mucosal barrier, bacterial invasion and proliferation, and gas formation within the bowel wall (Fig. 10-3) [213,298,306,330]. Surgical specimens from early stages of the disease show mucosal edema, hemorrhage, and superficial ulceration with very little inflammation or cellular response. By the 2nd or 3rd day, after progression to pneumatosis and transmural necrosis of the bowel wall, bacterial proliferation and the acute inflammatory reaction become more prominent [287,309].

There has been much investigation and little agreement on the importance of various perinatal events in the causation of NEC [213,298,306,331]. Except for immaturity and possibly polycythemia, other factors originally thought to predispose to NEC have been found on further study to occur with equal frequency in control populations of infants.* Maternal complications of pregnancy, labor, and delivery and neonatal respiratory distress syndrome are thought to be unrelated to the development of NEC, whereas evidence linking NEC to birth asphyxia, hypotension, hypothermia, use of vascular catheters, exchange transfusion, feeding history, abnormalities of gut motility, neonatal achlorhydria, and the presence of patent ductus arteriosus is often contradictory. Each of these conditions, singly or together, may act as a stress leading

to mucosal injury, but none has been consistently associated with NEC [334,335]. NEC has occurred among apparently healthy infants with no known predisposing risk factors [282,280,318,330]. Several studies (discussed later) suggest that early feeding of premature neonates may play a causal role in NEC. Dvorak and coworkers [330] showed that maternal milk may be protective compared with artificial formulas in a neonatal rat model of NEC; similar to human NEC, artificial feeding of maternal milk reduced the incidence and severity of NEC injury, and IL-10 expression was significantly increased when neonates were fed maternal milk.

Some epidemiologic observations suggest that NEC is an infectious contagious disease of nosocomial origin. The temporal clustering of cases at institutions, the association of some outbreaks with single infectious agents or alterations in bowel flora, and the possible beneficial effects of breast-feeding, oral nonabsorbable antibiotics, or infection control measures in reducing the incidence of disease suggest a possible nosocomial cause. The evidence linking NEC to a specific infectious agent is often circumstantial or open to alternative interpretation. Evidence suggests that NEC is the end response of the immature gastrointestinal tract to multiple factors acting alone or in concert to produce mucosal injury, with colonization or invasion by the microorganisms representing only one part of the continuum of that disease process [266,272,273,280].

MICROBIOLOGY

Descriptions of sporadic outbreaks of NEC in NICUs have led to a search for transmissible agents, including bacterial, viral, and fungal pathogens [341a].* Predominance

*References [243,287,280,309,318,327-329].

*References [213,299,304,310,311,320,332,333].

*References [288,298,306,213,330,336-341].

of a single organism in stool, blood, bowel wall, or peritoneal cavity of infants during epidemics of NEC has implicated numerous agents, including *Klebsiella* species [285,295,312,329,342–344], *Enterobacter* species [345], *E. coli* [346,347], *Pseudomonas* species [348,349], *Salmonella* species [350], *S. epidermidis* and other coagulase-negative staphylococci [351], *S. aureus* [352], rotavirus [353,354], coronavirus [355], coxsackievirus B2 [356], and *Torulopsis glabrata* [357].

The analogous pathology of necrotizing enteritis caused by *Clostridium septicum* [36,72] and *Clostridium perfringens* in domestic animals [297,344,358], older children, and adults [359] favored suggestions that *Clostridium* species might act as a primary pathogen in NEC.* Several reports provided evidence that *C. perfringens* [297,361–364], *Clostridium difficile* [333,365], or *Clostridium butyricum*, acting alone [366,367] or synergistically with *Klebsiella* [368], was able to evoke NEC. Subsequent studies indicated, however, that these species were often acquired from the nursery environment [369,370] and could frequently be recovered from healthy neonates [359,370–376]. Clostridial cytotoxin, which had been recovered from the stool of infants involved in an outbreak of NEC [333,365], has also been found in the stool of 90% of normal infants [333,364,369,375–377]. The role of *Clostridium* species in NEC remains unclear [341,360].

The δ -toxins, hemolysins of coagulase-negative staphylococci [378] and *S. aureus*, have also been proposed as possible primary toxins capable of producing NEC in infants. Frequent colonization by δ -toxin staphylococci and higher levels of toxin production by associated strains causing NEC [379] and one outbreak with δ -toxin-producing *S. aureus* strains have been reported [352]. Prospective studies have documented significant shifts in

aerobic bacterial bowel flora within 72 hours before onset of clinical NEC [380]; the observed shift results from pre-clinical changes in the intestinal environment. This shift suggests that bacteria isolated at the time of onset were present because of possible intraluminal changes and are not directly involved in NEC.

Pending further experimental or epidemiologic observations, the weight of evidence indicates that although bacteria or bacterial toxins may play a primary or secondary role in the pathogenesis of NEC, the occasional association of this condition with a single organism probably reflects patterns of intestinal colonization prevalent in the nursery at the time of an outbreak [213,298,306,330]. Despite intensive efforts to identify a specific infectious agent or toxin in the cause of NEC, convincing reports implicating the same pathogen in more than one outbreak have not appeared [341].

CLINICAL MANIFESTATIONS

Signs of NEC usually develop in the first 7 days of life [291,318,381], and 50% or more of cases are recognized within 5 days of birth [282,309,312,382,383]. Small immature newborns often develop illness later, during the 2nd to 8th week [314,316], whereas low-risk term infants may become ill shortly after delivery, as early as the first 24 hours [319].

NEC is a disease with a wide spectrum of manifestations, ranging from a mild gastrointestinal disturbance to a fulminant course characterized by early bowel perforation, peritonitis, sepsis, and shock [306,384,385]. A staging system (Table 10–4) taking these clinical variations into account may be useful in guiding patient evaluation and therapy [306,386]. The apparent stage of disease for an individual infant usually can be defined on the 2nd day of illness. An infant who exhibits only mild systemic and

*References [213,298,306,330,341,360].

TABLE 10–4 Modified Bell Staging Criteria and Recommendations for Therapy for Necrotizing Enterocolitis

Stage	Signs			
	Systemic	Intestinal	Radiologic	Treatment
IA (suspected)	Temperature instability, apnea, bradycardia, lethargy	Elevated residuals, mild abdominal distention, emesis guaiac-positive stools	Normal, mild ileus	NPO, antibiotics for 3 days
IB (suspected)	Same as IA	Frank rectal blood	Same as IA	Same as IA
IIA (definite), mild	Same as IA and IB	Same as IB plus absent bowel sounds \pm abdominal tenderness	Dilation, ileus, pneumatosis intestinalis	NPO, antibiotics for 7–14 days if examination is normal in 24–48 hr
IIB (definite), moderate	Same as IIA with mild metabolic acidosis, mild thrombocytopenia	Same as IIA with definite abdominal tenderness \pm abdominal cellulitis or right lower quadrant mass	Same as IIA plus portal gas \pm ascites	NPO, antibiotics for 14 days
IIIA (advanced), bowel intact	Same as IIB plus hypotension, bradycardia, severe apnea, respiratory/metabolic acidosis, disseminated intravascular coagulation, neutropenia	Same as IIB plus peritonitis, marked tenderness, abdominal distention	Same as IIB with ascites	Same as IIB plus 200 mL/kg fluid, inotropic agents, assisted ventilation, paracentesis
IIIB (advanced), bowel perforated	Same as IIIA	Same as IIIA	Same as IIIA plus pneumoperitoneum	Same as IIIA plus surgery

NPO, nothing by mouth.

Adapted from Walsh MC, Kliegman RM. Necrotizing enterocolitis: treatment based on staging criteria. *Pediatr Clin North Am* 33:179, 1986.

intestinal signs 24 to 48 hours after onset is unlikely to develop a more serious illness [306].

The classic presentation of NEC includes a triad of abdominal distention, retention of gastric contents, and gastrointestinal bleeding.* These findings are often preceded or accompanied by signs consistent with sepsis, such as lethargy, poor feeding, temperature instability, apnea, and bradycardia. Diarrhea is variable, rarely observed in some series [306], but common in others [316,347]. Progression of bowel wall necrosis leading to perforation, peritonitis, and sepsis is reflected in deteriorating vital signs accompanied by persistent acidosis [387], clotting disorders, and circulatory collapse. Redness, induration, and edema of the anterior abdominal wall are commonly described in the advanced stages of NEC. In the absence of aggressive medical and surgical intervention, the course is rapidly downhill when late signs appear.

DIAGNOSIS

Radiographic signs of NEC are largely nonspecific [385], and interobserver variability in the interpretation of films is substantial [387]. Radiographic examination of the abdomen remains the most reliable aid, however, in establishing a diagnosis of NEC [282,302,309,387,388]. Ileus with generalized bowel dilation and abdominal distention are the earliest radiologic findings. Increasing distention, separation of loops by peritoneal fluid or edema of the bowel wall, a gasless abdomen, pneumatosis intestinalis, and hepatic or portal air occur as NEC worsens. A persistent single dilated loop of bowel remaining relatively unchanged in shape and position in serial films is strongly suggestive [389–391] but not diagnostic [390,392] of localized bowel ischemia with impending perforation.

If free air or ascites is absent on initial abdominal examination, supine and left lateral decubitus films should be obtained every 6 to 8 hours until improvement or definitive surgery or invasive diagnostic measures have ruled out the presence of perforation. When perforation occurs, it is usually within 1 day after diagnosis [327], but may be delayed for 5 or 6 days [393]. Although the presence of pneumoperitoneum [244,265,275] or intraperitoneal fluid generally indicates perforation, its absence does not exclude perforation [221,243,244]. In one study [327], only 63% of infants with NEC and proven perforation had free air, 21% had ascites, and 16% had neither free air nor ascites.

When plain films are normal or equivocal, other studies may be diagnostic. A metrizamide gastrointestinal series may show intestinal perforation or abnormalities of the bowel wall, mucosa, or lumen [245,273,394]. Real-time ultrasonography may reveal portal venous and hepatic parenchymal gas in standard radiographs [327,395,396]. Serial abdominal transillumination with a fiberoptic light has been described as a bedside method for early detection of ascites or pneumoperitoneum, although its sensitivity compared with standard radiographic methods has not been determined [277].

A rapid and direct means of establishing the presence of intestinal necrosis or perforation is by abdominal paracentesis [269,397,398]. This procedure is unnecessary in

infants to rule out NEC or in infants improving on medical therapy. The procedure is generally reserved for infants suspected, on the basis of clinical, radiographic, and laboratory findings, to have intestinal gangrene. When performed properly, paracentesis is safe and accurate, as described by Kosloske [399]: The abdomen is palpated to locate any masses or enlarged viscera. After an antiseptic skin preparation, a small needle (22-gauge or 25-gauge) is inserted carefully in the flank, at a 45-degree angle. It is advanced slowly and aspirated gently until free flow of 0.5 mL or more of peritoneal fluid is obtained. Any volume less than 0.5 mL is considered a dry tap and cannot be accurately interpreted. The color and appearance of the fluid are noted, and the fluid is transported immediately to the laboratory in an approved anaerobic container for Gram staining and for aerobic and anaerobic cultures. A positive paracentesis consists of brown fluid or bacteria on the unspun fluid.

The accuracy of paracentesis in determining the need for an operation is 90% to 95% [269,398]. False-positive results are rare; false-negative results are quite common. Patients with a dry tap should be closely observed under medical therapy with continuing serial paracenteses until indications for or against surgical intervention are clearly defined. Infants with a positive result should undergo exploratory surgery immediately.

Thrombocytopenia and disseminated intravascular coagulation are the most common hematologic complications [282,312,340,397,399–401], particularly in the presence of bowel gangrene or perforation [287,401,402]. Platelet-activating factor has been used to assist in the staging of NEC [403]; a cutoff level of 10.2 ng/mL had a positive predictive value of 100% in identifying infants with stage II or III NEC. Leukopenia and absolute granulocytopenia, apparently caused by margination of white blood cells rather than bone marrow depletion [404], also have occurred during early stages of the illness [399,400]. A low absolute granulocyte count persisting for 2 to 3 days is associated with a poor prognosis. Hemolytic anemia has been reported in association with NEC related to *C. perfringens* [363]. No consistent urinary abnormalities have been described for NEC, although increased lactate excretion, reflecting heightened enteric bacterial activity, may occur [405]. Increased amounts of fecal-reducing substances have been found in almost three fourths of formula-fed premature infants during early stages of NEC, before the onset of abdominal distention, poor feeding, or emesis [406]. Although not readily available, levels of growth factors in urine have been found to be much higher in children with stage II and III NEC [407]; such an analysis might identify children at higher risk of complications and the need for surgical intervention.

The evaluation of patients with NEC should include culture of blood and, when appropriate and clinically safe, cerebrospinal fluid, urine, and stool. The likelihood of bacteremia accompanying NEC depends on the severity of bowel involvement; the reported incidence has ranged from 10% to 67% among symptomatic infants. Combined data from several large studies showed positive blood cultures in about one third of newborns with NEC [287,288,339,393]. The usual organisms have been *E. coli*, *Klebsiella* species, *S. aureus*, and *Pseudomonas*

*References [292,303,316,318,320,329,337,342,391,396].

species, whereas enterococci and anaerobic bacteria were isolated occasionally. A spectrum of organisms similar to those causing sepsis have been isolated from the peritoneal fluid [287,295,316,340,393]. Meningitis may accompany bacteremia, occurring in approximately 1% of NEC cases [319,408].

TREATMENT

Early and aggressive treatment must be initiated for any infant suspected to have NEC [298,306]. The modified Bell staging system of NEC may guide diagnostic studies, management, antibiotics, and surgical consultation and intervention (see Table 10-4). Umbilical catheters should be removed whenever possible, oral feedings should be stopped, and nasogastric tube drainage should be instituted. Fluid and electrolyte deficits and maintenance require rigorous attention; blood, plasma, or colloid infusions are often necessary for volume expansion and maintenance of tissue perfusion.

After appropriate cultures are obtained, parenteral antibiotic therapy should be started with clindamycin and gentamicin or an extended-spectrum cephalosporin and ampicillin. In nurseries where coagulase-negative staphylococcal colonization or infection is prevalent, initial therapy with vancomycin may replace ampicillin [409]. β -lactam and β -lactamase inhibitor combinations (e.g., piperacillin plus tazobactam) can replace gentamicin, ampicillin, and clindamycin, covering anaerobic, gram-negative enteric aerobic, and many gram-positive pathogens. Gentamicin and vancomycin dosages should be modified as necessary on the basis of serum levels. Despite anecdotal evidence that oral nonabsorbable aminoglycosides prevent gastrointestinal perforation in infants with NEC [410], later controlled studies did not corroborate this finding [411]; their use is not routinely recommended. The need for inclusion of clindamycin to provide activity against anaerobic bacteria in the management of NEC has been questioned [412], based on the observation that anaerobic bacteria are late bowel colonizers, present in relatively much lower numbers in infants than in adult subjects, and that anaerobes other than *Clostridium* species have been infrequently identified in NEC.

After immediate treatment has been started, follow-up studies should be instituted. These include serial examinations with measurement of abdominal girth; testing of stools for blood; levels of serum electrolytes, blood glucose, and arterial blood gases; complete blood cell count and platelet count; urine-specific gravity; and supine and left lateral decubitus abdominal radiographs. These tests should be considered every 6 to 8 hours until the infant's clinical condition stabilizes. Attention to vital functions should be provided as necessary on the basis of clinical, laboratory, or radiographic studies. Parenteral nutritional support through a central or peripheral vein must be started as soon as possible.

Early recognition and prompt initiation of medical therapy may reduce the need for surgery. Generally accepted criteria for surgical exploration are a deteriorating clinical condition despite appropriate medical therapy, signs of peritonitis, presence of free air within the abdomen, or a positive paracentesis result. The principles of surgical preparation and management have been discussed by

several investigators [289,398,413,414]. In addition to laparotomy with removal of necrotic bowel, closed peritoneal drainage has been proposed as an alternative in very small infants, with a resultant survival of more than 50% [415].

PREVENTION

The first observations implicating bacterial proliferation as a factor in pathogenesis of NEC prompted efforts at suppression of gut flora with topical antibiotics in the hope of preventing NEC. Attempts to prevent NEC by giving oral kanamycin or gentamicin prophylactically in the first hours of life, before any signs of bowel involvement are recognized, have generated contradictory data. In controlled clinical trials, a significant reduction in the incidence of NEC in treated premature infants was shown in some trials [313,416-419], whereas in others, investigators were unable to show any protective effect [419,420]. Studies of vancomycin [421] have shown a significant reduction in NEC in high-risk infants. Previous studies revealed selective growth of resistant organisms in bowel flora [313,420,422] and evidence of significant systemic absorption of aminoglycoside antibiotics [411,423,424], suggesting that oral aminoglycoside prophylaxis is not free of potential risks. Potential risk factors have not been examined in vancomycin trials, however. Until additional evidence is presented indicating clear-cut benefits from the use of oral aminoglycosides or vancomycin, it does not seem that either agent should be used routinely for prevention of NEC in premature infants. Epidemiologic evidence that early use of parenteral ampicillin and aminoglycoside therapy may delay or decrease the risk of NEC has not been confirmed in controlled studies [329].

Oral probiotics have been suggested to alter the bowel flora of the infant to reduce the incidence and severity of NEC. Infants fed breast milk and a product including *Lactobacillus acidophilus* and *Bifidobacterium infantis* had a reduced incidence and severity of NEC compared with infants fed breast milk alone [335]. A recent meta-analysis of 11 trials of probiotics concluded that the risk for NEC and death was significantly lower, but the risk for sepsis did not differ significantly from those neonates not receiving probiotics [335a]. These results seem to confirm the significant benefits of or probiotic oral supplements, with a 30% reduction in the incidence of NEC.

Excessive or accelerated feedings have been associated with increased frequency of endemic NEC [425], and some clinicians have recommended a schedule of slow advancement of daily feeding volumes limited to about 20 mL/kg/day. Infants with NEC are more likely to have been fed earlier, to have received full-strength formulas sooner, and to have received larger feeding volumes and increments, and stress and associated respiratory problems may make such infants more vulnerable to NEC [244,292-297,426]. Prior studies of the use of a feeding regimen employing prolonged periods of bowel rest in high-risk infants showed the regimen to be successful in preventing NEC in some nurseries [298], but totally without value in others [213,300]. Later studies added additional support for standardized feeding schedules in low birth weight infants (500 to 2500 g); all used maximum volumes no greater than 20 mL/kg/day, with feeding

beginning at 24 to 72 hours after birth, depending on birth weight and gestational age [307,308]. Synthetic formulas and breast milk have been successful.

Carrion and Egan [427] suggested that relative hypochlorhydria of the neonate may contribute to NEC and found that hydrochloric acid supplements (0.01 to 0.02 1.0 N HCl/mL of formula) significantly reduced NEC rates and lowered gastric pH. Additional studies have shown that standardized feedings begun at a median of 4 days after onset of NEC can be associated with an abbreviated time until institution of full enteral feedings, a reduced incidence of the use of central catheters and catheter infections, and ultimately a shorter hospital stay [428].

Many NEC “epidemics” in NICUs, lasting 2 weeks to 7 months, have been reported from centers worldwide [329,340,341,429,430]. Although the microbiologic agents associated with these outbreaks have varied, institution of strict infection control measures was often useful in bringing about a significant decrease in the incidence of NEC; the reasons for success are less clear. Results have been sufficiently impressive, however, to recommend that enforcement of bedside enteric precautions, together with cohorting of infants and staff, be instituted when two or more cases of NEC occur in a nursery [306,341,431].

The use of human breast milk has been claimed, largely on the basis of experimental evidence, to exert a protective effect against the development of NEC. There are no prospective, controlled studies showing any benefit from the feeding of colostrum or breast milk to human neonates. A study showing the protective effect of an orally administered IgA-IgG preparation suggests a possible way to provide benefits of high levels of functionally active antibodies in the gastrointestinal tract [432]. Orally administered probiotics have been suggested as a possible prophylactic measure in NEC based on an assumption that an accentuated and inappropriate inflammatory response to colonizing pathogenic flora in the premature gut plays a role in the initiation of NEC [433]. Bifidobacteria and lactobacilli are commonly found in breast-fed infants [434] and may be the reason in part for the protective effects of breast milk. A single large, multicenter, randomized controlled trial of *Bifidobacterium bifidum* and *L. acidophilus* was completed and suggested a lower incidence of NEC (stage II or greater) in infants receiving these probiotics [435]. This study showed a lower incidence of death or NEC in recipients of probiotics (4 of 217 probiotic infants versus 20/217 control infants). No adverse effects of the probiotics, such as sepsis, flatulence, or diarrhea, were noted. This large study seems to confirm several smaller studies with similar results [436,437,438].

PROGNOSIS

The mortality rate of NEC is difficult to determine because mild cases of suspected NEC are probably more common than is recognized [306,376,439]. In studies in which analysis has been limited to infants with “definite NEC,” mortality figures range from 20% to 40%.* Several longitudinal studies have shown a significant improvement in outcome [291,302,387,440]. A poor prognosis has

been linked with very low birth weight, associated congenital defects, bacterial sepsis, disseminated intravascular coagulation, intestinal perforation, and persistent hemodynamic or respiratory instability [311,336,381,393]. Surgical intervention, generally reserved for the sickest infants with more extensive bowel involvement, is also associated with higher mortality rates [311,381,387,398,439].

Infants who survive the acute phase of illness generally do well, although NEC may recur in 5% to 10% [291,316,440,441]. In addition to surgical complications (e.g., short bowel syndrome, anastomotic leaks, fistula formation), enteric strictures are probably the most common delayed complication in surviving infants, occurring in 4% to 20%. Usually found at sites of ischemia and necrosis in terminal ileum or colon [282,309,442], these strictures often become apparent within a few weeks, but may be delayed 18 months. When multiple strictures occur, the intervening normal bowel may form an enterocyst [388,443]. Clinically, strictures manifest as frequent episodes of abdominal distention, often with vomiting, obstipation, or hematochezia. Diagnosis is confirmed by gastrointestinal contrast studies. Surgery with removal of the stenotic site is necessary to effect a cure.

Long-term follow-up of low birth weight infants with severe NEC (i.e., Bell stages II and III) has documented higher rates of subnormal body weight (15% to 39%) and head circumference (30%) in addition to significant neurodevelopmental impairment (83%) [444]. Clinical observations suggest that infants with bowel resection for NEC are at increased risk of sepsis, occurring 1 week to 3 years (mean 4 months) later. Almost all had had a central venous catheter in place for parenteral nutrition at the time of infection. Enteric bacilli were responsible for more than 40% of bacteremias, whereas only 20% were caused by staphylococcal species, which are the usual causes of catheter sepsis. Several infants had two or more episodes of sepsis, and 2 of 19 died as a direct consequence of infection [427].

ENDOCARDITIS

Neonatal bacterial endocarditis, previously uncommon, has been recognized more frequently in recent years. About 60 cases that meet clinical and bacteriologic criteria sufficient to establish this diagnosis were reported in the literature before the mid-1980s [77,445–472]. The prolonged survival of critically ill infants, including infants with complex congenital heart disease, and the increased use of intravascular catheters, together with advances in the diagnostic sensitivity and availability of echocardiography, may be responsible for an increased recognition of endocarditis. In a 35-year review of 76 cases of endocarditis in children, 10% of patients were younger than 1 year; the youngest patient was 1 month old [473]. Of patients, 62 (83%) had congenital heart disease, and 77% had had prior surgery. Central venous catheters were additional significant risk factors. At the University of New Mexico in a level III nursery with 3200 to 3500 admissions annually, 12 cases of endocarditis occurred in children younger than 3 months [474]. Organisms isolated from these 10 cases included *S. aureus* (6 cases), *K. pneumoniae* (1 case), *Enterobacter cloacae* (2 cases),

*References [269,291,298,213,316–320,381,385,387].

Candida species (1 case), α -hemolytic streptococci (1 case), and coagulase-negative staphylococci (1 case). Three patients had congenital heart disease with early surgical intervention; all had surgically implanted catheters or intravenous access devices, one had NEC, and one had an associated osteomyelitis.

Etiologic agents of bacterial endocarditis in newborns have been identified by isolation from blood cultures or morphologic characteristics of organisms entrapped within valvular vegetations examined at autopsy. The causative organisms have included *S. aureus* in 36 infants*; streptococci in 6 infants [425,447,475]; *S. epidermidis* [455,456,476] and GBS [449,459,463] each in 5 infants; *S. pneumoniae* [447,475], *P. aeruginosa* [448,454], and *S. marcescens* [452,464] each in 2 infants; and *Neisseria gonorrhoeae* [447], *S. faecalis* [477], *Streptococcus salivarius* [71], and mixed α -hemolytic streptococcus, *K. pneumoniae*, and *P. mirabilis* [455] each in 1 infant. Despite widespread cardiovascular involvement associated with congenital syphilis, there is no conclusive evidence that this disease produces valvular heart lesions in infected infants [58]. *Candida* endocarditis has become increasingly prevalent, particularly associated with the use of central venous catheters.

Factors that predispose a newborn to endocarditis are not well understood, although intravascular catheters are associated with endocarditis. In contrast to older children, in whom congenital heart disease is often associated with endocarditis [478], cardiac anomalies were found in only nine of the reported cases in neonates in series before 1994.* Bacteremia arising from an infected umbilical stump [446–448], conjunctivitis [447], and skin lesions [447,469] were the presumed sources of valvular involvement in six infants; the invasive organisms associated with these conditions and with neonatal endocarditis in general can infect normal heart valves [480]. Nevertheless, the greater frequency of bacterial and fungal [468,481–488] endocarditis in newborns in recent years, particularly in association with prematurity or placement of central vessel catheters, or both, indicates that other, more complex mechanisms may also be operative in some cases [454–457,471,476,481,489].

Observations in laboratory animals and autopsy studies of adults have shown that damage to the intracardiac endothelium with formation of a sterile platelet-fibrin thrombus at the site of the injury is often the initiating event in a patient with endocarditis [486]. Endocardial trauma caused by placement of cardiac catheters, disseminated intravascular coagulation, and various nonspecific stresses associated with prematurity, such as hypotension and hypoxia, has been implicated in the genesis of thrombi.* Nonbacterial thrombotic endocarditis or verrucous endocarditis usually remains uninfected and is described as an incidental finding at autopsy [454,471,491,492]. With bacteremia, implantation of organisms may lead to valvular infection. Whether this mechanism or direct bacterial invasion is primarily responsible for valvulitis is unknown. A similar pathogenesis has been postulated for formation of mycotic aortic aneurysms in newborns [467,493,494].

Endocarditis should be suspected in any neonate, particularly a premature infant, with an indwelling vascular catheter, evidence of sepsis, and new or changing heart murmurs. When these findings are accompanied by persistent bacteremia or signs of congestive heart failure in the absence of underlying heart disease, the diagnosis must be considered seriously. Although Janeway lesions [471], a generalized petechial rash [466,477,481], and splinter hemorrhages [477] have been seen, murmurs characteristic of semilunar valve insufficiency, Osler nodes, Roth spots, arthritis, and other findings typical of valvular infection in adults and older children have not been observed in neonates. Multiple septic emboli with involvement of the skin, bones, viscera, and central nervous system are common findings, however.*

Two-dimensional echocardiography has proved to be an invaluable rapid, noninvasive method for diagnosing endocarditis [456–459,465,486,495]. Although it cannot differentiate between infected and sterile vegetations and other valvular lesions (discussed later), imaging is quite specific, and false-positive readings are uncommon. Less certainty can be placed on a negative report. Despite detection of vegetations 2 mm in diameter with echocardiography, the number of false-negative examinations is significant [465,471,496]; in one series, two of three infants with thrombotic valvular lesions 3 to 7 mm in diameter had normal two-dimensional echocardiograms [471]. A diagnosis of bacterial endocarditis should be considered in any infant with a compatible history and physical findings regardless of the results obtained by echocardiography. Widespread use of new techniques, such as transesophageal echocardiography, which provides detailed views of the mitral and tricuspid valves, and color flow Doppler imaging, which can identify areas of turbulence as blood passes over vegetations or through narrowed valve leaflets, may greatly improve diagnostic accuracy in the future [495].

When endocarditis is suspected, specimens of blood, cerebrospinal fluid, and urine obtained by catheterization or suprapubic aspiration should be sent for bacterial and fungal culture. Because blood drawn from a central catheter often contains organisms colonizing the line but not present in the systemic circulation, at least two peripheral venous blood cultures should be obtained before antimicrobial therapy is initiated. Volumes of 1 to 5 mL, depending on the infant's size, should be adequate [486].

Routine laboratory studies are helpful in supporting a diagnosis of endocarditis in the newborn. The leukocyte count, differential count, and platelet count are usually indicative of sepsis, rather than cardiac valve infection in particular. Microhematuria has been reported, although rarely [471,477]. A chest radiograph should be obtained to determine signs of cardiac failure or pulmonary or pleural space infection. CT or MRI of the brain can be helpful in an infant with neurologic signs, particularly if left-sided endocarditis or a right-to-left shunt exists. Baseline determinations of inflammatory markers are useful and can be used for assessing the efficacy of the therapy; erythrocyte sedimentation rate and C-reactive protein level have been used.

*References [455–457,460,463–465,467,468,475–487,489–510].

*References [77,450,452,455,459,460,465,479].

*References [455,471,476,480,490,491].

*References [455,457,459,460,464–466,468–474,478,479,481,485,487,489,508–510].

Intravenous therapy with a penicillinase-resistant penicillin and an aminoglycoside should be started after appropriate cultures have been obtained. In nurseries where MRSA or methicillin-resistant *S. epidermidis* infections have been a problem, vancomycin should be substituted initially for the penicillin antibiotic [486,495,496]. If endocarditis caused by *Enterococcus* species is suspected, ampicillin should be added or substituted for the penicillinase-resistant penicillin. After the infecting organism is isolated and antibiotic susceptibilities have been determined, specific antimicrobial therapy can be instituted.

Four to 8 weeks of parenteral treatment is usually adequate, depending on the susceptibility of the organism, response to therapy assessed clinically by reduction or elimination of the observed vegetations, and laboratory response. C-reactive protein level often normalizes 2 to 3 weeks before erythrocyte sedimentation rate, and blood cultures are usually sterile after 3 to 5 days of effective therapy. *Candida* species may persist for weeks, however, despite the use of active antifungal drugs. Dosage and efficacy should be monitored weekly with clinical and bacteriologic response with or without serum antibiotic and bactericidal levels [495,497]. Determination of serum bactericidal titers (Schlichter test) is of uncertain value and has never been validated in neonatal endocarditis [486,495,498]. Efficacy of treatment may also be monitored with serial echocardiograms taken until vegetations remain stable in size or disappear [465,466,468,476,481].

Intravascular catheters must be removed whenever possible, and the tip of the removed catheter should be cultured [486,499]. Extremely large or mobile vegetations occluding an outflow tract or posing a high risk of embolism may have to be removed surgically [463,465,468]. In infants with right-sided endocarditis, demonstration of decreased pulmonary blood flow through the use of ventilation-perfusion scan can be valuable in confirming the presence of emboli, particularly if there is clinical evidence of increasing respiratory effort and diminished peripheral oxygen saturation [463].

With the availability of echocardiography, improved clinical awareness, and early diagnosis, prognosis has improved. Although there were infrequent survivors before 1973 [480], the first survivors with proven endocarditis were reported in 1983 [465,466,471]. Approximately two thirds of subsequent cases have been cured. Death is usually the result of overwhelming sepsis, often in conjunction with cardiac failure. Early reconstructive surgery for infants who fail medical management may be helpful, but has been reported in only a few cases [500,501].

Inspection of the heart at autopsy has shown the mitral valve to be infected, alone or in combination with other valves, in about half of patients. The tricuspid valve was involved in 12 infants, the pulmonary valve in 7, the aortic valve in 6, infected mural thrombi in 12, and an unspecified site in 3. Microscopic examination of valve cusps has revealed the characteristic lesions of endocarditis, with multiple small, confluent, friable vegetations composed principally of bacteria and thrombi surrounded by inflammatory exudate [77,469,471]. On gross inspection, these vegetations are easily confused with noninflammatory lesions, such as those of nonbacterial thrombotic endocarditis, blood cysts [502], developmental valvular defects

[503], or hemangiomas or other vascular anomalies [504]. Cases described as fetal endocarditis in the literature are almost certainly examples of these types of lesions [503,505,506].

PERICARDITIS

Purulent pericarditis is a very unusual complication of neonatal sepsis. Approximately 20 cases of proven bacterial origin have been reported within the past 50 years [18,507–518]. No single causative agent has predominated. *S. aureus* was responsible for seven cases [18,507–509]; *E. coli* was isolated from three patients [509,514,515]; *Haemophilus influenzae* was found in two cases [510,517]; and *Salmonella wichita* [511], *Klebsiella* [518], and *P. aeruginosa* [516] were isolated from single cases. One early review of *Pseudomonas* sepsis described suppurative pericarditis in four neonates [512]. Another report on the recovery of *Pseudomonas* from the pericardium of premature infants dying of septicemia and meningitis during a nursery outbreak is difficult to evaluate because details of clinical and autopsy findings were not provided [519]. Cases caused by *Candida* species and *Mycoplasma hominis* have also been described [481,485,520]. The causes of a pericardial effusion in three fetuses with multiple congenital anomalies, myocardial hypertrophy, and pericarditis are uncertain. Although the inflammatory exudate found at autopsy contained polymorphonuclear leukocytes in addition to lymphocytes, no evidence of bacterial infection was found [521].

Virtually every infant with pericarditis has associated septic foci; pneumonia and multiple pulmonary abscesses are the most common sites. Involvement of pericardium may occur by direct extension from adjoining lung abscesses or by hematogenous spread of bacteria [508]. The presence of infectious processes elsewhere is sufficiently frequent to warrant the suggestion that pericarditis should be suspected in all infants who develop clinical signs of “heart failure” or a sudden increase in the size of the cardiac silhouette during the course of a purulent infection such as meningitis, pneumonia, or omphalitis [510,522].

Neonates with bacterial pericarditis generally present with signs and symptoms suggesting sepsis and respiratory distress. Poor feeding, listlessness, emesis, or abdominal distention may be seen in the presence of tachypnea, tachycardia, and cyanosis of various degrees. More specific signs of cardiac involvement become apparent with the accumulation of increasing amounts of pericardial effusion. The clinical findings of cardiac tamponade are extremely subtle and difficult to differentiate from the findings of myocardial disease with right-sided heart failure. A rapid pulse, quiet precordium, muffled heart sounds, neck vein distention, and hepatomegaly are findings common to both entities. More specific signs of tamponade, such as narrow pulse pressure or respiratory variations in pulse volume of more than 20 mm Hg (i.e., pulsus paradoxus), are technically difficult to obtain in neonates without an arterial catheter in place. A pericardial friction rub is absent in more than 50% of older infants and children and in most neonates with purulent pericarditis.

Rapid enlargement of the cardiac silhouette, a globular heart shape with widening of the base on tilting, and

diminished cardiac pulsation on fluoroscopic examination are of little value in differentiating pericardial effusion from cardiac dilation [523]. The early ST segment elevation and subsequent T wave inversion seen on electrocardiogram reflect subepicardial damage or inflammation and are similar to changes seen with primary myocarditis. Diminution in the amplitude of the QRS complex by fluid surrounding the heart is not a constant finding. Confirmation of the presence of a pericardial effusion is usually obtained by two-dimensional echocardiography [507,523]. In some cases, CT or MRI can also be helpful in delineating the extent of a pericardial effusion [524]; occasionally, pericardial fluid is found incidental to the chest or abdominal scanning. Additional causes of neonatal pericardial effusion other than purulent disease include viral pericarditis [525], intrapericardial teratoma [526], maternal lupus [527], immune and nonimmune [528] fetal hydrops, congenital diaphragmatic defects [529], chylopericardium [530], and central venous catheter perforation of the right atrium [531].

A definitive diagnosis of purulent pericarditis can be made only by obtaining fluid at surgery or through needle aspiration. Care and experience are necessary to facilitate aspiration while avoiding the risks of cardiac puncture or laceration [521]. Accurate monitoring of needle position can usually be obtained through CT guidance, with echocardiographic or fluoroscopic imaging, or by attaching the exploring electrode (V lead) of an electrocardiograph to the needle and by looking for injury current if contact is made with the epicardial surface of the heart.

When fluid is obtained, it should be sent for analysis to the laboratory in a laboratory-approved container appropriate for aerobic and anaerobic culture, mycobacteria, and fungi. In addition to cell count and protein and glucose levels, Gram and acid-fast stains should be performed with cultures for bacteria, viruses, mycobacteria, and fungi. Rapid identification of bacterial antigens by latex agglutination or by counterimmunoelectrophoresis of pericardial fluid, urine, or serum may also help to establish an etiologic diagnosis [532]. Specialized testing for rare bacterial, viral, or fungal organisms by molecular methods may be available by request in specialized (research) or reference laboratories.

Purulent pericarditis is a medical and surgical emergency. Therapy must be directed toward relief of the cardiac tamponade through adequate pericardial drainage and toward resolution of the infection. Both modes of treatment are essential for successful therapy for bacterial pericarditis in the newborn. No infant with suppurative pericarditis has recovered when treated by antibiotics alone [508]. Although repeated needle aspirations or catheter drainage [533] may be sufficient, the frequent occurrence of loculations of pus, particularly with staphylococcal infection, suggests that open surgical pericardiostomy is the method of choice to achieve adequate drainage.

Cultures of blood, and, when clinically indicated, cerebrospinal fluid and urine should be obtained before instituting antimicrobial therapy. Initial therapy should be based on results of Gram stain or antigen detection tests of the pericardial fluid. If no organisms can be identified, treatment can be started with penicillinase-resistant penicillin and an aminoglycoside (or extended-spectrum

cephalosporin) until definitive culture and susceptibility data are available. In nurseries where MRSA infection has been a problem, vancomycin should be substituted for penicillin [497].

The prognosis of neonatal purulent pericarditis is very poor. Only three survivors had been reported [507,509,518] before the 1990s. Treatment of these patients consisted of needle aspirations, drainage, and systemic antibiotic therapy, and in one case, treatment was combined with local instillation.

MEDIASTINITIS

Purulent mediastinitis has been reported in 11 infants younger than 6 weeks of age, although it is likely that a great many more cases occur than have been reported in the literature, and that it is an occasional complication of cardiothoracic surgery performed in the neonatal period. Six reported patients acquired mediastinal abscess through blood-borne dissemination of organisms [534–536] or by extension from a focus of infection in an adjacent retropharyngeal abscess [537], pleural or pulmonary abscess [508,535], or vertebral osteomyelitis [538]. One infant developed infection as a complication of surgery for esophageal atresia [525]. *S. aureus* was the causative organism in four infants, and *S. pneumoniae*, *Clostridium* species, and mixed *S. aureus* and *E. coli* were causative in one infant each. Organisms were not identified in four cases.

Traumatic perforation of the posterior pharynx or esophagus, often the result of resuscitative efforts in infants involving endotracheal or gastric intubation, produces a potential site for entry of microorganisms [539–546]. Retropharyngeal abscess [547], an infected pseudodiverticulum, or pyopneumothorax may occur as a consequence; purulent mediastinitis has been reported three times as a complication [536,540,541], but it is probably more common than is reported. At least one case of mediastinitis has occurred as the result of overly vigorous passage of a nasogastric tube through an atretic esophageal pouch [540]. Low (intrathoracic) perforations are said to have a higher risk of mediastinitis and abscess formation than perforations in the cervical region [542].

Early symptoms are nonspecific and are similar to symptoms of any septic process in a neonate. As purulent fluid accumulates in the mediastinum, it places increasing pressure on the esophagus, trachea, and tributaries of the superior vena cava and thoracic duct, bringing about rapid development of dysphagia, dyspnea, neck vein distention, and facial cyanosis or edema. To maintain a patent tracheal airway, an affected infant lies in an arched position with head extended in a manner very similar to that seen in neonates with congenital vascular ring. A halting, inspiratory, staccato type of breathing, probably because of pain, is also characteristic. Ultimately, the abscess may point on the anterior chest wall or in the suprasternal notch.

Usually, mediastinitis is first suspected when widening of the mediastinum is observed on a chest radiograph obtained for evaluation of respiratory distress. Forward displacement of the trachea and larynx may accompany these findings when retropharyngeal abscess is associated with mediastinitis. Infection after traumatic perforation of the esophagus or

pharynx is often accompanied by pneumomediastinum with or without a pneumothorax [540,542].

Contrast studies performed to define the cause of respiratory or feeding difficulties in infants with mediastinitis may result in flow of radiopaque fluid into an esophageal laceration, mimicking the findings of an atresia, duplication, or diverticulum of the esophagus [542,543]. In such cases, endoscopy often shows a mucosal tear, confirming the diagnosis [543,544].

Treatment should be directed toward establishment of drainage and relief of pressure on vital structures through a mediastinotomy and placement of drainage tubes. A tracheostomy or endotracheal tube may be necessary for maintenance of an adequate airway. Initial empirical antimicrobial therapy with clindamycin (or metronidazole), ampicillin, and an aminoglycoside (or extended-spectrum cephalosporin) and β -lactam and β -lactamase inhibitor combination should be started after cultures of the blood and all other clinically indicated cultures have been obtained. More limited empirical antibiotic therapy can be provided with a β -lactam and β -lactamase inhibitor combination alone, such as piperacillin plus tazobactam, ampicillin plus sulbactam, or ticarcillin plus clavulanate. Specific therapy can subsequently be determined by the results of bacteriologic studies of the cultures or purulent fluid obtained at surgery.

ESOPHAGITIS

The esophagus is infrequently a focus for infection of the fetus or newborn [548]. Esophageal atresia is associated with congenital rubella (see Chapter 28). Severe esophagitis has also been reported in neonates with congenital cytomegalovirus infection [549]. The esophagus may be involved in infants with congenital Chagas disease, identified by signs of dysphagia, regurgitation, and megaesophagus [550]. Esophageal disease may follow mediastinitis in the neonate (discussed earlier). Only occasional cases of bacterial esophagitis in a neonate have been reported; a 940-g infant boy developed signs of sepsis on the 5th day of life and died 5 hours later [551]. Premortem blood cultures were positive for *Bacillus* species. Examination at autopsy revealed histologic evidence of esophagitis with pseudomembranous necrosis of squamous epithelium and many gram-positive bacilli. No other focus of infection was evident.

INFECTIONS OF ENDOCRINE ORGANS

Endocrine glands other than the adrenal are rarely involved in fetal or neonatal infection. Nelson [552] reported neonatal suppurative thyroiditis in a term Laotian infant. The infant presented with a left anterior neck mass at 3 days of age. At surgery, a cystic mass within the left lobe of the thyroid was identified. Purulent material within the mass grew viridans streptococci and nonhemolytic streptococci.

Orchitis caused by *S. enteritidis* was described in a 10-week-old neonate [553]. The infant presented with symptoms of sepsis and diarrhea, subsequently developing unilateral scrotal swelling and erythema on the 5th day after onset of illness. Ultrasound examination of the testis showed a patchy increased echo intensity; the diagnosis

was confirmed at exploratory surgery to rule out testicular torsion. Three other cases of infection of the testes caused by *Salmonella* species in infants younger than 3 months have been described [552].

INFECTIONS OF THE SALIVARY GLANDS

Neonatal infections of salivary glands are uncommon; although rare, involvement of the parotid is the most frequent [554–559], and submandibular gland infection is infrequent [554–556]. Most infections are caused by *S. aureus* [553–556], but *E. coli* [556], *P. aeruginosa* [556], and GBS (see Chapter 12) have also been implicated in suppurative parotitis. Oral anaerobic bacteria, including *Bacteroides* species and *Peptostreptococcus* species, may be found in mixed or isolated infections in more than half of cases [557]. Infections of the salivary glands occur more frequently in premature and male infants [556,559] and most commonly manifest during the 2nd week of life. The oral cavity is the probable portal of entry for the infecting organism. Blood-borne bacteria may invade the salivary glands, however. Dehydration with resultant decreased salivary flow may be a predisposing cause in some infants.

The clinical manifestations of salivary gland infection include fever, anorexia, irritability, and failure to gain weight. There may be swelling, tenderness, or erythema over the involved gland. Purulent material may be expressed from the ductal opening with or without gentle pressure over the gland.

The diagnosis is made by culture, Gram stain, or both of the pus exuding from the duct or by percutaneous aspiration of a fluctuant area. If microscopic examination of the Gram stain does not suggest a responsible pathogen, initial antibiotic therapy should be directed against *S. aureus*, *E. coli*, and *P. aeruginosa* (i.e., penicillinase-resistant penicillin or vancomycin plus an aminoglycoside or extended-spectrum cephalosporin with activity against *Pseudomonas* organisms). If there is a strong suspicion of involvement with anaerobic bacteria (i.e., negative aerobic cultures or failure to respond to therapy directed at aerobic pathogens), consideration should be given to adding or substituting antibiotics appropriate for anaerobic bacteria (e.g., clindamycin, metronidazole in combination with other antibiotics, or a β -lactam and β -lactamase antibiotic alone). The duration of therapy should extend throughout the period of inflammation and 3 to 5 days after signs of local inflammation have disappeared. Incision and drainage often may be required; surgical drainage should be considered if there is not a prompt response to therapy within 72 hours or if the gland becomes fluctuant. When considering incision and drainage, careful attention to preservation of the function of overlying motor branches of cranial nerve VII is important.

INFECTIONS OF THE SKIN AND SUBCUTANEOUS TISSUE

Bacterial infections of the skin of the newborn may manifest as maculopapular rash, vesicles, pustules, bullae, abscesses, cellulitis, impetigo, erythema multiforme, and petechiae or purpura. In a review of 2836 neonatal

infections in Finland, only 6 were characterized as cellulitis [561]. Most infections of skin are caused by *S. aureus*, including bullous impetigo, chronic furunculosis, scalded skin syndrome, and breast abscesses (see Chapter 14). Cellulitis frequently accompanied by adenitis and bacteremia may be caused by GBS (see Chapter 12). Cutaneous infections caused by many other bacteria are discussed in this section; however, most microorganisms that cause disease in the neonate may produce cutaneous infections, and those infections are discussed in other chapters when relevant. For additional information on bacterial infections of the skin, the reader is referred to the text by Solomon and Esterly [562] and the reviews by Swartz and Weinberg [563] and Frieden [568]. Excellent color photographs are included in the *Color Atlas of Pediatric Dermatology* by Weinberg and coworkers [569].

Methicillin (oxacillin)-resistant (i.e., MRSA) infection is responsible for about 50% of cutaneous infections and is the frequent portal of entry for cutaneous infections in neonates [564]. MRSA cutaneous infections have included cellulitis and abscess formation, but may progress more frequently and rapidly to septicemia and disseminated infections to involve bones, joints, lung, central nervous system, and the cardiac endothelium [565]. In a 4-year survey (2001-2005) of staphylococcal infections in one center, 89 neonates were infected with staphylococci, and 61 (68.5%) infections were caused by MRSA; 30% of MRSA and methicillin-susceptible *S. aureus* infections manifested with pustular lesions, most often in the groin [566]. Cellulitis or abscess developed with 20% of MRSA and 32% of methicillin-susceptible *S. aureus* lesions, and 13% of lesions developed into an invasive infection (bacteremia, urinary tract infection). Also, infants with MRSA were more likely to have a mother with a history of skin infection (21% versus 4%). Other investigators have attempted to eradicate high colonization and disease rates in neonatal nurseries with the use of bacteriostatic agents applied to the umbilical cord. In one study with a high rate of MRSA impetigo, 0.3% triclosan (Bacti-Stat) was not effective in slowing the progress of the epidemic [567]. In this epidemic, 22 infants were affected, and all but 2 of the affected infants were circumcised, suggesting that the colonization of the surgical site facilitated invasion of the skin.

PATHOGENESIS

The skin of the newborn has unique characteristics, including absent microflora at birth; the presence of vernix caseosa; a less acid pH than that of older children; and often the presence of surgical wounds, including the severed umbilical cord, a circumcision site, and catheter wounds. The infant is immediately exposed to other infants, personnel, and the nosocomial environment. After the staphylococcal pandemic of the 1950s, information on the colonization of the skin, predisposing factors responsible for neonatal skin infection, bacterial transmission in the nursery, the inflammatory response of the skin to bacterial invasion, virulence factors of staphylococci, and methods of prevention of cross-infection became available. These studies are described in part in Chapters 14 and 35 and have been reviewed elsewhere [568,569].

Cutaneous bacterial infection may be a primary event or the result of systemic infection [570]. Septicemic embolic infection may occur at widely separated sites, whereas local infections often occur at a site with an identifiable predisposing cause. Procedures resulting in breaks in the cutaneous continuity, such as forceps abrasions or wounds at fetal electrodes or at venipuncture sites, may be readily identified. The necrotic umbilical cord is a site for proliferation of microorganisms that may invade local tissues.

Infection of the circumcision site remains a concern because it is the most common surgical procedure in infants in the United States. Speert [571] found that cleanliness was frequently disregarded by professional circumcisers in the 19th century. Operators were frequently uneducated, were dirty, and often spat on their instruments. Erysipelas, tetanus, and diphtheria have long been recognized as complications of unsterile surgical technique performed on newborns. In a now obsolete and prohibited part of the Orthodox Jewish circumcision ritual, the operator applied his lips to the fresh circumcision wound and sucked a few drops of blood. Such practices were responsible for transmission of syphilis and tuberculosis in neonates in the past. In one report [570], a 4-month-old infant presented with a penile ulcer, bilateral inguinal adenopathy, and a draining inguinal sinus caused by *Mycobacterium tuberculosis* after the "barber" spat on his razor before circumcision. Reports of 43 cases of tuberculosis associated with circumcision had been published by 1916 [569].

Subsequent case reports of severe infection after circumcision include bacteremia related to GBS [571], local infection and fatal staphylococcal pneumonia [572], staphylococcal scalded skin syndrome [573,574], necrotizing fasciitis [575], and bullous impetigo [576]. Two reports of necrotizing fasciitis after Plastibell circumcision emphasize severe infection as a potential risk of this procedure [577]. One infection caused by *S. aureus* and *Klebsiella* species was associated with prolonged convalescent and multiple surgical repairs, whereas a second infant survived staphylococcal necrotizing fasciitis after 14 days of intravenous antibiotic treatment.

The incidence of infection after elective circumcision was investigated at the University of Washington Hospital [578] from 1963-1972. Infection, defined as the presence of pus or erythema, occurred in 0.41% of 5521 infants and was more frequently associated with the use of a disposable plastic bell (Plastibell, 0.72%) than with the use of a metal clamp (Gomco, 0.14%). Wound cultures were infrequently available, and the microbiologic diagnosis was uncertain for most infants. Circumcision infection is uncommon, but local spread of infection may be devastating and lead to systemic infection.

Intrapartum fetal monitoring with scalp electrodes and intrauterine pressure catheters and measurements of fetal blood gases through scalp punctures have been associated with infections related to herpesvirus (see Chapter 26), *Mycoplasma* (see Chapter 20), and various aerobic and anaerobic bacteria. Bacterial infections have included pustules, abscesses, and fasciitis [579-583]. Infection rates are relatively low (0.1% to 4.5%) [579,580]; however, severe infections, including fasciitis, meningitis, and osteomyelitis, have occurred as severe complications. A review [581]

of causative organisms in fetal scalp monitor infections found that 61% of infections were polymicrobial, involving anaerobic bacteria, aerobic gram-positive cocci, and gram-negative bacilli.

A multitude of specific virulence factors may be important determinants of disease. Some phage types of *S. aureus* are responsible for local tissue damage and systemic disease; other staphylococci elaborate toxins that result in bullae and other cutaneous pathology. Groups A and B streptococci are responsible for cellulitis and impetigo in infants. *P. aeruginosa* may invade and proliferate in small blood vessels, causing local necrosis and eschar formation (i.e., ecthyma gangrenosum). Infections with *Clostridium* species cause disease in devitalized tissues such as the umbilical stump [584]. Similarly, organisms usually considered commensals, such as diphtheroids, might be responsible for infection of the cord and fetal membranes [585].

MICROBIOLOGY

The skin of the infant is colonized initially by microorganisms present in the maternal birth canal. The skin of infants delivered by cesarean section is usually sterile at birth. After birth, microorganisms may be transferred to the skin during handling by the parents and nursery personnel. The prevalent organisms on the skin during the first few days of life include coagulase-negative staphylococci, diphtheroids, and gram-negative enteric bacilli (including *E. coli*) [586,587]. The umbilicus, genitalia, and adjacent skin areas (groin and abdomen) are colonized first; organisms then spread to the nose, throat, conjunctivae, and other body sites. Organisms present in the nursery environment colonize neonatal skin after a few days in the nursery. *S. aureus*, GBS, and various species of gram-negative bacilli may be present, but the microbiologic flora differs among nurseries and from time to time in the same nursery. Use of soaps and antiseptic solutions modifies the flora on the skin of the newborn. Hexachlorophene decreases colonization with staphylococci and diphtheroids, but gram-negative organisms are unaffected or may increase after use of this agent [588].

EPIDEMIOLOGY

Infant boys are more susceptible to skin infections caused by *S. aureus* than girls. Thompson and coworkers [589] showed that boys were colonized more frequently in every body site cultured, including the nose, groin, rectum, and umbilicus. Their review of studies indicated that in England, the United States, and Australia, approximately 50% more boys had skin lesions than girls. Although the incidence of breast abscesses is equal in boys and girls during the first 2 weeks of life, such abscesses are more frequent thereafter in girls [590]. The reason for this pattern is unclear, but Rudoy and Nelson [590] hypothesized that physiologic breast enlargement may play a role. Hormone production in the female infant after the 2nd week might account for the increase in abscesses of the breast.

Infections caused by MRSA involving the skin of children and neonates have markedly increased. The *mecA* gene responsible for resistance to oxacillin and nafcillin

is often closely linked to a gene responsible for skin invasion. Before 1997, epidemic MRSA infections occurred in neonatal units involving infections of the respiratory tract, nasopharynx, gastrointestinal tract, eye, blood, wounds, or umbilicus [591]; these infections were usually restricted to single nurseries and involved a single genetic variant of MRSA. Since 1990, MRSA infections acquired in the community have been reported with increased frequency [592], including in infants 2 weeks old. Of these infections, 91% have involved the skin and soft tissues. In contrast to typical nosocomial MRSA, community-acquired MRSA organisms have frequently remained susceptible to trimethoprim-sulfamethoxazole and clindamycin. At the University of New Mexico, continued surveillance of MRSA-colonized and MRSA-infected infants in NICUs showed that by 2003, more than half of all isolates were community acquired.

Seasonal variation in the frequency of neonatal skin infections has been reported by Evans and coworkers [593], who conducted a series of studies at Harlem Hospital in New York. The prevalence of *S. aureus*, *E. coli*, and streptococci in the nares and umbilicus of infants was lowest in the autumn and usually highest in the summer or spring. No seasonal variation was observed for *S. epidermidis* or *Enterobacter* species. The investigators concluded that seasonal differences must be considered in investigations of bacterial colonization of the newborn skin, and that high humidity may favor gram-negative colonization.

The time of onset of skin lesions associated with sepsis may be early (during the 1st week of life) or late (several weeks or months after birth). Disease acquired in the nursery usually becomes apparent after 5 days of age. Many skin lesions do not appear until after the infant has left the nursery; the observed incidence of skin disease caused by bacteria should include surveillance of infants in the home during the 1st month of life. Physicians responsible for neonatal care must be alert to the unusual occurrence of skin lesions. The introduction of a new and virulent bacterium, an alteration in technique, or the use of contaminated materials must be considered as possible causes of an increased incidence of such infections.

CLINICAL MANIFESTATIONS

Infants who have skin infections that remain localized that are not invasive or part of a systemic infection have few general signs of disease, such as fever, alteration in feeding habits, vomiting, or diarrhea. These signs may be present when significant tissue invasion occurs, as in abscesses or extensive cellulitis. Cutaneous manifestations that result from infectious diseases are listed in Table 10-5.

Among the common and least specific lesions are maculopapular rashes; these rashes may be caused by viruses (measles, rubella, or enteroviruses), fungi (*Candida* species), or bacteria (streptococci or staphylococci), or they may be unassociated with any infectious process. Erythema multiforme lesions have been observed in cases of sepsis related to *S. aureus* [594], streptococci [560], and *P. aeruginosa* [595]. Virtually any rash may be associated with bacterial infection. In an outbreak of sepsis caused by *Achromobacter* in premature infants [596], illness was

TABLE 10-5 Manifestations and Etiologies of Infections of the Skin in Newborns

Clinical Manifestation	Etiologic Agent	
	Bacterial	Nonbacterial
Maculopapular rash	<i>Treponema pallidum</i> * [†] <i>Listeria monocytogenes</i> <i>Streptococcus</i> * <i>Staphylococcus</i>	Measles virus* Rubella virus* Enteroviruses* Molluscum contagiosum [653] <i>Candida</i> species*
Cellulitis (erysipelas)	Groups A and B streptococci Achromobacter species [626]	
Impetigo	Groups A and B streptococci [627]* <i>Staphylococcus aureus</i> * <i>Escherichia coli</i>	
Erythema multiforme	β-hemolytic streptococci [624] <i>S. aureus</i> * <i>Pseudomonas aeruginosa</i> [625]	
Vesicular or bullous lesions	<i>S. aureus</i> * <i>P. aeruginosa</i> <i>T. pallidum</i> <i>Haemophilus influenzae</i> type b [628] <i>L. monocytogenes</i> [629]	Herpes simplex virus* [†] Cytomegalovirus* Varicella virus* [†] Variola virus [†] Coxsackieviruses* <i>Candida</i> species* <i>Aspergillus</i> species* <i>Drosophila larvae</i> [654] <i>Sarcoptes scabiei</i> [655]
Pustular rashes	<i>S. aureus</i> * <i>L. monocytogenes</i> * <i>H. influenzae</i> [631]	
Ecthyma gangrenosum	<i>P. aeruginosa</i> [632–634]	
Abscesses and wound infections	<i>S. aureus</i> * <i>Staphylococcus epidermidis</i> * β-hemolytic streptococci [637] Group B streptococci [638] <i>E. coli</i> [639–641] <i>Klebsiella</i> species [642] <i>Proteus mirabilis</i> [643] <i>P. aeruginosa</i> [644] <i>Salmonella</i> species [645] <i>Serratia marcescens</i> [647] <i>H. influenzae</i> [648] <i>Haemophilus parainfluenzae</i> [649] <i>Corynebacterium vaginalis</i> [650] <i>Neisseria gonorrhoeae</i> [658] <i>Gardnerella vaginalis</i> [651] <i>Bacteroides</i> species [652]	<i>Mycoplasma hominis</i> * <i>Candida albicans</i> [656]
Petechiae, purpura, and ecchymoses	Gram-positive cocci* and gram-negative bacilli* associated with sepsis <i>L. monocytogenes</i> [629] <i>Streptococcus pneumoniae</i> [629] <i>T. pallidum</i> * [†]	Rubella virus* [†] Cytomegalovirus* [†] Herpes simplex virus* [†] Coxsackievirus B* [†] <i>Toxoplasma gondii</i> * [†]

*See appropriate chapter for further discussion.

[†]Including infections acquired in utero.

marked by respiratory distress, including apnea and cyanosis, but was characterized by a rash consisting of indurated, erythematous lesions with sharply defined borders that began on the cheeks or chest and spread rapidly to adjacent areas.

Cellulitis, erysipelas, and impetigo are usually associated with streptococcal infection (group A or B) [597], although impetigo caused by *S. aureus* or *E. coli* has also been reported in infants. Vesicles, commonly associated with infections by herpesviruses, also are seen occasionally during early stages of skin lesions caused by *S. aureus*, *H. influenzae* [598], *L. monocytogenes* [599], and *P. aeruginosa*. GBS [600], *S. aureus*, *P. aeruginosa*, herpes simplex virus, and *T. pallidum* may also be responsible for bullous lesions. Pustules commonly occur in staphylococcal diseases, but also occur in infections caused by *L. monocytogenes* and, rarely, in skin infections with *H. influenzae* [601].

Ecthyma gangrenosum is a local manifestation of infection with *P. aeruginosa* [602,603]. Lesions begin as a vesicular eruption on a wide erythematous base. Vesicles rupture and form an indurated black eschar followed by larger, sharply demarcated, painless necrotic areas, resulting from a small vessel vasculitis with necrosis of the adjacent tissue. The organisms are present in purulent material underlying the necrotic membrane. These lesions are particularly more common adjacent to nose, lip, ear, mouth, and perineum, resulting in avascular necrosis and loss of tissue. *P. aeruginosa* may be grown in pure culture from blood and lesions. Among 48 infants described in one outbreak, lesions appeared within the first 2 weeks of life; most infants died within 3 days of onset [604]. Ecthyma is relatively specific for *Pseudomonas* infections, but similar or identical lesions have rarely been described in infections owing to *S. aureus*, *Aeromonas hydrophila*, *S. marcescens*, *Aspergillus* species, or *Mucor* species [605].

Many infants with *Candida* infections have cutaneous manifestations. Baley and Silverman [606] described 18 infants with systemic candidiasis; 8 had a burnlike truncal erythema, and 9 other infants had typical candidal diaper rashes or maculopapular rashes of the axillae or neck.

Abscesses of the skin and subcutaneous tissue are usually caused by *S. aureus* and, less frequently, by group A streptococci, GBS [607,608], or gram-negative enteric bacilli [609–617]. Community-acquired MRSA organisms are even more likely to produce skin infections with abscess formation. Organisms that colonize the skin over an area that has been disrupted by an abrasion or other wound may invade the subcutaneous tissue and produce an abscess. *Haemophilus* species [618–620], *Gardnerella vaginalis* [621], *Bacteroides* species [622], molluscum contagiosum [623], *Drosophila* myiasis [624], scabies [625], and *Candida* [626] are examples of diverse causes of cutaneous abscesses; virtually any bacterial, fungal, or parasitic agent that is normally or transiently on skin may become a pathogen. *E. coli*, *Klebsiella* species, *P. aeruginosa* [605,610,627], *N. gonorrhoeae* [628], and *Bacteroides fragilis* [629] have caused wound infections in infants whose scalps were lacerated by forceps, fetal electrodes, or instruments used for obtaining blood from the scalp in utero. An extensive outbreak of systemic disease caused by *S. marcescens* in an NICU in Puerto Rico included wound infections at the site of intravenous infusions.

A cephalhematoma may become infected during sepsis or from manipulation of the cephalhematoma, such as through diagnostic or therapeutic needle puncture [630] or by puncture from a fetal monitor. Infections may be caused by *Bacteroides* species [622], *E. coli* [609,610], and *P. aeruginosa* [631]. The infection may be associated with meningitis [631] or with osteomyelitis of the underlying skull [609,610].

S. aureus is the most frequent etiologic agent in breast abscess, but gram-negative enteric bacilli may be becoming more common [611,615,616]. Of 36 cases with mastitis seen in Dallas, Texas, during a 16-year period, 32 cases were caused by *S. aureus*, 1 was caused by *E. coli*, and 2 were caused by *Salmonella* species; both *E. coli* and *S. aureus* were isolated from one abscess [632]. At Children's Hospital in Boston from 1947–1983, 41 cases of mastitis in neonates were managed [633]. *S. aureus* was responsible for 29 of 34 cases with an identifiable bacterial pathogen. All cases occurred in term infants during weeks 1 to 5 of life. Bilaterality and extramammary foci were rare. One third of infants were febrile, and most had elevated white blood cell counts (>15,000 cells/mm³). Other reports have identified GBS [608] and *P. mirabilis* [613] as causes of breast abscesses. Brook [616] found that 5 of 14 breast abscesses contained anaerobic bacteria (i.e., *Bacteroides* species and *Peptostreptococcus*), but *S. aureus*, GBS, or enteric bacteria predominated; anaerobic bacteria occurred alone in only 2 of 14 cases.

Paronychia may occur in neonates after injury to the cuticle. The lesion is usually caused by *S. aureus* or β -hemolytic streptococci [592]. The authors of a report on an outbreak of paronychia in a Kuala Lumpur nursery suggest but do not prove that the lesions were caused by an anaerobic *Veillonella* species [634].

Omphalitis is defined by the presence of erythema or serous or purulent discharge from the umbilical stump or periumbilical tissues. A review by Cushing [635] provided a useful discussion of the pathophysiology, microbiology, diagnosis, and management of omphalitis. The incidence of infection is more frequent in infants with low birth weight and infants with complications of delivery. A survey of infants born at the Royal Woman's Hospital in Brisbane, Australia [636], identified an incidence of approximately 2% among term infants. The mean age of infants at presentation of omphalitis was 3.2 days. Perhaps because hexachlorophene bathing was used, gram-negative bacilli were more frequently associated with infection than gram-positive cocci. Microbiologic results are difficult to interpret, however, because swabs of the site of infection do not exclude surface contaminants, unless cultures are taken with extreme care and precision.

A series from the United States [637] found that periumbilical fasciitis was more frequent in boys, but did not find that umbilical catheterization, low birth weight, or septic delivery was associated with a high risk; overall, the incidence of omphalitis was equal in boys and girls. In this series, omphalitis manifested as discharge, cellulitis, or fasciitis; gram-positive organisms were found in 94% of cultures, and gram-negative bacteria were found in 64%. *S. aureus* was the most frequent isolate, with *E. coli* and *Klebsiella* species the next most common. Group A

streptococci have been responsible for nursery outbreaks that may include an indolent form of omphalitis characterized by erythema and oozing of the umbilical stump for days to weeks, accompanied by pustular lesions of the abdominal wall in some cases [638]. Neonatal tetanus usually occurs as a result of contamination of the umbilical wound by *Clostridium tetani* at delivery.

Acute necrotizing fasciitis is a bacterial infection of subcutaneous tissue and fascial sheath [614,639,640]. Infection can arise in an operative wound or in a focal infection, such as a breast abscess, or there may be no apparent predisposing cause. Necrotizing fasciitis has been reported after circumcision [577] and as a complication of insertion of a fetal monitor [641]. The trunk and extremities are the areas most commonly involved; inflammation spreads rapidly along fascial planes, producing thrombosis and extensive necrosis, with infarcts developing in overlying skin. Vesicles and bullae appear, and the skin may become blue-gray or black. Myositis and bacteremia may accompany fasciitis. Staphylococci, GBS [642], *E. coli*, *P. aeruginosa*, anaerobic bacteria [643], and mixtures of gram-positive and gram-negative bacteria have been associated with this disease. The bacteria are present in skin lesions, deep fascia, and, in some cases, blood. The mortality is high despite the use of fasciotomy, wide débridement, and antibiotics.

Perirectal abscesses may occur in newborns. In contrast to older children, most newborns with perirectal abscess do not have underlying immunodeficiency, although infants with acquired or congenital immunodeficiency often present with this condition. The most common causes of perirectal abscess are *S. aureus*, *E. coli*, or other enteric bacilli [644,645]; however, anaerobic bacteria can also be involved. *S. aureus* and enteric bacilli may be more common in infants and newborns [645]. Recent rectal surgery for conditions such as Hirschsprung disease or imperforate anus (myotomy or rectal dilation) may be predisposing causes in infants; as in older children, neutropenia may be associated with an increased risk for perirectal abscess.

Otitis externa is uncommon in newborns. Victorin [646] described an outbreak of neonatal infections in which *P. aeruginosa* was cultured from seven infants with suppurative of the auditory canal. The author suggested that this outbreak was caused by contaminated bath water used in the nursery.

DIAGNOSIS

The appearance of a skin lesion alone may be sufficiently typical to suspect certain etiologic agents (e.g., ecthyma gangrenosum), but more often, the appearance is nonspecific. A microbiologic diagnosis should be sought to provide specific therapy. The lesion and the surrounding tissue should be cleaned with 70% ethanol to prevent contamination from organisms that colonize the surface. If crusts are present, they should be lifted with a sterile swab to provide drainage, and cultures should be obtained from the base of the lesion.

Vesicles and pustules can be aspirated with a needle (20-gauge to 25-gauge) attached to a syringe, or they can be opened and exudate can be collected on a sterile

swab. Generally, swabs are not preferred for specimen collection because swab materials bind or inactivate bacterial organisms. Aspiration of abscesses is important; more than one aspiration may be required because the suppurative focus may not be easily distinguished from the surrounding inflammatory tissue. Aspiration of the leading edge or point of maximal inflammation of an area of cellulitis may be valuable and should be performed if no other suppurative or purulent sites are available for culture. A small needle (25-gauge or 26-gauge) should be attached to a tuberculin or other small-volume syringe filled with 0.25 to 0.50 mL of sterile nonbacteriostatic saline; the needle should be inserted into the area of soft tissue to be sampled, with continuous, gentle aspiration applied to the syringe. If no fluid is returned to the syringe, a small amount of fluid should be injected and immediately aspirated back into the syringe. Collected material may be sent to the laboratory in the syringe for Gram stain and culture, or, alternatively, the contents may be washed into a tube of bacteriologic broth medium for transport and subsequent culture.

If swabs are used, care must be taken that the material does not dry before it is plated on bacteriologic media. Swabs preferentially should be directly inoculated or rinsed in bacteriologic media and immediately transported to the microbiology laboratory. Alternatively, they may be refrigerated or placed in appropriate transport media if more than a few hours will elapse before inoculation of media in the laboratory. Whenever sufficient material is available (on swabs or in liquid), several slides should be prepared for Gram staining.

It is often difficult to distinguish petechiae from vascular dilation. Pressure with a glass slide on the border of the lesion is a simple and reliable method for detecting extravasation of red blood cells. If the lesion disappears on pressure, it is probably caused by dilation of small vessels, whereas persistence of the lesion after application of pressure indicates extravasation of red blood cells. Bacteria may be present in petechial lesions that occur in infants with bacterial sepsis. Blood obtained by aspiration or gentle scraping with a scalpel at the center of the petechiae may reveal the causative organism on Gram stain or culture.

DIFFERENTIAL DIAGNOSIS

Sclerema neonatorum, milia, and erythema toxicum are noninfectious lesions that are often confused with infections of the skin [647]. Bullous and purpuric lesions may be caused by noninfectious disorders, including mast cell diseases (e.g., urticaria pigmentosa), histiocytosis X, acrodermatitis enteropathica, dermatitis herpetiformis, epidermolysis bullosa, congenital porphyria [562], and pemphigus vulgaris [648]. A syndrome of generalized erythroderma, failure to thrive, and diarrhea has been associated with various forms of immunodeficiency [649].

Sclerema neonatorum is a diffuse, spreading, waxy hardness of the skin and subcutaneous tissue that occurs during the first weeks of life [628,650]. The subcutaneous tissue seems to be bound to underlying muscle and bone. This condition is usually seen on the thighs, buttocks, and trunk. Although associated with sepsis in some infants,

sclerema also affects infants with dehydration, acidosis, and shock. Most evidence supports the hypothesis that sclerema is a manifestation of shock and insufficiency of the peripheral circulation. When it occurs in infants with generalized infection, sclerema is associated with a poor prognosis. In a review of cases of sepsis at The New York Hospital, sclerema was detected in 6 of 71 infants, 5 of whom died [651].

Milia are yellow or pearly white papules that are 1 mm in diameter and usually found scattered over the cheeks, forehead, and nose [647,652]. The lesion is a small cyst formed from retention of sebum in sebaceous glands. Because the cyst is capped by a shiny surface of epidermis, it may be confused with a small pustule. Milia are common; Gordon [652] estimated that 40% of healthy newborns have milia. The lesions are common in the first few weeks of life. These cysts may be distinguished from staphylococcal pustules by aspiration and Gram stain of the material.

Erythema toxicum consists of several types of lesions, including 1- to 3-mm, yellow-white papules or pustules on an erythematous base, erythematous macules, or diffuse erythema. These lesions are usually present on the trunk, but may involve the head and neck and extremities as well. Most lesions appear within the first hours of life and are uncommon after 2 days of age. Erythema toxicum is uncommon in low birth weight or premature infants [653]. Affected infants have no signs of systemic illness or local irritation. A smear of the contents of pustules reveals the presence of eosinophils and an absence of bacteria. Other noninfectious pustular lesions of newborns include neonatal pustular melanosis, which is marked by a mixed infiltrate that has a predominance of neutrophils [654], and infantile acropustulosis, which is characterized by an eosinophilic infiltration of the skin [655,656].

Bullae may occur on the skin of the wrist or forearm and usually are caused by trauma [657]. Sucking of the extremity by the infant is believed to cause the bullae, which contain sterile serous fluid. Purpura may be caused by noninfectious conditions, including trauma; erythroblastosis fetalis; or, less frequently, coagulation disorders, maternal drug ingestion, congenital leukemia, and congenital Letterer-Siwe disease.

Diaper rash is primarily a contact dermatitis associated with soiling of the skin by urine and stool [658–660]. The rash may occur as a mild erythema or scaling, a sharply demarcated and confluent erythema, or discrete shallow ulcerations. A beefy red, confluent rash with raised margins, satellite (e.g., folliculitis) oval lesions, or discrete vesicular-pustular lesions indicates secondary invasion by *C. albicans* or *S. aureus*. Systemic infectious illnesses that manifest as disseminated rashes (e.g., herpes, varicella, syphilis) may be characterized by early typical lesions in the diaper area.

TREATMENT

The treatment of localized skin lesions consists of the use of local antiseptic materials, systemic antimicrobial agents, and appropriate incision and drainage or débridement. Hexachlorophene (3% detergent emulsion) and chlorhexidine (4% solution) are valuable in cleaning

small, abraded areas and discrete pustular lesions. Because of concern over its neurotoxicity and cutaneous absorption, hexachlorophene should not be used on large open areas of skin (see Chapter 14).

Systemic antibiotics should be considered for therapy whenever there is significant soft tissue infection with abscess or cellulitis. The specific antibiotic choice should be made on the basis of the microbiology of the lesion; streptococci may be treated effectively with penicillin G, ampicillin, or extended-spectrum cephalosporins (i.e., cefotaxime or ceftriaxone), whereas staphylococci generally must be treated with penicillinase-resistant penicillins or vancomycin. Infections owing to gram-negative enteric bacilli may be treated with aminoglycosides or extended-spectrum cephalosporins based on the results of susceptibility testing. Infections owing to *Pseudomonas* organisms can be effectively treated with aminoglycosides or ceftazidime.

Local heat and moist dressings over areas of abscess formation may facilitate localization or spontaneous drainage. Indications for incision and drainage of abscesses in infants are the same as for the indications in older children and adults.

PREVENTION

Prevention of local skin infections is best provided by appropriate routine hygiene, maintenance of the integrity of skin (i.e., avoidance of drying, trauma, or chemical contact), frequent diaper changes, and hygienic care of the umbilicus or other wounds or noninfectious skin inflammation. The following measures of skin care are recommended by the Committee of the Fetus and Newborn of the American Academy of Pediatrics [660] to prevent infection:

1. The first bath should be postponed until the infant is thermally stable.
2. Nonmedicated soap and water should be used; sterile sponges (not gauze) soaked in warm water may be used.
3. The buttocks and perianal should be cleaned with fresh water and cotton or with mild soap and water at diaper changes.
4. Ideally, agents used on the newborn skin should be dispensed in single-use containers.
5. No single method of cord care has proved to be superior, and none is endorsed [660].

Cord care may include application of alcohol, triple dye (i.e., brilliant green, proflavine hemisulfate, and crystal violet), or antimicrobial agents such as bacitracin. Alcohol hastens drying of the cord, but is probably not effective in preventing cord colonization and omphalitis. A randomized study of triple dye, povidone-iodine, silver sulfadiazine, and bacitracin ointment showed comparability in antimicrobial control [661].

During nursery outbreaks, the Centers for Disease Control and Prevention recommends the judicious use of hexachlorophene bathing [662]. Daily hexachlorophene bathing of the diaper area [663] and umbilical cord care with 4% chlorhexidine solution [664] have shown efficacy for prevention of staphylococcal disease (see Chapter 14).

CONJUNCTIVITIS AND OTHER EYE INFECTIONS

Conjunctivitis in the newborn usually results from one of four causes: infection with *N. gonorrhoeae*, infection with *S. aureus*, inclusion conjunctivitis caused by *Chlamydia trachomatis*, or chemical conjunctivitis caused by silver nitrate solution [665,666]. Less commonly, other microorganisms have been implicated as a cause of conjunctivitis, including group A streptococci, GBS, *S. pneumoniae*, *H. influenzae* (nontypable [591] and group b [667]), *P. aeruginosa*, *Moraxella* (*Neisseria*) *catarrhalis* [668], *Neisseria meningitidis* [669], *Corynebacterium diphtheriae* [670], *Pasteurella multocida* [671], *Clostridium* species [672], herpes simplex virus, echoviruses, *M. hominis*, and *Candida* species. In addition to meningococcal infections, other *Neisseria* species can be confused with gonococcal infections; *Neisseria cinerea* has been reported to cause conjunctivitis that was indistinguishable from gonococcal infection [673].

An epidemic of erythromycin-resistant *S. aureus* conjunctivitis affected 25 of 215 newborns during a 10-month period; control of the epidemic was achieved by identification of staff carriers and substitution of silver nitrate prophylaxis for erythromycin [674]. The major causes of conjunctivitis in neonates are discussed in Chapters 14 and 15. Cultures of the conjunctivae of neonates with purulent conjunctivitis and from the comparable eyes of a similar number of infants chosen as controls revealed significant differences, suggesting causality for viridans streptococci, *S. aureus*, *E. coli*, and *Haemophilus* species [675,676] MRSA infections may have an increased propensity to infect the eye. Conjunctivitis owing to MRSA was reported as the first case in an epidemic that eventually involved 14 neonates in an NICU including multiple invasive infections and two deaths [677]. MRSA has also caused dacryocystitis [678] and keratitis [679] in neonates.

Compared with chemical (e.g., silver nitrate) conjunctivitis, other noninfectious causes for conjunctivitis occur only rarely. Eosinophilic pustular folliculitis has been described since 1970 [680] although this disease usually occurs after 3 months of age, some infants younger than 4 to 6 weeks have been described. These infants present with recurrent crops of pruritic papules primarily affecting the scalp and brow. Biopsy specimens reveal folliculitis with a predominant eosinophilic infiltrate; most infants also have a leukocytosis and eosinophilia. Other acute or chronic cutaneous conditions may also manifest as conjunctival or periocular inflammation, such as seborrhea, atopic dermatitis, acropustulosis of infancy, and erythema toxicum (see “Infections of the Skin and Subcutaneous Tissue”).

In a review by Hammerschlag [681], the incidence of the two major pathogens ranged from 17% to 32% for *C. trachomatis* and 0% to 14.2% for *N. gonorrhoeae* in four U.S. studies. In other developed countries such as England [682], investigators found 8 cases of gonococcal infection and 44 cases of chlamydial infection among 86 newborns with ophthalmia neonatorum; in Denmark [683], investigators found that 72% of infants with conjunctivitis at 4 to 6 days after birth had positive cultures, but 70% were caused by staphylococci (*S. aureus* and *S. epidermidis*), and chlamydiae were isolated from only 2 of 300 newborns.

The incidence and microbiology of neonatal conjunctivitis depend on the incidence of transmissible infections in the maternal genital tract or the nursery and the use and efficacy of chemoprophylaxis. In Nairobi, Kenya, in a hospital where ocular prophylaxis had been discontinued, the incidence of gonococcal and chlamydial ophthalmitis was 3.6 and 8.1 cases per 100 live births [684]; in Harare, Zimbabwe, in a hospital where prophylaxis also was not used, the most common cause of conjunctivitis was *S. aureus* [685]. The introduction of tetracycline ointment for prophylaxis at Bellevue Hospital in New York City led to an overall increase in conjunctivitis associated with an increase in the incidence of gonococcal infection [686] because of the emergence of tetracycline resistance among gonococci.

Infections related to *P. aeruginosa* warrant special attention. Although uncommon, pseudomonal conjunctivitis may be a devastating disease if not recognized and treated appropriately [686]. The infection is usually acquired in the nursery, and the first signs of conjunctivitis appear between the 5th and 18th days of life. At first, the clinical manifestations are localized to the eye and include edema and erythema of the lid and purulent discharge. In some children, the conjunctivitis progresses rapidly, with denuding of the corneal epithelium and infiltration with neutrophils. With extension of the corneal infiltration, perforation of the cornea may occur. The anterior chamber may fill with fibrinous exudate, and the iris can adhere to the cornea. Subsequent invasion of the cornea by small blood vessels (pannus) is characteristic of pseudomonal conjunctivitis. The late ophthalmic complications may be followed by bacteremia and septic foci in other organs [687].

Pseudomonal eye infections in neonates can occur in epidemic form, with subsequent high rates of mortality and ophthalmic morbidity. Burns and Rhodes [687] reported a series of eye infections caused by *P. aeruginosa* in premature infants with purulent conjunctivitis rapidly progressing to septicemia, shock, and death in four infants. Five other children with conjunctivitis alone survived, but one child required enucleation. Drewett and coworkers [688] described a nursery outbreak of pseudomonal conjunctivitis believed to be caused by contaminated resuscitation equipment; of 14 infected infants, 1 became blind, and 1 had severe corneal opacities. Rapidity of the course of this infection is indicated in a case report of a 10-day-old infant who developed a corneal ulcer with perforation within 2 days after first observation of a purulent discharge [689]. An outbreak of four cases of *Pseudomonas* conjunctivitis in premature infants occurred within 2 weeks at the American University of Beirut Medical Center [690]; no cause for the outbreak was found.

A review by Lohrer and Belohradsky [691] of bacterial endophthalmitis in neonates emphasizes the importance of *P. aeruginosa* in invasive bacterial eye infections ranging from keratitis to panophthalmitis. The literature review included 16 cases of invasive eye infections in neonates; 13 were caused by *P. aeruginosa*, and the others were cases of endophthalmitis caused by GBS and *S. pneumoniae*. Other opportunistic gram-negative pathogens associated with outbreaks of infections in nurseries may also include conjunctivitis as a part of the infection syndrome. In a report by Christensen and coworkers [692], multiply antibiotic-resistant *S. marcescens* was responsible for 15 cases of pneumonia, sepsis, and meningitis and for

20 cases of conjunctivitis, cystitis, and wound infection over a 9-month period in an NICU.

Dacryocystitis may complicate a congenital lacrimal sac distention (i.e., dacryocystocele) which may appear as early as the during the first week of life with edema and erythema of the lower lid. Purulent material emerged from the puncta after moderate pressure over the lacrimal sac; *S. marcescens* was grown from the material.

The physician responsible for management of a child with purulent conjunctivitis must consider the major causes of the disease and must be alert to rare pathogens. In hospitals that practice Credé method (i.e., silver nitrate application), purulent conjunctivitis during the first 48 hours of life is almost always caused by chemical toxicity [694]. After the first 2 days, the pus of an exudative conjunctivitis must be carefully examined by Gram stain for the presence of gram-negative intracellular diplococci, gram-positive cocci in clusters, and gram-negative bacilli. Appropriate cultures should be used for isolation of the organisms concerned. If the smears are inconclusive and no pathogens are isolated on appropriate media, and if conjunctivitis persists, a diagnosis of inclusion or chlamydial infection is likely [693,694].

The treatment of staphylococcal and gonococcal conjunctivitis is discussed in Chapters 14 and 15. Chlamydial conjunctivitis is reviewed in Chapter 19.

If infection with *Pseudomonas* species is suspected, treatment should be started at once with an effective parenteral antibiotic, such as an aminoglycoside (e.g., tobramycin, amikacin, or gentamicin) with or without an anti-pseudomonal penicillin or ceftazidime (see Chapter 35) and with a locally applied ophthalmic ointment. The use of subconjunctival gentamicin or other antipseudomonal aminoglycoside is of uncertain value; however, if the cornea seems to be extensively involved, there is a risk of rapid development of endophthalmitis, and the subconjunctival injection of antibiotics should be considered in consultation with an ophthalmologist. If the diagnosis is confirmed, this regimen is continued until the local signs of *Pseudomonas* infection resolve.

Recommendations for ocular chemoprophylaxis are discussed in Chapters 15 and 19. Additional information is available in the 2009 edition of the *Report of the Committee on Infectious Diseases* published by the American Academy of Pediatrics [695].

REFERENCES

- [1] F.M. Murphy, C.J. Baker, Solitary hepatic abscess: a delayed complication of neonatal bacteremia, *Pediatr. Infect. Dis. J.* 7 (1988) 414.
- [2] B. Guillois, et al., Staphylococci pleuro-pulmonaire néonatale avec abcès hépatiques multiples, *Ann. Pediatr.* 36 (1989) 681.
- [3] L.P. Dehner, J.M. Kissane, Pyogenic hepatic abscesses in infancy and childhood, *J. Pediatr.* 74 (1969) 763.
- [4] H.T. Wright Jr., Personal communication, (1987).
- [5] T.J. Moss, T.J. Pysker, Hepatic abscess in neonates, *Am. J. Dis. Child.* 135 (1981) 726.
- [6] M.J. Chusid, Pyogenic hepatic abscess in infancy and childhood, *Pediatrics* 62 (1978) 554.
- [7] P. Dineen, Personal communication, Cornell University Medical College, New York, NY.
- [8] T.V. Bilfinger, et al., Pyogenic liver abscesses in nonimmunocompromised children, *South. Med. J.* 79 (1986) 37.
- [9] K.C. Beutow, S.W. Klein, R.B. Lane, Septicemia in premature infants: the characteristics, treatment, and prevention of septicemia in premature infants, *Am. J. Dis. Child.* 110 (1965) 29.
- [10] E.C. Dunham, Septicemia in the newborn, *Am. J. Dis. Child.* 45 (1933) 229.
- [11] W. Nyhan, M.D. Fousek, Septicemia of the newborn, *Pediatrics* 22 (1958) 268.
- [12] S.P. Gotoff, R.E. Behrman, Neonatal septicemia, *J. Pediatr.* 76 (1970) 142.
- [13] J.R. Hamilton, A. Sass-Kortsak, Jaundice associated with severe bacterial infection in young infants, *J. Pediatr.* 63 (1963) 121.
- [14] P. Hänninen, P. Terhe, A. Toivanen, Septicemia in a pediatric unit: a 20-year study, *Scand. J. Infect. Dis.* 3 (1971) 201.
- [15] E. Potter, *Pathology of the Fetus and Infant*, third ed., Year Book Medical Publishers, Chicago, 1975.
- [16] W.A. Silverman, W.E. Homan, Sepsis of obscure origin in the newborn, *Pediatrics* 3 (1949) 157.
- [17] R.T. Smith, E.S. Platau, R.A. Good, Septicemia of the newborn: current status of the problem, *Pediatrics* 17 (1956) 549.
- [18] W.M. Gersony, G.H. McCracken Jr., Purulent pericarditis in infancy, *Pediatrics* 40 (1967) 224.
- [19] J.H.M. Axton, Amoebic proctocolitis and liver abscess in a neonate, *S. Afr. Med. J.* 46 (1972) 258.
- [20] T. Botman, P.J. Ruys, Amoebic appendicitis in newborn infant, *Trop. Geogr. Med.* 15 (1963) 221.
- [21] Y.W. Brans, R. Ceballos, G. Cassady, Umbilical catheters and hepatic abscesses, *Pediatrics* 53 (1974) 264.
- [22] H.J. Cohen, S. Dresner, Liver abscess following exchange transfusion for erythroblastosis fetalis, *Q. Rev. Pediatr.* 16 (1961) 148.
- [23] J. deBeaujeu, et al., Abcès hépatique à forme tumorale chez un nourrisson, *Pediatric 23* (1968) 363.
- [24] W. Heck, F. Rehbein, B. Reismann, Pyogene Leberabszesse im Säuglingsalter, *Z. Kinderchir. (Suppl. 1)* (1966) 49.
- [25] S.R. Kandall, A.B. Johnson, L.M. Gartner, Solitary neonatal hepatic abscess, *J. Pediatr.* 85 (1974) 567.
- [26] T. Kutsunai, Abscess of the liver of umbilical origin in infants: report of two cases, *Am. J. Dis. Child.* 51 (1936) 1385.
- [27] C.M. Madsen, N. Secouris, Solitary liver abscess in a newborn, *Surgery* 47 (1960) 1005.
- [28] C. Martin, et al., Abcès gazeux du foie avec coagulopathie chez le nouveau-né: guérison (à propos de 2 observations), *Bord. Med.* 5 (1972) 1181.
- [29] M.D. Pouyanne, Abcès du foie à staphylococques chez un nouveau-né, compliqué de suppuration sous-phrénique puis de péritonite à évolution subaiguë, *Guerison J. Med. Bordeaux* 130 (1953) 929.
- [30] L.J. Pyrtrek, S.A. Bartus, Hepatic pyemia, *N. Engl. J. Med.* 272 (1965) 551.
- [31] K. Sharma, R. Kumar, Solitary abscess of the liver in a newborn infant, *Surgery* 61 (1967) 812.
- [32] J.W. Williams, et al., Liver abscess in newborn: complication of umbilical vein catheterization, *Am. J. Dis. Child.* 125 (1973) 111.
- [33] D.W. Beaven, Staphylococcal peritonitis in the newborn, *Lancet* 1 (1958) 869.
- [34] J.R. Fraga, B.A. Javate, S. Venkatesan, Liver abscess and sepsis due to *Klebsiella pneumoniae* in a newborn, *Clin. Pediatr.* 13 (1974) 1081.
- [35] A.A. Tariq, N.A. Rudolph, E.J. Levin, Solitary hepatic abscess in a newborn infant: a sequel of umbilical vein catheterization and infusion of hypertonic glucose solutions, *Clin. Pediatr.* 16 (1977) 577.
- [36] P. Cushman, O.C. Ward, Solitary liver abscess in a neonate: complication of umbilical vein catheterization, *Ir. J. Med. Sci.* 147 (1978) 374.
- [37] H. Wiedersberg, P. Pawlowski, Pyelophlebitis nach Nabelvenenkatheterismus, *Monatsschr. Kinderheilkd.* 128 (1980) 128.
- [38] F. Gonzalez Rivera, M. Montoro Burgos, A. Cabrera Molina, Absceso hepático en un recién nacido, *An. Esp. Pediatr.* 23 (1985) 59.
- [39] P.W. Nars, L. Klco, C.P. Fliegel, Successful conservative management of a solitary liver abscess in a premature baby, *Helv. Paediatr. Acta* 38 (1983) 489.
- [40] L.R. Larsen, J. Raffensperger, Liver abscess, *J. Pediatr. Surg.* 14 (1979) 329.
- [41] F. Montoya, et al., Abcès du foie chez un nouveau-né guérison après ponction percutanée sous contrôle échographique, *Pediatric 38* (1983) 547.
- [42] S.M. Murphy, C.J. Baker, Solitary hepatic abscess: a delayed complication of neonatal bacteremia, *Pediatr. Infect. Dis. J.* 7 (1988) 414.
- [43] D. Anagnostakis, et al., Risk of infection associated with umbilical vein catheterization: a prospective study in 75 newborn infants, *J. Pediatr.* 86 (1975) 759.
- [44] J. Sabbaj, V. Sutter, S.M. Finegold, Anaerobic pyogenic liver abscess, *Ann. Intern. Med.* 77 (1972) 629.
- [45] W.W. Meyer, J. Lind, Postnatal changes in the portal circulation, *Arch. Dis. Child.* 41 (1966) 606.
- [46] J. Hageman, et al., Congenital tuberculosis: critical reappraisal of clinical findings and diagnostic procedures, *Pediatrics* 66 (1980) 980.
- [47] M.R. Hughesdon, Congenital tuberculosis, *Arch. Dis. Child.* 21 (1946) 121.
- [48] M.F. Cantwell, et al., Brief report: congenital tuberculosis, *N. Engl. J. Med.* 330 (1994) 1051.
- [49] N. Abughal, et al., Congenital tuberculosis, *Pediatr. Infect. Dis. J.* 13 (1994) 738.
- [50] B. Simma, et al., Bacille Calmette-Guérin-associated hepatitis, *Eur. J. Pediatr.* 150 (1991) 423.
- [51] T.N. Lide, Congenital tularemia, *Arch. Pathol.* 43 (1947) 165.
- [52] J.C. Regan, A. Litvak, C. Regan, Intrauterine transmission of anthrax, *JAMA* 80 (1923) 1769.
- [53] H.T. Hicks, H. French, Typhoid fever and pregnancy with special reference to foetal infection, *Lancet* 1 (1905) 1491.
- [54] M. Sarram, et al., Intrauterine fetal infection with *Brucella melitensis* as a possible cause of second-trimester abortion, *Am. J. Obstet. Gynecol.* 119 (1974) 657.
- [55] A. Brim, A bacteriologic study of 100 stillborn and dead newborn infants, *J. Pediatr.* 15 (1939) 680.

- [56] E. Madan, M.P. Meyer, A.J. Amortegui, Isolation of genital mycoplasmas and *Chlamydia trachomatis* in stillborn and neonatal autopsy material, Arch. Pathol. Lab. Med. 112 (1988) 749.
- [57] E.H. Oppenheimer, J.B. Hardy, Congenital syphilis in the newborn infant: clinical and pathological observations in recent cases, Johns Hopkins Med. J. 129 (1971) 63.
- [58] J.H. Stokes, H. Beerman, N.R. Ingraham, Modern Clinical Syphilology: Diagnosis, Treatment, Case Study, third ed., WB Saunders, Philadelphia, 1944.
- [59] A. Venter, et al., Liver function in early congenital syphilis. Does penicillin cause a deterioration? J. Pediatr. Gastroenterol. Nutr. 12 (1991) 310.
- [60] S. Lindsay, J.W. Luke, Fetal leptospirosis (Weil's disease) in a newborn infant: case of intrauterine fetal infection with report of an autopsy, J. Pediatr. 34 (1949) 90.
- [61] V. Topciu, et al., Voie transplacentaire dans un cas de leptospirose humaine, Gynecol. Obstet. 65 (1966) 617.
- [62] H.O. Gsell Jr., et al., Intrauterine leptospirosis pomona: 1st reported case of an intrauterine transmitted and cured leptospirosis, Dtsch. Med. Wochenschr. 96 (1971) 1263.
- [63] H.H.W. Cramer, Abortus beim Leptospirosis canicola, Arch. Gynecol. 177 (1950) 167.
- [64] H. Chung, et al., Transplacental or congenital infection of leptospirosis: clinical and experimental observations, Chin. Med. J. 82 (1963) 777.
- [65] P.C. Fuchs, A.A. Oyama, Neonatal relapsing fever due to transplacental transmission of *Borrelia*, JAMA 208 (1969) 690.
- [66] P. Yagupsky, S. Moses, Neonatal *Borrelia* species infection (relapsing fever), Am. J. Dis. Child. 139 (1985) 74.
- [67] P.C. Fuchs, Personal communication, 1973.
- [68] K. Weber, et al., *Borrelia burgdorferi* in a newborn despite oral penicillin for Lyme borreliosis during pregnancy, Pediatr. Infect. Dis. J. 7 (1988) 286.
- [68a] Lyme disease, in L.K. Pickering et al., (Ed.), Red Book 2006 Report of the Committee on Infectious Diseases, twenty-seventh ed., American Academy of Pediatrics, Elk Grove Village, IL, 2006, pp. 428–433.
- [69] A.C. Steere, Lyme disease, N. Engl. J. Med. 321 (1989) 586.
- [70] J.K. Lipinski, et al., Falciform ligament abscess in the infant, J. Pediatr. Surg. 20 (1985) 556.
- [71] K. Betke, H. Richarz, Nabelsepsis mit Pyelphlebitis, multiplen Leberabszessen, Lungenabszessen, und Osteomyelitis. Ausgang in Heilung, Monatsschr. Kinderheilkd. 105 (1957) 70.
- [72] R.I.K. Elliott, The ductus venosus in neonatal infection, Proc. R. Soc. Med. 62 (1969) 321.
- [73] C.G. McKenzie, Pyogenic infection of liver secondary to infection in the portal drainage area, BMJ 4 (1964) 1558.
- [74] K. Menzel, H. Buttenberg, Pyelophlebitis mit multiplen Leberabszessen als Komplikation mehrfacher Sondierung der Nabelvene, Kinderarztl. Prax. 40 (1972) 14.
- [75] S. Sarrut, J. Alain, F. Alison, Les complications précoces de la perfusion par la veine ombilicale chez le premature, Arch. Fr. Pediatr. 26 (1969) 651.
- [76] B. Santerne, et al., Diagnostic et traitement d'une abcédation hépatique néo-natale multifocale par l'échographie, Presse. Med. 16 (1987) 12.
- [77] J. Scott, Iatrogenic lesions in babies following umbilical vein catheterization, Arch. Dis. Child. 40 (1965) 426.
- [78] A. Shaw, S. Pierog, "Ectopic" liver in the umbilicus: an unusual focus of infection in a newborn infant, Pediatrics 44 (1969) 448.
- [79] J.E. Morison, Umbilical sepsis and acute interstitial hepatitis, J. Pathol. Bacteriol. 56 (1944) 531.
- [80] J. Bernstein, A.K. Brown, Sepsis and jaundice in early infancy, Pediatrics 29 (1962) 873.
- [81] R.G.F. Parker, Jaundice and infantile diarrhea, Arch. Dis. Child. 33 (1958) 330.
- [82] J.L. Gwinn, F.A. Lee, Radiologic case of the month: pyogenic liver abscess, Am. J. Dis. Child. 123 (1972) 50.
- [83] D.J. Martin, Neonatal disorders diagnosed with ultrasound, Clin. Perinatol. 12 (1985) 219.
- [84] V.M. Pineiro-Carrero, J.M. Andres, Morbidity and mortality in children with pyogenic liver abscess, Am. J. Dis. Child. 143 (1989) 1424.
- [85] K.H. Caron, Magnetic resonance imaging of the pediatric abdomen, Semin. Ultrasound CT MR 12 (1991) 448.
- [86] R.A. Halvorsen Jr., et al., Hepatic abscess: sensitivity of imaging tests and clinical findings, Gastrointest. Radiol. 13 (1988) 135.
- [87] J.C. Weinreb, et al., Imaging the pediatric liver: MRI and CT, AJR Am. J. Roentgenol. 147 (1986) 785.
- [88] M.D. Cohen, Clinical utility of magnetic resonance imaging in pediatrics, Am. J. Dis. Child. 140 (1986) 947.
- [89] M.J. Diamant, et al., Percutaneous aspiration and catheter drainage of abscesses, J. Pediatr. 108 (1986) 204.
- [90] Z. Rubinstein, et al., Ultrasound and computed tomography in the diagnosis and drainage of abscesses and other fluid collections, Isr. J. Med. Sci. 19 (1983) 1050.
- [91] T.B. Reynolds, Medical treatment of pyogenic liver abscess, Ann. Intern. Med. 96 (1982) 373.
- [92] R. Loh, G. Wallace, Y.H. Thong, Successful non-surgical management of pyogenic liver abscess, Scand. J. Infect. Dis. 19 (1987) 137.
- [93] C.M. Keidl, M.J. Chusid, Splenic abscesses in childhood, Pediatr. Infect. Dis. J. 8 (1989) 368.
- [94] A. Bowen, Acute gallbladder dilatation in a neonate: emphasis on ultrasonography, J. Pediatr. Gastroenterol. Nutr. 3 (1984) 304.
- [95] J.F. Goldthorn, D.W. Thomas, A.D. Ramos, Hydrops of the gallbladder in stressed premature infants, Clin. Res. 28 (1980) 122A.
- [96] P.W. Brill, P. Winchester, M.S. Rosen, Neonatal cholelithiasis, Pediatr. Radiol. 12 (1982) 285.
- [97] J. Callahan, et al., Cholelithiasis in infants: association with total parenteral nutrition and furosemide, Radiology 143 (1982) 437.
- [98] M.S. Keller, et al., Spontaneous resolution of cholelithiasis in infants, Radiology 157 (1985) 345.
- [99] W.J. Schirmer, E.R. Grisoni, M.W.L. Gauderer, The spectrum of cholelithiasis in the first year of life, J. Pediatr. Surg. 24 (1989) 1064.
- [100] D. Debray, et al., Cholelithiasis in infancy: a study of 40 cases, J. Pediatr. 122 (1993) 38.
- [101] J. Neu, A. Arvin, R.L. Ariagno, Hydrops of the gallbladder, Am. J. Dis. Child. 134 (1980) 891.
- [102] E.A. Leichty, et al., Normal gallbladder appearing as abdominal mass in neonates, Am. J. Dis. Child. 136 (1982) 468.
- [103] K.J. Peevy, H.J. Wiseman, Gallbladder distension in septic neonates, Arch. Dis. Child. 57 (1982) 75.
- [104] T. Dutta, et al., Gallbladder disease in infancy and childhood, Prog. Pediatr. Surg. 8 (1975) 109.
- [105] R.L. Saldanha, C.A. Stein, A.E. Kopelman, Gallbladder distention in ill preterm infants, Am. J. Dis. Child. 137 (1983) 1179.
- [106] M. El-Shafie, C.L. Mah, Transient gallbladder distention in sick premature infants: the value of ultrasonography and radionuclide scintigraphy, Pediatr. Radiol. 16 (1986) 468.
- [107] N. Modi, A.J. Keay, Neonatal gallbladder distention, Arch. Dis. Child. 57 (1982) 562.
- [108] J.B. Amodio, et al., Neonatal hydrops of the gallbladder: evaluation by cholescintigraphy and ultrasonography, N. Y. State J. Med. 85 (1985) 565.
- [109] J.P. McGahan, H.E. Phillips, K.L. Cox, Sonography of the normal pediatric gallbladder and biliary tract, Radiology 144 (1982) 873.
- [110] J.O. Haller, Sonography of the biliary tract in infants and children, AJR Am. J. Roentgenol. 157 (1991) 1051.
- [111] K.J. Guthrie, G.L. Montgomery, Infections with *Bacterium enteritidis* in infancy with the triad of enteritis, cholecystitis, and meningitis, J. Pathol. Bacteriol. 49 (1939) 393.
- [112] W. Faller, J.E. Berkelhamor, J.R. Esterly, Neonatal biliary tract infection coincident with maternal methadone therapy, Pediatrics 48 (1971) 997.
- [113] P.N. Jamieson, D.G. Shaw, Empyema of gallbladder in an infant, Arch. Dis. Child. 50 (1975) 482.
- [114] L.A. Arnsperger, J.G. Martin, H.O. Krempin, Acute noncalculous cholecystitis in children: report of a case in a 17-day-old infant, Am. J. Surg. 100 (1960) 103.
- [115] J.L. Ternberg, J.P. Keating, Acute acalculous cholecystitis: complication of other illnesses in childhood, Arch. Surg. 110 (1975) 543.
- [116] R.F. Crystal, R.L. Fink, Acute acalculous cholecystitis in childhood: a report of two cases, Clin. Pediatr. 10 (1971) 423.
- [117] A.E. Robinson, et al., Cholecystitis and hydrops of the gallbladder in the newborn, Radiology 122 (1977) 749.
- [118] W.H. Snyder Jr., L. Chaffin, L. Oettinger, Cholelithiasis and perforation of the gallbladder in an infant, with recovery, JAMA 149 (1952) 1645.
- [119] M.E. Washburn, P.J. Barcia, Uncommon cause of a right upper quadrant abdominal mass in a newborn: acute cholecystitis, Am. J. Surg. 140 (1980) 704.
- [120] W.A. Thurston, E.N. Kelly, M.M. Silver, Acute acalculous cholecystitis in a premature infant treated with parenteral nutrition, Can. Med. Assoc. J. 135 (1986) 332.
- [121] R. Pieretti, A.W. Auld, C.A. Stephens, Acute cholecystitis in children, Surg. Obstet. Gynecol. 140 (1975) 16.
- [122] B.A. Hanson, G.H. Mahour, M.M. Woolley, Diseases of the gallbladder in infancy and childhood, J. Pediatr. Surg. 6 (1971) 277.
- [123] P.A. Dewan, K.B. Stokes, J.R. Solomon, Paediatric acalculous cholecystitis, Pediatr. Surg. Int. 2 (1987) 120.
- [124] J. Denes, et al., Die Frühgeborenen-Appendicitis, Z. Kinderchir. 5 (1968) 400.
- [125] V.C. Traynelis, E.E. Hrabovsky, Acalculous cholecystitis in the neonate, Am. J. Dis. Child. 139 (1985) 893.
- [126] R.D. Aach, Cholecystitis in childhood, in: R.D. Feigin, J.D. Cherry (Eds.), Textbook of Pediatric Infectious Diseases, second ed., WB Saunders, Philadelphia, 1987, pp. 742–743.
- [127] R. Wyllie, J.F. Fitzgerald, Bacterial cholangitis in a 10-week-old infant with fever of undetermined origin, Pediatrics 65 (1980) 164.
- [128] D.M.O. Becroft, Biliary atresia associated with prenatal infection by *Listeria monocytogenes*, Arch. Dis. Child. 47 (1972) 656.
- [129] B.E. Favara, D.R. Akers, R.A. Franciosi, Adrenal abscess in a neonate, J. Pediatr. 77 (1970) 682.
- [130] C. Mondor, et al., Nonsurgical management of neonatal adrenal abscess, J. Pediatr. Surg. 23 (1988) 1048.
- [131] A. François, et al., Abscès surrénalien neonatal à *Proteus mirabilis*, Arch. Fr. Pediatr. 48 (1991) 559.
- [132] J.R. Lizardo-Barahona, J. Nieto-Zermño, E. Bracho-Blanchet, Absceso adrenal en el recién nacido: informe de un caso y revisión de la literatura, Bol. Med. Hosp. Infant. Mex. 47 (1990) 401.
- [133] J.M. Torres-Simon, et al., Absceso suprarrenal en el recién nacido, An. Esp. Pediatr. 31 (1989) 601.
- [134] O. Zamir, et al., Adrenal abscess: a rare complication of neonatal adrenal hemorrhage, Pediatr. Surg. Int. 2 (1987) 117.

- [135] J.M. Van de Water, E.W. Fonkalsrud, Adrenal cysts in infancy, *Surgery* 60 (1966) 1267.
- [136] W.J. Blankenship, et al., Suprarenal abscess in the neonate: a case report and review of diagnosis and management, *Pediatrics* 55 (1975) 239.
- [137] D.M. Gibbons, et al., Abdominal flank mass in the neonate, *J. Urol.* 119 (1978) 671.
- [138] G.O. Atkinson Jr., et al., Adrenal abscess in the neonate, *Radiology* 155 (1985) 101.
- [139] A. Carty, P. Stanley, Bilateral adrenal abscesses in a neonate, *Pediatr. Radiol.* 1 (1973) 63.
- [140] K.M. Walker, W.F. Coyer, Suprarenal abscess due to group B Streptococcus, *J. Pediatr.* 94 (1979) 970.
- [141] R. Camilleri, et al., Abcédation d'un hématoome surrénal avec hypertension artérielle, *Arch. Fr. Pediatr.* 41 (1984) 705.
- [142] K. Rajani, S.R. Shapiro, B.W. Goetsman, Adrenal abscess: complication of supportive therapy of adrenal hemorrhage in the newborn, *J. Pediatr. Surg.* 15 (1980) 676.
- [143] R.G. Wells, J.R. Sty, N.B. Hodgson, Suprarenal abscess in the neonate: technetium-99m glucoheptonate imaging, *Clin. Nucl. Med.* 11 (1986) 32.
- [144] B.A. Cadarso, A.M.R. Mialdea, Absceso adrenal en la epoca neonatal, *An. Esp. Pediatr.* 21 (1984) 706.
- [145] S. Ohta, et al., Neonatal adrenal abscess due to *Bacteroides*, *J. Pediatr.* 93 (1978) 1063.
- [146] B.A. Bekdash, M.S. Slim, Adrenal abscess in a neonate due to gas-forming organisms: a diagnostic dilemma, *Z. Kinderchir.* 32 (1981) 184.
- [147] M. Gross, P.K. Kottmeier, K. Waterhouse, Diagnosis and treatment of neonatal adrenal hemorrhage, *J. Pediatr. Surg.* 2 (1967) 308.
- [148] V. Vigi, et al., Suprarenal abscess in a newborn, *Helv. Paediatr. Acta* 36 (1981) 263.
- [149] S. Suri, et al., Adrenal abscess in a neonate presenting as a renal neoplasm, *Br. J. Urol.* 54 (1982) 565.
- [150] C.A. Mittelstaedt, et al., The sonographic diagnosis of neonatal adrenal hemorrhage, *Radiology* 131 (1979) 453.
- [151] A. Rey, et al., Hemorragia suprarenal encapsulada en el recién nacido: estudio de ocho casos, *An. Esp. Pediatr.* 21 (1984) 238.
- [152] J. Black, D.I. Williams, Natural history of adrenal hemorrhage in the newborn, *Arch. Dis. Child.* 48 (1973) 173.
- [153] O. Iklof, W. Mortensson, B. Sandstedt, Suprarenal haematoma versus neuroblastoma complicated by haemorrhage: a diagnostic dilemma in the newborn, *Acta Radiol. Diagn.* 27 (1986) 3.
- [154] S.J. White, et al., Sonography of neuroblastoma, *Am. J. Radiol.* 141 (1983) 465.
- [155] W.E. Etherington-Wilson, Appendicitis in newborn: report of a case 16 days old, *Proc. R. Soc. Med.* 38 (1945) 186.
- [156] P. Puri, B. O'Donnell, Appendicitis in infancy, *J. Pediatr. Surg.* 13 (1978) 173.
- [157] B. Landaa, Diagnosis of appendicitis in young children, *Tidsskr. Nor. Laegeforen.* 68 (1948) 335.
- [158] G. Reuter, I. Krause, Beitrag zur Problematik der Appendizitis des Neugeborenen, *Kinderarztl. Prax.* 47 (1975) 289.
- [159] W.J. Norris, Appendicitis in children, *West J Surg Obstet Gynecol* 54 (1946) 183.
- [160] J.M. Parsons, B.G. Miscall, C.K. McSherry, Appendicitis in the newborn infant, *Surgery* 67 (1970) 841.
- [161] W. Schaupp, E.G. Clausen, P.K. Ferrier, Appendicitis during the first month of life, *Surgery* 48 (1960) 805.
- [162] E.G. Stanley-Brown, Acute appendicitis during the first five years of life, *Am. J. Dis. Child.* 108 (1964) 134.
- [163] W.H. Snyder Jr., L. Chaffin, Appendicitis during the first 2 years of life: report of 21 cases and review of 447 cases from the literature, *Arch. Surg.* 64 (1952) 549.
- [164] I.A. Fields, M.J. Naiditch, P.E. Rothman, Acute appendicitis in infants, *Am. J. Dis. Child.* 93 (1957) 287.
- [165] W. Dick, H.J. Hirt, W. Vogel, Die Appendizitis im Säuglings- und Kleinkindesalter, *Fortschr. Med.* 94 (1976) 125.
- [166] R.E. Gross, *The Surgery of Infancy and Childhood: Its Principles and Techniques*, WB Saunders, Philadelphia, 1953.
- [167] J.L. Grosfeld, M. Weinberger, H.W. Clatworthy Jr., Acute appendicitis in the first two years of life, *J. Pediatr. Surg.* 8 (1973) 285.
- [168] J.S. Janik, H.V. Firor, Pediatric appendicitis: a 20-year study of 1, 640 children at Cook County (Illinois) Hospital, *Arch. Surg.* 114 (1979) 717.
- [169] C.D. Benson, J.J. Coury Jr., D.R. Hagge, Acute appendicitis in infants: fifteen-year study, *Arch. Surg.* 64 (1952) 561.
- [170] M. Massad, et al., Neonatal appendicitis: case report and a review of the English literature, *Z. Kinderchir.* 41 (1986) 241.
- [171] G.R. Schorlemmer, C.A. Herbst Jr., Perforated neonatal appendicitis, *South. Med. J.* 76 (1983) 536.
- [172] F. Diess, *Die Appendizitis im Basler Kinderspital*. Basle Dissertations, 1908, Case No. 36, p 63. Cited in reference 138 and in Abt IA: Appendicitis in infants, *Arch. Pediatr.* 34 (1917) 641.
- [173] R.H. Bartlett, A.J. Eraklis, R.H. Wilkinson, Appendicitis in infancy, *Surg. Gynecol. Obstet.* 130 (1970) 99.
- [174] N.R.G. Broadbent, J.L. Jardine, Acute appendicitis in a premature infant: a case report, *Aust. N. Z. J. Surg.* 40 (1971) 362.
- [175] L.R. Bryant, et al., Appendicitis and appendiceal perforation in neonates, *Am. Surg.* 36 (1970) 523.
- [176] R.D.G. Creery, Acute appendicitis in the newborn, *BMJ* 1 (1953) 871.
- [177] R.P. Hardman, D. Bowerman, Appendicitis in the newborn, *Am. J. Dis. Child.* 105 (1963) 99.
- [178] F. Klimt, G. Hartmann, Appendicitis perforata mit tiefsitzenden Dünnarmverschluss beim Neugeborenen, *Paediatr. Prax.* 1 (1962) 271.
- [179] G. Kolb, E.L. Schaeffer, Über Appendizitis mit Perforation in den ersten Lebenswochen, *Kinderarztl. Prax.* 1 (1955) 1.
- [180] R.E. Liechti, W.H. Snyder Jr., Acute appendicitis under age 2, *Am. Surg.* 29 (1963) 92.
- [181] S.C. Meigher, A.W. Lucas, Appendicitis in the newborn: case report, *Ann. Surg.* 136 (1952) 1044.
- [182] J.F. Meyer, Acute gangrenous appendicitis in a premature infant, *J. Pediatr.* 41 (1952) 343.
- [183] R. Neve, N.F. Quenville, Appendicitis with perforation in a 12-day-old infant, *Can. Med. Assoc. J.* 94 (1966) 447.
- [184] M.A. Nilforoushan, Fever and ascites in a newborn, *Clin. Pediatr.* 14 (1975) 878.
- [185] M. Nuri, W.C. Hecker, W. Duckert, Beitrag zur Appendizitis im Neugeborenenalter, *Z. Kinderheilkd.* 91 (1964) 1.
- [186] G.F. Parkhurst, S.C. Wagoner, Neonatal acute appendicitis, *N. Y. State J. Med.* 69 (1969) 1929.
- [187] S.J. Phillips, B. Cohen, Acute perforated appendicitis in newborn children, *N. Y. State J. Med.* 71 (1971) 985.
- [188] A.L. Smith, R.A. MacMahon, Perforated appendix complicating rhesus immunization in a newborn infant, *Med. J. Aust.* 2 (1969) 602.
- [189] J. Tabrisky, R. Westerfeld, J. Cavanaugh, Appendicitis in the newborn, *Am. J. Dis. Child.* 111 (1966) 557.
- [190] H. Vinz, U. Erben, H. Winkelvoss, Neugeborenen peritonitis, *Bruns. Beitr. Klin. Chir.* 215 (1967) 321.
- [191] R.H. Walker, Appendicitis in the newborn infant, *J. Pediatr.* 51 (1958) 429.
- [192] C.D. Morehead, P.W. Houck, Epidemiology of *Pseudomonas* infections in a pediatric intensive care unit, *Am. J. Dis. Child.* 124 (1972) 564.
- [193] J.Q. Trojanowski, et al., Fatal postoperative acute appendicitis in a neonate with congenital heart disease, *J. Pediatr. Surg.* 16 (1981) 85.
- [194] A. Ayalon, et al., Acute appendicitis in a premature baby, *Acta Chir. Scand.* 145 (1979) 285.
- [195] W. Schellerer, K. Schwemmler, R. Decker, Perforierte Appendizitis bei einem Frühgeborenen im Alter von 14 Tagen, *Z. Kinderchir.* 9 (1971) 434.
- [196] E.S. Golladay, et al., Intestinal obstruction from appendiceal abscess in a newborn infant, *J. Pediatr. Surg.* 13 (1978) 175.
- [197] V. Hemalatha, L. Spitz, Neonatal appendicitis, *Clin. Pediatr.* 18 (1979) 621.
- [198] P. Tucci, et al., Congenital uretero-pelvic junction obstruction associated with unsuspected acute perforated appendicitis in a neonate, *J. Urol.* 120 (1978) 247.
- [199] G.L. Fowkes, Neonatal appendicitis, *BMJ* 1 (1978) 997.
- [200] M.S. Kwong, M. Dinner, Neonatal appendicitis masquerading as necrotizing enterocolitis, *J. Pediatr.* 96 (1980) 917.
- [201] W.L. Shaul, Clues to the early diagnosis of neonatal appendicitis, *J. Pediatr.* 98 (1981) 473.
- [202] R. Grussner, et al., Appendicitis in childhood, *Monatsschr. Kinderheilkd.* 133 (1985) 158.
- [203] H.A. Lassiter, M.H. Werner, Neonatal appendicitis, *South. Med. J.* 76 (1983) 1173.
- [204] M.J. Carol, et al., Appendicitis neonatal: a report of a new case, *An. Esp. Pediatr.* 20 (1984) 80.
- [205] N.M.A. Bax, et al., Perforation of the appendix in the neonatal period, *J. Pediatr. Surg.* 15 (1980) 200.
- [206] W.L. Buntain, Neonatal appendicitis mistaken for necrotizing enterocolitis, *South. Med. J.* 75 (1982) 1155.
- [207] J.J. Heydenrych, D.F. DuToit, Unusual presentations of acute appendicitis in the neonate: a report of 2 cases, *S. Afr. Med. J.* 62 (1982) 1003.
- [208] Radiological case of the month: neonatal appendicitis with perforation, *Am. J. Dis. Child.* 145 (1991) 111.
- [209] O.P. Pathania, et al., Fatal neonatal perforation of appendix, *Indian Pediatr.* 17 (1982) 1166.
- [210] N.K. Arora, et al., Neonatal appendicitis: a rare cause of surgical emergency in preterm babies, *Indian Pediatr.* 28 (1991) 1330.
- [211] M.N. Srouji, J. Chatten, C. David, Pseudodiverticulitis of the appendix with neonatal Hirschsprung disease, *J. Pediatr.* 93 (1978) 988.
- [212] J. Arliss, L.O. Holgersen, Neonatal appendiceal perforation and Hirschsprung's disease, *J. Pediatr. Surg.* 25 (1990) 694.
- [213] R.M. Kliegman, A.A. Fanaroff, Necrotizing enterocolitis, *N. Engl. J. Med.* 310 (1984) 1093.
- [214] M.N. Srouji, B.E. Buck, Neonatal appendicitis: ischemic infarction in incarcerated inguinal hernia, *J. Pediatr. Surg.* 13 (1978) 177.
- [215] P. Charif, Perforated appendicitis in premature infants: a case report and review of the literature, *Johns Hopkins Med. J.* 125 (1969) 92.
- [216] H.H. Stone, S.L. Sanders, J.D. Martin Jr., Perforated appendicitis in children, *Surgery* 69 (1971) 673.
- [217] J.E. Dennis, et al., Nosocomial *Rhizopus* infection (zygomycosis) in children, *J. Pediatr.* 96 (1980) 824.
- [218] K. Buschard, A. Kjaeldgaard, Investigation and analysis of the position, fixation, length, embryology of the vermiform appendix, *Acta Chir. Scand.* 139 (1973) 293.
- [219] W.R. Jones, M.D. Kaye, R.M.Y. Ing, The lymphoid development of the fetal and neonatal appendix, *Biol. Neonate* 20 (1972) 334.
- [220] G.M. Smith, Inflammatory changes in the appendix during early infancy, *Am. J. Dis. Child.* 1 (1911) 299.

- [221] W.B. Hill, C.C. Mason, Prenatal appendicitis with rupture and death, *Am. J. Dis. Child.* 29 (1925) 86.
- [222] W.J. Corcoran, Prenatal rupture of the appendix, *Am. J. Dis. Child.* 39 (1930) 277.
- [223] W.F. Jackson, A case of prenatal appendicitis, *Am. J. Med. Sci.* 127 (1904) 710.
- [224] E.W. Kümmell, Cited in Etherington-Wilson WE. Appendicitis in newborn: report of a case 16 days old, *Proc. R. Soc. Med.* 38 (1945) 186.
- [225] L.W. Martin, P.M. Glen, Prenatal appendiceal perforation: a case report, *J. Pediatr. Surg.* 21 (1986) 73.
- [226] R.H. Wilkinson, R.H. Bartlett, A.J. Eraklis, Diagnosis of appendicitis in infancy: the value of abdominal radiographs, *Am. J. Dis. Child.* 118 (1969) 687.
- [227] T.M. Holder, L.L. Leape, The acute surgical abdomen in the neonate, *N. Engl. J. Med.* 278 (1968) 605.
- [228] J.H.T. Chang, The use of antibiotics in pediatric abdominal surgery, *Pediatr. Infect. Dis.* 3 (1984) 195.
- [229] M.J. Bell, Peritonitis in the newborn—current concepts, *Pediatr. Clin. North Am.* 32 (1985) 1181.
- [230] A.J. Barson, A postmortem study of infection in the newborn from 1976 to 1988, in: J. deLouvois, D. Harvey (Eds.), *Infection of the Newborn*, John Wiley & Sons, New York, 1990, pp. 13–34.
- [231] E.W. Fonkalsrud, D.G. Ellis, H.W. Clatworthy Jr., Neonatal peritonitis, *J. Pediatr. Surg.* 1 (1966) 227.
- [232] J.R. Lloyd, The etiology of gastrointestinal perforations in the newborn, *J. Pediatr. Surg.* 4 (1969) 77.
- [233] W.S. McDougal, R.J. Izant, R.M. Zollinger Jr., Primary peritonitis in infancy and childhood, *Ann. Surg.* 181 (1975) 310.
- [234] P.P. Rickham, Peritonitis in the neonatal period, *Arch. Dis. Child.* 30 (1955) 23.
- [235] J. Denes, J. Leb, Neonatal peritonitis, *Acta Paediatr. Acad. Sci. Hung.* 10 (1969) 297.
- [236] R. Daum, U. Schütze, H. Hoffman, Mortality of preoperative peritonitis in newborn infants without intestinal obstruction, *Prog. Pediatr. Surg.* 13 (1979) 267.
- [237] U. Schütze, K.H. Fey, G. Hess, Die Peritonitis im Neugeborenen-, Säuglings-, und Kindesalter, *Munch. Med. Wochenschr.* 116 (1974) 1201.
- [238] J. Prevot, G. Grosdidier, M. Schmitt, Fatal peritonitis, *Prog. Pediatr. Surg.* 13 (1979) 257.
- [239] B. Singer, B. Hammar, Neonatal peritonitis, *S. Afr. Med. J.* 46 (1972) 987.
- [240] O.J. Hensley, C.A. Hart, R.W.I. Cooke, Serious infection in a neonatal intensive care unit: a two-year survey, *J. Hyg. (Camb)* 95 (1985) 289.
- [241] M.A. Valdes-Dapeña, J.B. Arey, The causes of neonatal mortality: an analysis of 501 autopsies on newborn infants, *J. Pediatr.* 77 (1970) 366.
- [242] M.B. Duggan, M.S. Khwaja, Neonatal primary peritonitis in Nigeria, *Arch. Dis. Child.* 50 (1975) 130.
- [243] M.J. Bell, Perforation of the gastrointestinal tract and peritonitis in the neonate, *Surg. Gynecol. Obstet.* 160 (1985) 20.
- [244] A.G. Birtch, A.G. Coran, R.E. Gross, Neonatal peritonitis, *Surgery* 61 (1967) 305.
- [245] M. Lacheretz, et al., Péritonite néo-natale par perforation gastrique: à propos de 21 observations, *Chirurgie* 109 (1983) 887.
- [246] D.L. Mollitt, J.J. Tepas, J.L. Talbert, The microbiology of neonatal peritonitis, *Arch. Surg.* 123 (1988) 176.
- [247] J.E.S. Scott, Intestinal obstruction in the newborn associated with peritonitis, *Arch. Dis. Child.* 38 (1963) 120.
- [248] H.E. Thelander, Perforation of the gastrointestinal tract of the newborn infant, *Am. J. Dis. Child.* 58 (1939) 371.
- [249] G. Dinari, H. Haimov, M. Geiffman, Umbilical arteritis and phlebitis with scrotal abscess and peritonitis, *J. Pediatr. Surg.* 6 (1971) 176.
- [250] I. Forshall, Septic umbilical arteritis, *Arch. Dis. Child.* 32 (1957) 25.
- [251] E.G. Chadwick, S.T. Shulman, R. Yogeve, Peritonitis as a late manifestation of group B streptococcal disease in newborns, *Pediatr. Infect. Dis.* 2 (1983) 142.
- [252] T.M. Reyna, Primary group B streptococcal peritonitis presenting as an incarcerated inguinal hernia in a neonate, *Clin. Pediatr.* 25 (1987) 422.
- [253] W. Serlo, E. Heikkinen, K. Kouvalainen, Group A streptococcal peritonitis in infancy, *Ann. Chir. Gynaecol.* 74 (1985) 183.
- [254] D.E. Johnson, et al., *Candida* peritonitis in a newborn infant, *J. Pediatr.* 97 (1980) 298.
- [255] M. Kaplan, et al., Necrotizing bowel disease with *Candida* peritonitis following severe neonatal hypothermia, *Acta Paediatr. Scand.* 79 (1990) 876.
- [256] K.M. Butler, M.A. Bench, C.J. Baker, Amphotericin B as a single agent in the treatment of systemic candidiasis in neonates, *Pediatr. Infect. Dis. J.* 9 (1990) 51.
- [257] L. MacDonald, C.J. Baker, C. Chenoweth, Risk factors for candidemia in a children's hospital, *Clin. Infect. Dis.* 26 (1998) 642.
- [258] I.A. Abt, Fetal peritonitis, *Med. Clin. North Am.* 15 (1931) 611.
- [259] E.Y. Pan, et al., Radiographic diagnosis of meconium peritonitis: a report of 200 cases including six fetal cases, *Pediatr. Radiol.* 13 (1983) 199.
- [260] L.O. Hølgersen, The etiology of spontaneous gastric perforation of the newborn: a reevaluation, *J. Pediatr. Surg.* 16 (1981) 608.
- [261] P.P. Rickham, Neugeborenen-peritonitis, *Langenbecks Arch. Klin. Chir.* 292 (1959) 427.
- [262] R. Fowler, Primary peritonitis: changing aspects 1956–1970, *Aust. Paediatr. J.* 7 (1971) 73.
- [263] C.L. Wells, M.A. Maddaus, R.L. Simmons, Proposed mechanisms for the translocation of intestinal bacteria, *Rev. Infect. Dis.* 10 (1988) 958.
- [264] R. Wilson, et al., Short communication: age at onset of necrotizing enterocolitis: an epidemiologic analysis, *Pediatr. Res.* 16 (1982) 82.
- [265] J.M. Caralaps-Riera, B.D. Cohn, Bowel perforation after exchange transfusion in the neonate: review of the literature and report of a case, *Surgery* 68 (1970) 895.
- [266] R.J. Touloukian, A. Kadar, R.P. Spencer, The gastrointestinal complications of neonatal umbilical venous exchange transfusion: a clinical and experimental study, *Pediatrics* 51 (1973) 36.
- [267] R.M. Freedman, et al., A half century of neonatal sepsis at Yale: 1928 to 1978, *Am. J. Dis. Child.* 135 (1981) 140.
- [268] N.T. Griscom, et al., Diagnostic aspects of neonatal ascites: report of 27 cases, *AJR Am. J. Roentgenol.* 128 (1977) 961.
- [269] A.M. Kosloske, J.R. Lilly, Paracentesis and lavage for diagnosis of intestinal gangrene in neonatal necrotizing enterocolitis, *J. Pediatr. Surg.* 13 (1978) 315.
- [270] U. Töllner, F. Pohlandt, Aszitespunktion zur Differentialdiagnose beim akuten Abdomen des Neugeborenen, *Klin. Paediatr.* 196 (1984) 319.
- [271] J.B.J. McKendry, W.K. Lindsay, M.C. Gerstein, Congenital defects of the lymphatics in infancy, *Pediatrics* 19 (1959) 21.
- [272] W. Lees, J.E. Mitchell, Bile peritonitis in infancy, *Arch. Dis. Child.* 41 (1966) 188.
- [273] M.D. Cohen, T.R. Weber, J.L. Grosfeld, Bowel perforation in the newborn: diagnosis with metrizamide, *Radiology* 150 (1984) 65.
- [274] D.L. Rosenfeld, C.E. Cordell, N. Jadeja, Retrocardiac pneumomediastinum: radiographic finding and clinical implications, *Pediatrics* 85 (1989) 92.
- [275] E.D. Wind, G.P. Pillari, W.J. Lee, Lucent liver in the newborn: a roentgenographic sign of pneumoperitoneum, *JAMA* 237 (1977) 2218.
- [276] R. Pochaczewsky, D. Bryk, New roentgenographic signs of neonatal gastric perforation, *Radiology* 102 (1972) 145.
- [277] S.S. Gellis, M. Finegold, Picture of the month: pneumoperitoneum demonstrated by transillumination, *Am. J. Dis. Child.* 130 (1976) 1237.
- [278] S. Thomas, C. Sainsbury, J.F. Murphy, Pancurionium belly, *Lancet* 2 (1984) 870.
- [279] S.H. Ein, C.A. Stephens, B.J. Reilly, The disappearance of free air after pediatric laparotomy, *J. Pediatr. Surg.* 20 (1985) 422.
- [280] B. Emanuel, P. Zlotnik, J.G. Raffensperger, Perforation of the gastrointestinal tract in infancy and childhood, *Surg. Gynecol. Obstet.* 146 (1978) 926.
- [281] K. Opitz, Beitrag zur Klinik und Pathologie der Nabelschnurinfektionen, *Arch. Kinderheilkd.* 150 (1955) 174.
- [282] V.Y.H. Yu, D.I. Tudehope, G.J. Gill, Neonatal necrotizing enterocolitis. 1. Clinical aspects. 2. Perinatal risk factors, *Med. J. Aust.* 1 (1977) 685 688.
- [283] R.C. Holman, J.K. Stehr-Green, M.T. Zelasky, Necrotizing enterocolitis mortality in the United States, 1979–85, *Am. J. Public Health* 79 (1989) 987.
- [284] N.N. Finer, R.R. Moriarty, Reply, letter to the editor, *J. Pediatr.* 96 (1980) 170.
- [285] L.S. Book, et al., Necrotizing enterocolitis in low-birth-weight infants fed an elemental formula, *J. Pediatr.* 87 (1975) 602.
- [286] J.A. O'Neill Jr., M.T. Stahlman, H.C. Meng, Necrotizing enterocolitis in the newborn: operative indications, *Ann. Surg.* 182 (1975) 274.
- [287] T.D. Moore (Ed.), *Necrotizing Enterocolitis in the Newborn Infant: Report of the Sixty-Eighth Ross Conference on Pediatric Research*, Ross Laboratories, Columbus, OH, 1975.
- [288] N.L. Virnig, J.W. Reynolds, Epidemiological aspects of neonatal necrotizing enterocolitis, *Am. J. Dis. Child.* 128 (1974) 186.
- [289] R.J. Touloukian, Neonatal necrotizing enterocolitis: an update on etiology, diagnosis, and treatment, *Surg. Clin. North Am.* 56 (1976) 281.
- [290] M.J. Bell, J.L. Ternberg, R.J. Bower, The microbial flora and antimicrobial therapy of neonatal peritonitis, *J. Pediatr. Surg.* 15 (1980) 569.
- [291] R.M. Kliegman, A.A. Fanaroff, Neonatal necrotizing enterocolitis: a nine-year experience. I. Epidemiology and uncommon observations, *Am. J. Dis. Child.* 135 (1981) 603.
- [292] H.I. Goldman, Feeding and necrotizing enterocolitis, *Am. J. Dis. Child.* 134 (1980) 553.
- [293] E.G. Brown, A.Y. Sweet, Preventing necrotizing enterocolitis in neonates, *JAMA* 240 (1978) 2452.
- [294] L.S. Book, J.J. Herbst, A. Jung, Comparison of fast- and slow-feeding rate schedules to the development of necrotizing enterocolitis, *J. Pediatr.* 89 (1976) 463.
- [295] M.J. Bell, et al., Epidemiologic and bacteriologic evaluation of neonatal necrotizing enterocolitis, *J. Pediatr. Surg.* 14 (1979) 1.
- [296] A.I. Eidelman, R.J. Inwood, Marginal comments: necrotizing enterocolitis and enteral feeding. Is too much just too much, *Am. J. Dis. Child.* 134 (1980) 545.
- [297] R.M. Kliegman, et al., Clostridia as pathogens in neonatal necrotizing enterocolitis, *J. Pediatr.* 95 (1979) 287.
- [298] E.G. Brown, A.Y. Sweet, Neonatal necrotizing enterocolitis, *Pediatr. Clin. North Am.* 29 (1982) 1149.
- [299] W.P. Kanto Jr., et al., Perinatal events and necrotizing enterocolitis in premature infants, *Am. J. Dis. Child.* 141 (1987) 167.
- [300] S.G. Ostertag, et al., Early enteral feeding does not affect the incidence of necrotizing enterocolitis, *Pediatrics* 77 (1986) 275.
- [301] C.R.B. Merritt, J.P. Goldsmith, M.J. Sharp, Sonographic detection of portal venous gas in infants with necrotizing enterocolitis, *AJR Am. J. Roentgenol.* 143 (1984) 1059.
- [302] D. Cikrit, et al., Significance of portal venous air in necrotizing enterocolitis: analysis of 53 cases, *J. Pediatr. Surg.* 20 (1985) 425.
- [303] G.C. Maguire, et al., Infections acquired by young infants, *Am. J. Dis. Child.* 135 (1981) 693.
- [304] V.Y.H. Yu, et al., Perinatal risk factors for necrotizing enterocolitis, *Arch. Dis. Child.* 59 (1984) 430.

- [305] R.D. Uauy, et al., National Institute of Child Health and Human Development Neonatal Research Network. Necrotizing enterocolitis in very low birth weight infants: biodemographic and clinical correlates, *J. Pediatr.* 119 (1991) 630.
- [306] M.C. Walsh, R.M. Kliegman, Necrotizing enterocolitis: treatment based on staging criteria, *Pediatr. Clin. North Am.* 33 (1986) 179.
- [307] M.D. Kamitsuka, M.K. Horton, M.A. Williams, The incidence of necrotizing enterocolitis after introducing standardized feeding schedules for infants between 1250 and 2500 grams and less than 35 weeks of gestation, *Pediatrics* 105 (2000) 379.
- [308] S.F. Rayyis, et al., Randomized trial of slow versus fast feed advancements on the incidence of necrotizing enterocolitis in very low birth weight infants, *J. Pediatr.* 134 (1999) 293.
- [309] T.V. Santulli, et al., Acute necrotizing enterocolitis in infancy: a review of 64 cases, *Pediatrics* 55 (1975) 376.
- [310] R.M. Kliegman, et al., Epidemiologic study of necrotizing enterocolitis among low-birth-weight infants: absence of identifiable risk factors, *J. Pediatr.* 100 (1982) 440.
- [311] B.J. Stoll, et al., Epidemiology of necrotizing enterocolitis: a case control study, *J. Pediatr.* 96 (1980) 447.
- [312] I.D. Frantz III, et al., Necrotizing enterocolitis, *J. Pediatr.* 86 (1975) 259.
- [313] E.A. Egan, et al., A prospective controlled trial of oral kanamycin in the prevention of neonatal necrotizing enterocolitis, *J. Pediatr.* 89 (1976) 467.
- [314] R. Wilson, et al., Age at onset of necrotizing enterocolitis: an epidemiologic analysis, *Pediatr. Res.* 16 (1982) 82.
- [315] P. Gerard, et al., Mortality in 504 infants weighing less than 1501 g at birth and treated in four neonatal intensive care units of South-Belgium between 1976 and 1980, *Eur. J. Pediatr.* 144 (1985) 219.
- [316] V.Y.H. Yu, et al., Necrotizing enterocolitis in very low birthweight infants: a four-year experience, *Aust. Paediatr. J.* 20 (1984) 29.
- [317] S.R. Palmer, A. Biffin, H.R. Gamsu, Outcome of neonatal necrotising enterocolitis: results of the BAPM/CDSC surveillance study, 1989–84, *Arch. Dis. Child.* 64 (1989) 388.
- [318] E. de Gamarra, et al., Necrotizing enterocolitis in full-term neonates, *Biol. Neonate* 44 (1983) 185.
- [319] E.H. Thilo, R.A. Lazarte, J.A. Hernandez, Necrotizing enterocolitis in the first 24 hours of life, *Pediatrics* 73 (1984) 476.
- [320] R. Wilson, et al., Risk factors for necrotizing enterocolitis in infants weighing more than 2, grams at birth: a case-control study, *Pediatrics* 71 (1983) 19.
- [321] Necrotizing enterocolitis, Editorial, *Lancet* 1 (1977) 459.
- [322] J.A. Bisquera, T.R. Cooper, C.L. Berseth, Impact of necrotizing enterocolitis on length of stay and hospital charges in very low birth weight infants, *Pediatrics* 109 (2002) 423.
- [323] S.R. Hintz, et al., Neurodevelopmental and growth outcomes of extremely low birth weight infants after necrotizing enterocolitis, *Pediatrics* 115 (2005) 696.
- [324] A.A. Fanaroff, M. Hack, M.C. Walsh, The NICHD Neonatal Research Network: changes in practice and outcomes during the first 15 years, *Semin. Perinatol.* 27 (2003) 281.
- [325] R. Holman, et al., Necrotizing enterocolitis hospitalizations among neonates in the United States, *Pediatr. Perinatol. Epidemiol.* 20 (2006) 498.
- [326] P.W. Lin, B.J. Stoll, Necrotising enterocolitis, *Lancet* 127 (2006) 1271.
- [327] E.E. Frey, et al., Analysis of bowel perforation in necrotizing enterocolitis, *Pediatr. Radiol.* 17 (1987) 380.
- [328] R.M. Kliegman, W.B. Pittard, A.A. Fanaroff, Necrotizing enterocolitis in neonates fed human milk, *J. Pediatr.* 95 (1979) 450.
- [329] M. Guinan, et al., Epidemic occurrence of neonatal necrotizing enterocolitis, *Am. J. Dis. Child.* 133 (1979) 594.
- [330] B. Dvorak, et al., Maternal milk reduces the severity of necrotizing enterocolitis and increases intestinal IL-10 in a neonatal rat model, *Pediatr. Res* 55 (2003) 426.
- [331] H.C. Lin, et al., Oral probiotics reduce the incidence and severity of necrotizing enterocolitis in very low birthweight infants, *Pediatrics* 115 (2005) 1–4.
- [332] A.M. Kosloske, Pathogenesis and prevention of necrotizing enterocolitis: a hypothesis based on personal observation and a review of the literature, *Pediatrics* 74 (1984) 1086.
- [333] R.P. Gaynes, et al., The role of host factors in an outbreak of necrotizing enterocolitis, *Am. J. Dis. Child.* 138 (1984) 1118.
- [334] V.K.M. Han, et al., An outbreak of *Clostridium difficile* necrotizing enterocolitis: a case for oral vancomycin therapy? *Pediatrics* 71 (1983) 935.
- [335] Neonatal necrotizing enterocolitis: current concepts and controversies, *J. Pediatr.* 17 (Suppl) (1990) S1.
- [335a] G. Deshpande, et al., Updated meta-analysis of probiotics for preventing necrotizing enterocolitis in preterm neonates, *Pediatrics* 125 (2010) 921.
- [336] J.Z. Jona, Advances in neonatal surgery, *Pediatr. Clin. North Am.* 45 (1998) 605.
- [337] M.E. Milner, et al., Risk factors for developing and dying from necrotizing enterocolitis, *J. Pediatr. Gastroenterol. Nutr.* 5 (1986) 359.
- [338] T.E. Wiswell, C.T. Hankins, Twins and triplets with necrotizing enterocolitis, *Am. J. Dis. Child.* 142 (1988) 1004.
- [339] M. DeCurtis, et al., A case control study of necrotizing enterocolitis occurring over 8 years in a neonatal intensive care unit, *Eur. J. Pediatr.* 146 (1987) 398.
- [340] R.M. Kliegman, Neonatal necrotizing enterocolitis: implications for an infectious disease, *Pediatr. Clin. North Am.* 26 (1979) 327.
- [341] L.S. Book, et al., Clustering of necrotizing enterocolitis: interruption by infection-control measures, *N. Engl. J. Med.* 297 (1977) 984.
- [341a] M.J. Morowitz, et al., Redefining the role of intestinal microbes in the pathogenesis of necrotizing enterocolitis, *Pediatrics* 125 (2010) 777.
- [342] H.A. Rotbart, M.J. Levin, How contagious is necrotizing enterocolitis? *Pediatr. Infect. Dis.* 2 (1983) 406.
- [343] S.A. Roback, et al., Necrotizing enterocolitis, *Arch. Surg.* 109 (1974) 314.
- [344] M.D. Stanley, D.M. Null Jr., R.A. deLemos, Relationship between intestinal colonization with specific bacteria and the development of necrotizing enterocolitis, *Pediatr. Res.* 11 (1977) 543.
- [345] H.R. Hill, C.E. Hunt, J.M. Matsen, Nosocomial colonization with *Klebsiella*, type 26, in a neonatal intensive-care unit associated with an outbreak of sepsis, meningitis, and necrotizing enterocolitis, *J. Pediatr.* 85 (1974) 415.
- [346] J. Powell, et al., Necrotizing enterocolitis: epidemic following an outbreak of *Enterobacter cloacae* type 3305573 in a neonatal intensive care unit, *Am. J. Dis. Child.* 134 (1980) 1152.
- [347] M.E. Speer, et al., Fulminant neonatal sepsis and necrotizing enterocolitis associated with a "nonenteropathogenic" strain of *Escherichia coli*, *J. Pediatr.* 89 (1976) 91.
- [348] A.H. Cushing, Necrotizing enterocolitis with *Escherichia coli* heat-labile enterotoxin, *Pediatrics* 71 (1983) 626.
- [349] A. Henderson, J. Maclaurin, J.M. Scott, *Pseudomonas* in a Glasgow baby unit, *Lancet* 2 (1969) 316.
- [350] J.A. Waldhausen, T. Herendeen, H. King, Necrotizing colitis of the newborn: common cause of perforation of the colon, *Surgery* 54 (1963) 365.
- [351] H. Stein, et al., Gastroenteritis with necrotizing enterocolitis in premature babies, *BMJ* 2 (1972) 616.
- [352] J.A. Gruskay, et al., *Staphylococcus epidermidis*-associated enterocolitis, *J. Pediatr.* 109 (1986) 520.
- [353] G.D. Overturf, et al., Neonatal necrotizing enterocolitis associated with delta toxin producing methicillin-resistant *S. aureus*, *Pediatr. Infect. Dis. J.* 9 (1990) 88.
- [354] H.A. Rotbart, et al., Neonatal rotavirus-associated necrotizing enterocolitis: case control study and prospective surveillance during an outbreak, *J. Pediatr.* 112 (1988) 87.
- [355] H.A. Rotbart, et al., Confirmatory testing of Rotazyme results in neonates, *J. Pediatr.* 107 (1985) 289.
- [356] S. Rousset, et al., Intestinal lesions containing coronavirus-like particles in neonatal necrotizing enterocolitis: an ultrastructural analysis, *Pediatrics* 73 (1984) 218.
- [357] F.E. Johnson, et al., Association of fatal coxsackie B2 viral infection and necrotizing enterocolitis, *Arch. Dis. Child.* 52 (1977) 802.
- [358] *Clostridium septicum* and neutropenic enterocolitis. Editorial, *Lancet* 2 (1987) 608.
- [359] S.M. Finegold, Anaerobic Bacteria in Human Diseases, Academic Press, New York, 1977.
- [360] G. Lawrence, P.D. Walker, Pathogenesis of enteritis necroticans in Papua New Guinea, *Lancet* 1 (1976) 125.
- [361] R.M. Kliegman, The role of clostridia in the pathogenesis of neonatal necrotizing enterocolitis, in: S.P. Borriello (Ed.), *Clostridia in Gastrointestinal Disease*, CRC Press, Boca Raton, FL, 1985, pp. 68–92.
- [362] P. Volsted-Pedersen, et al., Necrotising enterocolitis of the newborn—is it gas gangrene of the bowels? *Lancet* 2 (1976) 715.
- [363] A.M. Kosloske, J.A. Ulrich, H. Hoffman, Fulminant necrotising enterocolitis associated with clostridia, *Lancet* 2 (1978) 1014.
- [364] S. Warren, J.R. Schreiber, M.F. Epstein, Necrotizing enterocolitis and hemolysis associated with *Clostridium perfringens*, *Am. J. Dis. Child.* 138 (1984) 686.
- [365] J.L. Blakey, et al., Enteric colonization in sporadic neonatal necrotizing enterocolitis, *J. Pediatr. Gastroenterol. Nutr.* 4 (1985) 591.
- [366] W.J. Cashore, et al., Clostridia colonization and clostridial toxin in neonatal necrotizing enterocolitis, *J. Pediatr.* 98 (1981) 308.
- [367] R. Sturm, et al., Neonatal necrotizing enterocolitis associated with penicillin-resistant, toxigenic *Clostridium butyricum*, *Pediatrics* 66 (1980) 928.
- [368] F.M. Howard, et al., Outbreak of necrotising enterocolitis caused by *Clostridium butyricum*, *Lancet* 2 (1977) 1099.
- [369] A.J. Zedd, et al., Nosocomial *Clostridium difficile* reservoir in a neonatal intensive care unit, *Pediatr. Infect. Dis.* 3 (1984) 429.
- [370] I.J. Al-Jumaili, et al., Incidence and origin of *Clostridium difficile* in neonates, *J. Clin. Microbiol.* 19 (1984) 77.
- [371] M.F. Smith, et al., Clinical and bacteriological findings in necrotizing enterocolitis: a controlled study, *J. Infect.* 2 (1980) 23.
- [372] L. Gothefors, I. Blenkarn, *Clostridium butyricum* and necrotising enterocolitis, *Lancet* 1 (1978) 52.
- [373] M. Laverdière, et al., Clostridia in necrotising enterocolitis, *Lancet* 2 (1978) 377.
- [374] M.C. Kelsey, A.J. Vince, Clostridia in neonatal feces, *Lancet* 2 (1979) 100.
- [375] A.D. Kindley, P.J. Roberts, W.H. Tulloch, Neonatal necrotising enterocolitis, *Lancet* 1 (1977) 649.
- [376] A.H. Lishman, et al., *Clostridium difficile* isolation in neonates in a special care unit: lack of correlation with necrotizing enterocolitis, *Scand. J. Gastroenterol.* 19 (1984) 441.
- [377] C.M. Westra-Meijer, et al., Quantitative study of the aerobic and anaerobic faecal flora in neonatal necrotizing enterocolitis, *Arch. Dis. Child.* 58 (1983) 523.
- [378] D.F.M. Thomas, et al., Clostridial toxins in neonatal necrotizing enterocolitis, *Arch. Dis. Child.* 59 (1984) 270.
- [379] D.W. Scheifele, G.L. Bjornson, Delta toxin activity in coagulase-negative staphylococci from the bowel of neonates, *J. Clin. Microbiol.* 26 (1988) 279.
- [380] D.W. Scheifele, et al., Delta-like toxin produced by coagulase-negative staphylococci is associated with neonatal necrotizing enterocolitis, *Infect. Immun.* 55 (1988) 2268.

- [381] C. Hoy, et al., Quantitative changes in faecal microflora preceding necrotizing enterocolitis in premature neonates, *Arch. Dis. Child.* 65 (1990) 1057.
- [382] E.H. Dykes, W.H. Gilmour, A.F. Azmy, Prediction of outcome following necrotizing enterocolitis in a neonatal surgical unit, *J. Pediatr. Surg.* 20 (1985) 3.
- [383] E.R. Wayne, J.D. Burrington, J. Hutter, Neonatal necrotizing enterocolitis: evolution of new principles in management, *Arch. Surg.* 110 (1975) 476.
- [384] R. Wilson, et al., Age at onset of necrotizing enterocolitis, *Am. J. Dis. Child.* 136 (1982) 814.
- [385] J.A. Richmond, V. Mikity, Benign form of necrotizing enterocolitis, *AJR Am. J. Roentgenol.* 123 (1975) 301.
- [386] J.A. Barnard, R.B. Cotton, W. Lutin, Necrotizing enterocolitis: variables associated with the severity of the disease, *Am. J. Dis. Child.* 139 (1985) 375.
- [387] M.J. Bell, et al., Neonatal necrotizing enterocolitis: therapeutic decisions based upon clinical staging, *Ann. Surg.* 187 (1978) 1.
- [388] R. Buras, et al., Acidosis and hepatic portal venous gas: indications for surgery in necrotizing enterocolitis, *Pediatrics* 78 (1986) 273.
- [389] A. Daneman, S. Woodward, M. de Silva, The radiology of neonatal necrotizing enterocolitis (NEC): a review of 47 cases and the literature, *Pediatr. Radiol.* 7 (1978) 70.
- [390] A.G. Mata, R.M. Rosengart, Interobserver variability in the radiographic diagnosis of necrotizing enterocolitis, *Pediatrics* 66 (1980) 68.
- [391] J.F. Johnson, L.H. Robinson, Localized bowel distension in the newborn: a review of the plain film analysis and differential diagnosis, *Pediatrics* 73 (1984) 206.
- [392] T. Leonard Jr., J.F. Johnson, P.G. Pettett, Critical evaluation of the persistent loop sign in necrotizing enterocolitis, *Radiology* 142 (1982) 385.
- [393] M.M. Weinstein, The persistent loop sign in neonatal necrotizing enterocolitis: a new cause, *Pediatr. Radiol.* 16 (1986) 71.
- [394] R.M. Kliegman, A.A. Fanaroff, Neonatal necrotizing enterocolitis: a nine-year experience. II. Outcome assessment, *Am. J. Dis. Child.* 135 (1981) 608.
- [395] M.S. Keller, H.S. Chawla, Neonatal metrizamide gastrointestinal series in suspected necrotizing enterocolitis, *Am. J. Dis. Child.* 139 (1985) 713.
- [396] S. Lindley, et al., Portal vein ultrasonography in the early diagnosis of necrotizing enterocolitis, *J. Pediatr. Surg.* 21 (1986) 530.
- [397] S.W. Malin, et al., Echogenic intravascular and hepatic microbubbles associated with necrotizing enterocolitis, *J. Pediatr.* 103 (1983) 637.
- [398] R.R. Ricketts, The role of paracentesis in the management of infants with necrotizing enterocolitis, *Am. Surg.* 52 (1986) 61.
- [399] A.M. Kosloske, Surgery of necrotizing enterocolitis, *World J. Surg.* 9 (1985) 277.
- [400] C.C. Patel, Hematologic abnormalities in acute necrotizing enterocolitis, *Pediatr. Clin. North Am.* 24 (1977) 579.
- [401] J.J. Hutter Jr., W.E. Hathaway, E.R. Wayne, Hematologic abnormalities in severe neonatal necrotizing enterocolitis, *J. Pediatr.* 88 (1976) 1026.
- [402] P.E. Hyman, C.E. Abrams, R.D. Zipser, Enhanced urinary immunoreactive thromboxane in neonatal necrotizing enterocolitis: a diagnostic indicator of thrombotic activity, *Am. J. Dis. Child.* 141 (1987) 688.
- [403] D.W. Scheifele, E.M. Olson, M.R. Pendray, Endotoxemia and thrombocytopenia during neonatal necrotizing enterocolitis, *Am. J. Clin. Pathol.* 83 (1985) 227.
- [404] S.S. Rabinowitz, et al., Platelet-activating factor in infants at risk for necrotizing enterocolitis, *J. Pediatr.* 138 (2001) 81.
- [405] R.D. Christensen, et al., Granulocyte transfusion in neonates with bacterial infection, neutropenia, and depletion of mature marrow neutrophils, *Pediatrics* 70 (1982) 1.
- [406] J. Garcia, F.R. Smith, S.A. Cucinell, Urinary d-lactate excretion in infants with necrotizing enterocolitis, *J. Pediatr.* 104 (1984) 268.
- [407] L.S. Book, J.J. Herbst, A. Jung, Carbohydrate malabsorption in necrotizing enterocolitis, *Pediatrics* 57 (1976) 201.
- [408] S. Scott, et al., Effect of necrotizing enterocolitis on urinary epidermal growth factor levels, *Am. J. Dis. Child.* 145 (1991) 804.
- [409] R.M. Kliegman, M.C. Walsh, The incidence of meningitis in neonates with necrotizing enterocolitis, *Am. J. Perinatol.* 4 (1987) 245.
- [410] D.W. Scheifele, et al., Comparison of two antibiotic regimens in neonates with necrotizing enterocolitis, *Clin. Invest. Med.* 8 (1985) A183.
- [411] M.J. Bell, et al., Neonatal necrotizing enterocolitis: prevention of perforation, *J. Pediatr. Surg.* 8 (1973) 601.
- [412] T.N. Hansen, et al., A randomized controlled study of oral gentamicin in the treatment of neonatal necrotizing enterocolitis, *J. Pediatr.* 97 (1980) 836.
- [413] R.B. Faix, T.Z. Polley, T.H. Grasel, A randomized, controlled trial of parental clindamycin in neonatal necrotizing enterocolitis, *J. Pediatr.* 112 (1988) 271.
- [414] J.D. Burrington, Necrotizing enterocolitis in the newborn infant, *Clin. Perinatol.* 5 (1978) 29.
- [415] M.J. Ghory, C.A. Sheldon, Newborn surgical emergencies of the gastrointestinal tract, *Surg. Clin. North Am.* 65 (1985) 1083.
- [416] S.H. Ein, et al., A 13-year experience with peritoneal drainage under local anesthesia for necrotizing enterocolitis perforation, *J. Pediatr. Surg.* 25 (1990) 1034.
- [417] L.J. Grylack, J.W. Scanlon, Oral gentamicin therapy in the prevention of neonatal necrotizing enterocolitis: a controlled double-blind trial, *Am. J. Dis. Child.* 132 (1978) 1192.
- [418] E.A. Egan, et al., Additional experience with routine use of oral kanamycin prophylaxis for necrotizing enterocolitis in infants under 1500 grams, *J. Pediatr.* 90 (1977) 331.
- [419] V.E. Brantley, I.M. Hiatt, T. Hegyi, The effectiveness of oral gentamicin in reducing the incidence of necrotizing enterocolitis (NEC) in treated and control infants, *Pediatr. Res.* 14 (1980) 592.
- [420] M.P. Rowley, G.W. Dahlenburg, Gentamicin in prophylaxis of neonatal necrotizing enterocolitis, *Lancet* 2 (1978) 532.
- [421] R. Boyle, et al., Alterations in stool flora resulting from oral kanamycin prophylaxis of necrotizing enterocolitis, *J. Pediatr.* 93 (1978) 857.
- [422] P.R.F. Dear, D.E.M. Thomas, Oral vancomycin in preventing necrotizing enterocolitis, *Arch. Dis. Child.* 63 (1988) 1390.
- [423] M.M. Conroy, R. Anderson, K.L. Cates, Complications associated with prophylactic oral kanamycin in preterm infants, *Lancet* 1 (1978) 613.
- [424] A.M. Bhat, R.G. Meny, Alimentary absorption of gentamicin in preterm infants, *Clin. Pediatr.* 23 (1984) 683.
- [425] L.J. Grylack, J. Boehnert, J.W. Scanlan, Serum concentration of gentamicin following oral administration to preterm newborns, *Dev. Pharmacol. Ther.* 5 (1982) 47.
- [426] D.M. Anderson, R.M. Kleigman, Relationship of neonatal alimentation practices to the occurrence of endemic neonatal necrotizing enterocolitis, *Am. J. Perinatol.* 8 (1991) 62.
- [427] R.E. McKeown, et al., Role of delayed feeding and of feeding increments in necrotizing enterocolitis, *J. Pediatr.* 121 (1992) 764.
- [428] V. Carrion, E.A. Egan, Prevention of neonatal necrotizing enterocolitis, *J. Pediatr. Gastroenterol. Nutr.* 11 (1990) 317.
- [429] B. Bohnhorst, et al., Early feeding after necrotizing enterocolitis in preterm infants, *J. Pediatr.* 143 (2003) 484.
- [430] C.L. Anderson, et al., A widespread epidemic of mild necrotizing enterocolitis of unknown cause, *Am. J. Dis. Child.* 138 (1984) 979.
- [431] A.R. Gerber, et al., Increased risk of illness among nursery staff caring for neonates with necrotizing enterocolitis, *Pediatr. Infect. Dis. J.* 4 (1985) 246.
- [432] G.A. Little (Ed.), American Academy of Pediatrics Committee on the Fetus and Newborn: Guidelines for Perinatal Care, second ed., American Academy of Pediatrics, Elk Grove Village, IL, 1988, pp. 182-183.
- [433] E.D. Claud, W.A. Walker, Hypothesis: inappropriate colonization of the premature intestine can cause neonatal necrotizing enterocolitis, *FASEB J.* 15 (2001) 1398.
- [434] M.A. Schell, M. Karmirantzou, B. Snel, et al., The genome sequence of *Bifidobacterium longum* reflects its adaptation to the human gastrointestinal tract, *Proc. Natl. Acad. Sci. U. S. A.* 99 (2002) 14422.
- [435] H.C. Lin, et al., Oral probiotics prevent necrotizing enterocolitis in very low birth weight preterm infants: a multicenter, randomized, controlled trial, *Pediatrics* 122 (2008) 693.
- [436] A. Bin-Nun, et al., Oral probiotics prevent necrotizing enterocolitis in very low birth weight neonates, *J. Pediatr.* 147 (2005) 192.
- [437] C.-H. Lin, et al., Oral probiotics reduce the incidence and severity of necrotizing enterocolitis in very low birth weight infants, *Pediatrics* 115 (2005) 1.
- [438] G. Deshpande, S. Rao, S. Patole, Probiotics for prevention of necrotizing enterocolitis in preterm neonates with very low birthweight: a systematic review of randomised controlled trials, *Lancet* 369 (2007) 1614.
- [439] J.C. Leonidas, R.T. Hall, Neonatal pneumatis coli: a mild form of neonatal necrotizing enterocolitis, *J. Pediatr.* 89 (1976) 456.
- [440] M.A. Eibl, et al., Prevention of necrotizing enterocolitis in low-birth-weight infants by IgA-IgG feeding, *N. Engl. J. Med.* 319 (1988) 1.
- [441] J.N. Schullinger, et al., Neonatal necrotizing enterocolitis: survival, management, and complications: a 25-year study, *Am. J. Dis. Child.* 136 (1981) 612.
- [442] S. Abbasi, et al., Long-term assessment of growth, nutritional status and gastrointestinal function in survivors of necrotizing enterocolitis, *J. Pediatr.* 104 (1984) 550.
- [443] J.S. Janik, S.H. Ein, K. Mancor, Intestinal structure after necrotizing enterocolitis, *J. Pediatr. Surg.* 16 (1981) 438.
- [444] T.I. Ball, J.B. Wyly, Enterocyst formation: a late complication of neonatal necrotizing enterocolitis, *AJR Am. J. Roentgenol.* 147 (1986) 806.
- [445] M.C. Walsh, R.M. Kliegman, M. Hack, Severity of necrotizing enterocolitis: influence on outcome at 2 years of age, *Pediatrics* 84 (1989) 808.
- [446] L.C. Blieden, et al., Bacterial endocarditis in the neonate, *Am. J. Dis. Child.* 124 (1972) 747.
- [447] I.C. Lewis, Bacterial endocarditis complicating septicemia in an infant, *Arch. Dis. Child.* 29 (1954) 144.
- [448] D. Macaulay, Acute endocarditis in infancy and early childhood, *Am. J. Dis. Child.* 88 (1954) 715.
- [449] D.R. Shanklin, The pathology of prematurity, in: *Prematurity and the Obstetrician*, Appleton-Century-Crofts, New York, 1969p. 471.
- [450] H. Steinitz, L. Schuchmann, G. Wegner, Leberzirrhose, Meningitis, und Endocarditis ulceropolyposa bei einer Neugeborenensepsis durch B-Streptokokken (*Streptokokkus agalactiae*), *Arch. Kinderheilkd.* 183 (1971) 382.
- [451] D.H. Johnson, A. Rosenthal, A.S. Nadas, Bacterial endocarditis in children under 2 years of age, *Am. J. Dis. Child.* 129 (1975) 183.
- [452] G. Mendelsohn, G.M. Hutchins, Infective endocarditis during the first decade of life: an autopsy review of 33 cases, *Am. J. Dis. Child.* 133 (1979) 619.
- [453] R. Liersch, et al., Gegenwärtige Merkmale der bakteriellen Endokarditis im Kindersalter, *Z. Kardiol.* 66 (1977) 501.
- [454] J. Colville, I. Jeffries, Bilateral acquired neonatal Erb's palsy, *Ir. Med. J.* 68 (1975) 399.
- [455] P.S. Szymchek, A.N. Krauss, P. Winchester, Endocarditis following intracardiac placement of umbilical venous catheters in neonates, *J. Pediatr.* 90 (1977) 287.
- [456] K. Edwards, et al., Bacterial endocarditis in 4 young infants: is this complication on the increase? *Clin. Pediatr.* 16 (1977) 607.

- [457] G.A. McGuiness, R.M. Schieken, G.F. Maguire, Endocarditis in the newborn, *Am. J. Dis. Child.* 134 (1980) 577.
- [458] R.L. Bender, et al., Echocardiographic diagnosis of bacterial endocarditis of the mitral valve in a neonate, *Am. J. Dis. Child.* 131 (1977) 746.
- [459] N.R. Lundström, G. Björkhem, Mitral and tricuspid valve vegetations in infancy diagnosed by echocardiography, *Acta Paediatr. Scand.* 68 (1979) 345.
- [460] A.G. Weinberg, W.P. Laird, Group B streptococcal endocarditis detected by echocardiography, *J. Pediatr.* 92 (1978) 335.
- [461] C.W. Barton, et al., A neonatal survivor of group B beta-hemolytic streptococcal endocarditis, *Am. J. Perinatol.* 1 (1984) 214.
- [462] B.N. Agarwala, Group B streptococcal endocarditis in a neonate, *Pediatr. Cardiol.* 9 (1988) 51.
- [463] B. Chattapadhyay, Fatal neonatal meningitis due to group B streptococci, *Postgrad. Med. J.* 51 (1975) 240.
- [464] E.T. Cabacungan, G. Tetting, D.Z. Friedberg, Tricuspid valve vegetation caused by group B streptococcal endocarditis: treatment by "vegetectomy," *J. Perinatol.* 13 (1993) 398.
- [465] H.H. Kramer, et al., Current clinical aspects of bacterial endocarditis in infancy, childhood, and adolescence, *Eur. J. Pediatr.* 140 (1983) 253.
- [466] R.E. Kavey, et al., Two-dimensional echocardiography assessment of infective endocarditis in children, *Am. J. Dis. Child.* 137 (1983) 851.
- [467] K.E. Ward, et al., Successfully treated pulmonary valve endocarditis in a normal neonate, *Am. J. Dis. Child.* 137 (1983) 913.
- [468] D.K. Nakayama, et al., Management of vascular complications of bacterial endocarditis, *J. Pediatr. Surg.* 21 (1986) 636.
- [469] P. Morville, et al., Intérêt de l'échocardiographie dans le diagnostic des endocardites néonatales: à propos de trois observations, *Ann. Pediatr.* 32 (1985) 389.
- [470] J.F. Eliaou, et al., Endocardite infectieuse en période néo-natale, *Pediatric* 38 (1983) 561.
- [471] J.E. McCartney, A case of acute ulcerative endocarditis in a child aged three and a half weeks, *J. Pathol. Bacteriol.* 25 (1922) 277.
- [472] D.G. Oelberg, et al., Endocarditis in high-risk neonates, *Pediatrics* 71 (1983) 392.
- [473] G. Gossius, P. Gunnes, K. Rasmussen, Ten years of infective endocarditis: a clinicopathologic study, *Acta Med. Scand.* 217 (1985) 171.
- [474] G.J. Noel, J.E. O'Loughlin, P.J. Edelson, Neonatal *Staphylococcus epidermidis* right-sided endocarditis: description of five catheterized infants, *Pediatrics* 82 (1988) 234.
- [475] M.J. Bannon, Infective endocarditis in neonates. Letter to the editor, *Arch. Dis. Child.* 63 (1998) 112.
- [476] Personal communication to Michael Marcy.
- [477] E.S. Giddings, Two cases of endocarditis in infants, *Can. Med. Assoc. J.* 35 (1936) 71.
- [478] S.S. Soo, D.L. Boxman, *Streptococcus faecalis* in neonatal infective endocarditis, *J. Infect.* 23 (1991) 209.
- [479] C. O'Callaghan, P. McDougall, Infective endocarditis in neonates, *Arch. Dis. Child.* 63 (1988) 53.
- [480] D. Prandstraller, A.M. Marata, F.M. Picchio, *Staphylococcus aureus* endocarditis in a newborn with transposition of the great arteries: successful treatment, *Int. J. Cardiol.* 14 (1987) 355.
- [481] T. Zakrzewski, J.D. Keith, Bacterial endocarditis in infants and children, *J. Pediatr.* 67 (1965) 1179.
- [482] L. Weinstein, J.J. Schlesinger, Pathoanatomic, pathophysiologic and clinical correlations in endocarditis, *N. Engl. J. Med.* 291 (Pt 1) (1974) 832.
- [483] P. Morand, et al., Endocardite tricuspidienne calcifiée du nourisson, *Arch. Mal. Coeur* 66 (1973) 901.
- [484] J.L. Luke, R.P. Bolande, S. Gross, Generalized aspergillosis and *Aspergillus* endocarditis in infancy: report of a case, *Pediatrics* 31 (1963) 115.
- [485] E.L. Wiley, G.M. Hutchins, Superior vena cava syndrome secondary to *Candida* thrombophlebitis complicating parenteral alimentation, *J. Pediatr.* 91 (1977) 977.
- [486] T.J. Walsh, G.M. Hutchins, Postoperative *Candida* infections of the heart in children: clinicopathologic study of a continuing problem of diagnosis and therapy, *J. Pediatr. Surg.* 15 (1980) 325.
- [487] R. Faix, et al., Successful medical treatment of *Candida parapsilosis* endocarditis in a premature infant, *Am. J. Perinatol.* 7 (1990) 272.
- [488] P.N. Zenker, et al., Successful medical treatment of presumed *Candida* endocarditis in critically ill infants, *J. Pediatr.* 119 (1991) 472.
- [489] P.J. Sanchez, J.D. Siegel, J. Fishbein, *Candida* endocarditis: successful medical management in three preterm infants and review of the literature, *Pediatr. Infect. Dis. J.* 10 (1991) 239.
- [490] D.D. Millard, S.T. Shulman, The changing spectrum of neonatal endocarditis, *Clin. Perinatol.* 15 (1988) 587.
- [491] W.R. Morrow, J.E. Haas, D.R. Benjamin, Nonbacterial endocardial thrombosis in neonates: relationship to persistent fetal circulation, *J. Pediatr.* 100 (1982) 117.
- [492] H. Kronsbein, Pathogenesis of endocarditis verrucosa simplex in the newborn, *Beitr. Pathol.* 161 (1977) 82.
- [493] H.F. Krous, Neonatal nonbacterial thrombotic endocarditis, *Arch. Pathol. Lab. Med.* 103 (1979) 76.
- [494] J. Bergsland, et al., Mycotic aortic aneurysms in children, *Ann. Thorac. Surg.* 37 (1984) 314.
- [495] C.A. Bullaboy, et al., Neonatal mitral valve endocarditis: diagnosis and successful management, *Clin. Pediatr.* 29 (1990) 398.
- [496] R.S. Baltimore, Infective endocarditis in children, *Pediatr. Infect. Dis. J.* 11 (1992) 907.
- [497] R.L. Popp, Echocardiography and infectious endocarditis, *Curr. Clin. Topics. Infect. Dis.* 4 (1983) 98.
- [498] C. Watanakunakorn, Treatment of infections due to methicillin-resistant *Staphylococcus aureus*, *Ann. Intern. Med.* 97 (1982) 376.
- [499] J.S. Wolfson, M.N. Swartz, Serum bactericidal activity as a monitor of antibiotic therapy, *N. Engl. J. Med.* 312 (1985) 968.
- [500] G.I. Cooper, C.C. Hopkins, Rapid diagnosis of intravascular catheter-associated infection by direct Gram staining of catheter segments, *N. Engl. J. Med.* 312 (1985) 1142.
- [501] M.J. Perelman, et al., Aortic root replacement for complicated bacterial endocarditis in an infant, *J. Pediatr. Surg.* 24 (1989) 1121.
- [502] R.M.R. Tulloh, E.D. Silove, L.D. Abrams, Replacement of an aortic valve cusp after neonatal endocarditis, *Br. Heart J.* 64 (1990) 204.
- [503] S.A. Levinson, A. Learner, Blood cysts on the heart valves of newborn infants, *Arch. Pathol.* 14 (1932) 810.
- [504] P. Gross, Concept of fetal endocarditis: a general review with report of an illustrative case, *Arch. Pathol.* 31 (1941) 163.
- [505] B.E. Favara, R.A. Franciosi, L.J. Butterfield, Disseminated intravascular and cardiac thrombosis of the neonate, *Am. J. Dis. Child.* 127 (1974) 197.
- [506] J.G. Begg, Blood-filled cysts in the cardiac valve cusps in foetal life and infancy, *J. Pathol. Bacteriol.* 87 (1964) 177.
- [507] T.R. Thompson, et al., Umbilical artery catheterization complicated by mycotic aortic aneurysm in neonates, *Adv. Pediatr.* 27 (1980) 275.
- [508] L. Charaf, et al., A case of neonatal endocarditis, *Acta Pediatr. Scand.* 79 (1990) 704.
- [509] A.J. Schaffer, M.E. Avery, *Diseases of the Newborn*, third ed., WB Saunders, Philadelphia, 1971, p. 252.
- [510] N. Neimann, et al., Les péricardites purulentes néonatales: à propos du premier cas guéri, *Arch. Fr. Pediatr.* 22 (1965) 238.
- [511] A.M. Collier, J.D. Connor, W.L. Nyhan, Systemic infection with *Hemophilus influenzae* in very young infants, *J. Pediatr.* 70 (1967) 539.
- [512] B. McKinlay, Infectious diarrhea in the newborn caused by an unclassified species of *Salmonella*, *Am. J. Dis. Child.* 54 (1937) 1252.
- [513] H. Chiari, Zur Kenntnis der Pyozyaneus-infektion bei Säuglingen, *Zentralbl. Allg. Pathol.* 38 (1926) 483.
- [514] F. Jaiyesimi, A.A. Abioye, A.U. Anita, Infective pericarditis in Nigerian children, *Arch. Dis. Child.* 54 (1979) 384.
- [515] R.J. Wynn, Neonatal *E. coli* pericarditis, *J. Perinatol. Med.* 7 (1979) 23.
- [516] J. Kachaner, J.M. Nouaille, A. Batisse, Les cardiomegalies massives du nouveau-né, *Arch. Fr. Pediatr.* 34 (1977) 297.
- [517] J.P.A. Graham, A. Martin, *B. pyocyaneus* pericarditis occurring four days after birth, *Cent. Afr. J. Med.* 1 (1955) 101.
- [518] W.E. Feldman, Bacterial etiology and mortality of purulent pericarditis in pediatric patients: review of 162 cases, *Am. J. Dis. Child.* 133 (1979) 641.
- [519] R.J. Morgan, et al., Surgical treatment of purulent pericarditis in children, *Thorac. Cardiovasc. Surg.* 85 (1983) 527.
- [520] C.H. Jellard, G.M. Churcher, An outbreak of *Pseudomonas aeruginosa* (*pyocyanea*) infection in a premature baby unit, with observations on the intestinal carriage of *Pseudomonas aeruginosa* in the newborn, *J. Hyg.* 65 (1967) 219.
- [521] T.C. Miller, S.I. Baman, W.W. Albers, Massive pericardial effusion due to *Mycoplasma hominis* in a newborn, *Am. J. Dis. Child.* 136 (1982) 271.
- [522] L. Shenker, et al., Fetal pericardial effusion, *Am. J. Obstet. Gynecol.* 160 (1989) 1505.
- [523] G.G. Cayler, H. Taybi, H.D. Riley Jr., Pericarditis with effusion in infants and children, *J. Pediatr.* 63 (1963) 264.
- [524] G.R. Noren, E.L. Kaplan, N.A. Staley, Nonrheumatic inflammatory diseases, in: F.H. Adams, G.C. Emmanouilides (Eds.), *Moss' Heart Disease in Infants, Children, and Adolescents*, third ed., Williams & Wilkins, Baltimore, 1983, pp. 585-594.
- [525] K.S. Kanarek, J. Coleman, Purulent pericarditis in a neonate, *Pediatr. Infect. Dis. J.* 10 (1991) 549.
- [526] J.D. Cherry, Enteroviruses, polioviruses (poliomyelitis), coxsackieviruses, echoviruses, and enteroviruses, in: R.D. Feigin, J.D. Cherry (Eds.), *Textbook of Pediatric Infectious Diseases*, third ed., WB Saunders, Philadelphia, 1992, pp. 1705-1752.
- [527] J.T. Zerella, D.C.E. Halpe, Intrapericardial teratoma-neonatal cardiorespiratory distress amenable to surgery, *J. Pediatr. Surg.* 15 (1980) 961.
- [528] N. Doshi, B. Smith, B. Kliensky, Congenital pericarditis due to maternal lupus erythematosus, *J. Pediatr.* 96 (1980) 699.
- [529] P. Sasidharan, et al., Nonimmune hydrops fetalis: case reports and brief review, *J. Perinatol.* 12 (1992) 338.
- [530] J.M.B. deFonseca, M.R.Q. Davies, K.D. Bolton, Congenital hydropericardium associated with the herniation of part of the liver into the pericardial sac, *J. Pediatr. Surg.* 22 (1987) 851.
- [531] A.J. Jafa, et al., Antenatal diagnosis of bilateral congenital chylothorax with pericardial effusion, *Acta Obstet. Gynaecol. Scand.* 64 (1985) 455.
- [532] R.H. Mupaneunda, H.R. Mackanje, A life-threatening complication of percutaneous central venous catheters in neonates, *Am. J. Dis. Child.* 146 (1992) 1414.
- [533] P.H. Dennehy, New tests for the rapid diagnosis of infection in children, in: S. Aronoff et al., (Ed.), *Advances in Pediatric Infectious Diseases*, vol. 8, Mosby-Year Book, St. Louis, 1993, pp. 91-129.
- [534] B. Zeevi, et al., Interventional cardiac procedures, *Clin. Perinatol.* 15 (1988) 633.

- [535] S. Achenbach, Mediastinalabszess bei einem 3 Wochen alten Säugling, Arch. Kinderheilkd. 74 (1924) 193.
- [536] H.E. Grewe, M. Martini Pape, Die eitrige Mediastinitis im frühen Säugling-salter, Kinderarztl. Prax. 32 (1964) 305.
- [537] M. Weichsel, Mediastinitis in a newborn, Proc. Rudolf Virchow Med. Soc. City N. Y. 22 (1963) 67.
- [538] G. Weber, Retropharyngeal und Mediastinalabszess bei einem 3 Wochen alten Säugling, Chirurg 21 (1950) 308.
- [539] A.A. deLorimier, D. Haskin, F.S. Massie, Mediastinal mass caused by vertebral osteomyelitis, Am. J. Dis. Child. 111 (1966) 639.
- [540] J.L. Talbert, et al., Traumatic perforation of the hypopharynx in infants, J. Thorac. Cardiovasc. Surg. 74 (1977) 152.
- [541] M. Grunebaum, et al., Iatrogenic transmural perforation of the oesophagus in the preterm infant, Clin. Radiol. 31 (1980) 257.
- [542] T. Sands, M. Glasson, A. Berry, Hazards of nasogastric tube insertion in the newborn infant, Lancet 2 (1989) 680.
- [543] I.H. Krasna, et al., Esophageal perforation in the neonate: an emerging problem in the newborn nursery, J. Pediatr. Surg. 22 (1987) 784.
- [544] J. Topsis, H.Y. Kinas, S.R. Kandall, Esophageal perforation—a complication of neonatal resuscitation, Anesth. Analg. 69 (1989) 532.
- [545] Y. Vandenplas, et al., Cervical esophageal perforation diagnosed by endoscopy in a premature infant, J. Pediatr. Gastroenterol. Nutr. 8 (1989) 390.
- [546] R.J. Touloukian, et al., Traumatic perforation of the pharynx in the newborn, Pediatrics 59 (1977) 1019.
- [547] D.E. Johnson, et al., Management of esophageal and pharyngeal perforation in the newborn infant, Pediatrics 70 (1982) 592.
- [548] M. Coulthard, D. Isaacs, Neonatal retropharyngeal abscess, Pediatr. Infect. Dis. J. 10 (1991) 547.
- [549] A. Bittencourt, Congenital Chagas disease, Am. J. Dis. Child. 130 (1976) 97.
- [550] P.H. Azimi, J. Willert, A. Petru, Severe esophagitis in a newborn, Pediatr. Infect. Dis. J. 15 (1996) 385.
- [551] T.J. Walsh, N.J. Belitsos, S.R. Hamilton, Bacterial esophagitis in immunocompromised patients, Arch. Intern. Med. 146 (1986) 1345.
- [552] A.J. Nelson, Neonatal suppurative thyroiditis, Pediatr. Infect. Dis. 2 (1983) 243.
- [553] R. Berner, et al., *Salmonella enteritidis* orchitis in a 10-week old boy, Acta Pediatr. 83 (1994) 922.
- [554] H.N. Sanford, I. Shmigelsky, Purulent parotitis in the newborn, J. Pediatr. 26 (1945) 149.
- [555] B.H. Shulman, Acute suppurative infections of the salivary glands in the newborn, Am. J. Dis. Child. 80 (1950) 413.
- [556] W.A.B. Campbell, Purulent parotitis in the newborn: report of a case, Lancet 2 (1951) 386.
- [557] D. Leake, R. Leake, Neonatal suppurative parotitis, Pediatrics 46 (1970) 203.
- [558] I. Brook, E.H. Frazier, D.H. Thompson, Aerobic and anaerobic microbiology of acute suppurative parotitis, Laryngoscope 101 (1991) 170.
- [559] R.B. David, E.J. O'Connell, Suppurative parotitis in children, Am. J. Dis. Child. 119 (1970) 332.
- [560] W.W. Banks, et al., Neonatal submandibular sialadenitis, Am. J. Otolaryngol. 1 (1980) 261.
- [561] M. Saarinen, et al., Spectrum of 2,836 cases of invasive bacterial or fungal infections in children: results of prospective nationwide five-year surveillance in Finland. Finnish Pediatric Invasive Infection Study Group, Clin. Infect. Dis. 21 (1995) 1134-1144.
- [562] L.M. Solomon, N.B. Esterly, Neonatal Dermatology, WB Saunders, Philadelphia, 1973.
- [563] M.N. Swartz, A.N. Weinberg, Bacterial diseases with cutaneous involvement, in: T.B. Fitzpatrick Jr. (Ed.), Dermatology in General Medicine, McGraw-Hill, New York, 1971.
- [564] Personal communication to Michael Marcy.
- [565] Personal communication to Michael Marcy.
- [566] R.M. Fortunov, et al., Community acquired *Staphylococcus aureus* infections in term and near-term previously healthy neonates, Pediatrics 118 (2006) 874.
- [567] A.B. Zafar, et al., Use of 0.3% triclosan (Bacti-Stat) to eradicate an outbreak of methicillin-resistant *Staphylococcus aureus* in a neonatal nursery, Am J Infect Control 23 (1995) 200.
- [568] I.J. Frieden, Blisters and pustules in the newborn, Curr. Probl. Pediatr. 19 (1989) 553.
- [569] S. Weinberg, M. Leider, L. Shapiro, Color Atlas of Pediatric Dermatology, McGraw-Hill, New York, 1975.
- [570] H.I. Maibach, G. Hildick-Smith (Eds.), Skin Bacteria and Their Role in Infection, McGraw-Hill, New York, 1965.
- [571] H. Speert, Circumcision of the newborn: an appraisal of its present status, Obstet. Gynecol. 2 (1953) 164.
- [572] S.H. Annabil, A. Al-Hifi, T. Kazi, Primary tuberculosis of the penis in an infant, Tubercle 71 (1990) 229.
- [573] T.G. Cleary, S. Kohl, Overwhelming infection with group B beta-hemolytic streptococcus associated with circumcision, Pediatrics 64 (1979) 301.
- [574] L. Sauer, Fatal staphylococcal bronchopneumonia following ritual circumcision, Am. J. Obstet. Gynecol. 46 (1943) 583.
- [575] D. Annunziato, L.M. Goldblum, Staphylococcal scalded skin syndrome, Am. J. Dis. Child. 132 (1978) 1187.
- [576] G.S. Breuer, S. Walfisch, Circumcision complications and indications for ritual recircumcision—clinical experience and review of the literature, Isr. J. Med. Sci. 23 (1987) 252.
- [577] J.R. Woodside, Necrotizing fasciitis after neonatal circumcision, Am. J. Dis. Child. 134 (1980) 301.
- [578] J. Stranko, M.E. Ryan, A.M. Bowman, Impetigo in newborn infants associated with a plastic bell clamp circumcision, Pediatr. Infect. Dis. 5 (1986) 597.
- [579] S.F. Siddiqi, P.M. Taylor, Necrotizing fasciitis of the scalp, Am. J. Dis. Child. 136 (1982) 226.
- [580] D.M. Okada, A.W. Chow, V.T. Bruce, Neonatal scalp abscess and fetal monitoring: factors associated with infection, Am. J. Obstet. Gynecol. 129 (1977) 185.
- [581] L. Cordero, C.W. Anderson, E.H. Hon, Scalp abscess: a benign and infrequent complication of fetal monitoring, Am. J. Obstet. Gynecol. 146 (1983) 126.
- [582] M.M. Wagener, et al., Septic dermatitis of the neonatal scalp and maternal endomyometritis with intrapartum internal fetal monitoring, Pediatrics 74 (1984) 81.
- [583] I. Brook, Microbiology of scalp abscesses in newborns, Pediatr. Infect. Dis. J. 11 (1992) 766.
- [584] J.C. Bogdan, R.H. Rapkin, Clostridia infection in the newborn, Pediatrics 58 (1976) 120.
- [585] W.F. Fitter, D.J. DeSa, H. Richardson, Chorioamnionitis and funisitis due to *Corynebacterium kutscheri*, Arch. Dis. Child. 55 (1979) 710.
- [586] I. Sarkany, C.C. Gaylarde, Skin flora of the newborn, Lancet 1 (1967) 589.
- [587] H.E. Evans, S.O. Akpata, A. Baki, Factors influencing the establishment of neonatal bacterial flora. I. The role of host factors, Arch. Environ. Health 21 (1970) 514.
- [588] I. Sarkany, L. Arnold, The effect of single and repeated applications of hexachlorophene on the bacterial flora of the skin of the newborn, Br. J. Dermatol. 82 (1970) 261.
- [589] D.J. Thompson, et al., Excess risk of staphylococcal infection and disease in newborn males, Am. J. Epidemiol. 84 (1966) 314.
- [590] R.C. Rudoy, J.D. Nelson, Breast abscess during the neonatal period: a review, Am. J. Dis. Child. 129 (1975) 1031.
- [591] A.C. Reboli, J.F. John, A.H. Levkoff, Epidemic methicillin-gentamicin-resistant *Staphylococcus aureus* in a neonatal intensive care unit, Am. J. Dis. Child. 143 (1989) 34.
- [592] J.E. Fergie, K. Purcell, Community-acquired methicillin-resistant *Staphylococcus aureus* infection in south Texas children, Pediatr. Infect. Dis. J. 20 (2001) 860.
- [593] H.E. Evans, et al., Flora in newborn infants: annual variation in prevalence of *Staphylococcus aureus*, *Escherichia coli*, and streptococci, Arch. Environ. Health 26 (1973) 275.
- [594] H.J. Starr, P.B. Holliday Jr., Erythema multiforme as a manifestation of neonatal septicemia, J. Pediatr. 38 (1951) 315.
- [595] J.L. Washington, R.E.L. Fowler, G.J. Guarino, Erythema multiforme in a premature infant associated with sepsis due to *Pseudomonas*, Pediatrics 39 (1967) 120.
- [596] J.F. Foley, et al., *Achromobacter* septicemia—fatalities in prematures. I. Clinical and epidemiological study, Am. J. Dis. Child. 101 (1961) 279.
- [597] T.K. Belgaumkar, Impetigo neonatorum congenita due to group B beta-hemolytic streptococcus infection. Letter to the editor, J. Pediatr. 86 (1975) 982.
- [598] F. Halal, et al., Congenital vesicular eruption caused by *Hemophilus influenzae* type b, Pediatrics 62 (1978) 494.
- [599] M.O. Martin, et al., Les signes cutanés des infections bactériennes néonatales, Arch. Fr. Pediatr. 42 (1985) 471.
- [600] A. Kline, E. O'Connell, Group B streptococcus as a cause of neonatal bullous skin lesions, Pediatr. Infect. Dis. J. 12 (1993) 165.
- [601] N. Khuri-Bulos, K. McIntosh, Neonatal *Haemophilus influenzae* infection: report of eight cases and review of the literature, Am. J. Dis. Child. 129 (1975) 57.
- [602] D.A. Bray, Ecthyma gangrenosum: full thickness nasal slough, Arch. Otolaryngol. 98 (1973) 210.
- [603] R.W. Heffner, G.F. Smith, Ecthyma gangrenosum in *Pseudomonas* septicemia, Am. J. Dis. Child. 99 (1960) 524.
- [604] S.P. Ghosal, et al., Noma neonatorum: its aetiopathogenesis, Lancet 1 (1978) 289.
- [605] G.P. Bodey, et al., Infections caused by *Pseudomonas aeruginosa*, Rev. Infect. Dis. 5 (1983) 279.
- [606] J.E. Baley, R.A. Silverman, Systemic candidiasis: cutaneous manifestations in low birth weight infants, Pediatrics 82 (1988) 211.
- [607] L. Cordero Jr., E.H. Hon, Scalp abscess: a rare complication of fetal monitoring, J. Pediatr. 78 (1971) 533.
- [608] J.D. Nelson, Bilateral breast abscess due to group B streptococcus, Am. J. Dis. Child. 130 (1976) 567.
- [609] H.L. Levy, J.F. O'Connor, D. Ingall, Bacteremia, infected cephalhematoma, and osteomyelitis of the skull in a newborn, Am. J. Dis. Child. 114 (1967) 649.
- [610] S.S. Ellis, et al., Osteomyelitis complicating neonatal cephalhematoma, Am. J. Dis. Child. 127 (1974) 100.
- [611] H. Stetler, et al., Neonatal mastitis due to *Escherichia coli*, J. Pediatr. 76 (1970) 611.
- [612] H.H. Balfour Jr., et al., Complications of fetal blood sampling, Am. J. Obstet. Gynecol. 107 (1970) 288.
- [613] M.A. McGuigan, R.P. Lipman, Neonatal mastitis due to *Proteus mirabilis*, Am. J. Dis. Child. 130 (1976) 1296.
- [614] H.D. Wilson, K.C. Haltalin, Acute necrotizing fasciitis in childhood, Am. J. Dis. Child. 125 (1973) 591.
- [615] V.F. Barry, M. Beezley, Infant mastitis due to gram-negative organisms, Am. J. Dis. Child. 124 (1972) 736.
- [616] I. Brook, The aerobic and anaerobic microbiology of neonatal breast abscess, Pediatr. Infect. Dis. J. 10 (1991) 785.

- [617] Centers for Disease Control, Nosocomial *Serratia marcescens* infections in neonates—Puerto Rico., MMWR Morb. Mortal. Wkly. Rep. 23 (1974) 183.
- [618] J.K. Todd, F.W. Bruhn, Severe *Haemophilus influenzae* infections: spectrum of disease, Am. J. Dis. Child. 129 (1975) 607.
- [619] S.H. Zinner, et al., Puerperal bacteremia and neonatal sepsis due to *Hemophilus parainfluenzae*: report of a case with antibody titers, Pediatrics 49 (1972) 612.
- [620] M.S. Platt, Neonatal *Hemophilus vaginalis* (*Corynebacterium vaginalis*) infection, Clin. Pediatr. 10 (1971) 513.
- [621] P.M. Leighton, B. Bulleid, R. Taylor, Neonatal cellulitis due to *Gardnerella vaginalis*, Pediatr. Infect. Dis. 1 (1982) 339.
- [622] Y.H. Lee, R.B. Berg, Cephalhematoma infected with *Bacteroides*, Am. J. Dis. Child. 121 (1971) 77.
- [623] M.J. Mandel, R.J. Lewis, Molluscum contagiosum of the newborn, Br. J. Dermatol. 84 (1970) 370.
- [624] J.M. Clark, W.R. Weeks, J. Tatton, *Drosophila* myiasis mimicking sepsis in a newborn, West. J. Med. 136 (1982) 443.
- [625] B.R. Burns, R.M. Lampe, G.H. Hansen, Neonatal scabies, Am. J. Dis. Child. 133 (1979) 1031.
- [626] O.J. Hensley, C.A. Hart, R.W.I. Cooke, *Candida albicans* skin abscesses, Arch. Dis. Child. 59 (1984) 479.
- [627] D.F. Turbeville, et al., Complications of fetal scalp electrodes: a case report, Am. J. Obstet. Gynecol. 122 (1975) 530.
- [628] Centers for Disease Control, Gonococcal scalp-wound infection—New Jersey., MMWR Morb. Mortal. Wkly. Rep. 24 (1975) 115.
- [629] I. Brook, Osteomyelitis and bacteremia caused by *Bacteroides fragilis*, Clin. Pediatr. 19 (1980) 639.
- [630] R.T. Mohon, et al., Infected cephalhematoma and neonatal osteomyelitis of the skull, Pediatr. Infect. Dis. 5 (1986) 253.
- [631] S.M. Cohen, B.W. Miller, H.W. Orris, Meningitis complicating cephalhematoma, J. Pediatr. 30 (1947) 327.
- [632] M. Walsh, K. McIntosh, Neonatal mastitis, Clin. Pediatr. 25 (1986) 395.
- [633] W.H. Langewisch, An epidemic of group A, type 1 streptococcal infections in newborn infants, Pediatrics 18 (1956) 438.
- [634] D. Sinniah, B.R. Sandiford, A.E. Dugdale, Subungual infection in the newborn: an institutional outbreak of unknown etiology, possibly due to *Veillonella*, Clin. Pediatr. 11 (1972) 690.
- [635] A.H. Cushing, Omphalitis: a review, Pediatr. Infect. Dis. 4 (1985) 282.
- [636] H. McKenna, D. Johnson, Bacteria in neonatal omphalitis, Pathology 7 (1977) 11.
- [637] W.H. Mason, et al., Omphalitis in the newborn infant, Pediatr. Infect. Dis. J. 8 (1989) 521.
- [638] C.C. Geil, W.K. Castle, E.A. Mortimer Jr., A. Group, streptococcal infections in newborn nurseries, Pediatrics 46 (1970) 849.
- [639] D.P. Bliss, P.J. Healey, J.H.T. Waldbraussen, Necrotizing fasciitis after Plastibell circumcision, J. Pediatr. 131 (1997) 459.
- [640] A.M. Kosloske, et al., Cellulitis and necrotizing fasciitis of the abdominal wall in pediatric patients, J. Pediatr. Surg. 16 (1981) 246.
- [641] W.F. Gee, J.S. Ansell, Neonatal circumcision: a 10-year overview: with comparison of the Gomco clamp and the Plastibell device, Pediatrics 58 (1976) 824.
- [642] G.N. Goldberg, R.C. Hansen, P.J. Lynch, Necrotizing fasciitis in infancy: report of three cases and review of the literature, Pediatr. Dermatol. 2 (1984) 55.
- [643] R.S. Ramamurthy, G. Srinivasan, N.M. Jacobs, Necrotizing fasciitis and necrotizing cellulitis due to group B streptococcus, Am. J. Dis. Child. 131 (1977) 1169.
- [644] R.W. Krieger, M.J. Chusid, Perirectal abscess in childhood, Am. J. Dis. Child. 133 (1979) 411.
- [645] M. Arditi, R. Yogev, Perirectal abscess in infants and children: report of 52 cases and review of the literature, Pediatr. Infect. Dis. J. 9 (1990) 411.
- [646] L. Victorin, An epidemic of otitis in newborns due to infection with *Pseudomonas aeruginosa*, Acta. Paediatr. Scand. 56 (1967) 344.
- [647] E.J. Laubo, A.S. Paller, Common skin problems during the first year of life, Pediatr. Clin. North Am. 41 (1994) 1105.
- [648] M.T. Glover, D.J. Atherton, R.J. Levinsky, Syndrome of erythroderma, failure to thrive, and diarrhea in infancy: a manifestation of immunodeficiency, Pediatrics 81 (1988) 66.
- [649] L.S. Prod'hom, et al., Care of the seriously ill neonate with hyaline membrane disease and with sepsis (sclerema neonatorum), Pediatrics 53 (1974) 170.
- [650] W.E. Hughes, M.L. Hammond, Sclerema neonatorum, J. Pediatr. 32 (1948) 676.
- [651] G.H. McCracken Jr., H.R. Shinefield, Changes in the pattern of neonatal septicemia and meningitis, Am. J. Dis. Child. 112 (1966) 33.
- [652] I. Gordon, Miliary sebaceous cysts and blisters in the healthy newborn, Arch. Dis. Child. 24 (1949) 286.
- [653] J.A. Carr, et al., Relationship between toxic erythema and infant maturity, Am. J. Dis. Child. 112 (1966) 219.
- [654] P. Merlob, A. Metzker, S.H. Reisman, Transient neonatal pustular melanosis, Am. J. Dis. Child. 136 (1982) 521.
- [655] G. Kahn, A.M. Rywlin, Acropustulosis of infancy, Arch. Dermatol. 115 (1979) 831.
- [656] A.W. Lucky, J.S. McGuire, Infantile acropustulosis with eosinophilic pustules, J. Pediatr. 100 (1982) 428.
- [657] W.F. Murphy, A. Langley, Common bullous lesions—presumably self-inflicted—occurring in utero in the newborn infant, Pediatrics 32 (1963) 1099.
- [658] W.L. Weston, A.T. Lane, J.A. Weston, Diaper dermatitis: current concepts, Pediatrics 66 (1980) 532.
- [659] Nappy rashes, Editorial, BMJ 282 (1981) 420.
- [660] J.C. Hauth, G.B. Merenstein (Eds.), Guidelines for Perinatal Care, fourth ed., American Academy of Pediatrics and American College of Obstetricians and Gynecologists, Elk Grove Village, IL, 1997.
- [661] I.M. Gladstone, et al., Randomized study of six umbilical cord care regimens, Clin. Pediatr. 27 (1988) 127.
- [662] Centers for Disease Control, National nosocomial infections study report: nosocomial infections in nurseries and their relationship to hospital infant bathing practices—a preliminary report, Centers for Disease Control, Atlanta, 1974, pp. 9–23.
- [663] H.M. Gezon, M.J. Schaberg, J.O. Klein, Concurrent epidemics of *Staphylococcus aureus* and group A streptococcus disease in a newborn nursery—control with penicillin G and hexachlorophene bathing, Pediatrics 51 (1973) 383.
- [664] S. Seeberg, et al., Prevention and control of neonatal pyoderma with chlorhexidine, Acta Paediatr. Scand. 73 (1984) 498.
- [665] A.R. de Toledo, J.W. Chandler, Conjunctivitis of the newborn, Infect. Dis. Clin. North Am. 6 (1992) 807.
- [666] J.P. Whitcher, Neonatal ophthalmia. Have we advanced in the last 20 years? Int. Ophthalmol. Clin. 30 (1990) 39.
- [667] D.D. Millard, R. Yogev, *Haemophilus influenzae* type b: a rare case of congenital conjunctivitis, Pediatr. Infect. Dis. 7 (1988) 363.
- [668] D.T. McLeod, F. Ahmad, M.A. Calder, *Branhamella catarrhalis* (beta lactamase positive) ophthalmia neonatorum, Lancet 2 (1984) 647.
- [669] M. Ellis, et al., Neonatal conjunctivitis associated with meningococcal meningitis, Arch. Dis. Child. 67 (1992) 1219.
- [670] M.J. Naiditch, A.G. Bower, Diphtheria: a study of 1433 cases observed during a ten year period at Los Angeles County Hospital, Am. J. Med. 17 (1954) 229.
- [671] M.S. Khan, S.E. Stead, Neonatal *Pasteurella multocida* conjunctivitis following zoonotic infection of mother, J. Infect. Dis. 1 (1979) 289.
- [672] I. Brook, W.J. Martin, S.M. Finegold, Effect of silver nitrate application on the conjunctival flora of the newborn, and the occurrence of clostridial conjunctivitis, J. Pediatr. Ophthalmol. Strabismus 15 (1978) 179.
- [673] P. Bourbeau, V. Holla, S. Piemontese, Ophthalmia neonatorum caused by *Neisseria cinerea*, J. Clin. Microbiol. 28 (1990) 1640.
- [674] K. Hedberg, et al., Outbreak of erythromycin-resistant staphylococcal conjunctivitis in a newborn nursery, Pediatr. Infect. Dis. J. 9 (1990) 268.
- [675] M.J. Paentice, G.R. Hutchinson, D. Taylor-Robinson, A microbiological study of neonatal conjunctivitis, Br. J. Ophthalmol. 61 (1977) 9.
- [676] K.I. Sandstrom, et al., Microbial causes of neonatal conjunctivitis, J. Pediatr. 105 (1984) 706.
- [677] G. Regev-Yochav, et al., Methicillin-resistant *Staphylococcus aureus* in a neonatal intensive care unit, Emerg. Infect. Dis. 11 (2005) 453.
- [678] T. Rutar, Vertically acquired community methicillin-resistant *Staphylococcus aureus* dacryocystitis in a neonate, J. AAPOS 13 (2009) 79–81.
- [679] J.H. Kim, et al., Outbreak of Gram-positive bacterial keratitis associated with epidemic keratoconjunctivitis in neonates and infants, Eye 23 (2009) 1059–1065.
- [680] A.M. Duarte, et al., Eosinophilic pustular folliculitis in infancy and childhood, Am. J. Dis. Child. 147 (1993) 197.
- [681] M.R. Hammerschlag, Conjunctivitis in infancy and childhood, Pediatr. Rev. 5 (1984) 285.
- [682] J. Wincelaus, et al., Diagnosis of ophthalmia neonatorum, BMJ 295 (1987) 1377.
- [683] I.L. Molgaard, P.B. Nielsen, J. Kaern, A study of the incidence of neonatal conjunctivitis and of its bacterial causes including *Chlamydia trachomatis*, Acta Ophthalmol. 62 (1984) 461.
- [684] M. Laga, et al., Epidemiology of ophthalmia neonatorum in Kenya, Lancet 2 (1986) 1145.
- [685] K.J. Nathoo, A.S. Latif, J.E.S. Trijssenaar, Aetiology of neonatal conjunctivitis in Harare, Cent. Afr. J. Med. 30 (1984) 123.
- [686] S. Stenson, R. Newman, H. Fedukowicz, Conjunctivitis in the newborn: observations on incidence, cause, and prophylaxis, Ann. Ophthalmol. 13 (1981) 329.
- [687] R.P. Burns, D.H. Rhodes Jr., *Pseudomonas* eye infection as a cause of death in premature infants, Arch. Ophthalmol. 65 (1981) 517.
- [688] S.E. Drewett, et al., Eradication of *Pseudomonas aeruginosa* infection from a special-care nursery, Lancet 1 (1972) 946.
- [689] G.A. Cole, D.P. Davies, D.J. Austin, *Pseudomonas* ophthalmia neonatorum: a cause of blindness, BMJ 281 (1980) 440.
- [690] E.I. Traboulsi, et al., *Pseudomonas aeruginosa* ophthalmia neonatorum, Am. J. Ophthalmol. 98 (1984) 801.
- [691] R. Lohrer, B.H. Belohradsky, Bacterial endophthalmitis in neonates, Eur. J. Pediatr. 146 (1987) 354.
- [692] G.D. Christensen, et al., Epidemic *Serratia marcescens* in a neonatal intensive care unit: importance of the gastrointestinal tract as a reservoir, Infect. Control 3 (1982) 127.
- [693] H. Nishida, H.M. Risemberg, Silver nitrate ophthalmic solution and chemical conjunctivitis, Pediatrics 56 (1975) 3368.
- [694] S.S. Kripke, B. Golden, Neonatal inclusion conjunctivitis: a report of three cases and a discussion of differential diagnosis and treatment, Clin. Pediatr. 11 (1972) 261.
- [695] L.K. Pickering, et al., Red Book: 2009 Report of the Committee on Infectious Diseases, 28th ed., Elk Grove Village, IL, American Academy of Pediatrics, 2009.