

POSTER PRESENTATION

Open Access

Epidemiological description of the demographic and HIV disease characteristics of HIV patients who are in care but not on treatment in Spain

P Domingo^{1*}, P Arazo², JL Casado³, MA Castaño⁴, E Ferrer⁵, MA Goenaga⁶, R Granados⁷, F Pulido⁸, J Rodriguez⁹, R Rubio⁸, J Sanz¹⁰, P Viciana¹¹

From Tenth International Congress on Drug Therapy in HIV Infection Glasgow, UK. 7-11 November 2010

Background

It is estimated that around 15% of HIV infected patients are being followed in HIV units for disease evolution and need of treatment. Data on the demographics and HIV disease status of these patients are scarce.

Methods

Cross sectional analysis within 12 HIV units in Spain with the aim of describing the characteristics of patients in care but not on treatment. We collected data on demographics, HIV disease status as well as the prevalence of non-HIV co-morbidities that could qualify for antiretroviral treatment initiation regardless immunological status according to current Spanish treatment guidelines.

Results

Data from 865 patients naïve to ARV treatment were collected. Mean age 37 y, most of them were male (83%) and Caucasian (85%). Predominant HIV risk factor was MSM (56%). Median time from HIV diagnosis 2.3 y (IQR 1.0 to 5.1). Current median CD4 cell count was 602 /mm³ (IQR 422 to 729) and mean HIV-1 RNA levels were 4.6 log.

The prevalence of key HIV and non-HIV features according to different CD4 strata can be seen in Table 1

Conclusions

In our cohort of patients who are under care but not on treatment, a considerable amount of them could benefit from treatment initiation according to current guide-

Table 1

	CD4<350 (n=112)	CD4 350 - 500 (n=261)	CD4>500 (n=492)
>55 years of age	1.8%	3.1%	2.6%
HIV RNA >100,000 cop/mL	20.5%	13.4%	8.1%
CD4 percentage < 14%	19.6%	3.1%	1.2%
HCV coinfection	25.0%	14.2%	14.0%
HBV coinfection requiring treatment†	0%	0.4%	0%
Liver cirrhosis	4.5%	1.1%	1.6%
HIV associated nephropaty	0%	0.8%	0%
10 y CHD>20% (Framinghan score)‡	Not available	Not available	0.8%

^{†:} According to Spanish guidelines ‡: Data not available for 30% of the subjects

Full list of author information is available at the end of the article



¹Hospital de Sa Santa Creu i Sant Pau, c/ Sant Antoni Maria Claret, 167, Barcelona. Spain

lines. When monitoring to what extent treatment guidelines are being followed, non HIV comorbidities should also be taken into account. Among these patients, HCV coinfection is the most prevalent comorbidity. Some of these patients could also benefit from generalisation of cardiovascular risk evaluation in HIV units.

Author details

¹Hospital de Sa Santa Creu i Sant Pau, c/ Sant Antoni Maria Claret, 167, Barcelona, Spain. ²Hospital Miguel Servet, Zaragoza, Spain. ³Hospital Ramón y Cajal, Madrid, Spain. ⁴Hospital Carlos Haya, Málaga, Spain. ⁵Ciudad Univ Bellvitge, Barcelona, Spain. ⁶Hospital Donostia, Donosti, Spain. ⁷Hospital Dr Negrín, Las Palmas, Spain. ⁸Hospital 12 de Octubre, Madrid, Spain. ⁹Hospital Univ Virgen Macarena, Seville, Spain. ¹⁰Hospital Príncipe de Asturias, Madrid, Spain. ¹¹Hospital Univ Virgen del Rocío, Seville, Spain.

Published: 8 November 2010

doi:10.1186/1758-2652-13-S4-P53

Cite this article as: Domingo et al.: Epidemiological description of the demographic and HIV disease characteristics of HIV patients who are in care but not on treatment in Spain. Journal of the International AIDS Society 2010 13(Suppl 4):P53.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit

