

Adapting Virtual Reality-Based Mental Health Interventions for Equitable Global Use: Proof of Concept

Ana F. Trueba, Ph.D., Cristina Crespo-Andrade, Ph.D., Maria Sol Garces, Ph.D., Camila Alejandra Merino Larrea, B.A., Nicolas A. Alvarez Frank, B.A., Boyu Ren, Ph.D., Hailey V. Cray, M.P.H., Julia Kimball, B.A., Rebecca Dickinson, B.Sc., B.A., Ipsit V. Vahia, M.D.

Among low and middle-income countries (LMICs), mental health challenges are compounded by poverty and limited access to mental health services, particularly among older adults (1–3). There is a need for effective and innovative interventions tailored to these populations. Mindfulness interventions can be applied trans-diagnostically and have been shown as feasible and effective in LMICs (4).

However, mindfulness can be challenging to engage with, especially at the start of the practice (5). Notably, minoritized and disadvantaged populations tend to engage less in mindfulness (6), and older adults may also have difficulty with these interventions (7). One approach to improving the effectiveness of mindfulness among older adults may be by using virtual reality (VR). VR has been shown to facilitate a greater state of mindfulness when compared to audio-only mindfulness interventions (8).

Low-cost VR hardware and expanding content availability may facilitate their implementation to older adults in LMICs. In this study, we aimed to demonstrate whether a manualized low-cost VR mindfulness intervention is feasible, tolerable, and effective among older adults with and without mild cognitive impairment (MCI) in Quito, Ecuador. This feasibility study has significant implications for determining the scalability of mindfulness-based interventions as well as for demonstrating how advanced technologies like VR can be made much more accessible.

METHODS

We conducted 10 weekly sessions of VR-assisted mindfulness for older adults in Quito, Ecuador. This study was approved by the Institutional Review Board at Universidad San Francisco de Quito. The study used low-cost VR DesTeK VS VR headsets (costing \$70) requiring a smartphone to run VR apps. For this study, we specifically selected the free and publicly available “Sites in VR” application as it offers a variety of static nature scenes ideal for guided meditation. This approach leverages the

widespread use of smartphones in LMICs (9) and removes the need for WIFI or more complex VR headsets.

We recruited 28 older adults ages 61 or older with subclinical anxiety and depression through social media. Inclusion criteria were normal cognitive functioning or MCI determined using the Montreal Cognitive Assessment (MoCA), being able to give written consent, and being a Spanish speaker. The MoCA cut-off score was set at 26 based on the original MoCA and the use of the Spanish MoCA in other Latin American countries (10). Exclusion criteria were moderate to severe cognitive impairment, blindness or severe vision problems, deafness or hearing problems, history of dizziness, and current psychosis or mania.

Tailoring the Intervention for the Study Population and Location

We used a Spanish-translated standardized mindfulness script to deliver the mindfulness intervention. We elected to use static 360° VR content using the “Sites in VR” app. Because the gyroscope sensor required to use the VR headset is not always available in smartphones sold in Latin America, we used devices obtained in the United States. Gyroscope sensors allow the mobile device to detect and measure an individual’s orientation, rotational movement, and the position of the device in space. These sensors are essential for full immersion in the VR environment. Following the conclusion of the study, all devices were returned to study staff.

Intervention Sessions

We offered participants a choice between home visits or clinic visits to include those with transportation issues. Nineteen participants preferred home visits, and nine attended clinic visits. In each session, we selected a nature scene from the “Sites in VR” app, then placed the smartphone in the headset and adjusted the fit for each participant. Participants were seated while using the headset to

minimize the risk of falls, accidents, and motion sickness while eliminating the need for hand controllers to “move” in the virtual space.

We delivered a practical form of Acceptance and Commitment Therapy using guided visualization to emphasize the principles of cognitive diffusion and emotional acceptance (11, 12). In each 45-min session, the therapist guided participants through a nature scene in VR for 15 min. Participants were asked to imagine what they were smelling, hearing, touching, and tasting while in the VR environment to facilitate full immersion in the experience. Study participants were also prompted to imagine how moving around the VR environment might feel. Processing and debriefing sessions followed the exercise. All sessions were video and audio recorded to ensure consistency and to assess participants’ responses.

Outcomes

Attrition and Acceptability. We documented attrition and reasons for dropout and used standardized open-ended questions to document experiences after every intervention session and any physical symptoms or discomfort.

If participants reported dizziness when asked about general discomfort, we administered the Spanish version of the Dizziness Handicap Inventory (DHI).

Clinical Outcomes. Though our population was subclinical, we evaluated anxiety and depression symptoms to evaluate the general impact on psychological health, using the Spanish versions of the Geriatric Depression Scale (GDS-15) (13) and the Generalized Anxiety Disorder (GAD-7) (14). In addition, we assessed perceived stress, trait mindfulness, and behavior activation using Spanish versions of the Perceived Stress Scale (PSS) (15), the Mindfulness Attention Awareness Scale (MAAS) (16), and the Behavioral Activation Scale for Depression (BADSD) (17). All the scales were administered at three time points: pre-screening, post-intervention, and 1-month post-intervention. We used the MoCA (18) scores to assess cognitive status.

Statistical Analysis

We applied a Bonferroni Correction to set the p -value cut-off for significance at 0.00 and tested the normality of the data using the Shapiro-Wilk test. For scales that were not normally distributed (GAD and GDS), we used the Wilcoxon Signed Ranked test to examine changes from pre- to post-intervention and 1-month follow-up. We used linear mixed models with random intercept to examine changes across time points for normally-distributed outcome data (PSS, BADSD, and MAAS).

RESULTS

Attrition and Acceptability

We recruited a total of 28 older adults aged 61 to 81. The minimum age to participate was 61, but there was no upper

cut-off. The average age of our sample was 68 years ($SD = 6.5$) and women ($N = 22$) comprised the majority. Six participants (21%) dropped out over the course of the study: five were for reasons unrelated to the study, and one discontinued study participation after the third intervention session due to dizziness and discomfort with the VR headset. Accordingly, study staff administered the DHI to this participant. No other participants reported physical symptoms during the intervention.

Our sample was highly educated with all participants reporting at least a high school degree and 21 reporting an undergraduate degree or higher. The majority of our sample ($N = 25$) identified as “Mestizo,” a person of mixed European and indigenous background, three identified as White, and 13 reported being married. Seventeen (60%) of the participants had MoCA scores at or below 26, suggesting MCI.

Only three participants had a formal diagnosis of a depressive or anxiety disorder and none were receiving any additional psychological or psychiatric treatment at the time of study participation. The data of all participants were pooled for analyses as all participants completed the same intervention and outcome measures.

Psychological Measures


We noted that GDS scores significantly decreased using the Wilcoxon Signed ranked test pre- to post-intervention, and pre- to 1-month follow-up (see Figure 1 and Table 1). We did not find a significant change in GAD-7 scores but there was a trend reduction from pre-intervention to 1 month follow-up.

Using linear-mixed modeling, we found that BADSD had a trend increase from pre-intervention to post- ($B = 8.58$, $t(26) = 2.29$ $p = 0.030$) and significantly increased from pre-intervention to 1-month follow-up $B = 9.44$, $t(28) = 3.05$ $p = 0.005$. The MAAS scores significantly increased from pre- to post-intervention ($B = 10.75$, $t(32) = 4.76$ $p < 0.001$) and pre- to 1-month follow-up ($B = 6.71$, $t(21) = 3.44$ $p = 0.002$). The PSS score did not significantly change across any assessment.

DISCUSSION

Our aim for this study was to demonstrate the feasibility of using low-cost, and minimally resource-intensive VR-delivered mindfulness in an older non-English speaking sample in Ecuador—a low-income country. We successfully demonstrated proof-of-concept for this approach, finding that it was acceptable, well tolerated, and with initial evidence of positive clinical impact. The attrition rate of this study was 21%, similar to that found in other mindfulness-based interventions for older adults (19).

Furthermore, 17 participants (60%) had MCI, which is consistent with literature demonstrating that VR interventions can be deployed in this population (20, 21).

FIGURE 1. Psychological outcomes. + $p \leq 0.05$, ** $p < 0.006$, *** $p < 0.001$.+ $p \leq 0.05$, ** $p < 0.006$, *** $p < 0.001$

Although the absence of a control group limits our ability to determine if VR enhanced mindful states, we found that participants had significantly higher trait mindfulness scores post-intervention and at 1-month post-intervention. It is possible that this finding represents the effect of more sustained attention to mindfulness tasks, as reported in studies in younger adults (22–25). We noted that the intervention led to significant decreases in depression scores and increases in behavior activation scores. These results are both consistent with existing literature on the beneficial impact of mindfulness on mood

(26, 27). In contrast, anxiety and perceived stress scores did not significantly change, which may be at least in part due to our participants' low baseline anxiety and stress scores (28).

Our study has several limitations. Our sample was one of convenience and biased towards those with access to social media. The absence of a control group, and a non-clinical sample means that we cannot reliably infer the direct clinical impact of this intervention on symptoms. We also cannot quantify a differential impact from VR. A larger controlled study will need to replicate our findings

TABLE 1. Psychological outcomes.^a

	N	Minimum	Maximum	Mean	Std. deviation
GDS pre-intervention	28	0.00	7.00	2.43	1.85
GDS post-intervention	22	0.00	4.00	1.36**	1.33
GDS follow-up	22	0.00	4.00	1.18***	1.22
BADS pre-intervention	28	65.00	130.00	101.36	17.14
BADS post-intervention	22	87.00	131.00	111.73**	11.30
BADS follow-up	22	80.00	137.00	110.85 ⁺	15.61
MAAS pre-intervention	28	29.00	85.00	61.43	13.64
MAAS post-intervention	22	51.00	90.00	69.36**	11.03
MAAS follow-up	22	45.00	89.00	73.41***	11.08
GAD7 pre-intervention	28	1.00	17.00	6.36	3.85
GAD7 post-intervention	22	0.00	11.00	5.27	2.85
GAD7 follow-up	22	0.00	15.00	4.68 ⁺	3.92
PSS pre-intervention	28	6.00	27.00	17.423	5.71
PSS post-intervention	22	6.00	27.00	15.73	5.66
PSS follow-up	22	6.00	28.00	15.73	5.86

^a BADS, Behavioral Activation Scale for Depression; GAD-7, Generalized Anxiety Disorder; GDS, Geriatric Depression Scale; MAAS, Mindfulness Attention Awareness Scale; PSS, Perceived Stress Scale.

+ $p \leq 0.05$, ** $p < 0.006$, *** $p < 0.001$.

and ascertain their generalizability. Additionally, the Sites in VR app used in this study necessitates a smartphone with a gyroscope, which is more expensive and may limit scalability to an extent. However, our approach represents a novel use of VR technology adapted for scaling to LMICs. This proof of concept along with a preliminary signal towards clinical efficacy has greater implications for how digital tools can be adapted for more equitable use globally. While our findings are modest, they are an important first step in this crucial direction.

Clinical Implications

Our study has clinical implications for older adults facing economic, language, and cultural barriers. Our results support the feasibility of broadening access and implementation of mindfulness interventions to diverse demographics. In essence, our findings suggest that VR-delivered mindfulness interventions are a viable, accessible, and cost-effective tool that could improve the well-being and mental health of older adults, especially in resource-limited settings.

AUTHOR AND ARTICLE INFORMATION

McLean Hospital, Belmont, Massachusetts, USA (Trueba, Ren, Cray, Kimball, Dickinson, Vahia); Department of Psychiatry, Harvard Medical School, Boston, Massachusetts, USA (Trueba, Ren, Vahia); Department of Psychology, Universidad San Francisco de Quito, Quito, Ecuador (Trueba, Crespo-Andrade, Garcés, Larrea, Alvarez Frank).

Send correspondence to Dr. Trueba (atruebayepz@mclean.harvard.edu).

An unrestricted gift from the Eric Warren Goldman Charitable Trust and the McLean Technology and Aging Lab. An intramural research grant awarded by Universidad San Francisco de Quito specifically for this study.

Dr. Vahia receives current research support from the National Institute on Aging, the National Institute of Mental Health, the Once Upon a Time Foundation, and the Harvard Dean's Initiative on Aging.

Dr. Vahia has performed scientific consultation for Otsuka. None of this work is related to the current study.

The data that support the findings of this study are available from the corresponding author upon reasonable request.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2024 The Authors. Psychiatric Research and Clinical Practice published by Wiley Periodicals LLC on behalf of American Psychiatric Association.

Received October 13, 2023; revised March 15, 2024; accepted March 17, 2024.

REFERENCES

- Salazar-Xirinachs JM, García-Buchaca R, Cecchini S, Shaw S. Envejecimiento en América Latina y el Caribe: inclusión y derechos de las personas mayores [Internet]. Cepal.org. 2022 [citado el 9 de agosto de 2023]. Available from: https://repositorio.cepal.org/bitstream/handle/11362/48568/4/S2201042_en.pdf
- Bonilla-Sierra P, Vargas-Martínez A.-M, Davalos-Batallas V, Leon-Larios F, Lomas-Campos M.-M. Chronic diseases and associated factors among older adults in Loja, Ecuador. *Int J of Environ Res Public Health*. 2020;17(11):4009. <https://doi.org/10.3390/ijerph17114009>
- Quijada Y, Villagrán L, Vaccari Jiménez P, Reyes C, Gallardo LD. Social Inequality and mental health in Chile, Ecuador, and Colombia. *Lat Am Perspect*. 2019;46(6):92–108. <https://doi.org/10.1177/0094582X18803682>
- Patel V, Naslund JA, Wood S, Patel A, Chauvin JJ, Agrawal R, et al. EMPOWER: toward the global dissemination of psychosocial interventions. *Focus*. 2022;20(3):301–6. <https://doi.org/10.1176/appi.focus.20220042>
- Lomas T, Cartwright T, Edginton T, Ridge D. A qualitative analysis of experiential challenges associated with meditation practice. *Mindfulness*. 2015;6(4):848–60. <https://doi.org/10.1007/s12671-014-0329-8>
- Olano HA, Kachan D, Tannenbaum SL, Mehta A, Annane D, Lee DJ. Engagement in mindfulness practices by U.S. adults: socio-demographic barriers. *J Altern Complement Med*. 2015;21(2):100–2. <https://doi.org/10.1089/acm.2014.0269>

7. Morone NE, Greco CM. Adapting mindfulness meditation for the older adult. *Mindfulness*. 2014;5(5):610–2. <https://doi.org/10.1007/s12671-014-0297-z>
8. Yildirim C, O'Grady T. The efficacy of a virtual reality-based mindfulness intervention. In: 2020 IEEE International Conference on Artificial Intelligence and Virtual Reality (AIVR). IEEE; 2020. <https://doi.org/10.1109/aivr50618.2020.00035>
9. Bastawrous A, Armstrong MJ. Mobile health use in low- and high-income countries: an overview of the peer-reviewed literature. *J R Soc Med*. 2013;106(4):130–42. <https://doi.org/10.1177/0141076812472620>
10. Aguilar-Navarro SG, Mimenza-Alvarado AJ, Palacios-García AA, Samudio-Cruz A, Gutiérrez-Gutiérrez LA, Ávila-Funes JA. Validity and reliability of the Spanish version of the Montreal Cognitive Assessment (MoCA) for the detection of cognitive impairment in Mexico. *Rev Colomb Psiquiatr*. 2018;47(4):237–43. <https://doi.org/10.1016/j.rcp.2017.05.003>
11. Seshadri A, Orth SS, Adaji A, Singh B, Clark MM, Frye MA, et al. Mindfulness-based cognitive therapy, acceptance and commitment therapy, and positive psychotherapy for major depression. *Am J Psychother*. 2021;74(1):4–12. <https://doi.org/10.1176/appi.psychotherapy.20200006>
12. Hayes SC, Strosahl K, Wilson KG. Acceptance and commitment therapy: the process and practice of Mindful Change. New York: The Guilford Press; 2016.
13. Erazo M, Fors M, Mullo S, González P, Viada C. Internal consistency of Yesavage Geriatric Depression Scale (GDS 15-item version) in Ecuadorian older adults. *Inquiry*. 2020;57:004695802097118. <https://doi.org/10.1177/0046958020971184>
14. Camargo L, Herrera-Pino J, Shelach S, Soto-Añari M, Porto MF, Alonso M, et al. Escala de ansiedad generalizada GAD-7 en profesionales médicos colombianos durante pandemia de COVID-19: validez de constructo y confiabilidad. *Rev Colomb Psiquiatr*. 2021;52(3):245–50. <https://doi.org/10.1016/j.rcp.2021.06.003>
15. Remor E, Rueda MCPL. La relación entre niveles de la actividad física y la experiencia de estrés y de síntomas de malestar físico. *Interamerican J Psych*. 2007;41(3):313–22.
16. Soler J, Tejedor R, Feliu-Soler A, Pascual JC, Cebolla A, Soriano J, et al. Propiedades psicométricas de la versión española de la escala Mindful Attention Awareness Scale (MAAS). *Actas Esp Psiquiatr*. 2012;40(1):19–26. Available from: <https://medes.com/publication/72182>
17. Sánchez-Angulo J, Barraca J, Mora EJ, Reyes-Ortega M. propiedades psicométricas de la escala de activación conductual para la depresión (BADS) en una muestra mexicana. *Clin Salud*. 2018;151–5. <https://doi.org/10.5093/clysa2018a22>
18. Delgado C, Araneda A, Behrens MI. Validación del instrumento Montreal Cognitive Assessment en español en adultos mayores de 60 años. *Neurología*. 2019;34(6):376–85. <https://doi.org/10.1016/j.nrl.2017.01.013>
19. Geiger PJ, Boggero IA, Brake CA, Caldera CA, Combs HL, Peters JR, et al. Mindfulness-based interventions for older adults: a review of the effects on physical and emotional well-being. *Mindfulness*. 2016;7(2):296–307. <https://doi.org/10.1007/s12671-015-0444-1>
20. Huygelier H, Schraepen B, van Ee R, Vanden Abeele V, Gillebert CR. Acceptance of immersive head-mounted virtual reality in older adults. *Sci Rep*. 2019;9(1):4519. <https://doi.org/10.1038/s41598-019-41200-6>
21. Manera V, Chapoulie E, Bourgeois J, Guerchouche R, David R, Ondrej J, et al. A feasibility study with image-based rendered virtual reality in patients with mild cognitive impairment and dementia. *PLoS One*. 2016;11(3):e0151487. <https://doi.org/10.1371/journal.pone.0151487>
22. Schachtner-Appel A, Sahyoun N, Chen J, Song H.-J. Formative research for an intervention to reduce plate waste and improve produce intake in schools. *Health Behav Policy Rev*. 2021;8(5):477–87. <https://doi.org/10.14485/hbpr.8.5.8>
23. Benham S, Enam N, Ivaturi S. A mindfulness program addressing sleep quality and stress: transition to a telehealth format for higher education students during COVID-19. *Int J Telerehabil*. 2022;14(1):e6439. <https://doi.org/10.5195/ijt.2022.6439>
24. Navarro-Haro MV, López-del-Hoyo Y, Campos D, Linehan MM, Hoffman HG, García-Palacios A, et al. Meditation experts try virtual reality mindfulness: a pilot study evaluation of the feasibility and acceptability of virtual reality to facilitate mindfulness practice in people attending a mindfulness conference. *PLoS One*. 2017;12(11):e0187777. <https://doi.org/10.1371/journal.pone.0187777>
25. Seabrook E, Kelly R, Foley F, Theiler S, Thomas N, Wadley G, et al. Understanding how virtual reality can support mindfulness practice: mixed methods study. *J Med Internet R*. 2020;22(3):e16106. <https://doi.org/10.2196/16106>
26. O'Connor M, Piet J, Hougaard E. The effects of mindfulness-based cognitive therapy on depressive symptoms in elderly bereaved people with loss-related distress: a controlled pilot study. *Mindfulness*. 2013;5(4):400–9. <https://doi.org/10.1007/s12671-013-0194-x>
27. Gable SL, Reis HT, Elliot AJ. Behavioral activation and inhibition in everyday life. *J Pers Soc Psychol*. 2000;78(6):1135–49. <https://doi.org/10.1037/0022-3514.78.6.1135>
28. Gallegos AM, Hoerger M, Talbot NL, Moynihan JA, Duberstein PR. Emotional benefits of mindfulness-based stress reduction in older adults: the moderating roles of age and depressive symptom severity. *Aging Ment Health*. 2013;17(7):823–9. <https://doi.org/10.1080/13607863.2013.799118>