



Correction to: Ephedra Herb extract activates/desensitizes transient receptor potential vanilloid 1 and reduces capsaicin-induced pain

Shunsuke Nakamori^{1,2} · Jun Takahashi^{1,2} · Sumiko Hyuga²  · Toshiko Tanaka-Kagawa³ · Hideto Jinno⁴ · Masashi Hyuga⁵ · Takashi Hakamatsuka⁵ · Hiroshi Odaguchi² · Yukihiro Goda⁵ · Toshihiko Hanawa² · Yoshinori Kobayashi^{1,2}

Published online: 1 March 2018
© The Author(s) 2018. This article is an open access publication

Correction to: J Nat Med (2017) 71:105–113
<https://doi.org/10.1007/s11418-016-1034-9>

The article Ephedra Herb extract activates/desensitizes transient receptor potential vanilloid 1 and reduces capsaicin-induced pain, written by Shunsuke Nakamori, Jun Takahashi, Sumiko Hyuga, Toshiko Tanaka-Kagawa, Hideto Jinno, Masashi Hyuga, Takashi Hakamatsuka, Hiroshi Odaguchi, Yukihiro Goda, Toshihiko Hanawa and Yoshinori Kobayashi, was originally published Online First without open access. After publication in volume 71, issue 1, page 105–113 the author decided to opt for Open Choice and to make the article an open access publication. Therefore, the copyright of the article has been changed to © The Author(s) 2018 and the article is forthwith distributed under the terms

of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, duplication, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, duplication, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The original article can be found online at <https://doi.org/10.1007/s11418-016-1034-9>.

✉ Sumiko Hyuga
hyuga-s@insti.kitasato-u.ac.jp

- ¹ Department of Pharmacognosy, School of Pharmacy, Kitasato University, 5-9-1 Shirokane, Minato-ku, Tokyo 108-8641, Japan
- ² Oriental Medicine Research Center, Kitasato University, 5-9-1, Shirokane, Minato-ku, Tokyo 108-8642, Japan
- ³ Department of Biochemical Toxicology, Yokohama University of Pharmacy, 601 Matano-cho, Totsuka-ku, Yokohama 245-0066, Japan
- ⁴ Faculty of Pharmacy, Meijo University, 150 Yagotoyama, Tempaku-ku, Nagoya 468-8503, Japan
- ⁵ National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan